

CONSTRUCTION OF OPEN SETS OF FREE k -TUPLES OF MATRICES

by A. MANDEL and J. Z. GONÇALVES*

(Received 10th August 1985)

0. Introduction

A k -tuple of members of a group is *free* if it freely generates a free group. In an earlier paper [1] the authors have shown:

Theorem 0. *Let F be a locally compact field with a nontrivial absolute value. Then the set of free k -tuples of $PGL(n, F)$ has nonvoid interior in $PGL(n, F)^k$, where the latter has the topology induced from that of F .*

In the present work we elaborate on the proof of the theorem above, in order to describe an explicit construction for open sets of free k -tuples. For this purpose, it is more convenient to work in $GL(n, F)$ than $PGL(n, F)$. The first step produces an open subset of F^{kn^2+kn} and an open map ϕ_k of it into $GL(n, F)^k$ whose image consists of free k -tuples; this, together with elementary facts about the quotient map $GL(n, F) \rightarrow PGL(n, F)$ yields Theorem 0. The second step consists of intersecting that set with a suitable affine copy of F^{nk^2} such that ϕ_k immerses that intersection into $GL(n, F)^k$. This provides open subsets of free k -tuples of $GL(n, F)$ described by exactly kn^2 parameters. It is not very hard to follow the computations in the proof on small examples over familiar fields; two such examples are included in the paper.

We introduce some notation. Throughout, F is a locally compact field with a nontrivial absolute value $x \mapsto |x|$. A product space F^m will be equipped with the norm $|(x_1, \dots, x_m)| = \max\{|x_1|, \dots, |x_m|\}$. As F is fixed, we use the simple notation $M_n = M_n(F)$, $GL_n = GL_n(F)$ and denote by D_n the set of diagonal matrices in M_n , and N_n the subset of D_n whose members have diagonal entries which are distinct and nonzero. We identify $M_n = F^{n^2}$ and $D_n = F^n$ (via $(x_1, \dots, x_n) \mapsto \text{diag}(x_1, \dots, x_n)$). Hence $GL_n \times N_n$ is an open and dense subset of F^{n^2+n} . Now we state our main results. Let k be a positive integer and define the map

$$\phi_k: GL_n^k \times D_n^k \rightarrow M_n^k$$

by

$$\phi_k((A^{(1)}, \dots, A^{(k)}), (D^{(1)}, \dots, D^{(k)})) = (A^{(1)}D^{(1)}(A^{(1)})^{-1}, \dots, A^{(k)}D^{(k)}(A^{(k)})^{-1}).$$

*This paper forms part of the Proceedings of the conference Groups–St Andrews 1985.

Theorem 1. *The restriction of ϕ_k to $GL_n^k \times N_n^k$ is an open map.*

An m -tuple ($m \geq n$) of members of F^n is said to be in *general position* provided each n of its members comprises a basis of F^n .

Theorem 2. *Let $W = (W^{(1)}, \dots, W^{(k)}) \in GL_n^k$. Suppose that the kn -tuple of columns of W is in general position. Then there exist a neighborhood \mathcal{A} of W in GL_n^k and a subset \mathcal{D} of D_n^k with nonempty interior such that $\phi_k(\mathcal{A} \times \mathcal{D})$ consists only of free k -tuples.*

Note that as N_n^k is dense in D_n^k , the combination of these two results yields Theorem 0. As in [1], the discovery of Theorem 2 as well as many ideas in its proof are inspired by Tits [5].

One can remove the degeneracy of ϕ_k , by restricting this map to a suitable subspace. For convenience of explicit constructions, let $P \in GL_n^k$ be a k -tuple of permutation matrices and denote by M_n^P the affine subspace of M_n^k which consists of k -tuples $(A^{(1)}, \dots, A^{(k)})$ such that for $i = 1, \dots, k$, $A^{(i)}$ has a 1 in each entry where $P^{(i)}$ has a 1.

Theorem 3. *Let $W, \mathcal{A}, \mathcal{D}$ be as in Theorem 2. Then there exist a k -tuple P of permutation matrices and an open set \mathcal{B} of GL_n^k such that the restrictions of ϕ_k to $(\mathcal{B} \cap M_n^P) \times \mathcal{D}$ is an open immersion and $\phi_k((\mathcal{B} \cap M_n^P) \times \mathcal{D}) = \phi_k(\mathcal{A} \times \mathcal{D})$.*

It is clear that the quotient map $GL_n \rightarrow PGL_n$ maps free k -tuples to free k -tuples. As this map is open, the theorems above yield open sets of free k -tuples in PGL_n . Actually, the open sets constructed in the proof of Theorem 2 are unions of fibres of this quotient map.

The authors have conjectured [1] that free k -tuples form (alternatively, contain) a dense open set in PGL_n^k . It was pointed to us by D. Sullivan that the conjectures fail for $PGL(2, \mathbb{C})$; this may be extracted from [4] where the interior of the set of free k -tuples of $PSL(2, \mathbb{C})$ is characterized.

1. The map ϕ_k

Proof of Theorem 1. It is enough to show that $\phi = \phi_1$ is an open map, as $\phi_k = (\phi, \dots, \phi)$. In order to prove that, we compute the differential and show that it is surjective at each point of $GL_n \times N_n$. The desired result is a standard consequence of the “inverse mapping theorem” when $F = \mathbb{R}$ or \mathbb{C} ; for non-archimedean F , see the remark following the proof.

Let us fix a point $(A, D) \in GL_n \times N_n$ and write $d\phi$ for the differential of ϕ at (A, D) . Using well known differentiation rules, we obtain:

$$d\phi(H, K) = HDA^{-1} - ADA^{-1}HA^{-1} + AK A^{-1}. \tag{1}$$

To show that $d\phi$ is surjective, we consider instead the linear map $T: M_n \times D_n \rightarrow M_n$ defined by $T(H, K) = A^{-1}d\phi(AH, K)$. Thus

$$T(H, K) = HD - DH + K. \tag{2}$$

Let E_{ij} be the matrix which has a 1 in position ij and 0 elsewhere. It is enough to show that each E_{ij} is in the image of T . Suppose that $D = \text{diag}(x_1, \dots, x_n)$, with the x_i 's distinct. Direct computations using (2) yield:

$$T((x_j - x_i)^{-1} E_{ij}, 0) = E_{ij},$$

if $i \neq j$ and

$$T(0, E_{ii}) = E_{ii}. \quad \square$$

Remark. A differential calculus for non-archimedean fields has been developed by Schikhof [3] with results that validate the proof above in that case. However, there seems to be a mistake in the proof of Theorem 2.3 of [3]. This can be bypassed by noting that proofs of the inverse mapping theorem using the notion of “strong derivative”, as in Nijenhuis [2] (we are grateful to Prof. E. L. Lima for pointing out this reference to us) remain valid in the non-archimedean setting. That extends the theorem at least to rational functions $F^m \rightarrow F^n$, which is enough for our purposes.

Now we remove the degeneracy of ϕ . This is done by first characterizing its fibres. That, together with Theorem 1 enables one to recognize the triple $(GL_n \times N_n, \phi, \text{Im } \phi)$ as a principal G -bundle, where G is the group of monomial matrices in GL_n (matrices with exactly one nonzero on each row and column). Then we proceed to the construction of local cross sections.

Lemma 1.1. *Let (A, D) and (\bar{A}, \bar{D}) be elements of $GL_n \times N_n$. Then $\phi(A, D) = \phi(\bar{A}, \bar{D})$ if and only if there exists a monomial matrix M such that $\bar{A} = AM$ and $\bar{D} = M^{-1}DM$. Further, there exists a neighborhood of (A, D) whose intersection with any fibre of ϕ has fixed D -component.*

Proof. Clearly $\phi(AM, M^{-1}DM) = \phi(A, D)$ if M is a monomial matrix. For the converse, suppose that $\phi(A, D) = \phi(\bar{A}, \bar{D})$. It follows that the diagonal entries of D and \bar{D} are the eigenvalues of $\phi(A, D)$. Since $D \in N_n$, there is a uniquely determined permutation matrix P such that $P^{-1}DP = \bar{D}$. One concludes that $\phi(\bar{A}P^{-1}, D) = \phi(A, D)$. Hence, for $i = 1, \dots, n$, the i th column of A , and that of $\bar{A}P^{-1}$ are eigenvectors of $\phi(A, D)$ corresponding to the same eigenvalue. As each eigenspace of $\phi(A, D)$ is 1-dimensional, each column of $\bar{A}P^{-1}$ is a (nonzero) scalar multiple of the corresponding column of A , whence $\bar{A}P^{-1} = AK$ for some diagonal matrix K . The result follows with $M = KP$.

As for the last assertion, we note first that the action of monomial matrices on N_n , given by conjugation, factors through the induced action of the symmetric group, hence has finite orbits. Thus, given (A, D) the assertion is validated by a product neighborhood $B \times C$, where C is a ball in N_n centred in D , of radius less than half the distance of D to the finite set $\{PDP^{-1} \mid P \text{ a non-identity permutation matrix}\}$. \square

Theorem 1.2. *Let GL_n^1 be the subset of matrices in GL_n whose main diagonal consists of ones only. Then the restriction of ϕ to $GL_n^1 \times N_n$ is an open immersion onto $\phi(GL_n \times N_n)$.*

Proof. Let $(A, D) \in GL_n^I \times N_n$. Choose a neighborhood of (A, D) in $GL_n \times N_n$ satisfying the conclusions of Lemma 1.1, and let B be its intersection with $GL_n^I \times N_n$. Suppose that (A', D') and $(A'', D'') \in B$ have the same image under ϕ . By choice of $B, D' = D''$, hence, by the Lemma, $A'' = A'K$, for some diagonal matrix K . Since A' and $A'' \in GL_n^I, A' = A''$, whence $\phi|_B$ is 1-1. It will now follow that $\phi|_{GL_n^I \times N_n}$ is an immersion from the fact that it is open, proved as follows:

Let GL'_n be the open subset of GL_n comprised of matrices with nonzero diagonal entries, and let $\xi: GL'_n \rightarrow GL_n^I$ be division of each row by the corresponding diagonal element. With $\iota: GL_n^I \rightarrow GL'_n$ denoting the inclusion, we have the following commutative diagram of maps:

$$\begin{array}{ccc}
 GL'_n \times N_n & \xrightarrow{\phi} & GL_n \\
 \downarrow \xi \times 1 & & \uparrow \phi \\
 GL_n^I \times N_n & \xrightarrow{\iota \times 1} & GL'_n \times N_n
 \end{array}$$

Since ϕ is open and ξ is continuous, it follows that $\phi \circ \iota \times 1 = \phi|_{GL_n^I \times N_n}$ is open.

To see that $\phi(GL_n^I \times N_n) = \phi(GL_n \times N_n)$, choose $(A, D) \in GL_n \times N_n$; choose a nonzero term in the determinant expansion of A and let M be the monomial matrix corresponding to that term. Then, $(AM^{-1}, MDM^{-1}) \in GL_n^I \times N_n$ and $\phi(AM^{-1}, MDM^{-1}) = \phi(A, D)$.

An obvious, but convenient consequence of this is stated here with the notation of Theorem 3.

Corollary 1.3. *Let $P = (P^{(1)}, \dots, P^{(k)})$ be a k -tuple of permutation matrices. Then the restriction of ϕ_k to $(GL_n^I)^k \times N_n^k$ is an open immersion onto $\phi_k(GL_n^k \times N_n^k)$.*

2. Construction of free k -tuples

The proof of Theorem 2 is a modification of the proof of Proposition 2.12 of Tits [5]. As in [1], the tool for proving a k -tuple to be free is the following criterion, due to Tits, and which has as ancestor Klein's Table-tennis Lemma.

Lemma 2.1. *Let G be a group acting on a set P and let $g = (g_1, \dots, g_k)$ be a k -tuple of members of G . Suppose that there exists a k -tuple (P_1, \dots, P_k) of subsets of P and $p \in P - (P_1 \cup \dots \cup P_k)$ such that, for all distinct $i, j, 1 \leq i, j \leq k$, and for all $n \in \mathbb{Z}^*$, $g_i^n(P_j \cup \{p\}) \subseteq P_i$. Then, g is free.*

This is the plan of the proof of Theorem 2. Beginning with the given k -tuple W , we choose a point $p \in F^n$ that will play the role suggested in Lemma 2.1. Next we choose suitable small disks in F^n , centred at the first and last column of each $W^{(i)}$ (for simplicity, all with the same radius); for each i , the union of all 1-dimensional subspaces through the two corresponding disks will yield the set P_i of Lemma 2.1. Then the set A is chosen as a convenient disk about W . Finally the set \mathcal{D} is chosen subject to two requirements: for any $(A, D) \in \mathcal{A} \times \mathcal{D}$, the k -tuple $g = \phi_k(A, D)$ satisfies:

- (a) the first (last) column of each $A^{(i)}$ is an eigenvector corresponding to the largest (smallest) eigenvalue of g_i , and
- (b) the relative sizes of the entries of members of \mathcal{D} satisfy computed constraints that guarantee that $g_i(P_j \cup \{p\})$ and $g_i^{-1}(P_j \cup \{p\})$ are contained in P_i , $i \neq j$. A simple observation will then show that the conditions of Lemma 2.1 are fulfilled for each $g \in \phi(\mathcal{A} \times \mathcal{D})$. Note that the P_i 's and p are the same for all g 's.

We need two simple preliminary results. Denote F^n by V . Recall that when $m \geq n$ an m -tuple of members of V is in *general position* if every n of its members are linearly independent.

Lemma 2.2. *The set G_m of m -tuples ($m \geq n$) of V in general position is open in V^m .*

Proof. Indeed, (V_1, \dots, V_m) is in general position if and only if for each $1 \leq i_1 < \dots < i_n \leq m$, $\det(V_{i_1}, \dots, V_{i_n}) \neq 0$. That is, G_m is the complement in $V^m = F^{nm}$ of a finite number of hypersurfaces. \square

If A is a matrix (vector), A_i denote its i th column (component).

Lemma 2.3. *Suppose that $A \in M$ and $x \in V$ are such that $(A_1, \dots, A_n, x) \in G_{n+1}$. Then, A is nonsingular and each component ^{n} of $A^{-1}x$ is nonzero.*

Proof. That A is nonsingular follows immediately from the definition of general position. Also, as $x = \sum_i A_i (A^{-1}x)_i$, general position of (A_1, \dots, A_n, x) implies that $(A^{-1}x)_i \neq 0$. \square

We have arrived at the objective of this section. Denote, whenever a set $S \subseteq F^r$ (some r) and α is a positive real, by

$$\mathcal{B}(S, \alpha) = \{x \in F^r \mid d(s, x) \leq \alpha \text{ for some } s \in S\}$$

the closed α -neighborhood of S .

Proof of Theorem 2. Regarding W as a list of column vectors, by hypothesis $W \in G_{nk}$. Choose a point $p \in V$ not lying on any of the hyperplanes spanned by columns of W ; thus $w = (W, p) \in G_{nk+1}$. Choose now positive reals $\gamma, \varepsilon, \delta$ such that:

- (a) $\mathcal{B}(w, \gamma) \subseteq G_{nk+1}$ (this is possible, by Lemma 2.2),
- (b) $\mathcal{B}(\mathcal{B}(w, \delta), \varepsilon) \subseteq \mathcal{B}(w, \gamma)$ (clearly, $\delta + \varepsilon \leq \gamma$ suffices but in working on non-archimedean examples, $\gamma = \varepsilon = \delta$ will also do).

Define now, for $i = 1, \dots, k$, $B_s = \mathcal{B}(W_1^{(s)}, \gamma)$, $C_i = \mathcal{B}(W_n^{(i)}, \gamma)$ and $\mathcal{A}^{(i)} = \mathcal{B}(W^{(i)}, \delta)$. Set now:

- (c) $\mathcal{A} = \mathcal{B}(W, \delta) = \mathcal{A}^{(1)} \times \dots \times \mathcal{A}^{(k)}$,
- (d) $P_i = \{x \in V \mid \lambda x \in B_i \cup C_i \text{ for some } \lambda \in F\}$.

Finally, we define $\mathcal{D} = \mathcal{D}^{(1)} \times \dots \times \mathcal{D}^{(k)}$, where

- (e) $\mathcal{D}^{(i)} = \{\text{diag}(\lambda_1, \dots, \lambda_n) \mid \mu_i^{-1} |\lambda_1| \geq |\lambda_2| \geq \dots \geq |\lambda_{n-1}| \geq \mu_i |\lambda_n| > 0\}$, where μ_1, \dots, μ_k are positive reals yet to be determined. To begin with, we just require that $\mu_i \geq 1$. The other restrictions on μ_i will be determined so that the following is true.
- (f) If $g = (g_1, \dots, g_k) \in \phi_k(\mathcal{A} \times \mathcal{B})$, then for $1 \leq i, j \leq k, i \neq j$, and $x \in P_j \cup \{p\}$, there exist $\alpha, \alpha' \in F^*$ such that $g_i(\alpha x) \in B_i$ and $g_i^{-1}(\alpha' x) \in C_i$.

Let us finish the proof of the Theorem under the assumption that (f) holds. We must show that each $g \in \phi_k(\mathcal{A} \times \mathcal{D})$ is free. Now, from (f) and the definition of the P_i , it follows clearly that $g_i^{-1}(P_j \cup \{p\}) \subseteq P_i$, whenever $i \neq j$. Also, since $\mu_i \geq 1$, if $D \in \mathcal{D}^{(i)}$, so does D^r for every positive r hence $(g_1^r, \dots, g_k^r) \in \phi_k(\mathcal{A} \times \mathcal{D})$ for each positive r . Applying the previous observation to $(g_1^r, \dots, g_k^r), r > 0$, shows that the conditions of Lemma 2.2 are fulfilled.

Incidentally, note that if some $\mu_i = 1$, we could obtain $g_i = I$, which cannot be a member of a free k -tuple. This shows that if the numbers $\mu_i \geq 1$ are such that (f) holds, indeed $\mu_i > 1, i = 1, \dots, k$.

Now we compute the μ_i 's. Let us denote, for $r = 1$ or n ,

$$\mathcal{L}_r = \{\text{diag}(\lambda_1, \dots, \lambda_n) \mid \lambda_r = 0, |\lambda_i| \leq 1 \text{ for } i = 1, \dots, n\} \subseteq D_n.$$

Now, for $1 \leq i, j \leq k, i \neq j$ and $r = 1$ or n , let

$$\beta_{i,j,r} = \sup\{ \|(ADA^{-1}x)/(A^{-1}x)_r\| \mid A \in \mathcal{A}^{(i)}, x \in B_j \cup C_j \cup \{p\}, D \in \mathcal{L}_r \}.$$

This is finite: when A, x are the domain of this expression, the columns of A occur together with x in a member of G_{nk+1} (see (a)), hence by Lemma 2.3 the denominator of $\|(ADA^{-1}x)/(A^{-1}x)_r\|$ never vanishes, and this is a continuous real valued function with compact domain.

Set now

$$(g) \quad \mu_i = \sup_{j \neq i} \left\{ \frac{\beta_{i,j,1}}{\varepsilon}, \frac{\beta_{i,j,n}}{\varepsilon}, 1 \right\}$$

(As remarked before, it will follow that $\mu_i > 1$, hence the presence of 1 in the expression defining μ_i is unnecessary. But we know no other "a priori" way of guaranteeing that $\mu_i \geq 1$).

Now we prove (f). Let $g_i = ADA^{-1}$, with $A \in \mathcal{A}^{(i)}, D \in \mathcal{D}^{(i)}$, and let $x \in P_j \cup \{p\}$. Let $\alpha = (D_{11}, (A^{-1}x)_1)^{-1}$. Then

$$g_i(\alpha x) = (ADA^{-1}x)/(D_{11}(A^{-1}x)_1) = A\bar{D}A^{-1}\bar{x},$$

where

$$\bar{D} = D_{11}^{-1}D = \text{diag}(1, \lambda_2, \dots, \lambda_n), \lambda_i = D_{ii}/D_{11}$$

and $\bar{x} = x/(A^{-1}x)_1$. Note that $\mu_i^{-1} > |\lambda_2| \geq |\lambda_3| \geq \dots \geq |\lambda_n|$. Further, $(A^{-1}\bar{x})_1 = 1$ whence

the identity $A_1 = A \operatorname{diag}(1, 0, \dots, 0)A^{-1}\bar{x}$. We have:

$$\|g_i(\alpha x) - A_1\| = \|A\bar{D}A^{-1}\bar{x} - A \operatorname{diag}(1, 0, \dots, 0)A^{-1}\bar{x}\| = \|\lambda_2 A\bar{D}A^{-1}\bar{x}\|,$$

where $\bar{D} = \lambda_2^{-1} \operatorname{diag}(0, \lambda_2, \dots, \lambda_n) \in \mathcal{L}_1$, yielding finally

$$\|g_i(\alpha x) - A_1\| = |\lambda_2| \| (A\bar{D}A^{-1}x) / (A^{-1}x)_1 \| \leq \mu_i^{-1} \beta_{i,j,1} \leq \varepsilon.$$

Therefore as $A \in \mathcal{A}^{(i)}$, $A_i \in \mathcal{B}(W_1^{(i)}, \delta)$ whence

$$g_i(\alpha x) \in \mathcal{B}(\mathcal{B}(W_1^{(i)}, \delta), \varepsilon) \subseteq B_i.$$

This completes the proof of the conclusion of (f) about g_i ; for g_i^{-1} , use the same argument with $\alpha' = D_{nn} / (A^{-1}x)_n$ and replace $A_1, W_1^{(i)}$ by $A_n, W_n^{(i)}$.

3. Examples

We shall follow here the proof of Theorem 2 in order to construct examples of open sets of free pairs, first over the reals, second over \mathbb{Q}_p , the p -adic completion of the field of rational numbers. The simplification suggested by Corollary 1.3 is to be followed so that we obtain an 8-parameter family of free pairs, where each parameter is allowed to vary within a certain interval. A computation of appropriately sized intervals, which ensure that the resulting pairs are free is the gist of the calculation.

3.1. Let $F = \mathbb{R}$, $n = k = 2$

$$W = \left(\begin{array}{cc|cc} 1 & 0 & 1/2 & 1 \\ 0 & 1 & 1 & 1/2 \end{array} \right),$$

$p = (1, 1)$, $\gamma = 1/6$. One readily verifies that $B(W, 1/6) \subseteq G_5$. In the notation of Corollary 1.3, with ξ suitably adapted to P ,

$$P = \left(\begin{array}{cc|cc} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{array} \right), \xi(B(W, 1/6)) = \left(\begin{array}{cc|cc} 1 & x_2 & 1/2 + x_3 & 1 \\ x_1 & 1 & 1 & 1/2 + x_4 \end{array} \right)$$

where $|x_1|, |x_2| \leq 1/5$, $-3/4 \leq x_3, x_4 \leq 3/10$. Recall that $\xi(B_1), \xi(C_1), \xi(B_2), \xi(C_2)$ are the projections of $\xi(B(W, 1/6))$ onto the first through fourth column, respectively.

Choosing $\delta = 1/11, \varepsilon = 5/66$ (so that $\varepsilon + \delta = 1/6$), we have, with $\mathcal{A} \sim B(W, \delta)$,

$$\xi(\mathcal{A}) = \left(\begin{array}{cc|cc} 1 & \varepsilon_2 & 1/2 + \varepsilon_3 & 1 \\ \varepsilon_1 & 1 & 1 & 1/2 + \varepsilon_4 \end{array} \right) \begin{array}{l} |\varepsilon_1|, |\varepsilon_2| \leq 1/10, \\ -1/8 \leq \varepsilon_2, \varepsilon_4 \leq 3/20 \end{array}$$

Note that $\mathcal{L}_1 = \{\lambda E_{22} \mid |\lambda| \leq 1\}$.

If $A \in \xi(\mathcal{A}^{(1)})$,

$$A = \begin{pmatrix} 1 & \varepsilon_2 \\ \varepsilon_1 & 1 \end{pmatrix}, (\det A) \cdot A^{-1} = \begin{pmatrix} 1 & -\varepsilon_2 \\ -\varepsilon_1 & 1 \end{pmatrix}, (\det A) \cdot AE_{22}A^{-1} = \begin{pmatrix} -\varepsilon_1\varepsilon_2 & \varepsilon_2 \\ \varepsilon_1 & 1 \end{pmatrix}$$

If $x \in B_2$,

$$\det A \cdot AE_{22}A^{-1}x = (\varepsilon_2(-\varepsilon_1(1/2 + x_3) + 1), \varepsilon_1(1/2 + x_3) + 1),$$

hence

$$|(\det A) \cdot AE_{22}A^{-1}x| \leq 1/2(1/2 + 3/10) + 1 = \frac{108}{100};$$

also

$$|\det(A) \cdot (A^{-1}x)_1| = |1/2 + x_3 - \varepsilon_2| \geq 1/2 - \frac{3}{14} - \frac{1}{10} = \frac{13}{70}.$$

Hence

$$\sup \left\{ \left| \frac{ADA^{-1}x}{(A^{-1}x)_1} \right| \mid A \in \xi(\mathcal{A}^{(1)}), D \in \mathcal{L}_1, x \in B_2 \right\} \leq \frac{108/13}{100/70} = \frac{378}{65}.$$

(A slightly more complicated exact calculation gives $\sup = \frac{72}{13}$).

Similarly, one finds that

$$\sup \left| \frac{ADA^{-1}x}{(A^{-1}x)_1} \right|$$

is bounded above by 90/92 when $x \in C_2$, and by 11/9 when $x = p$.

It follows that $\beta_{1,2,1} \leq 378/65$, and the same bound holds for $\beta_{1,2,2}$ (no coincidence—this reflects symmetries of the chosen set \mathcal{A}).

Thus one gets

$$\mu_1 = \left\{ \frac{\beta_{1,2,1}}{\varepsilon}, \frac{\beta_{1,2,2}}{\varepsilon}, 1 \right\} \leq \frac{24948}{325} < 77.$$

Similar calculations yield $\mu_2 = \frac{14916}{175} < 86$ (additional effort yields the exact values $\mu_1 = 4752/65, \mu_2 = 407/5$).

In conclusion, Theorem 2 guarantees that,

$$\phi_2(A^{(1)}, A^{(2)}, D^{(1)}, D^{(2)})$$

is a free pair whenever

$$A^{(1)} = \begin{pmatrix} 1 & \varepsilon_2 \\ \varepsilon_1 & 1 \end{pmatrix}, |\varepsilon_1|, |\varepsilon_2| \leq 1/10,$$

$$A^{(2)} = \begin{pmatrix} 1/2 + \varepsilon_2 & 1 \\ 1 & 1/2 + \varepsilon_4 \end{pmatrix}, \quad -1/8 \leq \varepsilon_3, \varepsilon_4 \leq 3/20,$$

$$D^{(1)} = \begin{pmatrix} \alpha_1 & 0 \\ 0 & \alpha_2 \end{pmatrix}; \quad |\alpha_1| > \mu_1, |\alpha_2| > 0,$$

$$D^{(2)} = \begin{pmatrix} b_1 & 0 \\ 0 & \beta_2 \end{pmatrix}; \quad |\beta_1| > \mu_2, |\beta_2| > 0,$$

and by Corollary 1.3 the image of the interior of this domain is an open set.

3.2. Let $F = \mathbb{Q}_p$

We indicate one set of parameters; the calculations, following the first example as paradigm, are easily completed due to the ultrametric property of $\| \cdot \|_p$, yielding exact values for μ_1 and μ .

Let

$$W = \left(\begin{array}{cc|cc} 1 & 0 & P & 1 \\ 0 & 1 & 1 & P \end{array} \right), \quad P = (1, 1),$$

with $\gamma = P^{-2}$. We take $\varepsilon = \delta = \gamma$. Then

$$\xi(B(W), \gamma) = \xi(\mathcal{A}) = \left(\begin{array}{cc|cc} 1 & \varepsilon_2 & P + \varepsilon_3 & 1 \\ \varepsilon_1 & 1 & 1 & P + \varepsilon_4 \end{array} \right), \quad |\varepsilon_i|_p \leq P^2.$$

One gets $\mu_1 = \mu_2 = P^3$. Thus one gets an open set of free pairs of $PGL(2, \mathbb{Q}_p)$ as $\phi_2(A^{(1)}, A^{(2)}, D^{(1)}, D^{(2)})$ (sort of) where

$$A^{(1)} = \begin{pmatrix} 1 & P^2 d_2 \\ P^2 d_1 & 1 \end{pmatrix}, \quad A^{(2)} = \begin{pmatrix} P + P^2 d_3 & 1 \\ 1 & P + P^2 d_4 \end{pmatrix} \quad |d_i|_p \leq 1, \quad 1 \leq i \leq 6.$$

$$D^{(1)} = \begin{pmatrix} 1 & 0 \\ 0 & P^2 d_5 \end{pmatrix}, \quad D^{(2)} = \begin{pmatrix} 1 & 0 \\ 0 & P^2 d_6 \end{pmatrix}.$$

4. What about Zariski Topology

It makes sense to ask whether an analogue of Theorem 0 exists in the sense of Zariski topology; namely, one would look for Zariski-open sets of free k -tuples. For fields with a non-trivial absolute value, those sets would be open and dense, in the topology hitherto considered. Thus, we would be looking for something even stronger than conjecture 2 of [1], which has been shown false for \mathbb{C} by Sullivan, as we remarked at the introduction. Indeed, the intended analogue of Theorem 0 is false in general, as we show now by elementary methods.

Here F will be a field with *infinitely many roots of unity*. The only topology considered herewith in the several spaces involved is the Zariski topology ascribed to them by their obvious identification as a constructible subset of some F^m .

Theorem 4.1. *The set $T_n = \{A \in GL_n \mid A^s = I \text{ for some } s \in \mathbb{N}^*\}$ is dense in GL_n .*

Corollary 4.2. *T_n^k is dense in GL_n^k .*

Corollary 4.3. *The set of free k -tuples has empty interior in GL_n^k .*

Since the product of dense sets is dense in the Zariski product topology, Theorem 4.1 implies the first corollary, and this entails the second one immediately. Notice that the existence of infinitely many roots of unity is also a necessary condition for Theorem 4.1 to hold: if a field has only r roots of 1, every member of T_n is a root of the polynomial (in n^2 variables) $(\det A)^r - 1$. However, one feels that Corollary 4.3 is true whenever F is infinite.

The proof of 4.1 requires the two lemmas below:

Lemma 4.4. *The set $T_n(F)$ is dense in the set of semisimple matrices of $GL(n, K)$, where K is the algebraic closure of F .*

Proof. Consider the morphism of varieties

$$\phi: GL(n, K) \times D_n(K) \rightarrow GL(n, K),$$

given by the now usual formula $\phi(A, D) = ADA^{-1}$. Its image is the set of semisimple matrices. The set $T_n(F)$ contains the image of $GL(n, F) \times D_n(R)$, where R is the set of roots of 1 in F , $D_n(R)$ is the set of diagonal matrices all whose diagonal entries are in R . Since R is infinite, it is dense in K , hence $D_n(R)$ is dense in $D_n(K)$; essentially by the same token $GL(n, F)$ is dense in $GL(n, K)$. It follows that $GL(n, K) \times D_n(R)$ is dense in the domain of ϕ ; by continuity, its image is dense in the image of ϕ . \square

Lemma 4.3. *For an algebraically closed field K , the set of matrices with only simple eigenvalues is open in $M_n(K)$.*

Proof. Let $X = (x_{ij})$ be a matrix of indeterminates over R and let $d(x_{ij}) \in K[x_{ij}]$ be the discriminant of the characteristic polynomial P_X of X . A polynomial has multiple roots if and only if its discriminant is zero. Hence a matrix A has multiple eigenvalues if and only if it is a zero of $d(x_{ij})$. \square

Proof of Theorem 4.1. Combining the two lemmas, $T_n(F)$ is dense in a subset of $GL(n, K)$ which contains an open set. As open sets are dense, $T_n(F)$ is dense in $GL(n, K)$; a fortiori, $T_n(F)$ is dense in $GL(n, F)$, since this has the topology induced from $GL(n, K)$. \square

REFERENCES

1. A. MANDEL and J. Z. GONÇALVES, Free k -tuples in linear groups, *Bull. Austr. Math. Soc.* **28** (1983), 152–157.
2. A. NIJENHUIS, Strong derivatives and inverse mappings, *Amer. Math. Monthly* **81** (1971), 969–981 and **83** (1976), 22.
3. W. H. SCHIKHOF, Differentiation in non-archimedean valued fields, *Nederl. Akad. Wetensch. Proc. Ser. A* **73** (1970), 35–44.
4. D. SULLIVAN, Quasi conformal homeomorphisms and dynamics II: Structural stability implies hyperbolicity for Kleinian groups, preprint, 1982.
5. J. TITS, Free subgroups in linear group, *J. Algebra* **20** (1972), 250–270.

INSTITUTO DE MATEMÁTICA E ESTATÍSTICA
UNIVERSIDADE DE SÃO PAULO
CAIXA POSTAL, 20.570—AG. IGUA TEMI
01000—SÃO PAULO—SP—BRAZIL