

# References

- [1] E. Abdalla and M. C. B. Abdalla, "Updating QCD in two-dimensions," *Phys. Rept.* **265**, 253 (1996) [arXiv:hep-th/9503002].
- [2] E. Abdalla, M. C. B. Abdalla and K. D. Rothe, *Nonperturbative Methods in Two-Dimensional Quantum Field Theory*, Singapore: World Scientific (2001).
- [3] A. Abrashkin, Y. Frishman and J. Sonnenschein, "The spectrum of states with one current acting on the adjoint vacuum of massless QCD<sub>2</sub>," *Nucl. Phys. B* **703**, 320 (2004) [arXiv:hep-th/0405165].
- [4] C. Adam, "Charge screening and confinement in the massive Schwinger model," *Phys. Lett. B* **394**, 161 (1997) [arXiv:hep-th/9609155].
- [5] G. S. Adkins, C. R. Nappi and E. Witten, "Static properties of nucleons in the Skyrme model," *Nucl. Phys. B* **228**, 552 (1983).
- [6] S. L. Adler, J. C. Collins and A. Duncan, "Energy momentum tensor trace anomaly in spin 1/2 quantum electrodynamics" *Phys. Rev. D* **15**, 1712 (1977).
- [7] I. Affleck, "On the realization of chiral symmetry in (1+1)-dimensions," *Nucl. Phys. B* **265**, 448 (1986).
- [8] O. Aharony, O. Ganor, J. Sonnenschein and S. Yankielowicz, "On the twisted G/H topological models," *Nucl. Phys. B* **399**, 560 (1993) [arXiv:hep-th/9208040].
- [9] O. Aharony, O. Ganor, J. Sonnenschein, S. Yankielowicz and N. Sochen, "Physical states in G/G models and 2-d gravity," *Nucl. Phys. B* **399**, 527 (1993) [arXiv:hep-th/9204095].
- [10] O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, "Large N field theories, string theory and gravity," *Phys. Rept.* **323**, 183 (2000) [arXiv:hep-th/9905111].
- [11] A. Y. Alekseev and V. Schomerus, "D-branes in the WZW model," *Phys. Rev. D* **60**, 061901 (1999) [arXiv:hep-th/9812193].
- [12] D. Altschuler, K. Bardakci and E. Rabinovici, "A construction of the  $c < 1$  modular invariant partition function," *Commun. Math. Phys.* **118**, 241 (1988).
- [13] L. Alvarez-Gaume, G. Sierra and C. Gomez, "Topics in conformal field theory," contribution to the Knizhnik Memorial Volume, L. Brink, *et al.*, World Scientific. In Brink, L. (ed.) *et al.: Physics and Mathematics of Strings* 16-184 (1989). Singapore.
- [14] F. Antonuccio and S. Dalley, "Glueballs from (1+1)-dimensional gauge theories with transverse degrees of freedom," *Nucl. Phys. B* **461**, 275 (1996) [arXiv:hep-ph/9506456].
- [15] A. Armoni, Y. Frishman and J. Sonnenschein, "The string tension in massive QCD(2)," *Phys. Rev. Lett.* **80**, 430 (1998) [arXiv:hep-th/9709097].
- [16] A. Armoni, Y. Frishman and J. Sonnenschein, "The string tension in two dimensional gauge theories," *Int. J. Mod. Phys. A* **14**, 2475 (1999) [arXiv:hep-th/9903153].
- [17] A. Armoni, Y. Frishman and J. Sonnenschein, "Massless QCD(2) from current constituents," *Nucl. Phys. B* **596**, 459 (2001) [arXiv:hep-th/0011043].
- [18] A. Armoni and J. Sonnenschein, "Mesonic spectra of bosonized QCD in two-dimensions models," *Nucl. Phys. B* **457**, 81 (1995) [arXiv:hep-th/9508006].
- [19] M. F. Atiyah and N. J. Hitchin, "The geometry and dynamics of magnetic monopole. M.B. Porter lectures" Princeton, USA: Princeton University Press (1988) 133p.
- [20] M. F. Atiyah, N. J. Hitchin, V. G. Drinfeld and Yu. I. Manin, "Construction of instantons," *Phys. Lett. A* **65**, 185 (1978).
- [21] F. A. Bais, "To be or not to be? Magnetic monopoles in non-abelian gauge theories," in 't Hooft, Hackensack, G. ed. *Fifty years of Yang-Mills Theory*, New Jersey World Scientific, C 2005.

- [22] A. P. Balachandran, "Solitons in nuclear and elementary physics. Proceedings of the Lewes workshop," World Scientific, 1984.
- [23] I. I. Balitsky and L. N. Lipatov, "The Pomeranchuk singularity in quantum chromodynamics," *Sov. J. Nucl. Phys.* **28**, 822 (1978) [*Yad. Fiz.* **28**, 1597 (1978)].
- [24] V. Baluni, "The Bose form of two-dimensional quantum chromodynamics," *Phys. Lett. B* **90**, 407 (1980).
- [25] T. Banks, "Lectures on conformal field theory," presented at the Theoretical Advanced Studies Institute, St. John's College, Santa Fe, N. Mex., Jul 5 – Aug 1, 1987. Published in Santa Fe: TASI 87:572.
- [26] T. I. Banks and C. M. Bender, "Anharmonic oscillator with polynomial self-interaction," *J. Math. Phys.* **13**, 1320 (1972).
- [27] K. Bardakci and M. B. Halpern, "New dual quark models," *Phys. Rev. D* **3**, 2493 (1971).
- [28] A. Bassetto, G. Nardelli and R. Soldati, *Yang-Mills Theories in Algebraic Noncovariant Gauges: Canonical Quantization and Renormalization*, Singapore World Scientific (1991).
- [29] R. J. Baxter, *Exactly Solved Models in Statistical Mechanics*, London: Academic press (1989).
- [30] K. Becker, M. Becker and J. H. Schwarz, *String Theory and M-Theory: A modern introduction*, Cambridge, UK: Cambridge University Press (2007).
- [31] N. Beisert, "The dilatation operator of  $N = 4$  super Yang-Mills theory" *Phys. Rept.* **405**, 1 (2005) [arXiv:hep-th/0407277].
- [32] A. A. Belavin, A. M. Polyakov, A. S. Shvarts and Yu. S. Tyupkin, "Pseudoparticle solutions of the Yang-Mills equations," *Phys. Lett. B* **59**, 85 (1975).
- [33] A. A. Belavin, A. M. Polyakov and A. B. Zamolodchikov, "Infinite conformal symmetry in two-dimensional quantum field theory," *Nucl. Phys. B* **241**, 333 (1984).
- [34] A. V. Belitsky, V. M. Braun, A. S. Gorsky and G. P. Korchemsky, "Integrability in QCD and beyond," *Int. J. Mod. Phys. A* **19**, 4715 (2004) [arXiv:hep-th/0407232].
- [35] D. E. Berenstein, J. M. Maldacena and H. S. Nastase, "Strings in flat space and pp waves from  $N = 4$  super Yang Mills," *JHEP* **0204**, 013 (2002) [arXiv:hep-th/0202021].
- [36] M. Bernstein and J. Sonnenschein, "A comment on the quantization of the chiral bosons" *Phys. Rev. Lett.* **60**, 1772 (1988).
- [37] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Field Theory*, New York: McGraw-Hill (1965), ISBN 0-07-005494-0.
- [38] G. Bhanot, K. Demeterfi and I. R. Klebanov, "(1+1)-dimensional large  $N$  QCD coupled to adjoint fermions," *Phys. Rev. D* **48**, 4980 (1993) [arXiv:hep-th/9307111].
- [39] D. Bernard and A. Leclair, "Quantum group symmetries and nonlocal currents in 2-D QFT," *Commun. Math. Phys.* **142**, 99 (1991).
- [40] K. M. Bitar and S. J. Chang, "Vacuum tunneling of gauge theory in minkowski space," *Phys. Rev. D* **17**, 486 (1978).
- [41] E. B. Bogomolny, "Stability of classical solutions," *Sov. J. Nucl. Phys.* **24**, 449 (1976) [*Yad. Fiz.* **24**, 861 (1976)].
- [42] V. M. Braun, S. E. Derkachov, G. P. Korchemsky and A. N. Manashov, "Baryon distribution amplitudes in QCD," *Nucl. Phys. B* **553**, 355 (1999) [arXiv:hep-ph/9902375].
- [43] V. M. Braun, G. P. Korchemsky and D. Mueller, "The uses of conformal symmetry in QCD," *Prog. Part. Nucl. Phys.* **51**, 311 (2003) [arXiv:hep-ph/0306057].
- [44] E. Brezin and J. L. Gervais, "Nonperturbative aspects in quantum field theory. Proceedings of Les Houches Winter Advanced Study Institute, March 1978," *Phys. Rept.* **49**, 91 (1979).
- [45] E. Brezin, C. Itzykson, J. Zinn-Justin and J. B. Zuber, "Remarks about the existence of nonlocal charges in two-dimensional models," *Phys. Lett. B* **82**, 442 (1979).
- [46] E. Brezin and S. R. Wadia, "The large  $N$  expansion in quantum field theory and statistical physics: From spin systems to two-dimensional gravity," Singapore, Singapore: World Scientific (1993) 1130 p
- [47] S. J. Brodsky, "Gauge theories on the light-front," *Braz. J. Phys.* **34**, 157 (2004) [arXiv:hep-th/0302121].

- [48] S. J. Brodsky, H. C. Pauli and S. S. Pinsky, "Quantum chromodynamics and other field theories on the light cone," *Phys. Rept.* **301**, 299 (1998) [arXiv:hep-ph/9705477].
- [49] S. J. Brodsky, Y. Frishman, G. P. Lepage and C. T. Sachrajda, "Hadronic wave functions at short distances and the operator product expansion," *Phys. Lett. B* **91**, 239 (1980).
- [50] S. J. Brodsky, Y. Frishman and G. P. Lepage, "On the application of conformal symmetry to quantum field theory," *Phys. Lett. B* **167**, 347 (1986).
- [51] S. J. Brodsky, P. Damgaard, Y. Frishman and G. P. Lepage, "Conformal symmetry: exclusive processes beyond leading order," *Phys. Rev. D* **33**, 1881 (1986).
- [52] R. C. Brower, W. L. Spence and J. H. Weis, "Effects of confinement on analyticity in two-dimensional QCD," *Phys. Rev. D* **18**, 499 (1978).
- [53] G. E. Brown, 'Selected papers, with commentary, of Tony Hilton Royle Skyrme,' Singapore: World Scientific (1994) 438 p. (World Scientific series in 20th century physics, 3).
- [54] R. N. Cahn, *Semisimple Lie Algebras and their Representations*, Menlo Park, USA: Benjamin/Cummings (1984) 158 p. (Frontiers In Physics, 59)
- [55] C. G. Callan, "Broken scale invariance in scalar field theory," *Phys. Rev. D* **2**, 1541 (1970).
- [56] C. G. Callan, N. Coote and D. J. Gross, "Two-dimensional Yang-Mills theory: a model of quark confinement," *Phys. Rev. D* **13**, 1649 (1976).
- [57] C. G. Callan, R. F. Dashen and D. J. Gross, "Toward a theory of the strong interactions," *Phys. Rev. D* **17**, 2717 (1978).
- [58] J. L. Cardy, "Boundary conditions, fusion rules and the Verlinde formula," *Nucl. Phys. B* **324**, 581 (1989).
- [59] J. L. Cardy, "Conformal invariance and statistical mechanics," Les Houches Summer School 1988:0169-246.
- [60] A. Casher, H. Neuberger and S. Nussinov, "Chromoelectric flux tube model of particle production," *Phys. Rev. D* **20**, 179 (1979).
- [61] A. Chodos, "Simple connection between conservation laws in the Korteweg-De Vries and sine-Gordon systems," *Phys. Rev. D* **21**, 2818 (1980).
- [62] E. Cohen, Y. Frishman and D. Gepner, "Bosonization of two-dimensional QCD with flavor," *Phys. Lett. B* **121**, 180 (1983).
- [63] S. R. Coleman, "Quantum sine-Gordon equation as the massive Thirring model," *Phys. Rev. D* **11**, 2088 (1975).
- [64] S. R. Coleman, "More about the massive Schwinger model," *Annals Phys.* **101**, 239 (1976).
- [65] S. R. Coleman, "The uses of instantons," *Subnucl. Ser.* **15**, 805 (1979).
- [66] S. Coleman, *Aspects of Symmetry*, selected Erice lectures of Sidney Coleman, Cambridge, UK, Cambridge Univ. Press (1985).
- [67] S. R. Coleman, The magnetic monopole fifty years later, In the *Unity of Fundamental Interactions*, ed. A. Zichichi: New York, Plenum (1983).
- [68] S. R. Coleman, R. Jackiw and L. Susskind, "Charge shielding and quark confinement in the massive Schwinger model," *Annals Phys.* **93**, 267 (1975).
- [69] S. Cordes, G. W. Moore and S. Ramgoolam, "Large N 2-D Yang-Mills theory and topological string theory," *Commun. Math. Phys.* **185**, 543 (1997) [arXiv:hep-th/9402107].
- [70] E. Corrigan, D. B. Fairlie, S. Templeton and P. Goddard, "A Green's function for the general selfdual gauge field," *Nucl. Phys. B* **140**, 31 (1978).
- [71] E. Corrigan and P. Goddard, "Construction of instanton and monopole solutions and reciprocity," *Annals Phys.* **154**, 253 (1984).
- [72] S. Dalley and I. R. Klebanov, "String spectrum of (1+1)-dimensional large N QCD with adjoint matter," *Phys. Rev. D* **47**, 2517 (1993) [arXiv:hep-th/9209049].
- [73] R. F. Dashen and Y. Frishman, "Four fermion interactions and scale invariance," *Phys. Rev. D* **11**, 2781 (1975).
- [74] R. F. Dashen, B. Hasslacher and A. Neveu, "Nonperturbative methods and extended hadron models in field theory. 2. Two-dimensional models and extended hadrons," *Phys. Rev. D* **10**, 4130 (1974).

- [75] G. Date, Y. Frishman and J. Sonnenschein, "The spectrum of multiflavor QCD in two-dimensions" *Nucl. Phys. B* **283**, 365 (1987).
- [76] G. F. Dell-Antonio, Y. Frishman and D. Zwanziger, "Thirring model in terms of currents: solution and light cone expansions," *Phys. Rev. D* **6**, 988 (1972).
- [77] P. Di Francesco, P. Mathieu, D. Senechal *Conformal Field Theory*, Series:Graduate Texts in Contemporary Physics. New York: Springer-Verlag (1997).
- [78] P. A. M. Dirac, "Theory Of Magnetic Monopoles," In *\*Coral Gables 1976, Proceedings, New Pathways In High-energy Physics, Vol. I\**, New York: Plenum Press, 1976, 1-14.
- [79] F. M. Dittes and A. V. Radyushkin, "Two loop contribution to the evolution of the pion wave function," *Phys. Lett. B* **134**, 359 (1984); M. H. Sarmadi, *Phys. Lett. B* **143**, 471 (1984); S. V. Mikhailov and A. V. Radyushkin, *Nucl. Phys. B* **254**, 89 (1985); G. R. Katz, *Phys. Rev. D* **31**, 652 (1985).
- [80] N. Dorey, T. J. Hollowood, V. V. Khoze, M. P. Mattis and S. Vandoren, "Multi-instanton calculus and the AdS/CFT correspondence in  $N = 4$ " *Nucl. Phys. B* **552**, 88 (1999) [arXiv:hep-th/9901128].
- [81] N. Dorey, T. J. Hollowood, V. V. Khoze and M. P. Mattis, "The calculus of many instantons," *Phys. Rept.* **371**, 231 (2002) [arXiv:hep-th/0206063].
- [82] M. R. Douglas, K. Li and M. Staudacher, "Generalized two-dimensional QCD," *Nucl. Phys. B* **420**, 118 (1994) [arXiv:hep-th/9401062].
- [83] A. V. Efremov and A. V. Radyushkin, "Factorization and asymptotical behavior of pion form-factor in QCD," *Phys. Lett. B* **94**, 245 (1980).
- [84] T. Eguchi and H. Ooguri, "Chiral bosonization on a Riemann surface," *Phys. Lett. B* **187**, 127 (1987).
- [85] S. Elitzur and G. Sarkissian, "D-branes on a gauged WZW model," *Nucl. Phys. B* **625**, 166 (2002) [arXiv:hep-th/0108142].
- [86] J. R. Ellis, Y. Frishman, A. Hanany and M. Karliner, "Quark solitons as constituents of hadrons," *Nucl. Phys. B* **382**, 189 (1992) [arXiv:hep-ph/9204212].
- [87] J. R. Ellis, Y. Frishman and M. Karliner, "Meson baryon scattering in QCD(2) for any coupling," *Phys. Lett. B* **566**, (2003) 201 [arXiv:hep-ph/0305292].
- [88] L. D. Faddeev, "How algebraic Bethe ansatz works for integrable model," arXiv:hep-th/9605187.
- [89] B. L. Feigin and D. B. Fuchs "Skew symmetric differential operators on the line and Verma modules over the Virasoro algebra" *Funct. Anal. Prilozhen* **16**, (1982) 47.
- [90] S. Ferrara, A. F. Grillo and R. Gatto, "Improved light cone expansion," *Phys. Lett. B* **36**, 124 (1971) [*Phys. Lett. B* **38**, 188 (1972)].
- [91] D. Finkelstein and J. Rubinstein, "Connection between spin, statistics, and kinks," *J. Math. Phys.* **9**, 1762 (1968).
- [92] R. Floreanini and R. Jackiw, "Selfdual fields as charge density solitons," *Phys. Rev. Lett.* **59**, 1873 (1987).
- [93] D. Friedan, "Introduction To Polyakov's string theory," Published in Les Houches Summer School 1982:0839.
- [94] D. Friedan, E. J. Martinec and S. H. Shenker, "Conformal invariance, supersymmetry and string theory," *Nucl. Phys. B* **271**, 93 (1986).
- [95] D. Friedan, Z. a. Qiu and S. H. Shenker, "Conformal invariance, unitarity and two-dimensional critical exponents," *Phys. Rev. Lett.* **52**, 1575 (1984).
- [96] Y. Frishman, A. Hanany and J. Sonnenschein, "Subtleties in QCD theory in two-dimensions," *Nucl. Phys. B* **429**, 75 (1994) [arXiv:hep-th/9401046].
- [97] Y. Frishman and M. Karliner, "Baryon wave functions and strangeness content in QCD in two-dimensions," *Nucl. Phys. B* **344**, 393 (1990).
- [98] Y. Frishman and M. Karliner, "Scattering and resonances in QCD(2)," *Phys. Lett. B* **541**, 273 (2002). Erratum-ibid. *B* **562**, 367, (2003). [arXiv:hep-ph/0206001].
- [99] Y. Frishman and J. Sonnenschein, "Bosonization of colored-flavored fermions and QCD in two-dimensions," *Nucl. Phys. B* **294**, 801 (1987).
- [100] Y. Frishman and J. Sonnenschein, "Gauging of chiral bosonized actions" *Nucl. Phys. B* **301**, 346 (1988).

- [101] Y. Frishman and J. Sonnenschein, "Bosonization and QCD in two-dimensions," *Phys. Rept.* **223**, 309 (1993) [arXiv:hep-th/9207017].
- [102] Y. Frishman and W. J. Zakrzewski, "Multibaryons in QCD in two-dimensions," *Nucl. Phys. B* **328**, 375 (1989).
- [103] Y. Frishman and W. J. Zakrzewski, "Explicit expressions for masses and bindings of multibaryons in QCD(2)," *Nucl. Phys. B* **331**, 781 (1990).
- [104] S. Fubini, A. J. Hanson and R. Jackiw, "New approach to field theory," *Phys. Rev. D* **7**, 1732 (1973).
- [105] O. Ganor, J. Sonnenschein and S. Yankielowicz, "The string theory approach to generalized 2-d Yang-Mills theory," *Nucl. Phys. B* **434**, 139 (1995) [arXiv:hep-th/9407114].
- [106] E. G. Gimon, L. A. Pando Zayas, J. Sonnenschein and M. J. Strassler, "A soluble string theory of hadrons," *JHEP* **0305**, 039 (2003) [arXiv:hep-th/0212061].
- [107] D. Gepner, "Nonabelian bosonization and multiflavor QED and QCD in two-dimensions," *Nucl. Phys. B* **252**, 481 (1985).
- [108] D. Gepner and E. Witten, "String theory on group manifolds," *Nucl. Phys. B* **278**, 493 (1986).
- [109] P. H. Ginsparg, "Applied conformal field theory," Published in Les Houches Summer School 1988:1-168, arXiv:hep-th/9108028.
- [110] P. Goddard, A. Kent and D. I. Olive, "Virasoro algebras and coset space models," *Phys. Lett. B* **152**, 88 (1985).
- [111] P. Goddard and D. I. Olive, "Kac-Moody and Virasoro algebras in relation to quantum physics," *Int. J. Mod. Phys. A* **1**, 303 (1986).
- [112] D. Gonzales and A. N. Redlich, "The low-energy effective dynamics of two-dimensional gauge theories," *Nucl. Phys. B* **256**, 621 (1985).
- [113] M. B. Green, J. H. Schwarz and E. Witten, "Superstring theory. Vol. 2: Loop amplitudes, anomalies and phenomenology," Cambridge, UK: Univ. Pr. (1987) 596 P. (Cambridge Monographs On Mathematical Physics)
- [114] C. Gomez, G. Sierra and M. Ruiz-Altaba, "Quantum groups in two-dimensional physics," *Cambridge, UK: Univ. Pr. (1996) 457 p*
- [115] D. J. Gross, "Two-dimensional QCD as a string theory," *Nucl. Phys. B* **400**, 161 (1993) [arXiv:hep-th/9212149].
- [116] D. J. Gross, I. R. Klebanov, A. V. Matytsin and A. V. Smilga, "Screening vs. Confinement in 1+1 Dimensions," *Nucl. Phys. B* **461**, 109 (1996) [arXiv:hep-th/9511104].
- [117] D. J. Gross and A. Neveu, "Dynamical symmetry breaking in asymptotically free field theories," *Phys. Rev. D* **10**, 3235 (1974).
- [118] D. J. Gross and W. Taylor, "Two-dimensional QCD is a string theory," *Nucl. Phys. B* **400**, 181 (1993) [arXiv:hep-th/9301068].
- [119] D. J. Gross and W. Taylor, "Twists and Wilson loops in the string theory of two-dimensional QCD," *Nucl. Phys. B* **403**, 395 (1993) [arXiv:hep-th/9303046].
- [120] I. G. Halliday, E. Rabinovici, A. Schwimmer and M. S. Chanowitz, "Quantization of anomalous two-dimensional models," *Nucl. Phys. B* **268**, 413 (1986).
- [121] M. B. Halpern, "Quantum solitons which are SU(N) fermions," *Phys. Rev. D* **12**, 1684 (1975).
- [122] G. 't Hooft, "A planar diagram theory for strong interactions," *Nucl. Phys. B* **72**, 461 (1974).
- [123] G. 't Hooft, "Magnetic monopoles in unified gauge theories" *Nucl. Phys. B* **79**, 276 (1974).
- [124] G. 't Hooft, "A two-dimensional model for mesons," *Nucl. Phys. B* **75**, 461 (1974).
- [125] G. 't Hooft, "Computation of the quantum effects due to a four-dimensional pseudoparticle," *Phys. Rev. D* **14**, 3432 (1976) [Erratum-ibid. *D* **18**, 2199 (1978)].
- [126] P. Horava, "Topological strings and QCD in two-dimensions," hep-th/9311156, talk given at NATO Advanced Research Workshop on New Developments in String Theory, Conformal Models and Topological Field Theory, Cargese, France, 12–21 May 1993.
- [127] K. Hornbostel, "The application of light cone quantization to quantum chromodynamics in (1+1)-dimensions," Ph.D. thesis, SLAC-R-333, Dec 1988.

- [128] K. Hornbostel, S. J. Brodsky and H. C. Pauli, "Light cone quantized QCD in (1+1)-dimensions," *Phys. Rev. D* **41**, 3814 (1990).
- [129] C. Imbimbo and A. Schwimmer, "The Lagrangian formulation of chiral scalars," *Phys. Lett. B* **193**, 455 (1987).
- [130] C. Itzykson and J. B. Zuber, *Quantum Field Theory, New York, USA: McGraw-Hill* (1980) 705 P.(International Series In Pure and Applied Physics)
- [131] R. Jackiw and C. Rebbi, "Vacuum periodicity in a Yang-Mills quantum theory," *Phys. Rev. Lett.* **37**, 172 (1976).
- [132] R. Jackiw, C. Nohl and C. Rebbi, "Conformal properties of pseudoparticle configurations", *Phys. Rev. D* **15**, 1642 (1977).
- [133] A. D. Jackson and M. Rho, "Baryons as chiral solitons," *Phys. Rev. Lett.* **51**, 751 (1983).
- [134] B. Julia and A. Zee, "Poles with both magnetic and electric charges in nonabelian gauge theory," *Phys. Rev. D* **11**, 2227 (1975).
- [135] N. Ishibashi, "The boundary and crosscap states in conformal field theories," *Mod. Phys. Lett. A* **4**, 251 (1989).
- [136] V. G. Kac, "Simple graded algebras of finite growth," *Funct. Anal. Appl.* **1**, 328 (1967).
- [137] L. P. Kadanoff, "Correlators along the line of two dimensional Ising model," *Phys. Rev.* **188**, 859 (1969).
- [138] M. Kaku, *Strings, Conformal Fields, and M-Theory*, New York, USA: Springer (2000) 531 p.
- [139] V. A. Kazakov, "Wilson loop average for an arbitrary contour in two-dimensional U(N) gauge theory," *Nucl. Phys. B* **179**, 283 (1981).
- [140] S. V. Ketov, *Conformal Field Theory*, Singapore, Singapore: World Scientific (1995) 486 p.
- [141] V. V. Khoze, M. P. Mattis and M. J. Slater, "The instanton hunter's guide to supersymmetric SU(N) gauge theory," *Nucl. Phys. B* **536**, 69 (1998) [arXiv:hep-th/9804009].
- [142] E. Kiritsis, *String Theory in a Nutshell*, Princeton, USA: Univ. Pr. (2007) 588 p.
- [143] V. G. Knizhnik and A. B. Zamolodchikov, "Current algebra and Wess-Zumino model in two dimensions," *Nucl. Phys. B* **247**, 83 (1984).
- [144] W. Krauth and M. Staudacher, "Non-integrability of two-dimensional QCD," *Phys. Lett. B* **388**, 808 (1996) [arXiv:hep-th/9608122].
- [145] E. A. Kuraev, L. N. Lipatov and V. S. Fadin, "The Pommeranchuk singularity in non-abelian gauge theories," *Sov. Phys. JETP* **45**, 199 (1977) [*Zh. Eksp. Teor. Fiz.* **72**, 377 (1977)]. "Multi-reggeon processes in the Yang-Mills theory," *Sov. Phys. JETP* **44**, 443 (1976) [*Zh. Eksp. Teor. Fiz.* **71**, 840 (1976)].
- [146] D. Kutasov, "Duality off the critical point in two-dimensional systems with nonabelian symmetries," *Phys. Lett. B* **233**, 369 (1989).
- [147] D. Kutasov, "Two-dimensional QCD coupled to adjoint matter and string theory," *Nucl. Phys. B* **414**, 33 (1994) [arXiv:hep-th/9306013].
- [148] D. Kutasov and A. Schwimmer, "Universality in two-dimensional gauge theory," *Nucl. Phys. B* **442**, 447 (1995) [arXiv:hep-th/9501024].
- [149] T. D. Lee and Y. Pang, "Nontopological solitons," *Phys. Rept.* **221**, 251 (1992).
- [150] G. P. Lepage and S. J. Brodsky, "Exclusive processes in quantum chromodynamics: evolution equations for hadronic wave functions and the form-factors of mesons," *Phys. Lett. B* **87**, 359 (1979).
- [151] L. N. Lipatov, *The Creation of Quantum Chromodynamics and the Effective Energy*, Bologna, Italy: Univ. Bologna (1998) 367 p.
- [152] J. H. Lowenstein and J. A. Swieca, "Quantum electrodynamics in two-dimensions," *Annals Phys.* **68**, 172 (1971).
- [153] M. Luscher, "Quantum nonlocal charges and absence of particle production in the two-dimensional nonlinear Sigma model," *Nucl. Phys. B* **135**, 1 (1978).
- [154] D. Lust and S. Theisen, "Lectures on string theory," *Lect. Notes Phys.* **346**, 1 (1989).
- [155] A. Maciocia, "Metrics on the moduli spaces of instantons over Euclidean four space," *Commun. Math. Phys.* **135**, 467 (1991).
- [156] G. Mack and A. Salam, "Finite component field representations of the conformal group," *Annals Phys.* **53**, 174 (1969).

- [157] V. G. Makhankov, Y. P. Rybakov and V. I. Sanyuk, *The Skyrme model: Fundamentals, methods, applications*, Berlin, Germany: Springer (1993) 265 p. (Springer series in nuclear and particle physics).
- [158] J. M. Maldacena, "The large N limit of superconformal field theories and supergravity," *Adv. Theor. Math. Phys.* **2**, 231 (1998) [*Int. J. Theor. Phys.* **38**, 1113 (1999)] [arXiv:hep-th/9711200].
- [159] S. Mandelstam, "Soliton operators for the quantized sine-Gordon equation," *Phys. Rev. D* **11**, 3026 (1975).
- [160] A. V. Manohar, "Large N QCD," arXiv:hep-ph/9802419, Published in \*Les Houches 1997, Probing the standard model of particle interactions, Pt. 2\* 1091-1169.
- [161] A. A. Migdal, "Recursion equations in gauge field theories," *Sov. Phys. JETP* **42**, 413 (1975) [*Zh. Eksp. Teor. Fiz.* **69**, 810 (1975)].
- [162] J. A. Minahan and K. Zarembo, "The Bethe-ansatz for  $N = 4$  super Yang-Mills," *JHEP* **0303**, 013 (2003) [arXiv:hep-th/0212208].
- [163] C. Montonen and D. I. Olive, "Magnetic monopoles as gauge particles?," *Phys. Lett. B* **72**, 117 (1977).
- [164] R. V. Moody, "Lie algebras associated with generalized Cartan matrices," *Bull. Am. Math. Soc.* **73**, 217 (1967).
- [165] M. Moshe and J. Zinn-Justin, "Quantum field theory in the large N limit: A review," *Phys. Rept.* **385**, 69 (2003) [arXiv:hep-th/0306133].
- [166] D. Mueller, "Constraints for anomalous dimensions of local light cone operators in  $\phi^4$  in six-dimensions theory," *Z. Phys. C* **49**, 293 (1991); *Phys. Rev. D* **49**, 2525 (1994).
- [167] W. Nahm, "A simple formalism for the BPS monopole," *Phys. Lett. B* **90**, 413 (1980).
- [168] N. K. Nielsen, "Gauge invariance and broken conformal symmetry," *Nucl. Phys. B* **97**, 527 (1975).
- [169] N. K. Nielsen, "The energy momentum tensor in a nonabelian quark gluon theory," *Nucl. Phys. B* **120**, 212 (1977).
- [170] S. P. Novikov, "Multivalued functions and functionals, An analogue to Morse theory," *Sov. Math. Dock.* **24**, 222 (1981)
- [171] T. Ohrndorf, "Constraints from conformal covariance on the mixing of operators of lowest twist," *Nucl. Phys. B* **198**, 26 (1982).
- [172] R. D. Peccei and H. R. Quinn, "Constraints imposed by CP conservation in the presence of instantons," *Phys. Rev. D* **16**, 1791 (1977).
- [173] M. E. Peskin and D. V. Schroeder, *An Introduction To Quantum Field Theory*, Reading, USA: Addison-Wesley (1995) 842 p.
- [174] J. Polchinski, *String Theory. Vol. 2: Superstring Theory and Beyond*, Cambridge, UK: Univ. Pr. (1998) 531 p.
- [175] A. M. Polyakov, "Particle spectrum in quantum field theory," *JETP Lett.* **20**, 194 (1974) [*Pisma Zh. Eksp. Teor. Fiz.* **20**, 430 (1974)].
- [176] A. M. Polyakov, "Hidden symmetry of the two-dimensional chiral fields," *Phys. Lett. B* **72**, 224 (1977).
- [177] A. M. Polyakov, "Quantum geometry of bosonic strings," *Phys. Lett. B* **103**, 207 (1981).
- [178] A. M. Polyakov, *Gauge Fields and Strings*, Chur, Switzerland: Harwood (1987) 301 p. (Contemporary concepts in Physics, 3).
- [179] A. M. Polyakov and P. B. Wiegmann, "Goldstone fields in two-dimensions with multivalued actions," *Phys. Lett. B* **141**, 223 (1984).
- [180] M. K. Prasad and C. M. Sommerfield, "An exact classical solution for the 't Hooft monopole and the Julia-Zee dyon," *Phys. Rev. Lett.* **35**, 760 (1975).
- [181] E. Rabinovici, A. Schwimmer and S. Yankielowicz, "Quantization in the presence of Wess-Zumino terms," *Nucl. Phys. B* **248**, 523 (1984).
- [182] R. Rajaraman, *An Introduction to Solitons and Instantons in Quantum Field Theory*, Amsterdam, Netherlands: North-holland (1982) 409p.
- [183] C. Rebbi and G. Soliani, *Solitons and particles*, World Scientific 1984.
- [184] B. E. Rusakov, "Loop averages and partition functions in  $U(N)$  gauge theory on two-dimensional manifolds," *Mod. Phys. Lett. A* **5**, 693 (1990).

- [185] T. Schafer and E. V. Shuryak, "Instantons in QCD," *Rev. Mod. Phys.* **70**, 323 (1998) [arXiv:hep-ph/9610451].
- [186] J. Schechter and H. Weigel, "The Skyrme model for baryons," arXiv:hep-ph/9907554.
- [187] T. D. Schultz, D. C. Mattis and E. H. Lieb, "Two-dimensional Ising model as a soluble problem of many fermions," *Rev. Mod. Phys.* **36**, 856 (1964).
- [188] M. A. Shifman, *Instantons in Gauge Theories*, Singapore, Singapore: World Scientific (1994) 488 p.
- [189] E. V. Shuryak, "The role of instantons in quantum chromodynamics. 1. Physical vacuum," *Nucl. Phys. B* **203**, 93 (1982).
- [190] J. S. Schwinger, "Gauge invariance and mass. 2," *Phys. Rev.* **128**, 2425 (1962).
- [191] N. Seiberg and E. Witten, "Monopoles, duality and chiral symmetry breaking in N=2 supersymmetric QCD," *Nucl. Phys. B* **431**, 484 (1994) [arXiv:hep-th/9408099].
- [192] N. Seiberg and E. Witten, "Monopole condensation, and confinement in N=2 supersymmetric Yang-Mills theory," *Nucl. Phys. B* **426**, 19 (1994) [Erratum-ibid. B **430**, 485 (1994)] [arXiv:hep-th/9407087].
- [193] Y. M. Shnir, *Magnetic Monopoles*, Berlin, Germany: Springer (2005) 532 p.
- [194] W. Siegel, "Manifest Lorentz invariance sometimes requires nonlinearity," *Nucl. Phys. B* **238**, 307 (1984).
- [195] T. H. R. Skyrme, "A Nonlinear theory of strong interactions," *Proc. Roy. Soc. Lond. A* **247**, 260 (1958).
- [196] T. H. R. Skyrme, "Particle states of a quantized meson field," *Proc. Roy. Soc. Lond. A* **262**, 237 (1961).
- [197] T. H. R. Skyrme, "A unified field theory of mesons and baryons," *Nucl. Phys.* **31**, 556 (1962).
- [198] Smirnov, F. A. "Form factors in completely integrable models of quantum field theory," *Adv. Ser. Math. Phys.* **14**:1–208 (1992).
- [199] J. Sonnenschein, "Chiral bosons" *Nucl. Phys. B* **309**, 752 (1988).
- [200] M. Spiegelglas and S. Yankielowicz, "G/G topological field theories by cosetting G(K)," *Nucl. Phys. B* **393**, 301 (1993) [arXiv:hep-th/9201036].
- [201] P. J. Steinhardt, "Baryons and baryonium in QCD in two-dimensions," *Nucl. Phys. B* **176**, 100 (1980).
- [202] M. Stone, *Bosonization*, Singapore: World Scientific (1994) 539 p.
- [203] H. Sugawara, "A Field theory of currents," *Phys. Rev.* **170**, 1659 (1968).
- [204] K. Symanzik, "Small distance behavior in field theory and power counting," *Commun. Math. Phys.* **18**, 227 (1970).
- [205] W. E. Thirring, "A soluble relativistic field theory," *Annals Phys.* **3**, 91 (1958).
- [206] C. B. Thorn, "Computing the Kac determinant using dual model techniques and more about the no-ghost theorem," *Nucl. Phys. B* **248**, 551 (1984).
- [207] S. B. Treiman, R. Jackiw and D. Gross *Current Algebra and its Applications* (Princeton, University Press, New Jersey, 1972).
- [208] A. I. Vainshtein, V. I. Zakharov, V. A. Novikov and M. A. Shifman, "ABC of instantons," *Sov. Phys. Usp.* **25**, 195 (1982) [*Usp. Fiz. Nauk* **136**, 553 (1982)].
- [209] S. Vandoren and P. van Nieuwenhuizen, "Lectures on instantons," arXiv:0802.1862 [hep-th].
- [210] E. P. Verlinde, "Fusion rules and modular transformations in 2d conformal field theory," *Nucl. Phys. B* **300**, 360 (1988).
- [211] E. P. Verlinde and H. L. Verlinde, "Chiral bosonization, determinants and the string partition function," *Nucl. Phys. B* **288**, 357 (1987).
- [212] M. S. Virasoro, "Subsidiary conditions and ghosts in dual resonance models," *Phys. Rev. D* **1**, 2933 (1970).
- [213] M. Wakimoto, "Fock representations of the affine Lie algebra A1(1)," *Commun. Math. Phys.* **104**, 605 (1986).
- [214] E. J. Weinberg and P. Yi, "Magnetic monopole dynamics, supersymmetry, and duality," *Phys. Rept.* **438**, 65 (2007) [arXiv:hep-th/0609055].



- [215] S. Weinberg, *The Quantum Theory of Fields*. Vol. 1: *Foundations*, Cambridge, UK: Univ. Pr. (1995) 609 p.
- [216] G. Veneziano, "U(1) Without instantons," *Nucl. Phys. B* **159**, 213 (1979).
- [217] J. Wess and B. Zumino, "Consequences of anomalous Ward identities," *Phys. Lett. B* **37**, 95 (1971).
- [218] F. Wilczek, "Inequivalent embeddings of SU(2) and instanton interactions," *Phys. Lett. B* **65**, 160.
- [219] K. G. Wilson, "Nonlagrangian models of current algebra," *Phys. Rev.* **179**, 1499 (1969).
- [220] E. Witten, "Some exact multipseudoparticle solutions of classical Yang-Mills theory," *Phys. Rev. Lett.* **38**, 121 (1977).
- [221] E. Witten, "Instantons, the Quark model, and the 1/N expansion," *Nucl. Phys. B* **149**, 285 (1979).
- [222] E. Witten, "Baryons in the 1/N expansion," *Nucl. Phys. B* **160**, 57 (1979).
- [223] E. Witten, "Large N chiral dynamics," *Annals Phys.* **128**, 363 (1980).
- [224] E. Witten, "Nonabelian bosonization in two dimensions," *Commun. Math. Phys.* **92**, 455 (1984).
- [225] E. Witten, "Global aspects of current algebra," *Nucl. Phys. B* **223**, 422 (1983).
- [226] E. Witten, "Current algebra, baryons, and quark confinement," *Nucl. Phys. B* **223**, 433 (1983).
- [227] E. Witten, "On holomorphic factorization of WZW and coset models," *Commun. Math. Phys.* **144**, 189 (1992).
- [228] E. Witten, "Two-dimensional gauge theories revisited," *J. Geom. Phys.* **9**, 303 (1992) [arXiv:hep-th/9204083].
- [229] C. N. Yang and R. L. Mills, "Conservation of isotopic spin and isotopic gauge invariance," *Phys. Rev.* **96**, 191 (1954).
- [230] C. N. Yang, "Some exact results for the many body problems in one dimension with repulsive delta function interaction," *Phys. Rev. Lett.* **19**, 1312 (1967).
- [231] I. Zahed and G. E. Brown, "The Skyrme model," *Phys. Rept.* **142**, 1 (1986).
- [232] A. B. Zamolodchikov, "Renormalization group and perturbation theory near fixed points in two-dimensional field theory," *Sov. J. Nucl. Phys.* **46**, 1090 (1987) [*Yad. Fiz.* **46**, 1819 (1987)].
- [233] A. B. Zamolodchikov, "Exact solutions of conformal field theory in two-dimensions and critical phenomena," *Rev. Math. Phys.* **1**, 197 (1990).
- [234] A. B. Zamolodchikov, "Thermodynamic Bethe ansatz in relativistic models, scaling three state Potts and Lee-Yang models," *Nucl. Phys. B* **342**, 695 (1990).
- [235] A. B. Zamolodchikov and A. B. Zamolodchikov, "Factorized S-matrices in two dimensions as the exact solutions of certain relativistic quantum field models," *Annals Phys.* **120**, 253 (1979).
- [236] A. B. Zamolodchikov, A. B. Zamolodchikov and I. M. Khalatnikov, "Physics Reviews, Vol. 10, pt. 4: Conformal field theory and critical phenomena in two-dimensional systems," London, UK: Harwood (1989) 269–433. (Soviet scientific reviews, Section A, 10.4)
- [237] B. Zwiebach, *A First Course in String Theory*, Cambridge, UK: Univ. Pr. (2004) 558 p

