

Molecular epidemiology of *Shigella flexneri* in a diarrhoea-endemic area of Lima, Peru

C. M. FERNANDEZ-PRADA¹*, M. M. VENKATESAN¹, A. A. FRANCO²,
C. F. LANATA³, R. B. SACK², A. B. HARTMAN¹ AND W. SPIRA²

¹ Walter Reed Army Institute of Research (WRAIR), Silver Spring, MD 20910, USA

² Johns Hopkins University, Baltimore, MD 21205, USA

³ Instituto de Investigacion Nutricional, Lima, Peru

(Accepted 30 September 2003)

SUMMARY

A year-long community-based study of diarrhoeal diseases was conducted in Canto Grande, a periurban community in Lima, Peru. In 109 (34%) houses out of 323 that were visited, at least one individual was detected with shigellosis. The frequency of the 161 shigella isolates obtained was as follows: 117 *S. flexneri* (73%), 21 *S. boydii* (13%), 15 *S. dysenteriae* (9%), and 8 *S. sonnei* (5%). Using a non-radioactive *ipaH* gene probe as a molecular epidemiological tool, a total of 41 *S. flexneri* strains were shown to be distributed in 25 intra-family comparisons by pairs (icp). Further subdivision, based on a comparison of the serotype, plasmid profile, antibiotic resistances and *ipaH* hybridization patterns indicated that Group I, with 11 icp (44%), had strains that were identical, Group II with 8 icp (32%), had strains that were different and Group III with 6 icp (24%), had strains with the same serotype and identical *ipaH* profiles but with differences in other markers. This data indicates that a diversity of shigella clones circulated in this community resulting from both clonal spread and horizontal transfer of genetic elements. Furthermore, *ipaH* profiling of isolates can be used not only to differentiate between closely related shigella strains but also with other parameters, help to understand the dynamics of the generation of new clones of pathogenic bacteria.

INTRODUCTION

Each year, more than 5 million children die in developing countries due to diarrhoeal diseases, and approximately 10–20% of these deaths are caused by bacillary dysentery produced principally by *Shigella* [1–3]. A recent study estimated that the annual number of shigella episodes throughout the world was 164·7 million, of which 163·2 million were in

developing countries with 1·1 million deaths [4]. Shigellosis is also a public-health problem in developed countries, but is restricted to closed groups such as the military, mental hospitals, day-care nurseries, prisons, or cruise-ships [2, 3]. In the United States more than 20 000 cases were reported to the Centers for Disease Control, Atlanta, in 2001 [5].

In developing countries, shigella infections are caused principally by *S. flexneri*, followed by *S. sonnei*; while in developed countries, they are caused principally by *S. sonnei*, followed by *S. flexneri* [4, 6]. Contaminated food, flies, unclean water as well as the faecal–oral route have contributed to the transmission and high incidence of shigella organisms, especially for the small inoculum size required for infection

* Author for correspondence: Dr C. Fernandez-Prada, Department of Bacterial Diseases, Division of Communicable Diseases and Immunology, Walter Reed Army Institute of Research, Bldg. 503, Rm 2N57, Silver Spring, MD 20910, USA.

The views expressed herein are those of the authors and do not necessarily represent those of the Department of the Army or the Department of Defense (para 4-3, AR 360-5).

[2, 3]. The 50% infectious dose (ID₅₀) of shigella strains for humans has been reported to be as low as 100–200 bacteria [7]. This explains why the disease can be easily spread by person-to-person contact as well as by contaminated food and water [3].

Shigella is not only pathogenic to humans, but also to non-human primates like gorillas, chimpanzees and monkeys. All virulent *Shigella* species and enteroinvasive *Escherichia coli* (EIEC) contain plasmids of 120–140 MDa, carrying genes that, in conjunction with genes present in 'pathogenicity islands' on the chromosome, allow these organisms to invade, multiply and spread from cell to cell to extend the foci of infection [2, 8]. Infection with virulent *Shigella* or EIEC results in the production of immune serum with antibodies to surface-located and secreted proteins of the bacteria [2, 9]. Among these, IpaA, IpaB, IpaC, IpaD, VirG and IpaH proteins are encoded by genes on the virulence plasmid and play a role in the pathogenesis of the disease.

The *ipaH* gene is unique in that multiple copies have been detected on the invasion plasmid and on the chromosome of all *Shigella* serotypes and EIEC [8, 10–12], but not on the chromosome of non-pathogenic *E. coli* strains [13]. *ipaH* was initially isolated from a λ gt11 expression library of the invasion plasmid (pWR100) of *S. flexneri* 5, strain M90T [14]. The first sequence that was analysed (*ipaH*_{7.8}) did not have a *Hind*III site within its structural gene [11]. When this sequence was used as a probe against *Hind*III-digested DNA on Southern blots of M90T, five copies were identified on pWR100, designated *ipaH*_{9.8}, *ipaH*_{7.8}, *ipaH*_{4.5}, *ipaH*_{2.5} and *ipaH*_{1.4}, with the subscripts indicating the sizes of the *Hind*III band which hybridized to the *ipaH*_{7.8} probe [11]. The use of oligonucleotide probes of the *ipaH*_{7.8} gene seems to be useful in genetic lineage and epidemiological studies of shigellae and EIEC [10, 15, 16] (Hartman AB, unpublished observations). The development of specific non-radioactive *ipaH*-related DNA probes to identify *Shigella* and EIEC has enabled investigators to determine the prevalence of these organisms in different populations, to identify mixed infection and rare serotypes, and to screen for *Shigella* and EIEC in large numbers of patients [16–18]. Additionally, the sensitivity and specificity of the polymerase chain reaction (PCR) for detection of DNA sequences specific to shigellae and EIEC, like the *ipaH*, *virF* and *ial* loci [15, 17–20] have indicated the importance of this technique in diagnostic and epidemiological shigella studies.

In the present study, a non-radioactive *ipaH* probe was evaluated as a molecular epidemiological tool to detect genetic polymorphisms between *S. flexneri* strains from an endemic area of shigellosis in Lima, Peru. This procedure has been coupled to several other epidemiological markers such as serotype, biochemical reactions, antibiotic susceptibility and plasmid profile.

MATERIALS AND METHODS

Epidemiological investigations

As part of a prospective, community-based study of diarrhoeal diseases conducted in Canto Grande, a periurban community of low socioeconomic conditions and poor hygienic conditions in Lima, Peru [21–23], all children <3 years of age from a representative sample of 400 households were visited twice per week at their households by trained field workers as described elsewhere [24, 25]. A stool specimen or a rectal swab was collected from any child who developed diarrhoea (defined as three or more liquid or semi-liquid stools passed in a 24-h period or with one bloody stool) and were placed in a clean cup with a lid or in Cary–Blair (BBL) and transported with cold packs to the laboratory, where they were processed within 24 h of collection [25]. Between February 1985 and February 1986, 37 index-children <2 years of age with diarrhoea and a shigella strain identified in their stool sample were entered into this study. After obtaining informed consent, all family members from the household of the index-child were asked to participate in this study, as well as all family members of up to nine households selected from the same neighbourhood of the index-child. The selection was made applying the following procedures. Once a shigella strain was identified in the laboratory, a study team visited the household of the index-child as soon as possible (<60 h from the time the index-child was cultured) and all available family members were interviewed. Three households with a family member who had contact with the index-child since his/her diarrhoea started were identified, as well as three other households with a family member who had been in contact with a family member of the index-child, but not with the index-child. Three additional households from the same neighborhood of the index-child family without any type of recent contact with any member of the index-child were identified by door-to-door survey, starting in the same block of households

of the index-child. The group of the index-child household and the three types of household controls constituted one family study of up to ten households each. A daily rectal swab was taken from all consenting individuals of any age from all these households during one week.

Bacteriological investigations

The rectal swab samples were directly inoculated in the field onto Salmonella–Shigella, MacConkey, or Hecktoen agar plates (Difco Laboratories, Detroit, MI, USA). The plates were maintained at room temperature and immediately transported to the Microbiology Laboratory of the Universidad Peruana Cayetano Heredia, Lima, Peru. The plates were incubated overnight at 37 °C. Non-lactose-fermenting colonies were inoculated onto tryptic sugar iron (TSI), lysine iron agar (LIA), and urea tubes and examined for motility, glucose fermentation, production of gas, indole, hydrogen sulphide, and hydrolysis of urea. Final identification was performed by agglutination with specific polyvalent antisera [Laboratorio Nacional de Referencia de Enteropatógenos (LANARE), Lima, Peru]. A total of 161 shigella strains were isolated from humans during the year of study. Only one colony per individual was immediately stored at –70 °C in brain heart infusion broth (BHI; Difco Laboratories) with 25% (v/v) glycerol. When needed, the stock cultures of bacteria were plated onto tryptic soy agar (TSA) plates supplemented with 0.05% Congo Red dye and 0.2% galactose and incubated overnight at 37 °C. Congo Red-positive (CR+) colonies were selected as invasives and used for further characterization of the strains. Forty-one *S. flexneri* strains from households with two or more individuals infected with this species were selected to further characterize and compare the strains as follows:

- (1) biochemical analysis using the API 20E system;
- (2) serological analysis where the strains identified as shigella were sent to LANARE, Lima, Peru, for their corresponding serotyping;
- (3) antibiotic susceptibility testing which was done on Muller–Hinton agar plates by the disk diffusion method of Bauer et al. [26], antimicrobial agents tested included: ampicillin (A), cephalothionin (Cp), chloramphenicol (C), doxycycline (D), erythromycin (E), gentamicin (G), kanamycin (K), nalidixic acid (NA), neomycin (N), streptomycin (S), tetracycline (T), trimethoprim–sulphamethoxazole (Tsx), and triple sulfa (Ts);
- (4) analysis of virulence using the Sereny test [27], where approximately 10⁸ cells were inoculated into the conjunctival sac of a guinea-pig, and strains that elicited keratoconjunctivitis within 48 h were considered Sereny positive;
- (5) plasmid profile analysis performed by the method of Birnboim and Doly [28], where plasmids were detected by electrophoresis in 0.8% agarose gels containing 0.5 µg of ethidium bromide/ml and photographed with UV light illumination using a polaroid camera (Fotodyne, Inc., New Berlin, WI, USA);
- (6) *ipaH* hybridization patterns using non-radioactive and radioactive *ipaH* probes.

Plasmid DNA was prepared from 2 ml cultures using the method of Birnboim and Doly [28] or from 100–200 ml cultures by a modified version of the method described by Cassie and Denarie [14]. Approximately 1 µg of purified DNA plasmid was digested with *Hind*III or *Sal*I (Boehringer Mannheim, Indianapolis, IN, USA) according to the manufacturer's instructions. DNA was electrophoresed on 0.8% agarose gels containing 0.5 µg of ethidium bromide/ml and photographed with UV light illumination using a polaroid camera (Fotodyne, Inc.). The separated DNA fragments were transferred to nitrocellulose membranes (Schleicher & Schuell, Keene, NH, USA) by the method of Southern [28], and hybridized with non-radiolabelled or radiolabelled synthetic *ipaH* oligoprobes. The non-radioactive *ipaH* oligoprobe used in this study has been previously described [16], with the genetic sequence 5'-d CtG GAG GAC ATT GCC CGG G-3', where 't' indicated a modification with suberyl-alkaline phosphatase (Molecular Biosystems, San Diego, CA, USA). Prehybridization was carried out in 5× SSPE [5× SSPE is 0.9 M NaCl, 50 mM NaPO₄ (pH 7.7), and 5 mM EDTA] with 1% SDS/0.5% polyvinylpyrrolidone (Sigma-Aldrich, St. Louis, MO, USA) containing 100 µg of yeast tRNA (Gibco-BRL, Life Technologies, Gaithersburg, MD, USA) per ml for 15 min at 42 °C; this was followed by hybridization of each filter in 3 ml of the same buffer for 30 min at 42 °C by using 5 µl of the *ipaH* oligoprobe (250 nm/ml). The filter was washed twice at 50 °C for 10 min each time in 1× SSPE/1% SDS for the first wash and in 1× SSPE/1% Triton X-100 for the second wash. The third wash in 1× SSPE was carried out at room temperature for 10 min. Detection of bound probe was accomplished by incubation of filters for 15 h

at room temperature in the dark in 5 ml of 0.1 M Tris-HCl (pH 9.5)/0.1 M NaCl/50 mM MgCl₂ as the alkaline phosphatase substrate with 0.17 mg of 5-bromo-4-chloro-3-indolylphosphate *p*-toluidine salt (BCIP) and 0.33 mg of nitroblue tetrazolium chloride (NBT) per ml. Substrate development was terminated by washing the filters in distilled water, then the filters were allowed to dry in the dark. The same *ipaH* oligoprobe was end labelled with [γ -³²P]dATP by using T4 polynucleotide kinase (5'-DNA terminus labelling kit, Gibco-BRL) as previously described [10, 13]. The filters were hybridized and further processed as outlined previously [13]. Briefly, hybridizations were carried out for 16 h at 37 °C in 1× SSPE, 0.2% SDS, 5× Denhardt's solution, 0.01% NaPPi and 100 µl/ml denatured calf-thymus DNA. After washing the filters several times with 1× SSPE/0.2% SDS/0.01% NaPPi, the filters were exposed to X-ray film xAR (Eastman Kodak Co., Rochester, NY, USA).

Numerical analysis

The hybridization profiles of 41 *S. flexneri* strains were obtained after restriction enzyme digestion of their plasmid with *Hind*III and *Sal*I. These profiles were used to generate a 0–1 character-state matrix for computer-based data storage and processing. One was scored when a band was present and 0 when absent. A dissimilarity matrix was derived from the character-state matrix by calculating Jaccard's distance [29]. Classification was performed with a clustering algorithm according to the unweighted-pair group method with arithmetic averages (UPGMA) and dendrograms were produced by using a computer program elaborated by E. Gerres (Laboratoire d'Ecologie Médicale et Pathologie Parasitaire, Faculté de Médecine Montpellier, France) [30].

Statistical analyses

Statistical analysis was done by Student's *t* test using the INSTAT statistical analysis package (Graph Pad Software, Inc., San Diego, CA, USA). Significance was taken as $P < 0.05$.

RESULTS

Distribution of *Shigella* in Canto Grande

During the year of the study, 37 family studies were initiated and a total of 323 houses were visited in Canto Grande, an endemic area of shigellosis. In 109

houses (34%), at least one individual was detected with shigellosis, and in 34 of these houses (31%), between 2 and 4 individuals were infected with *Shigella* spp. A total of 161 shigella isolates were obtained and the distribution by species was as follows: 117 *S. flexneri* (73%), 21 *S. boydii* (13%), 15 *S. dysenteriae* (9%), and 8 *S. sonnei* (5%). Between 1 and 10 isolates (mean 4 isolates per family study) and between 1 and 6 serotypes (mean 3 serotypes per family study) were detected per family study, indicating a diversity of strains circulating in this population. One of 37 family studies showed the highest variation with 7 isolates of 6 different serotypes of shigella that were isolated. These 6 serotypes in this family study included *S. flexneri* 1a, *S. flexneri* 2a (2 isolates), *S. flexneri* 3a, *S. flexneri* 4a, *S. flexneri* 6 and *S. boydii* 11.

Comparison of *S. flexneri* in households with multiple infections

Since *S. flexneri* strains were isolated at a higher frequency in this study than other shigella serogroups (73%, $P < 0.05$), this species was selected to compare households with multiple shigella infections. The isolates of each household were compared by pairs. With two shigella isolates from a household, one intra-family comparison per pair (icp) was obtained. In a household with three shigella isolates, three combinations per pair were obtained (3 icp). In this way, 41 *S. flexneri* strains from 19 households with multiple shigella infections generated 25 icp cases. These icp cases were further characterized and divided into three groups depending on their similarities or differences in serotype, antibiogram susceptibility and plasmid profile. Group I consisted of 11 icp cases (44%) that showed similar characteristic for all markers used. There was only one exception in this group. Strains 30VI01 and 30VI03 belonging to *S. flexneri* serotype 1a were different in one out of 13 antibiotics tested (chloramphenicol, see Table 1). Group II contained 8 icp cases (32%), which were distinguished by different serotypes, plasmid profiles and antibiotic susceptibilities. One case included in this group showed the same antibiotic profile for the 13 antibiotics tested (3NC206, *S. flexneri* 6 and 3NC207, *S. flexneri* 2a, Table 1). Group III consisted of 6 icp cases (24%) with the same serotype but with different plasmid profile and antibiotic susceptibility (see Table 1). This indicated that (i) there were different shigella clones circulating at the same time in Canto

Table 1. *Intra-family comparison by pairs (icp) of the S. flexneri strains from households with multiple infections*

icp	Isolate	Sero-type	<i>ipaH</i> pattern		API 20E	Antibiotic resistance*	Plasmid profile (MDa)
			<i>SalI</i>	<i>HindIII</i>			
Group I							
1	1NC101	5	H	h	0044100	E, S, Ts	140, 4, 2-6, 2§
	1NC103	5	H	h	0044100	E, S, Ts	140, 4, 2-6, 2
2	10VI00	6	J	j	0004102	E, D, T, A, S, Ts	140, 4-4, 4, 2-6, 2
	10VI05	6	J	j	0004102	E, D, T, A, S, Ts	140, 4-4, 4, 2-6, 2
3	10NC303	2a	C	c	0004100	D, T, C, S, Ts	140, 2-6, 2, 1-4
	10NC306	2a	D	d	0004100	D, T, C, S, Ts	140, 2-6, 2, 1-4
4	15VI00	3a	G	g	0004140	†	140, 3-4, 2-6, 2
	15VI04	3a	G	g	0004140		140, 3-4, 2-6, 2
5	16NC207	6	J	j	0004102	E, D, T, C, S, Ts	140, 4-4, 4, 1-8
	16NC208	6	J	j	0004102	E, D, T, C, S, Ts	140, 4-4, 4, 1-8
6	20NC307	1a	A	a	0004100	E, T, C, A, S, Ts	140, 2-6, 2
	20NC309	1a	A	a	0004100	E, T, C, A, S, Ts	140, 2-6, 2
7	23CI307	3a	G	g	0044100	E, A, Tsx, S, Ts	140, 34, 3-4, 2-6, 2, 1-4
	23CI308	3a	G	g	0044100	E, A, Tsx, S, Ts	140, 34, 3-4, 2-6, 2, 1-4
8	24NC304	2a	C	c	0004100	E, D, T, C, S, Ts	140, 2-6, 2
	24NC312	2a	C	c	0004100	E, D, T, C, S, Ts	140, 2-6, 2
9	26NC303	1a	A	a	0004100	T, C, A, S, Ts	140, 2-6, 2
	26NC305	1a	A	a	0004100	T, C, A, S, Ts	140, 2-6, 2
10	30VI01	1a	A	a	0004100	T, A, S, Ts	140, 2-6, 2
	30VI03	1a	A	a	0004100	T, C, A, S, Ts‡	140, 2-6, 2
11	31CI103	2a	C	c	0004102	D, T, C, A, S	140, 2-6, 2
	31CI105	2a	C	c	0004102	D, T, C, A, S	140, 2-6, 2
Group II							
12	3NC206	6	J	j	0004142	D, T, C, S, Ts	140, 4, 2-6, 2
	3NC207	2a	C	c	0004142	D, T, C, S, Ts	140, 2-6, 2
13	3NC302	2a	C	c	0004102	D, T, C, S, Ts	140, 2-6, 2
	3NC305	6	J	j	0004102	E, D, C, K, S, N	140, 4, 2-6, 1-8
14	5CI205	3a	G	g	0004100	E, D, T, C, A, K, Tsx, S, N, Ts	140, 3-4, 2-6, 2
	5CI206	2a	C	c	0004100	E, D, T, C, A, K, S, N, Ts	140, 4-8, 3-4, 2-6, 2
15	6NC104	6	J	l	0004100	A	140, 4, 2-6, 1-8, 1-4
	6NC108	3a	G	g	0004100	E, D, T, C, A, S, Ts	140, 3-4, 2-6, 2
16	20NC305	3a	G	g	0004140	E, D, T, C, A, S, Ts	140, 3-4, 2-6, 2
	20NC307	1a	A	a	0004100	E, T, C, A, S, Ts	140, 2-6, 2
17	20NC305	3a	G	g	0004140	E, D, T, C, A, S, Ts	140, 3-4, 2-6, 2
	20NC309	1a	A	a	0004100	E, T, C, A, S, Ts	140, 2-6, 2
18	25VI00	6	K	k	0004002	E, T, C, S, Ts	140, 4, 2-6, 1-8
	25VI08	1a	A	a	0004100	D, T, C, S, Ts	140, 2-6, 2
19	34VI00	1a	A	a	0004100	E, D, T, C, A, K, Tsx, S, N, Ts	140, 4-8, 3-4, 2-6, 2, 1-4
	34VI06	2a	E	e	0004100	D, T, C, S, Ts	140, 2-6, 2, 1-4
Group III							
20	10NC303	2a	C	c	0004100	D, T, C, S, Ts	140, 2-6, 2, 1-4
	10NC308	2a	D	d	0004100	D, T, C, Ts	140, 90, 2-6, 2
21	10NC306	2a	D	d	0004100	D, T, C, S, Ts	140, 2-6, 2, 1-4
	10NC308	2a	D	d	0004100	D, T, C, Ts	140, 90, 2-6, 2
22	13NC106	2a	F	c	0004100	D, T, C, S, Ts	140, 90, 2-6, 2
	13NC107	2a	F	c	0004100	E, D, T, C, Tsx, S	140, 2-6, 2
23	23CI307	3a	G	g	0044100	E, A, Tsx, S, Ts	140, 34, 3-4, 2-6, 2, 1-4
	23CI309	3a	G	g	0004140	S	140, 50, 3-4, 2-6, 2
24	23CI308	3a	G	g	0044100	E, A, Tsx, S, Ts	140, 34, 3-4, 2-6, 2, 1-4
	23CI309	3a	G	g	0004140	S	140, 50, 3-4, 2-6, 2
25	24CI204	1a	B	b	0044100	E, D, T, C, S, Ts	140, 90, 2-6, 2, 1-4
	24CI208	1a	B	b	0044100	E, D, T, C, A, S, Ts	140, 34, 2-6, 2

* Antibiotics: A, ampicillin; C, chloramphenicol; Cp, cephalotholin; D, doxycycline; E, erythromycin; G, gentamicin; K, kanamycin; NA, nalidixic acid; N, neomycin; S, streptomycin; T, tetracycline; Tsx, trimethoprim-sulphamethoxazole; Ts, triple sulfa.

† These two strains were sensitive to the 13 antibiotics tested.

‡ Due to the difference in only one antibiotic, these strains were included in Group I.

§ Sizes of the plasmids refer to the sizes of the supercoiled form of the plasmid.




Fig. 1. The invasion plasmids of the *S. flexneri* strains were isolated as described in the Materials and Methods section, and they were digested with *Hind*III (*a, b, d*) or *Sal*I (*c*). The samples were run on agarose gels and blotted to nitrocellulose. The *ipaH*_{7,8} oligoprobe was non-radiolabelled (*a*) or radiolabelled (*b–d*). (*a*) Lane 1, *S. flexneri* 5 strain M90T; lanes 2–9, *S. flexneri* 2a, strains 31CI105, 31CI103, 24NC312, 24NC304, 13NC107, 13NC106, 3NC302, 3NC207; lanes 10–13, *S. flexneri* 6 strains 3NC305, 3NC206, 10VI05, 10VI00. (*b*) Lanes 1–3, *S. flexneri* 3a strains 23CI307, 23CI308, 23CI309; lanes 4, 5, 7–9, *S. flexneri* 1a strains 24CI204, 24CI208, 25VI08, 30VI01, 34VI00; lane 6, *S. flexneri* 6 strain 25VI00; lane 10, *S. flexneri* 2a strain 34VI06. (*c*) Lane 1, *S. flexneri* 5 strain M90T; lanes 2–12, *S. flexneri* 2a strains 10NC303, 10NC306, 10NC308, 13NC106, 13NC107, 24NC304, 24NC312, 31CI105, 31CI103, 34VI06. (*d*) Lane 1, *S. flexneri* 5 strain M90T; lanes 2, 10, *S. flexneri* 6 strains 6NC104, 25VI00; lanes 3–5, 13, *S. flexneri* 2a strains 10NC303, 10NC306, 10NC308, 34VI06; lanes 6, 7, *S. flexneri* 3a strains 20NC305, 23CI309; lanes 8, 9, 11, 12, *S. flexneri* 1a strains 24CI204, 24CI208, 25VI08, 34VI00. Molecular-weight markers are indicated to the left of each panel and reflect the mobilities of λ *Hind*III and Φ X174 *Hae*III fragments.

Grande, and (ii) *S. flexneri* strains were identical in less than half (44%) of the households with multiple shigella infections.

ipaH hybridization profiles of *S. flexneri* strains from households with multiple shigella infections

The 25 icp cases distributed in three groups described above were compared utilizing their *ipaH* hybridization patterns. The laboratory strain M90T (*S. flexneri* 5) was included as a control for the hybridizations with the *ipaH* oligoprobe. The hybridization patterns

were similar using either non-radioactive (Fig. 1*a*) or radioactive (Fig. 1*b–d*) *ipaH* probes and these patterns are schematized in Figure 2(*a, b*) (*Sal*I and *Hind*III restriction enzyme-digested DNA respectively). The *ipaH* patterns of the three *S. flexneri* groups described previously were analysed. We arbitrarily used capital letters for *Sal*I patterns and lower-case letters for *Hind*III patterns.

Group I: In 10 out of 11 icp cases (91%) the *ipaH* patterns were identical (Table 1). Only 1 icp case (icp no. 3, 10NC303 and 10NC306 corresponding to serotype 2a) showed a difference in 1 out of 5 bands




Fig. 2. Schematic representation of the *ipaH* hybridization profiles in invasion plasmid DNA isolated from *S. flexneri* strains from households with multiple shigella infections. Invasion plasmid DNA was isolated as described in the Materials and Methods section and digested with *SalI* (a) or *HindIII* (b) prior to electrophoretic separation on 0.8% agarose gels and blotting to nitrocellulose. Samples were screened with the *ipaH* oligoprobe as described. Molecular-weight markers are indicated to the left of each panel and reflect the mobilities of λ *HindIII* and Φ X174 *HaeIII* fragments.

Table 2. Comparison of ipaH patterns, antibiotic-resistance and plasmid profiles of *S. flexneri* serotypes

Serotype	ipaH pattern		No. of cases	API 20E	Antibiotic resistance*													Plasmid profile (MDa)																
	SalI	HindIII			A	Cp	C	D	E	G	K	NA	N	S	T	Tsx	Ts	140	90	50	34	4.8	4.4	4.0	3.4	2.6	2.0	1.8	1.4					
1	A	a	8	0004100	7	7	0	2	3	0	1	0	0	1	8	8	1	8	8	8	1	8	8	8	8	8	8	8	8	1				
	B	b	2	0044100	1	2	0	2	0	0	0	0	0	0	2	2	0	2	2	2	1	2	2	2	2	2	2	2	1	1				
	C	c	4	0004100	1	4	0	4	3	0	1	0	0	1	4	4	0	4	4	4	—	—	—	—	—	—	—	—	—	—	2			
	C	c	3	0004102	2	3	0	3	0	0	0	0	0	0	3	3	0	1	3	3	—	—	—	—	—	—	—	—	—	—	—			
2	C	c	1	0004142	0	1	0	1	0	0	0	0	0	0	1	1	0	1	1	1	—	—	—	—	—	—	—	—	—	—	—			
	D	d	2	0004102	0	2	0	2	0	0	0	0	0	0	1	2	0	2	2	2	1	—	—	—	—	—	—	—	—	—	—			
	E	e	1	0004102	0	1	0	1	0	0	0	0	0	0	1	1	0	1	1	1	—	—	—	—	—	—	—	—	—	—	—	—		
	F	f	2	0004100	0	2	0	2	1	0	0	0	0	0	2	2	1	1	2	2	1	—	—	—	—	—	—	—	—	—	—	—		
3	G	g	4	0004140	1	1	0	1	1	0	0	0	0	0	2	1	0	1	4	4	—	—	—	—	—	—	—	—	—	—	—	—		
	G	g	2	0004100	2	2	0	2	2	0	1	0	1	2	2	1	2	2	2	2	—	—	—	—	—	—	—	—	—	—	—	—	—	
5	G	g	2	0044100	2	0	0	0	2	0	0	0	0	0	2	0	2	2	2	2	—	—	—	—	—	—	—	—	—	—	—	—	—	
	H	h	2	0044100	0	0	0	0	2	0	0	0	0	0	2	0	0	2	2	2	—	—	—	—	—	—	—	—	—	—	—	—	—	
6	J	j	6	0004102	2	4	0	6	5	0	1	0	1	6	5	0	5	6	6	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
	K	k	1	0004102	0	1	0	0	1	0	0	0	0	0	1	1	0	1	1	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—
1	J	j	1	0004100	1	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—

* Antibiotics: A, ampicillin; C, chloramphenicol; Cp, cephalotholin; D, doxycycline; E, erythromycin; G, gentamicin; K, kanamycin; NA, nalidixic acid; N, neomycin; S, streptomycin; T, tetracycline; Tsx, trimethoprim-sulphamethoxazole; Ts, triple sulfa.

when the DNA was digested either with *SalI* (Fig. 1c, lanes 2 and 3, these patterns are also illustrated in Fig. 2a, serotype 2 patterns C and D) or *HindIII* (Fig. 1d, lanes 3 and 4, also illustrated in Fig. 2b, patterns c and d).

Group II: These cases were variable in their serotypes, antibiotic susceptibility and plasmid profiles. These strains also showed differences in their *ipaH* hybridization patterns (Table 1, Fig. 2a, b).

Group III: There were 6 icp cases in this group that had the same serotype but with differences in antibiotic susceptibility and plasmid profiles. In 5 out of 6 icp cases (83%) the *ipaH* pattern was the same when either *HindIII* or *SalI* was used to digest the invasion plasmid (Table 1). Only 1 of the 6 icp cases showed a different *ipaH* hybridization profile. The *ipaH* profile of this icp case (10NC303 and 10NC308, *S. flexneri* 2a) has been described above for Group I.

A strong correlation was obtained particularly between the serotype of *S. flexneri* and *ipaH* patterns. These patterns are illustrated schematically in Figure 2(a, b). The distribution of the *ipaH* patterns, antibiotic susceptibility patterns and plasmid profiles of the 41 *S. flexneri* strains is further described in Table 2. With strains of serotype 1 (Fig. 2a, b), pattern A (*SalI*) and pattern a (*HindIII*) were observed in 8 out of 10 strains, while pattern B (*SalI*) and pattern b (*HindIII*) were observed in 2 strains from the same house (24CI204 and 24CI208) belonging to Group III (Table 1). Strains of serotype 2a which were more numerous (13 strains) exhibited greater variation in their *ipaH* hybridization patterns. Using *SalI*, 4 patterns (C–F) were observed (Fig. 2a) while with *HindIII*-digested DNA, three patterns were obtained (Fig. 2b). The majority of strains of serotype 2a (8/13) presented patterns C and c. Two strains of this serotype which gave patterns D and d, were isolated from the same house and exhibited differences in plasmid profiles and antibiotic susceptibility (Tables 1 and 2). Only one strain of serotype 2a in this study presented the patterns E and e (Fig. 2a, b). Pattern F (Fig. 2a) and pattern c (Fig. 2b) were observed in two strains of serotype 2a and these two strains showed differences in antibiotic susceptibility and plasmid profiles (Tables 1 and 2). In the case of serotype 3a, the eight strains presented the same pattern (G and g; Fig. 2a, b). Interestingly, three strains of this serotype (23CI307, 23CI308 and 23CI309) were isolated from the same house, of which two (23CI307 and 23CI308) were identical but one (23CI309) showed differences

in plasmid profile, antibiotic-susceptibility pattern and biochemical reactions. Only two strains of serotype 5 were obtained in this study with identical markers. When the *ipaH* patterns of these *S. flexneri* 5 strains were compared to the pattern obtained with a laboratory strain control (M90T-W), differences in only one band were obtained with either *SalI* or *HindIII* digestions (Fig. 2*a, b*). In the case of serotype 6, eight strains were studied. Two patterns were obtained with *SalI* (J and K) and three patterns with *HindIII* (j, k, l) digestions (Fig. 2*a, b*). The pattern J (*SalI*) was observed in seven strains of serotype 6 (Table 2); however, in one of these strains the pattern 'l' instead of the pattern 'j' was obtained when *HindIII* was used for DNA digestion. Only one strain of serotype 6 showed pattern K (*SalI*; Fig. 2*a*) and pattern k (*HindIII*; Fig. 2*b*).

Phylogenetic analysis

Based on *ipaH* patterns of hybridization and serotype a dendrogram representing cluster analysis was carried out as described in Figure 3. The *S. flexneri* strains used in this study clustered in four groups. The first cluster includes the patterns Aa, Gg, Kk, Jj and Jl, of serotypes 1a, 3a and 6. The second cluster includes only two isolates of *S. flexneri* serotype 1a, 24CI204 and 24CI208, which presented the pattern Bb. These two isolates of *S. flexneri* 1a were biochemically different compared to the other eight strains of this serotype. The third cluster corresponds to the patterns Cc, Dd, Ee and Fc of the 13 *S. flexneri* 2a isolates. Finally, the fourth cluster includes the two isolates of *S. flexneri* serotype 5 with pattern Hh and the strain control M90T-W.

For the Group I strains described in Table 1 in this study, 10 out of 11 icp cases showed a Jaccard's distance of 0 when the *ipaH* hybridization patterns were compared using numerical analysis. Only 1 icp case [strains 10NC303 and 10NC306 (*S. flexneri* 2a)] presented small differences in their respective *ipaH* patterns and had a Jaccard's distance of only 0.1 (Fig. 3, Table 1). All of the icp cases in Group II, with strains that presented different serotypes and *ipaH* hybridization profiles, had Jaccard's distances that varied between 0.31 and 0.5 (Fig. 3, Table 1). In Group III, 5 out of 6 icp cases presented a Jaccard's distance of 0. Curiously, the strains (10NC303 and 10NC308, *S. flexneri* 2a) of the only icp case with a different *ipaH* pattern in this group had a Jaccard's distance of 0.1 and were isolated from the same house of the

other icp case in Group I that showed differences in the *ipaH* profiles (Fig. 3, Table 1).

DISCUSSION

In this study we found that a diversity of shigella clones were circulating in the population of Canto Grande at the time of infection. Up to six different serotypes of *Shigella* were isolated from the household contacts of one index case. This complicated picture of the epidemiology of shigella has also been observed in other endemic areas of shigellosis, such as Bangladesh and Somalia [3]. A total of 41 *S. flexneri* strains distributed in 25 icp cases were divided in three groups when their respective serotypes, plasmid and anti-biogram profiles and *ipaH* hybridization profiles were compared.

Several methods have been used to study the epidemiology of shigella infections. Serology constitutes one of the most important analyses for species identification of shigella isolates and is based on the specificity of the O-antigen [31]. However, this technique shows limitations since it is time consuming, does not always provide reliable data, and needs extensive confirmation especially with non-typable strains [32]. Additionally, serological analysis cannot be considered alone since strains of the same serotype can be obtained from different geographical regions, which have been exposed to pressure-selection changes in both their phenotypic and genotypic properties [33]. Plasmid fingerprinting [33–40], antibiotic profiles [33, 34, 38, 40–42], analysis of the rRNA gene restriction patterns [43–47], phage typing [48, 49], genomic DNA analysis by restriction fragment length polymorphism (RFLPs) [20, 50], pulsed-field gel electrophoresis (PFGE) of DNA samples [32, 35, 45, 51] and distribution of insertion sequences [32] have provided alternative techniques for shigella epidemiology. Individually, each assay has limitations.

In this study we exploited the *ipaH* heterogeneity as a tool for distinguishing shigella strains in an epidemiological study of transmission in an endemic area of shigellosis. We found a strong correlation between *ipaH* patterns and serotypes of *S. flexneri*. Our results indicate that *ipaH* hybridization patterns can differentiate between *S. flexneri* strains and can complement serotyping. *ipaH* genes are detected in multiple copies on the chromosome and invasion plasmids of virulent and avirulent *Shigella* spp. and EIEC strains [10]. The RFLP *ipaH* patterns not only correlated with the species and the geographic origin of the




Fig. 3. Dendrogram representing cluster analysis of the *ipaH* hybridization profiles and serotypes of the *S. flexneri* strains from households with multiple shigella infections.

strains, but were also able to discriminate between shigella isolates that had previously been shown to have similar RFLP patterns using Shiga toxin or sRNA probes [10]. In our study, we were able to discriminate between isolates of the same serotype. For example, we found that three isolates of *S. flexneri* 2a (strains 10NC303, 10NC306 and 10NC308) from the same household presented differences in their respective *ipaH* hybridization profiles. The three icp cases generated with these isolates were either identical or had some differences in their plasmid and antibiotic profiles. This indicates that in endemic areas such as Canto Grande, the shigella strains are undergoing rapid changes. Mutations or rearrangements in a region of the invasion plasmid will generate a different *ipaH* profile, even though the antibiotic characteristics and the presence of other plasmids are conserved (Group I, case no. 3). Alternatively, the *ipaH* profile can be conserved but changes in plasmids and/or antibiotic susceptibility may occur (Group III, case no. 21). We do not know which event occurred first or whether both events leading to differences among isolates occurred simultaneously. These changes result in the generation of an isolate with distinct phenotypic and genetic characteristics (Group III, case no. 20). A limitation of this study was to select and save only one strain per individual. Later, in another study of shigellosis conducted in Lima and Canto Grande, three shigella colonies per plate of selective medium were stored. We found that an individual could be infected with two different species of shigella, *S. flexneri* and *S. sonnei* (Fernandez-Prada CM, unpublished observations), and in some cases where there was only one species isolated, up to three clonal variants were detected when their *ipaH* RFLP profiles and serology were compared (Hartman AB, unpublished observations). These findings indicate that several clones are circulating in endemic areas of shigellosis and that the use of *ipaH* RFLP analysis of shigella strains is a powerful tool in epidemiological studies.

The role of *ipaH* in pathogenesis is unclear. Using suicide vector-driven mutagenesis, deletions in individual copies of *ipaH*_{9,8}, *ipaH*_{7,8}, and *ipaH*_{4,5} genes have been made, and double and triple *ipaH* mutants have been constructed [52]. These *ipaH* mutants remain invasive in HeLa cells and behave like wild-type strains in plaque assay indicating that these genes may not be involved in the entry or dissemination process [52]. However, these mutations induce an exaggerated Sereny response in guinea-pig eyes suggesting that

ipaH may play a role in modulating the inflammatory response elicited by infection [52]. Infection of human monocyte-derived macrophages *in vitro* with virulent *S. flexneri* results in cell death with release of IL-1 β [53, 54]. Recently, we have found that the rapid exit of virulent shigella from the phagocytic vacuoles of monocyte/macrophage cells may be facilitated by *ipaH*_{7,8} [52]. In the absence of IpaH_{7,8}, reduced cytotoxicity and lower levels of IL-1 β secretion were observed [52].

Comparison of the plasmid profiles of the 41 *S. flexneri* strains studied here, indicated that the 2.6 and 2.0 MDa plasmids were present in 95 and 88% of the isolates respectively. As shown in Table 2, these plasmids were absent only in some strains of serotype 6. The presence of these small plasmids in *S. flexneri* strains has been previously reported and indicates that a correlation between plasmid profiles and serotypes for *S. flexneri* exists [36, 37, 39]. Additionally, we found that only the serotypes 5 and 6 of *S. flexneri* had the 4 MDa plasmid, while the 3.4 MDa plasmid was present in all 8 *S. flexneri* 3a, in 1 out of 13 *S. flexneri* 2a and in 1 out of 10 *S. flexneri* 1a strains. In *S. flexneri* strains of serotype 6, when the 2.0 MDa plasmid was not present a small plasmid of 1.8 MDa was observed instead. The *S. flexneri* strains analysed in this study were multi-drug resistant and contained several plasmids; however, we did not find a correlation between presence of a particular plasmid and antibiotic-resistance profile. Multi-drug resistance patterns were observed in isolates containing both large-sized plasmids and those containing only small-sized plasmids in addition to the 140 MDa invasion plasmid. Several reports have indicated the presence of large transmissible plasmids that harbour multiple resistance genes among isolates of shigella [33–36, 38, 40, 42]. Recently, Rajakumar et al. [55, 56] have identified a chromosomal region in *S. flexneri* strains that confers multi-antibiotic resistance. A 99-kb deletion within the chromosome of *S. flexneri* 2a (strain YSH6000) indicated that this region was responsible for susceptibility to ampicillin, streptomycin, tetracycline and chloramphenicol [56]. This 99-kb chromosomal multi-antibiotic resistance locus was highly similar in sequence and organization to a region of the *Shigella* R-plasmid, NR1, and it may have arose following integration of an NR1-like plasmid [55]. The possibility of integration of R-plasmids in the chromosome of shigella strains in Canto Grande is an open question and could be the explanation for the differences in antibiotic and plasmid profiles

observed between isolates of icp cases with the same serotype and similar *ipaH* hybridization profiles (Group III). Curiously, the strains 15VI00 and 15VI04 (see Table 1, icp no. 4, Group I) were sensitive to the 13 antibiotics tested and this may be explained by a spontaneous deletion of the integrated multi-drug resistance genes in the chromosome of shigella or by the loss of the non-integrated R-plasmid.

Shigellosis remains endemic throughout the world, especially in many developing countries where poor sanitary conditions, malnutrition, overcrowded situations and unclean water supplies contribute to the high incidence and persistence of infection [2]. Shigellosis is a disease that can be a major killer of infants, children, elderly people, malnourished individuals and people living in marginal conditions [2] as in the case of Canto Grande in Lima, Peru. In order to understand the epidemiology of shigella, especially in endemic areas, it is necessary to have effective discriminatory tools to differentiate between isolates of shigella. These epidemiological tools have to be both easy to use and less time-consuming. We found that the *ipaH* hybridization profiles together with the data generated from serology, plasmid patterns and antibiotic characteristics represent a useful epidemiological tool not only to differentiate between strains of *S. flexneri* but also to understand the dynamic of the generation of new clones of pathogenic bacteria.

ACKNOWLEDGEMENTS

This study was supported by the Programme for the Control of Diarrhoeal Diseases of the World Health Organization, and by the Applied Diarrhoeal Disease Research Project of the United States Agency for International Development. The authors thank the field and laboratory workers and supervisors of the study for their dedicated assistance, especially Dora Maurtua, Rina Meza, Nancy Quevedo and Cesar Guerrero. Thanks are also due to the Laboratorio Nacional de Referencia de Enteropatógenos (LANARE) in Lima, Peru for the serotyping of the shigella strains, and to Dr David Taylor for reading the manuscript.

REFERENCES

- Black RE, Lanata CF. Epidemiology of diarrhoeal diseases in developing countries. In: Blaser MJ, Smith PD, Ravdin JI, Greenberg HB, Guerrant RL, eds. Infections of the gastrointestinal tract. New York: Raven Press, 1995: 13–36.
- Hale TL. Bacillary dysentery. In: Topley and Wilson's microbiology and microbial infection, Vol. 3: Bacterial infections. Oxford University Press: London, 1998: 479–493.
- Shears P. Shigella infections. Ann Trop Med Parasitol 1996; **90**: 105–114.
- Kotloff KL, Winickoff JP, Ivanoff B, et al. Global burden of shigella infections: implications for vaccine development and implementation of control strategies. Bull World Health Org 1999; **77**: 651–666.
- CDC. Summary of notifiable diseases – United States, 2001. MMWR 2003; **50**: 1–164.
- Dutta S, Rajendran K, Roy S, et al. Shifting serotypes, plasmid profile analysis and antimicrobial resistance pattern of shigellae strains isolated from Kolkata, India during 1995–2000. Epidemiol Infect 2002; **129**: 235–243.
- Dupont HL. Inoculum size in shigellosis and implications for expected mode of transmission. J Infect Dis 1989; **159**: 1126–1128.
- Wei J, Goldberg MB, Burland V, et al. Complete genome sequence and comparative genomics of *Shigella flexneri* serotype 2a strain 2457T. Infect Immun 2003; **71**: 2775–2786.
- Oaks VE, Hale TL, Formal SB. The immune response to *Shigella* protein antigens in rhesus monkeys and humans infected with *Shigella* spp. Infect Immun 1986; **53**: 57–63.
- Buysee JM, Hartman AB, Strockbine N, Venkatesan MM. Genetic polymorphism of the *ipaH* multicopy antigen gene in *Shigella* spp. and enteroinvasive *Escherichia coli*. Microb Pathog 1995; **19**: 335–349.
- Hartman AB, Venkatesan MM, Oaks EV, Buysee JM. Sequence and molecular characterization of a multicopy invasion plasmid antigen gene, *ipaH*, of *Shigella flexneri*. J Bacteriol 1990; **172**: 1905–1915.
- Venkatesan MM, Buysee JM, Hartman AB. Sequence variation in two *ipaH* genes of *Shigella flexneri* 5 and homology to the LRG-like family of proteins. Mol Microbiol 1991; **5**: 2435–2445.
- Venkatesan MM, Buysee JM, Kopecko DJ. Use of *Shigella flexneri ipaC* and *ipaH* gene sequences for the general identification of *Shigella* spp. and enteroinvasive *Escherichia coli*. J Clin Microbiol 1989; **27**: 2687–2691.
- Buysee JM, Stover CK, Oaks EV, Venkatesan MM, Kopecko DJ. Molecular cloning of invasion plasmid antigen (*ipa*) genes from *Shigella flexneri*: analysis of *ipa* gene products and genetic mapping. J Bacteriol 1987; **169**: 2561–2569.
- Dutta S, Chatterjee A, Dutta P, et al. Sensitivity and performance characteristics of a direct PCR with stool samples in comparison to conventional techniques for diagnosis of *Shigella* and enteroinvasive *Escherichia coli* infection in children with acute diarrhoea in Calcutta, India. J Med Microbiol 2001; **50**: 667–674.
- Oberhelman RA, Kopecko DJ, Venkatesan MM, et al. Evaluation of alkaline phosphatase-labeled *ipaH* probe for diagnosis of shigella infections. J Clin Microbiol 1993; **31**: 2101–2104.

17. Echevarria P, Sethabutr O, Serichantalergs O, Lexomboon U, Tamura K. *Shigella* and enteroinvasive *Escherichia coli* infections in households of children with dysentery in Bangkok. *J Infect Dis* 1992; **165**: 144–147.
18. Sethabutr O, Echevarria P, Hoge CW, Bodhidatta L, Pitarangsi C. Detection of *Shigella* and enteroinvasive *Escherichia coli* by PCR in the stools of patients with dysentery in Thailand. *J Diarrhoeal Dis Res* 1994; **12**: 265–269.
19. Gaudio PA, Sethabutr O, Echeverria P, Hoge CW. Utility of a polymerase chain reaction diagnostic system in a study of the epidemiology of shigellosis among dysentery patients, family contacts, and well controls living in a shigellosis-endemic area. *J Infect Dis* 1997; **176**: 1013–1018.
20. Yavzori M, Cohen D, Bercovier H. Molecular epidemiology of *Shigella* infections in Israel. *Epidemiol Infect* 1992; **109**: 273–282.
21. Gil AI, Lanata CF, Butron B, Gabilondo A, Molina M, Bravo N. Incidence of *Vibrio cholerae* O1 diarrhea in children at the onset of a cholera epidemic in periurban Lima, Peru. *Pediatr Infect Dis J* 1996; **15**: 415–418.
22. Lopez de Romana G, Brown KH, Black RE. Health and growth of infants and young children in Huascar, Peru. *Ecol Food Nutr* 1987; **19**: 213–229.
23. Yeager BA, Lanata CF, Lazo F, Verastegui H, Black RE. Transmission factors and socioeconomic status as determinants of diarrhoeal incidence in Lima, Peru. *J Diarrhoeal Dis Res* 1991; **9**: 186–193.
24. Lanata CF, Black RE, Gilman RH, Lazo F, Del Aguila R. Epidemiologic, clinical and laboratory characteristics of acute vs. persistent diarrhea in periurban Lima, Peru. *J Pediatr Gastroenterol Nutr* 1991; **12**: 82–88.
25. Lanata CF, Black RE, Maurtua D, et al. Etiologic agents in acute vs. persistent diarrhea in children under three years of age in periurban Lima, Peru. *Acta Paediatr* 1992; **381** (Suppl): 32–38.
26. Bauer AM, Kirby WMM, Sherris JC, Turck M. Antibiotic susceptibility testing by standardized single disk method. *Am J Clin Pathol* 1966; **45**: 493–496.
27. Sereny B. Experimental shigella keratoconjunctivitis. *Acta Microbiol Acad Sci Hung* 1955; **2**: 293–296.
28. Maniatis T, Fritsch EF, Sambrook J. Molecular cloning. A laboratory manual. New York: Cold Spring Harbor Laboratory, 1982.
29. Jaccard P. Nouvelles recherches sur la distribution florale [New studies on floral distribution]. *Bull Soc Vaud Sci Nat* 1908; **44**: 223–270.
30. Gerres E, Poux M. Pratique de la classification automatique. L'exemple des *Leishmania* [Method of automatic classification. The example of *Leishmania*]. In: Rioux JA, ed. International Symposium on taxonomy and phylogeny of *Leishmania*. Montpellier, 1986: 27–40.
31. Lindberg AA, Karnell A, Weintraub A. The lipopolysaccharide of *Shigella* bacteria as a virulence factor. *Rev Infect Dis* 1991; **13** (Suppl 4): S279–S284.
32. Soldati L, Piffaretti JC. Molecular typing of *Shigella* strains using pulsed field gel electrophoresis and genome hybridization with insertion sequences. *Res Microbiol* 1991; **142**: 489–498.
33. Scerpella EG, Mathewson JJ, Dupont HL, Marani SK, Ericsson CD. *Shigella sonnei* strains isolated from US summer students in Guadalajara, Mexico, from 1986 to 1992. *J Clin Microbiol* 1994; **32**: 2549–2552.
34. Bratoeva MP, John JF, Barg NL. Molecular epidemiology of trimethoprim-resistant *Shigella boydii* serotype 2 strains from Bulgaria. *J Clin Microbiol* 1992; **30**: 1428–1431.
35. Brian MJ, Van R, Townsend I, Murray BE, Cleary TG, Pickering LK. Evaluation of the molecular epidemiology of an outbreak of multiply resistant *Shigella sonnei* in a day-care center by using pulsed-field gel electrophoresis and plasmid DNA analysis. *J Clin Microbiol* 1993; **31**: 2152–2156.
36. Gebre-Yohannes A, Drasar BS. Molecular epidemiology of plasmid patterns in *Shigella flexneri* types 1–6. *Epidemiol Infect* 1991; **107**: 321–334.
37. Haider K, Huq MI, Talukder KA, Ahmad QS. Electrophoretotyping of plasmid DNA of different serotypes of *Shigella flexneri* isolated in Bangladesh. *Epidemiol Infect* 1989; **102**: 421–428.
38. Lin SR, Chang SF. Drug resistance and plasmid profile of shigellae in Taiwan. *Epidemiol Infect* 1992; **108**: 87–97.
39. Litwin CM, Storm AL, Chipowsky S, Ryan KJ. Molecular epidemiology of *Shigella* infections: plasmids profiles, serotype correlation, and restriction endonuclease analysis. *J Clin Microbiol* 1991; **29**: 104–108.
40. Yagupsky P, Loeffelholz M, Bell K, Menegus MA. Use of multiple markers for investigation of an epidemic of *Shigella sonnei* infection in Monroe county, New York. *J Clin Microbiol* 1991; **29**: 2850–2855.
41. Finkelman Y, Yagupsky P, Fraser D, Dagan R. Epidemiology of *Shigella* infections in two ethnic groups in a geographic region in southern Israel. *Eur J Clin Microbiol Infect Dis* 1994; **13**: 367–373.
42. Lima AA, Sidrim JJ, Lima NL, Titlow W, Evans ME, Greenberg RN. Molecular epidemiology of multiple antibiotic-resistant *Shigella flexneri* in Fortaleza, Brazil. *J Clin Microbiol* 1997; **35**: 1061–1065.
43. Coimbra RS, Nicastro G, Grimont PAD, Grimont F. Computer identification of *Shigella* species by rRNA gene restriction patterns. *Res Microbiol* 2001; **152**: 47–55.
44. Hinojosa-Ahumada M, Swaminathan B, Hunter SB, et al. Restriction fragment length polymorphisms in rRNA operons for subtyping *Shigella sonnei*. *J Clin Microbiol* 1991; **29**: 2380–2384.
45. Liu PY, Lau YJ, Hu BS, et al. Analysis of clonal relationships among isolates of *Shigella sonnei* by different molecular typing methods. *J Clin Microbiol* 1995; **33**: 1779–1783.
46. Mendoza MC, Martin MC, Gonzalez-Hevia MA. Usefulness of ribotyping in a molecular epidemiology study of shigellosis. *Epidemiol Infect* 1996; **116**: 127–135.
47. Strockbine NA, Parsonnet J, Greene K, Kiehlauch JA, Wachsmuth IK. Molecular epidemiologic techniques

- in analysis of epidemic and endemic *Shigella dysenteriae* type 1 strains. *J Infect Dis* 1991; **163**: 406–409.
48. Bentley CA, Frost JA, Rowe B. Phage typing and drug resistance of *Shigella sonnei* isolated in England and Wales. *Epidemiol Infect* 1996; **116**: 295–302.
 49. Lerman Y, Yavzori M, Ambar R, Sechter I, Wiener M, Cohen D. Epidemic spread of *Shigella sonnei* shigellosis and evidence for development of immunity among children attending day-care centers in a communal settlement (Kibbutz). *J Clin Microbiol* 1994; **32**: 1092–1094.
 50. Preston MA, Borczyk AA. Genetic variability and molecular typing of *Shigella sonnei* strains isolated in Canada. *J Clin Microbiol* 1994; **32**: 1427–1430.
 51. Litwin CM, Leonard RB, Carroll KC, Drummond WK, Pavia AT. Characterization of endemic strains of *Shigella sonnei* by use of plasmid DNA analysis and pulsed-field gel electrophoresis to detect patterns of transmission. *J Infect Dis* 1997; **175**: 864–870.
 52. Fernandez-Prada CM, Hoover DL, Tall BD, Kopelowitz J, Hartman AB, Venkatesan MM. *Shigella flexneri* *IpaH*_{7,8} facilitates escape of virulent bacteria from the endocytic vacuoles of mouse and human macrophages. *Infect Immun* 2000; **68**: 3608–3619.
 53. Fernandez-Prada CM, Hoover DL, Tall B, Venkatesan MM. Human monocyte-derived macrophages infected with virulent *Shigella flexneri* in vitro undergo a rapid cytolytic event similar to oncosis but not apoptosis. *Infect Immun*. 1997; **65**: 1486–1496.
 54. Fernandez-Prada CM, Hoover DL, Tall B, Elliott S, Nataro J, Venkatesan MM. Hemolysin-positive enteroaggregative and cell-detaching *Escherichia coli* strains cause oncosis of human monocyte derived macrophages and apoptosis of murine J774 cells. *Infect Immun* 1998; **66**: 3918–3924.
 55. Rajakumar K, Bulach D, Davies J, Ambrose L, Sasakawa C, Adler B. Identification of a chromosomal *Shigella flexneri* multi-antibiotic resistance locus which shares sequence and organizational similarity with the resistance region of the plasmid NR1. *Plasmid* 1997; **37**: 159–168.
 56. Rajakumar K, Sasakawa C, Adler B. A spontaneous 99-kb chromosomal deletion results in multi-antibiotic susceptibility and an attenuation of contact haemolysis in *Shigella flexneri* 2a. *J Med Microbiol* 1996; **45**: 64–75.