

CONTRIBUTORS

- D. E. Beever *AFRC Institute for Grassland and Animal Production, Hurley, Maidenhead SL6 5LR*
- Nazeli Borlak *Department of Biochemistry and Physiology, University of Reading, Reading RG6 2AP*
- Tristan Corring *Station de Physiologie de la Nutrition, INRA, 78350 Jouy-en-Josas, France*
- G. G. Duthie *Rowett Research Institute, Bucksburn, Aberdeen AB2 9SB*
- J. France *AFRC Institute for Grassland and Animal Production, Hurley, Maidenhead SL6 5LR*
- Smita Ganatra *Milk Marketing Board, Thames Ditton, Surrey KT7 0EL*
- M. Gill *AFRC Institute for Grassland and Animal Production, Hurley, Maidenhead SL6 5LR*
- M. H. N. Golden *Tropical Metabolism Research Unit, University of the West Indies, Kingston, Jamaica*
- George K. Grimble *Department of Gastroenterology and Nutrition, Central Middlesex Hospital, Acton Lane, London NW10 7NS*
- M. I. Gurr *Milk Marketing Board, Thames Ditton, Surrey KT7 0EL*
- Barbara F. Harland *Department of Human Nutrition and Food, Howard University, Washington DC 20059, USA*
- Johein Harmeyer *Institute of Physiology, School of Veterinary Medicine, D-3000 Hannover 1, West Germany*
- Peter H. Holmes *Glasgow University Veterinary School, Bearsden Road, Glasgow G61 1QH*
- G. Hussey *Clinical Nutrition Unit, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT*
- A. A. Jackson *Department of Human Nutrition, University of Southampton SO9 3TU*
- W. P. T. James *Rowett Research Institute, Bucksburn, Aberdeen AB2 9SB*
- Catherine Juste *Station de Physiologie de la Nutrition, INRA, 78350 Jouy-en-Josas, France*
- Uwe Kollenkirchen *Institute of Physiology, School of Veterinary Medicine, D-3000 Hannover 1, West Germany*
- H. A. Lee *Department of Renal Medicine, St Mary's Hospital, Milton Road, Portsmouth PO3 6AD*
- Evelyne F. Lhoste *Station de Physiologie de la Nutrition, INRA, 78350 Jouy-en-Josas, France*
- A. R. Michell *Department of Veterinary Medicine and Animal Husbandry, Royal Veterinary College, Hawkshead Lane, North Mymms, Herts AL9 7TA*

viii *Contributors*

D. J. Millward *Nutrition Research Unit, London School of Hygiene and Tropical Medicine, 4 St Pancras Way, London NW1 2PE*

James J. Parkins *Glasgow University Veterinary School, Bearsden Road, Glasgow G61 1QH*

Gillian Price *Nutrition Research Unit, London School of Hygiene and Tropical Medicine, 4 St Pancras Way, London NW1 2PE*

D. D. Ramdath *Tropical Metabolism Research Unit, University of the West Indies, Kingston, Jamaica*

J. P. W. Rivers *Nutrition Research Unit, London School of Hygiene and Tropical Medicine, 4 St Pancras Way, London NW1 2PE*

David B. A. Silk *Department of Gastroenterology and Nutrition, Central Middlesex Hospital, Acton Lane, London NW10 7NS*

S. T. Talbot *Department of Renal Medicine, St Mary's Hospital, Milton Road, Portsmouth PO3 6AD*

Andrew Tomkins *Clinical Nutrition Unit, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT*

K. W. J. Wahle *Rowett Research Institute, Bucksburn, Aberdeen AB2 9SB*