

Rank-one systems, flexible classes and Shannon orbit equivalence

CORENTIN CORREIA

Université Paris Cité, Paris 75006, France
(e-mail: corentin.correia@imj-prg.fr)

(Received 12 July 2024 and accepted in revised form 21 October 2024)

Abstract. We build a Shannon orbit equivalence between the universal odometer and a variety of rank-one systems. This is done in a unified manner using what we call flexible classes of rank-one transformations. Our main result is that every flexible class contains an element which is Shannon orbit equivalent to the universal odometer. Since a typical example of flexible class is $\{T\}$ when T is an odometer, our work generalizes a recent result by Kerr and Li, stating that every odometer is Shannon orbit equivalent to the universal odometer. When the flexible class is a singleton, the rank-one transformation given by the main result is explicit. This applies to odometers and Chacon's map. We also prove that strongly mixing systems, systems with a given eigenvalue, or irrational rotations whose angle belongs to any fixed non-empty open subset of the real line form flexible classes. In particular, strong mixing, rationality or irrationality of the eigenvalues are not preserved under Shannon orbit equivalence.

Key words: orbit equivalence, rank-one systems, odometers, point spectrum, mixing properties

2020 Mathematics Subject Classification: 37A35 (Primary); 37A20 (Secondary)

Contents

1	Introduction	2
1.1	The universal odometer and a theorem of Kerr and Li [KL24]	3
1.2	Rank-one systems	4
1.3	A first extension of Kerr and Li's theorem	5
1.4	A modified strategy	6
1.5	Flexible classes	6
1.6	Further comments	8
1.7	Outline of the paper	9
2	Preliminaries	9
2.1	Basics of ergodic theory	9
2.2	Quantitative orbit equivalence	10



3	Rank-one systems	11
3.1	The cutting-and-stacking method	11
3.2	Flexible classes	16
4	Proof of Proposition 3.8	19
4.1	BSP systems	19
4.2	Irrational rotations	19
4.3	Systems with a given eigenvalue	21
4.4	Strongly mixing systems	22
5	From flexible classes to the universal odometer	22
5.1	The construction	23
5.2	First properties of this construction	29
5.3	Towards flexible classes	34
5.4	Equality of the orbits, universal odometer and quantitative control of the cocycles	35
5.5	Proof of Theorem 3.9	42
	Acknowledgements	50
	References	50

1. Introduction

At the level of ergodic probability measure-preserving bijections, *quantitative orbit equivalence* aims at bridging the gap between the well studied but very complicated relation of *conjugacy* and the trivial relation of *orbit equivalence*, which is equality of orbits up to conjugacy.

To be more precise, given two ergodic probability measure-preserving bijections S and T on a standard atomless probability space (X, \mathcal{A}, μ) , if S and some system $\Psi^{-1}T\Psi$ conjugate to T have the same orbits, then S and T are said to be *orbit equivalent* and the probability measure-preserving bijection $\Psi: X \rightarrow X$ is called an *orbit equivalence* between T and S . Dye's theorem [Dye59] states that if S and T are ergodic, then they are orbit equivalent.

To get an interesting theory, let us define the *cocycles* associated to Ψ , these are the integer-valued functions c_S and c_T defined by $Sx = \Psi^{-1}T^{c_S(x)}\Psi(x)$ and $Tx = \Psi S^{c_T(x)}\Psi^{-1}(x)$. *Shannon orbit equivalence* requires that there exists an orbit equivalence whose cocycles are Shannon, meaning that the partitions associated to c_S and c_T are both of finite entropy. For φ -integrable orbit equivalence, we ask that both integrals $\int_X \varphi(|c_S(x)|) d\mu(x)$ and $\int_X \varphi(|c_T(x)|) d\mu(x)$ are finite. In the particular case of a linear map φ , φ -integrable orbit equivalence exactly requires the integrability of the cocycles, and is simply called integrable orbit equivalence.

Belinskaya's theorem [Bel69] implies that integrable orbit equivalence is exactly flip-conjugacy (S and T are flip-conjugate if S is conjugate to T or T^{-1}). In fact, it only requires that one of the two cocycles is integrable. Carderi *et al* [CJLMT23] proved that this result is optimal, meaning that φ -integrable orbit equivalence never implies flip-conjugacy for a sublinear map φ . Moreover, φ -integrable orbit equivalence implies Shannon orbit equivalence when φ is asymptotically greater than \log . An impressive result of Kerr and Li [KL24] guarantees that these relations are not trivial: entropy is preserved




FIGURE 1. Schematic view of the interplay between the relations on ergodic bijections we have seen so far.

under Shannon orbit equivalence (and this is the only invariant of which we know). As a consequence, two transformations with different entropies can neither be Shannon orbit equivalent nor φ -integrably orbit equivalent for any φ greater than \log .

Historically, the question of preservation of entropy in quantitative orbit equivalence was asked in the more general setting of group actions. We do not give any definition in this setting, as the paper is only about probability measure-preserving bijections S , which can be seen as \mathbb{Z} -actions via $(n, x) \in \mathbb{Z} \times X \mapsto S^n x$. Austin [Aus16] showed that integrable orbit equivalence between actions of infinite finitely generated amenable groups preserves entropy. Kerr and Li [KL21, KL24] then generalized this result, replacing integrable orbit equivalence by Shannon orbit equivalence, and going beyond the amenable case using sofic entropy (see Figure 1).

1.1. *The universal odometer and a theorem of Kerr and Li [KL24].* In [CJLMT23], the statement about φ -integrable orbit equivalence in the sublinear case is the following. This gives a result on Shannon orbit equivalence since this is implied by φ -integrable orbit equivalence for φ greater than \log .

THEOREM. (Carderi *et al* [CJLMT23]) *Let $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a sublinear function. Let S be an ergodic probability measure-preserving transformation and assume that S^n is ergodic for some $n \geq 2$. Then, there is another ergodic probability measure-preserving transformation T such that S and T are φ -integrably orbit equivalent but not flip-conjugate.*

COROLLARY. (Carderi *et al* [CJLMT23]) *Let S be an ergodic probability measure-preserving transformation and assume that S^n is ergodic for some $n \geq 2$. Then, there is another ergodic probability measure-preserving transformation T such that S and T are Shannon orbit equivalent but not flip-conjugate.*

The proof is constructive and the resulting transformation T is built so that T^n is not ergodic. It is natural to wonder whether this statement holds for systems T without ergodic non-trivial powers. A well-known example of such a system is the universal odometer.

Question 1.1. Which systems are Shannon orbit equivalent to the universal odometer?

A first answer is given by Kerr and Li.

THEOREM. (Kerr and Li [KL24]) *Every odometer is Shannon orbit equivalent to the universal odometer.*




FIGURE 2. In this example, (\mathcal{R}_n) denotes the nested sequence of Rokhlin towers defining an odometer. Dividing \mathcal{R}_1 in two sub-towers and stacking them, this gives the next tower \mathcal{R}_2 . From \mathcal{R}_2 , \mathcal{R}_3 is defined by dividing in three sub-towers and stacking them.

Odometers are exactly probability measure-preserving bijections admitting a nested sequence of partitions of the space, each of them being a Rokhlin tower, and increasing to the σ -algebra \mathcal{A} , see Figure 2 (we refer the reader to the end of §3.1 for concrete examples with adding machines). Kerr and Li use this combinatorial specificity of these bijections to build an orbit equivalence between them.

1.2. Rank-one systems. The aim of the paper is to extend Kerr and Li's result to *rank-one bijections*. These are more general transformations admitting a nested sequence of Rokhlin towers increasing to the σ -algebra \mathcal{A} , but the towers do not necessarily partition the space. This means that from a tower to the next one, we need to add some parts of the space which are not covered by the previous tower, called *spacers*, so that the measure of the subset covered by the n th tower tends to 1 as n goes to $+\infty$. As illustrated in Figure 3, to get the next tower, the current one is subdivided in sub-towers which are stacked with optional spacers between them. The number of sub-towers is called the cutting parameter and the number of consecutive spacers between these sub-towers are the spacing parameters (see Definition 3.2). For example, an odometer admits a cutting-and-stacking construction with spacing parameters equal to zero at each step.

Rank-one systems all have entropy zero. They include systems with a *discrete spectrum* [Jun76], also called *compact* systems. Such systems are not weakly mixing and are completely classified up to conjugacy by their point spectrum [HVN42]. Examples include odometers and irrational rotations.

The family of rank-one systems is much richer than its subclass of discrete spectrum systems. Indeed, the latter are not weakly mixing, whereas there exist strongly mixing systems of rank one and also rank-one systems which are weakly mixing but not strongly




FIGURE 3. In this example, there are four spacers and the cutting parameter is three.

mixing (Chacon’s map was the first example of such a system and opened the study of rank-one systems). Rank-one systems can have irrational eigenvalues (that is, of the form $\exp(2i\pi\theta)$ with irrational numbers θ), which is the case of irrational rotations, whereas odometers only have rational eigenvalues. The reader may refer to the complete survey of Ferenczi [Fer97] about rank-one systems and, more generally, systems of finite rank.

The combinatorial structure of a general rank-one system does not differ too much from the structure of an odometer but the systems can have completely different properties; thus this class may extend the result of Kerr and Li and provide interesting flexibility results about Shannon orbit equivalence.

1.3. *A first extension of Kerr and Li’s theorem.* The construction of an orbit equivalence between the universal odometer S and any rank-one system T is a natural generalization of Kerr and Li’s method for the universal odometer and any odometer (see Remark 5.18). The difficulty is to quantify the cocycles.

At the beginning of our work, we first proved that the Shannon orbit equivalence established by Kerr and Li in [KL24] is actually a φ -integrable orbit equivalence for any $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\varphi(t) = o(t^{1/3})$. We then generalized this to rank-one systems called BSP, for ‘bounded-spacing-parameter’, see Definition 3.5. This notion of BSP systems was already introduced by Gao and Ziegler in [GZ19], using the symbolic definition of rank-one systems (in this paper, we will only consider the cutting-and-stacking definition of rank-one systems, which is often more appropriate for constructions in a measure-theoretic setting).

THEOREM A. *Every BSP rank-one system is φ -integrably orbit equivalent to the universal odometer for any $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfying $\varphi(t) = o(t^{1/3})$.*

Therefore, φ -integrable orbit equivalence, for a φ as in the above theorem, and Shannon orbit equivalence do not preserve weak mixing since Chacon’s map is a BSP rank-one system.

Now the goal is to get a result for systems of rank one outside the class of BSP systems. For this purpose, we find a more general framework with the notion of *flexible classes*,

and a general statement (Theorem B) implying Theorem A and other flexibility results (Theorems C, E and F). Theorem D is a refinement of Theorem C.

1.4. *A modified strategy.* We first have to understand why the quantification of the cocycles is more difficult to determine for general rank-one systems than for odometers (or even for BSP systems in Theorem A). In [KL24], the quantification of the cocycles relies on a series whose terms vanish to zero as the cutting parameters get increasingly larger. The key is then to get quickly increasing cutting parameters for the series to converge. To do so, it suffices to skip steps in the cutting-and-stacking process, that is, from the n th Rokhlin tower, we can directly build the $(n + k)$ th Rokhlin for k so big that the new cutting parameter is large enough. In other words, we can recursively choose the cutting parameters so that they increase quickly enough.

When the rank-one system is not an odometer, we need an asymptotic control on the spacing parameters (recall that they are zero for an odometer) for the cocycles to be well quantified. When skipping steps in the cutting-and-stacking method, the spacing parameters may increase too quickly, preventing us from quantifying the cocycles. As we will see in Lemma 3.6, we do not have this problem with BSP rank-one systems.

When the rank-one system is not BSP, skipping steps in the cutting-and-stacking construction is not relevant as it may improperly change the spacing parameters. In §5.3 (see Lemma 5.9), we will notice that the construction of Kerr and Li enables us to build the universal odometer S while we are building the rank-one system T , focusing only on the combinatorics behind the systems, whereas for Kerr and Li, T and its cutting-and-stacking settings are fixed and S is built from these data. This new strategy will enable us to have a result for systems of rank one outside the class of BSP systems, with the notion of flexible class.

1.5. *Flexible classes.* A flexible class (see Definition 3.7) is basically a class of rank-one systems satisfying a common property (e.g. the set of strongly mixing rank-one systems), with the following two requirements. We first ask for a sufficient condition, given by a set \mathcal{F}_C , on the first n cutting and spacing parameters (for all integers $n \geq 0$) for the underlying rank-one system to be in this class. Second, given a sequence of n cutting and spacing parameters in \mathcal{F}_C (they will be the first n parameters of a cutting-and-stacking construction), we require that it can be completed in a sequence of $n + 1$ parameters in \mathcal{F}_C , with infinitely many choices for the $(n + 1)$ th cutting parameters, and with the appropriate asymptotic control on the $(n + 1)$ th spacing parameters.

The idea is to inductively choose the parameters so that the cutting parameters increase quickly enough, with the appropriate asymptotics on the spacing parameters, and the underlying rank-one system has the desired property, namely the system is in the flexible class that we consider.

The general statement on flexible classes is the following.

THEOREM B. (See Theorem 3.9) *Let $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a map satisfying $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$. If \mathcal{C} is a flexible class, then there exists T in \mathcal{C} which is φ -integrably orbit equivalent to the universal odometer.*

A very interesting phenomenon is when a rank-one system T is flexible, meaning that $\{T\}$ is a flexible class. This first means that given the parameters of a cutting-and-stacking construction of T , it is possible to change the $(n + 1)$ th parameters so that they have the desired asymptotic control, and to inductively do so for every n so that the underlying rank-one system is again T . We do not know if every rank-one system is flexible. Second, Theorem B is an existence result and when a flexible class is a singleton $\{T\}$, this statement provides a concrete example of a rank-one system which is φ -integrably orbit equivalent to the universal odometer.

The following proposition gives examples of flexible classes.

- PROPOSITION 1.2. (See Proposition 3.8) (1) *Every BSP rank-one system is flexible.*
 (2) *For every non-empty open subset \mathcal{V} of \mathbb{R} , the set $\{R_\theta \mid \theta \in \mathcal{V} \cap (\mathbb{R} \setminus \mathbb{Q})\}$ is a flexible class.*
 (3) *For every irrational number θ , the class of rank-one systems which have $e^{2i\pi\theta}$ as an eigenvalue is flexible.*
 (4) *The class of strongly mixing rank-one systems is flexible.*

Proving that a BSP system is flexible is not difficult and we rely on the fact that bounded spacing parameters already have the desired asymptotics even though we skip steps in the cutting-and-stacking process for the cutting parameters to increase quickly enough (see §4.1). We use a construction by Drillick *et al* [DEJ+23] to prove Proposition 1.2 for irrational rotations (see §4.2). We also consider a construction by Danilenko and Vieprik [DV23] for the rank-one systems with a given eigenvalue (see §4.3). Finally, Ornstein [Orn72] gives the first example of strongly mixing rank-one systems and the fact that these systems form a flexible class follows from his construction (see §4.4).

Combined with Proposition 1.2, Theorem B provides four flexibility results. The first one is Theorem A stated above, which is a generalization of Kerr and Li’s theorem. The second one is another result with almost explicit examples of systems which are φ -integrably orbit equivalent to the universal odometer.

THEOREM C. *Let $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a map satisfying $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$. The set of irrational numbers θ whose associated irrational rotation is φ -integrably orbit equivalent to the universal odometer is dense in \mathbb{R} .*

The point spectrum of R_θ is exactly the circle subgroup generated by $\exp(2i\pi\theta)$ and the eigenvalues of the universal odometer are rational, so Theorem C implies that there exist two Shannon orbit equivalent systems (more specifically, φ -integrably orbit equivalent with $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$), with non-trivial point spectra and such that 1 is the only common eigenvalue.

The way we prove Theorem B will enable us to get the following refinement, its proof is written at the end of the paper.

THEOREM D. *For every map $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfying $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$, and for every non-empty open subset \mathcal{V} of \mathbb{R} , the set of irrational numbers $\theta \in \mathcal{V}$ whose associated irrational rotation is φ -integrably orbit equivalent to the universal odometer is uncountable.*

Question 1.3. Let us consider the set of irrational numbers θ whose associated irrational rotation is φ -integrably orbit equivalent to the universal odometer. Is this set conull, with respect to the Lebesgue measure, equal to the set of irrational numbers?

Finally, we get the following corollaries, providing implicit examples.

THEOREM E. For every map $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfying $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$ and for every irrational number θ , there exists a rank-one system which has $e^{2i\pi\theta}$ as an eigenvalue and which is φ -integrably orbit equivalent to the universal odometer.

THEOREM F. For every map $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfying $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$, there exists a strongly mixing rank-one system which is φ -integrably orbit equivalent to the universal odometer.

As $\exp(2i\pi\theta)$ is an eigenvalue of the irrational rotation of angle θ , and as we do not know if Theorem C holds for every irrational number θ , Theorem E then completes this statement with a weaker result for the remaining θ .

Theorem F implies that φ -integrable orbit equivalence, with $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$, and Shannon orbit equivalence do not preserve strong mixing. This is also a consequence of the result from [CJLMT23]. Indeed, if S is strongly mixing, then all its non-trivial powers are ergodic and the statements give some T with a non-trivial power which is not ergodic, so T is not strongly mixing. Here, Theorem F gives an example starting from a very non-strongly mixing system S (the universal odometer). Finally, note that strongly mixing systems are not BSP. This is a consequence of [GZ19, Theorem 1.3]: BSP rank-one systems are not topologically mixing, therefore they are not measure-theoretically strongly mixing.

1.6. Further comments. As they both preserve entropy, we may wonder whether there is a connection between Shannon orbit equivalence (or, more generally, φ -integrable orbit equivalence for φ greater than \log) and *even Kakutani equivalence*. Two ergodic probability measure-preserving bijections S and T , respectively acting on (X, μ) and (Y, ν) , are evenly Kakutani equivalent if there exist measurable subsets $A \subseteq X$ and $B \subseteq Y$ with equal measure, that is, $\mu(A) = \nu(B)$, such that the induced maps S_A and T_B are conjugate. Even Kakutani equivalence is an equivalence relation, contrarily to Shannon orbit equivalence and φ -integrable orbit equivalence *a priori* (except for linear maps φ , by Belinskaya's theorem). The theory of Ornstein, Rudolph and Weiss [ORW82] gives a complete classification up to even Kakutani equivalence among loosely Bernoulli (LB) systems and entropy is a complete invariant. Moreover, the class of LB systems is closed by even Kakutani equivalence, meaning that if S is LB and equivalent to T , then T is also LB.

Rank-one systems are zero-entropy and LB, and by Theorems A, C, E and F, some of them are Shannon orbit equivalent to the universal odometer.

Question 1.4. Does even Kakutani equivalence imply Shannon orbit equivalence or φ -integrable orbit equivalence for some φ ?

In a forthcoming paper, we will provide a new construction of orbit equivalence to prove that the converse is false: for every $\varepsilon > 0$, there exists a non-LB system which is $(x \mapsto x^{(1/2)-\varepsilon})$ -integrably orbit equivalent to the dyadic odometer. So $(x \mapsto x^{(1/2)-\varepsilon})$ -integrable orbit equivalence and Shannon orbit equivalence do not imply even Kakutani equivalence.

1.7. *Outline of the paper.* After a few preliminaries in §2, rank-one systems are defined in §3 using the cutting-and-stacking method. We also define the central notion of flexible classes of rank-one transformations. In §4, we prove Proposition 1.2 (Proposition 3.8 in §3), that is, we show that the classes mentioned in Theorem B (Theorem 3.9 in §3) are flexible. It remains to show that every flexible class admits an element which is φ -integrably orbit equivalent to the universal odometer (Theorem 3.9). In §5, we will describe the construction of Kerr and Li, generalized to rank-one systems, and establish that this is an orbit equivalence with some important properties preparing for the proof of Theorem 3.9. Theorems A, C, E and F directly follow from Proposition 3.8 and Theorem 3.9. We prove Theorem D at the end of the paper.

2. *Preliminaries*

2.1. *Basics of ergodic theory.* The probability space (X, \mathcal{A}, μ) is assumed to be standard and atomless. Such a space is isomorphic to $([0, 1], \mathcal{B}([0, 1]), \text{Leb})$, that is, there exists a bimeasurable bijection $\Psi: X \rightarrow [0, 1]$ (defined almost everywhere) such that $\Psi_*\mu = \text{Leb}$, where $\Psi_*\mu$ is defined by $\Psi_*\mu(A) = \mu(\Psi^{-1}(A))$ for every measurable set A . We consider maps $T: X \rightarrow X$ acting on this space and which are bijective, bimeasurable and *probability measure-preserving (p.m.p.)*, meaning that $\mu(T^{-1}(A)) = \mu(A)$ for all measurable sets $A \subseteq X$, and the set of these transformations is denoted by $\text{Aut}(X, \mathcal{A}, \mu)$, or simply $\text{Aut}(X, \mu)$, two such maps being identified if they coincide on a measurable set of full measure. In this paper, elements of $\text{Aut}(X, \mu)$ are called *transformations* or *(dynamical) systems*.

A measurable set $A \subseteq X$ is *T-invariant* if $\mu(T^{-1}(A)\Delta A) = 0$, where Δ denotes the symmetric difference. A transformation $T \in \text{Aut}(X, \mu)$ is said to be *ergodic* if every T -invariant set is of measure 0 or 1. If T is ergodic, then T is *aperiodic*, that is, $T^n(x) \neq x$ for almost every $x \in X$ and for every $n \in \mathbb{Z} \setminus \{0\}$, or equivalently the *T-orbit* of x , denoted by $\text{Orb}_T(x) := \{T^n(x) \mid n \in \mathbb{Z}\}$, is infinite for almost every $x \in X$.

Here, T is *weakly mixing* if

$$\frac{1}{n} \sum_{k=0}^n |\mu(A \cap T^{-k}(B)) - \mu(A)\mu(B)| \xrightarrow{n \rightarrow +\infty} 0$$

for every measurable sets A, B . Additionally, T is *strongly mixing* if

$$|\mu(A \cap T^{-n}(B)) - \mu(A)\mu(B)| \xrightarrow{n \rightarrow +\infty} 0$$

for every measurable set A, B . It is not difficult to prove that strong mixing implies weak mixing and that the latter implies ergodicity.

The notions of weak mixing and ergodicity can be translated in terms of eigenvalues. Denoting by $L^2(X, \mathcal{A}, \mu)$ the space of complex-valued and square-integrable functions defined on X , a complex number λ is an *eigenvalue* of T if there exists $f \in L^2(X, \mathcal{A}, \mu) \setminus \{0\}$ such that $f \circ T = \lambda f$ almost everywhere (f is then called an eigenfunction). An eigenvalue λ is automatically an element of the unit circle $\mathbb{T} := \{z \in \mathbb{C} \mid |z| = 1\}$. The *point spectrum* of T is then the set of all its eigenvalues. Notice that $\lambda = 1$ is always an eigenvalue since the constant functions are in its eigenspace. Finally, T is ergodic if and only if the constant functions are the only eigenfunctions with eigenvalue one, in other words, the eigenspace of $\lambda = 1$ is the line of constant functions (we say that it is a simple eigenvalue). If T is ergodic, it is weakly mixing if and only if the only eigenvalue of T is 1. For a complete survey on spectral theory for dynamical systems, the reader may refer to [VO16].

All the properties that we have introduced are preserved under conjugacy. Two transformations $S \in \text{Aut}(X, \mu)$ and $T \in \text{Aut}(Y, \nu)$ are *conjugate* if there exists a bimeasurable bijection $\Psi: X \rightarrow Y$ such that $\Psi_*\mu = \nu$ and $\Psi \circ S = T \circ \Psi$ almost everywhere. Some classes of transformations have been classified up to conjugacy, the two examples to keep in mind are the following. By Ornstein [Orn70], entropy is a total invariant of conjugacy among Bernoulli shifts, and Ornstein and Weiss [OW87] generalized this result for Bernoulli shifts of amenable groups. For more details about entropy, see [Dow11] for non-necessarily invertible transformations $T: X \rightarrow X$, and [KL17] more generally for actions of amenable groups. Finally, Halmos and von Neumann [HVN42] showed that two systems with discrete spectra are conjugate if and only if they have equal point spectra (a system has a discrete spectrum if the span of all its eigenfunctions is dense in $L^2(X, \mathcal{A}, \mu)$).

2.2. Quantitative orbit equivalence. The conjugacy problem in full generality is very complicated (see [FRW11]). We now give the formal definition of orbit equivalence, which is a weakening of the conjugacy problem.

Definition 2.1. Two aperiodic transformations $S \in \text{Aut}(X, \mu)$ and $T \in \text{Aut}(Y, \nu)$ are *orbit equivalent* if there exist a bimeasurable bijection $\Psi: X \rightarrow Y$ satisfying $\Psi_*\mu = \nu$, such that $\text{Orb}_S(x) = \text{Orb}_{\Psi^{-1}T\Psi}(x)$ for almost every $x \in X$. The map Ψ is called an *orbit equivalence* between S and T .

We can then define the *cocycles* associated to this orbit equivalence. These are measurable functions $c_S: X \rightarrow \mathbb{Z}$ and $c_T: Y \rightarrow \mathbb{Z}$ defined almost everywhere by

$$Sx = \Psi^{-1}T^{c_S(x)}\Psi(x) \quad \text{and} \quad Ty = \Psi S^{c_T(y)}\Psi^{-1}(y)$$

($c_S(x)$ and $c_T(y)$ are uniquely defined by aperiodicity).

Given a function $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$, a measurable function $f: X \rightarrow \mathbb{Z}$ is said to be *φ -integrable* if

$$\int_X \varphi(|f(x)|) d\mu < +\infty.$$

For example, integrability is exactly φ -integrability when φ is non-zero and linear. Then, a weaker quantification on cocycles is the notion of φ -integrability for a *sublinear* map

φ , meaning that $\lim_{t \rightarrow +\infty} \varphi(t)/t = 0$. Two transformations in $\text{Aut}(X, \mu)$ are said to be φ -integrably orbit equivalent if there exists an orbit equivalence between them whose associated cocycles are φ -integrable. Another form of quantitative orbit equivalence is Shannon orbit equivalence. We say that a measurable function $f: X \rightarrow \mathbb{Z}$ is *Shannon* if the associated partition $\{f^{-1}(n) \mid n \in \mathbb{Z}\}$ of X has finite entropy. Two transformations in $\text{Aut}(X, \mu)$ are *Shannon orbit equivalent* if there exists an orbit equivalence between them whose associated cocycles are Shannon.

3. Rank-one systems

3.1. *The cutting-and-stacking method.* Before the definition of a rank-one system (Definition 3.2), and for the definition of flexible classes (Definition 3.7), we need to define sequences of integers which will provide the combinatorial data of a rank-one system, namely the cutting and spacing parameters.

Definition 3.1. By a *cutting and spacing parameter*, we mean a tuple of the form

$$(q, (\sigma_{\cdot,0}, \dots, \sigma_{\cdot,q}))$$

with an integer $q \geq 2$ (the *cutting parameter*) and non-negative integers $\sigma_{\cdot,0}, \dots, \sigma_{\cdot,q}$ (the *spacing parameters*), and we denote by \mathcal{P} the set of all cutting and spacing parameters. We also define the set of finite sequences of cutting and spacing parameters:

$$\mathcal{P}^* := \bigcup_{n \in \mathbb{N}} \mathcal{P}^n.$$

Given a sequence of cutting and spacing parameters $\mathbf{p} = (q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{k \geq 0} \in \mathcal{P}^{\mathbb{N}}$ and an integer $n \geq 0$, the tuple $(q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))$ in \mathcal{P} is the *n*th *cutting and spacing parameter* of \mathbf{p} , and the tuple $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n}$ is the *projection* of \mathbf{p} on \mathcal{P}^{n+1} (it gives the first $n + 1$ cutting and spacing parameters). From \mathbf{p} , we also define three sequences:

- $(h_n)_{n \geq 0}$ the *height sequence* of \mathbf{p} , inductively defined by

$$\begin{cases} h_0 = 1, \\ h_{n+1} = q_n h_n + \sigma_n, \end{cases}$$

where h_n is called the *height of the n*th *tower*;

- $(\sigma_n)_{n \geq 0}$, with $\sigma_n := \sum_{i=0}^{q_n} \sigma_{n,i}$ (the number of new spacers at step n);
- $(Z_n)_{n \geq 0}$, with $Z_n := \max \{\sigma_{j,i} \mid 0 \leq j \leq n, 0 \leq i \leq q_j\}$,

and it is also possible to consider the finite sequences $(h_k)_{0 \leq k \leq n+1}$, $(\sigma_k)_{0 \leq k \leq n}$ and $(Z_k)_{0 \leq k \leq n}$ associated to a finite sequence of cutting and spacing parameters in \mathcal{P}^{n+1} .

The terminology ‘cutting’, ‘spacing’, ‘tower’, ‘height’, etc., is justified by Definition 3.2 and Figure 4. There are many definitions of rank-one systems (see [Fer97] for a complete survey and various facts in this section). In this paper, the goal is to use the combinatorial structure given by the cutting-and-stacking method (see Figure 4).




FIGURE 4. An example of cutting-and-stacking construction with $h_n = 5, q_n = 3, \sigma_{n,0} = 1, \sigma_{n,1} = 2, \sigma_{n,2} = 0, \sigma_{n,3} = 1$. We then have $h_{n+1} = 19$.

Definition 3.2. A transformation $T \in \text{Aut}(X, \mu)$ is of rank one if there exist:

- (1) a sequence of cutting and spacing parameters $\mathbf{p} = (q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_{n \geq 0} \in \mathcal{P}^{\mathbb{N}}$ satisfying

$$\sum_{n=0}^{+\infty} \frac{\sigma_n}{h_{n+1}} < +\infty, \tag{F}$$

where (h_n) and (σ_n) are the sequences associated to \mathbf{p} , as described in Definition 3.1;

- (2) measurable subsets of X , denoted by B_n (for every $n \geq 0$), $B_{n,i}$ (for every $n \geq 0$ and $0 \leq i \leq q_n - 1$) and $\Sigma_{n,i,j}$ (for every $n \geq 0, 0 \leq i \leq q_n$ and $1 \leq j \leq \sigma_{n,i}$; if $\sigma_{n,i} = 0$, then there are no $\Sigma_{n,i,j}$) such that for all $n \geq 0$:

- (a) $B_n, \dots, T^{h_n-1}(B_n)$ are pairwise disjoint;
- (b) $(B_{n,0}, B_{n,1}, \dots, B_{n,q_n-1})$ is a partition of B_n ;
- (c) $T^{h_n}(B_{n,i}) = \begin{cases} \Sigma_{n,i+1,1} & \text{if } \sigma_{n,i} > 0, \\ B_{n,i+1} & \text{if } \sigma_{n,i} = 0 \text{ and } i < q_n - 1; \end{cases}$
- (d) if $\sigma_{n,i} > 0$, then $T(\Sigma_{n,i,j}) = \begin{cases} \Sigma_{n,i,j+1} & \text{if } j < \sigma_{n,i}, \\ B_{n,i} & \text{if } j = \sigma_{n,i} \text{ and } i \leq q_n - 1; \end{cases}$
- (e) $B_{n+1} = \begin{cases} \Sigma_{n,0,1} & \text{if } \sigma_{n,0} > 0, \\ B_{n,1} & \text{if } \sigma_{n,0} = 0, \end{cases}$

and if the Rokhlin towers $\mathcal{R}_n := (T^k(B_n))_{0 \leq k \leq h_n-1}$ are increasing to the σ -algebra \mathcal{A} meaning that the σ -algebra generated by $\{T^k(B_n) \mid n \in \mathbb{N}, 0 \leq k \leq h_n - 1\}$ is \mathcal{A} up to null sets (since \mathcal{A} is standard, this also means that $\{T^k(B_n) \mid n \in \mathbb{N}, 0 \leq k \leq h_n - 1\}$ separates the points). Note that \mathcal{R}_0 is the tower with only one level B_0 . The sets $\Sigma_{n,i,j}$ are called the *spacers*. In this paper, we will usually write:

- $X_n := B_n \sqcup \dots \sqcup T^{h_n-1}(B_n)$ as the subset covered by the n th tower \mathcal{R}_n ;
- $\varepsilon_n := \mu((X_n)^c)$, where $(X_n)^c$ denotes the complement of the subset X_n of X .

Since X_n is increasing and \mathcal{R}_n increases to the atomless σ -algebra \mathcal{A} , we have $\mu(X_n) \xrightarrow{n \rightarrow +\infty} 1$. In other words ε_n tends to 0.

Before giving examples, the following lemmas give some easy properties on these systems to understand their combinatorial structure and the hypotheses required in the definition.

LEMMA 3.3. *Let (h_n) and (σ_n) be the sequences associated to $(q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_n \in \mathcal{P}^{\mathbb{N}}$ (see Definition 3.1). The following assertions are equivalent:*

- (1) *the series $\sum(\sigma_n/h_{n+1})$ converges (condition (F) in Definition 3.2);*
- (2) *the series $\sum(\sigma_n/q_0 \dots q_n)$ converges;*
- (3) *there exists a constant $M_0 \leq 1$ such that $h_{n+1} \underset{n \rightarrow +\infty}{\sim} (q_0 \dots q_n/M_0)$,*

and if one of these equivalent assertions is true, then $\sum_{n \geq 0} (\sigma_n/q_0 \dots q_n) = (1/M_0) - 1$.

Proof of Lemma 3.3. If the series $\sum(\sigma_n/q_0 \dots q_n)$ converges, so does the series $\sum(\sigma_n/h_{n+1})$ since h_{n+1} is greater or equal to $q_0 \dots q_n$. Now assume that the series $\sum(\sigma_n/h_{n+1})$ converges. Notice that we have

$$\frac{\sigma_n}{h_{n+1}} = \frac{h_{n+1} - q_n h_n}{h_{n+1}} = 1 - q_n \frac{h_n}{h_{n+1}}$$

and since the series is convergent, the product $\prod q_n (h_n/h_{n+1})$ converges to some $M_0 > 0$, that is, $q_0 \dots q_n/h_{n+1} \rightarrow M_0$. The constant M_0 is less than or equal to 1 since we have $h_{n+1} \geq q_n h_n$ for every $n \geq 0$. Finally, let us assume $q_0 \dots q_n/h_{n+1} \rightarrow M_0$. Notice that we have

$$\frac{\sigma_n}{q_0 \dots q_n} = \frac{h_{n+1} - q_n h_n}{q_0 \dots q_n} = \frac{h_{n+1}}{q_0 \dots q_n} - \frac{h_n}{q_0 \dots q_{n-1}},$$

so by telescoping consecutive terms, we get $\sum_{n \geq 0} (\sigma_n/q_0 \dots q_n) = \lim_{n \rightarrow \infty} (h_{n+1}/q_0 \dots q_n) - h_0 = (1/M_0) - 1$ and we are done for the equivalence between the three assumptions. □

LEMMA 3.4. *Let $T: X \rightarrow X$ be a bimeasurable bijection. Assume that T preserves a non-zero measure μ and it admits a sequence of Rokhlin towers as in Definition 3.2. The following hold:*

- (1) *the levels $T^k(B_n)$ of the n th Rokhlin tower \mathcal{R}_n have μ -measure $\mu(B_0)/q_0 \dots q_{n-1}$;*
- (2) *μ is finite if and only if condition (F) is satisfied. Furthermore, if μ is a probability measure (this implies that T is a rank-one system), then $\mu(B_0) = M_0$ and $h_{n+1} \leq (q_0 \dots q_n/M_0)$, where M_0 is given by Lemma 3.3.*

Proof of Lemma 3.4. For a fixed n , the levels of \mathcal{R}_n have the same measure by T -invariance of the measure μ . Moreover, the first level B_n is a disjoint union of q_n levels $B_{n,0}, \dots, B_{n,q_n-1}$ of \mathcal{R}_{n+1} . Then, it is clear by induction that a level of \mathcal{R}_n has measure $\mu(B_0)/q_0 \dots q_{n-1}$. Since the sequence $(X_n)_{n \geq 0}$ is increasing to X , and X_{n+1} is obtained from X_n by adding σ_n spacers, which are levels of \mathcal{R}_{n+1} , we get

$$\mu(X) = \mu(X_0) + \sum_{n \geq 0} \mu(X_{n+1} \setminus X_n) = \mu(B_0) + \sum_{n \geq 0} \frac{\mu(B_0)\sigma_n}{q_0 \dots q_n}, \tag{1}$$

so $\mu(B_0)$ is non-zero, and $\mu(X)$ is finite if and only if the sum $\sum_{n \geq 0} (\sigma_n/q_0 \dots q_n)$ is finite. Finally, let us assume that μ is a probability measure. This implies $\sum_{n \geq 0} (\sigma_n/q_0 \dots q_n) = (1/M_0) - 1$ and, using equation (1), we get $\mu(B_0) = M_0$. The measurable set X_n is the disjoint union of h_n levels of \mathcal{R}_n , so the inequality $h_n \leq (q_0 \dots q_{n-1}/M_1)$ follows from the fact that μ is a probability measure. \square

It is possible to build a finite measure-preserving transformation T of rank one with a given combinatorial setting $(q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_{n \geq 0} \in \mathcal{P}^{\mathbb{N}}$ satisfying the hypothesis (F). For instance, it suffices to build (X_n) as an increasing sequence of intervals of \mathbb{R}_+ , with $B_{n,i}$ and $\Sigma_{n,i,j}$ being subintervals of equal length and disjoint (for a fixed n), each on which T is defined as an affine map, and with $B_0 = [0, M_0]$. The convergence of the series $\sum (\sigma_n/h_{n+1})$ and Lemma 3.3 ensure that $X := \bigcup X_n$ is equal to $[0, 1]$ (up to a null set), so the Lebesgue measure on $[0, 1]$ is a probability measure preserved by T . Notice that if the series is divergent, we can set $B_0 = [0, 1]$ and this defines T on the set of positive real numbers endowed with the Lebesgue measure, so this is an infinite measure-preserving transformation.

Therefore, for every $(q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_{n \geq 0} \in \mathcal{P}^{\mathbb{N}}$ satisfying condition (F), there exists a rank-one system having a cutting-and-stacking construction with these cutting and spacing parameters, this fact will be used in this paper since it enables us to only take into account the combinatorics behind the systems.

The hypothesis on the Rokhlin towers \mathcal{R}_n aims not only to have $\varepsilon_n \rightarrow 0$ but also to define two isomorphic systems when they admit cutting-and-stacking constructions with the same cutting and spacing parameters. Moreover, if T admits such a construction with Rokhlin towers increasing to a sub- σ -algebra \mathcal{B} of \mathcal{A} , then T , seen as an element of $\text{Aut}(X, \mathcal{A}, \mu)$, is not necessarily a rank-one system but admits a rank-one system (T on the sub- σ -algebra \mathcal{B}) as a factor.

Two different families of cutting and spacing parameters do not necessarily define non-isomorphic systems. Indeed, in a construction of a rank-one system with parameters q_n and $\sigma_{n,i}$, one can decide to only consider a subsequence \mathcal{R}_{n_k} of Rokhlin towers. For example, the new cutting parameters will be $q_{n_k}q_{n_k+1} \dots q_{n_{k+1}-1}$ for $k \geq 0$.

The rank-one systems form a class of ergodic and zero entropy systems. The easiest examples of rank-one systems are the *irrational rotations*

$$R_\theta : z \in \mathbb{T} \mapsto e^{2i\pi\theta} z \in \mathbb{T}$$

for every irrational number θ , where \mathbb{T} is the unit circle endowed with its Haar measure. These systems are not weakly mixing. Moreover, they have discrete spectrum and the point spectrum of R_θ is $\{e^{in\theta} \mid n \in \mathbb{Z}\}$, so by the Halmos–von Neumann theorem [HVN42], R_θ and $R_{\theta'}$ are isomorphic if and only if $\theta = \theta' \bmod \mathbb{Z}$ or $\theta = -\theta' \bmod \mathbb{Z}$.

The *odometers* are rank one. These are exactly the rank-one systems without spacers (that is, $\sigma_{n,i} = 0$), so the Rokhlin towers are partitions of the space. Such a system is isomorphic to the adding machine S in the space $\prod_{n \geq 0} \{0, 1, \dots, q_n - 1\}$,

namely the addition by $(1, 0, 0, 0, \dots)$ with carry over to the right, defined for every $x \in \prod_{n \geq 0} \{0, 1, \dots, q_n - 1\}$ by

$$Sx = \begin{cases} (0, \dots, 0, x_i + 1, x_{i+1}, \dots) & \text{if } i := \min \{j \geq 0 \mid x_j \neq q_j - 1\} \text{ is finite,} \\ (0, 0, 0, \dots) & \text{if } x = (q_0 - 1, q_1 - 1, q_2 - 1, \dots), \end{cases}$$

and it preserves the product of uniform probability measures on each finite set $\{0, 1, \dots, q_n - 1\}$. Denote the cylinders of length k by

$$[x_0, \dots, x_{k-1}]_k := \left\{ y \in \prod_{n \geq 0} \{0, 1, \dots, q_n - 1\} \mid y_0 = x_0, \dots, y_{k-1} = x_{k-1} \right\}.$$

If S is the odometer on the space $\prod_{n \geq 0} \{0, 1, \dots, q_n - 1\}$, we can also set a partially defined map

$$\zeta_n : X \setminus [q_0 - 1, \dots, q_{n-1} - 1]_n \rightarrow X \setminus [0, \dots, 0]_n$$

which is the addition by

$$\underbrace{(0, \dots, 0, 1, 0, 0, \dots)}_{n-1 \text{ times}}$$

(so S and ζ_1 coincide on $X \setminus [q_0 - 1]_1$). Then, we have

$$B_n = \underbrace{[0, \dots, 0]_n}_{n \text{ times}}$$

$$B_{n,i} = \underbrace{[0, \dots, 0, i]_{n+1}}_{n \text{ times}}$$

and $B_{n,i} = \zeta_{n+1}^i(B_{n,0})$ for every $0 \leq i \leq q_n - 1$, so it provides a scale in B_n . Note that it is possible to recover the odometer S from these partially defined maps ζ_n (see Figure 5). In §5.1, the strategy will be to build S from partially defined maps ζ_n .

In the class of odometers, the number of occurrences of every prime factors in the set $\{q_n \mid n \geq 0\}$ form a total invariant of conjugacy. As for irrational rotations, it is a consequence of the Halmos–von Neumann theorem since odometers have a discrete spectrum and their eigenvalues are given by these occurrences. In particular, odometers have eigenvalues non-equal to 1 and are not weakly mixing; moreover, odometers and irrational rotations are not isomorphic. Notice that the Halmos–von Neumann theorem implies that the conjugacy classes among systems with a discrete spectrum coincide with the flip-conjugacy classes since the point spectrum of a system is a subgroup of \mathbb{T} . If every prime number has infinite multiplicity in the set $\{q_n \mid n \geq 0\}$, then the odometer is said to be *universal*. An odometer is *dyadic* if 2 is the only prime factor.

Chacon’s map is the first example of weakly mixing system which is not strongly mixing [Cha69] and was the starting point for the notion of rank-one systems. It is a rank-one transformation defined with cutting and spacing parameters $q_n = 3$, $\sigma_{n,0} = \sigma_{n,1} = \sigma_{n,3} = 0$, $\sigma_{n,2} = 1$.




FIGURE 5. Example of odometer with $q_0 = 3, q_1 = 2, q_2 = 3$.

3.2. *Flexible classes.* Now we introduce classes of rank-one systems to which the main result of this paper applies. First, let us consider cutting-and-stacking constructions whose spacing parameters have controlled asymptotics. Recall that $\mathcal{P}^{\mathbb{N}}$ is the set of sequences of cutting and spacing parameters. As introduced in Definition 3.1, (h_n) , (σ_n) and (Z_n) denote the sequences associated to a sequence in $\mathcal{P}^{\mathbb{N}}$: h_n is the height of the n th tower, σ_n the number of new spacers at step n and Z_n is the maximum number of spacers between two consecutive towers, over the first n steps.

Definition 3.5. A construction by cutting and stacking with cutting and spacing parameters $(q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_{n \geq 0} \in \mathcal{P}^{\mathbb{N}}$ is said to be *CSP (controlled-spacing-parameter)* if there exists a constant $C > 0$ such that $Z_n \leq Ch_n$ for all n . It is furthermore *BSP (bounded-spacing-parameter)* if $Z_n \leq C$ and $\sigma_{n,0} = \sigma_{n,q_n} = 0$ for all n . A rank-one system T is *BSP* if it admits a BSP cutting-and-stacking construction.

Odometers and Chacon’s map are examples of BSP rank-one systems. Moreover, BSP implies CSP. The interest in the BSP property is its stability after skipping steps in the cutting-and-stacking process, as stated in the following lemma.

LEMMA 3.6. *Given a BSP cutting-and-stacking construction, any subsequence of its Rokhlin towers still provides a BSP construction with the same constant C .*

Proof of Lemma 3.6. Let q_n and $\sigma_{n,i}$ be the cutting and spacing parameters of the BSP construction, C the bound for the spacing parameters $\sigma_{n,i}$, \mathcal{R}_n the associated towers and




FIGURE 6. Illustration of the proof of Lemma 3.6, spacing parameters from \mathcal{R}_n to \mathcal{R}_{n+2} with $q_n = q_{n+1} = 2$ (the coloured levels are the base and the roof of the towers).

\mathcal{R}_{n_k} a subsequence. Let k be an integer and assume $n_{k+1} = n_k + 2$. Denote by q'_{n_k} and $\sigma'_{n_k,i}$ the new cutting and spacing parameters from \mathcal{R}_{n_k} to $\mathcal{R}_{n_{k+1}}$. It is easy to show that $q'_{n_k} = q_{n_k} q_{n_{k+1}}$, $\sigma'_{n_k,0} = \sigma'_{n_k,q'_{n_k}} = 0$ and for every $1 \leq j \leq q_{n_{k+1}}$, $\sigma'_{n_k,(j-1)q_{n_k}+i}$ is equal to $\sigma_{n_k,i}$ if $1 \leq i \leq q_{n_k} - 1$ and $\sigma_{n_{k+1},j}$ if $i = q_{n_k}$ (see Figure 6). Thus, the non-zero spacing parameters from \mathcal{R}_{n_k} to $\mathcal{R}_{n_{k+1}}$ are of the form $\sigma_{n_k,i}$ or $\sigma_{n_{k+1},i}$ and they are all bounded above by C . For n_{k+1} bigger than $n_k + 2$, the result is now clear by induction. \square

If the parameters σ_{n,q_n} are non-zero, then skipping steps in the cutting-and-stacking process will cause an accumulation of spacers above the last columns and the new spacing parameters will not be bounded if the subsequence is properly chosen so that the jumps $n_{k+1} - n_k$ increase quickly enough. We have the same problem for $\sigma_{n,0}$ (accumulation of spacers at the bottom of the first columns), and hence the conditions $\sigma_{n,0} = \sigma_{n,q_n} = 0$ in the definition of BSP.

Lemma 3.6 has no reason to hold for CSP construction that are not BSP. Indeed, the spacing parameters from \mathcal{R}_{n_k} to $\mathcal{R}_{n_{k+1}}$ have to be compared with h_{n_k} , the height of \mathcal{R}_{n_k} . The comparison is easily obtained for the spacing parameters $\sigma_{n_k,i}$, $0 \leq i \leq q_{n_k}$, but for the other spacing parameters, we only know that they are bounded above by $Ch_{n_{k+1}}, Ch_{n_{k+2}}, \dots, Ch_{n_{k+1}-1}$.

In the following, we will see other important CSP examples by considering classes containing ‘nice’ cutting-and-stacking constructions, meaning that we will be able to properly choose the parameters to have the desired quantification of the cocycles for the orbit equivalence built in §5.1. By definition, every flexible class \mathcal{C} will be associated to some subset $\mathcal{F}_{\mathcal{C}}$ of \mathcal{P}^* , which can be considered as sufficient conditions that the cutting and spacing parameters have to satisfy at each step for the underlying transformation to

belong to \mathcal{C} . Recall that \mathcal{P}^* denotes the set of all finite sequences of cutting and stacking parameters.

Definition 3.7. A class \mathcal{C} of rank-one systems is said to be *flexible* if there exists a subset $\mathcal{F}_{\mathcal{C}}$ of \mathcal{P}^* satisfying the following properties:

- (1) there exists a constant $C > 0$ such that for all $(q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_{n \geq 0} \in \mathcal{P}^{\mathbb{N}}$ satisfying condition (F) in Definition 3.2, if $\mathcal{F}_{\mathcal{C}}$ contains every projection $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n} \in \mathcal{P}^{n+1}$ for $n \geq 0$, then these parameters define a CSP construction (with the constant C) and the underlying rank-one transformation is in \mathcal{C} ;
- (2) there exists a cutting and spacing parameter $(q_0, (\sigma_{0,0}, \dots, \sigma_{0,q_0}))$ in $\mathcal{F}_{\mathcal{C}}$ with $q_0 \geq 3$;
- (3) there is a constant $C' > 0$ such that for all $n \geq 1$, if $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n-1}$ is in $\mathcal{F}_{\mathcal{C}}$, then there are infinitely many integers q_n such that $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n}$ is in $\mathcal{F}_{\mathcal{C}}$ for some $\sigma_{n,0}, \dots, \sigma_{n,q_n}$ satisfying the inequality

$$\sigma_n \leq C' q_n h_{n-1},$$

where $(h_k)_{0 \leq k \leq n+1}$ and $(\sigma_k)_{0 \leq k \leq n}$ denote the finite sequences associated to the finite sequence $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n}$ of cutting and stacking parameters.

A rank-one system T is *flexible* if $\{T\}$ is a flexible class.

The third point of the definition aims to recursively choose the cutting parameters (and we want them to increase quickly enough) with an asymptotic control on $(\sigma_n)_n$, while the first point guarantees that it is possible to do so for the underlying system to be in the class \mathcal{C} . The second point is minor, but it is required for the initialization of the recursive construction of an odometer orbit equivalent to an element of our flexible class (see Lemma 5.10 and Remark 5.11). It also ensures that $\mathcal{F}_{\mathcal{C}}$ is not an empty set.

Notice that if a construction satisfies $Z_n \leq C h_{n-1}$ for all n , then it is in particular CSP and we get $\sigma_n \leq C(q_n + 1)h_{n-1} \leq 2C q_n h_{n-1}$ as in the third point of Definition 3.7.

We now give examples of flexible classes. The proof is given in §4.

PROPOSITION 3.8.

- (1) Every BSP rank-one system is flexible.
- (2) For every non-empty open subset \mathcal{V} of \mathbb{R} , the set $\{R_{\theta} \mid \theta \in \mathcal{V} \cap (\mathbb{R} \setminus \mathbb{Q})\}$ is a flexible class.
- (3) For every irrational number θ , the class of rank-one systems which have $e^{2i\pi\theta}$ as an eigenvalue is flexible.
- (4) The class of strongly mixing rank-one systems is flexible.

Theorems A, C, E and F follow from Proposition 3.8 and the following theorem which is the main result.

THEOREM 3.9. Let $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a map satisfying $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$. If \mathcal{C} is a flexible class, then there exists T in \mathcal{C} which is φ -integrably orbit equivalent to the universal odometer.

4. Proof of Proposition 3.8

In this section, we prove the four statements in Proposition 3.8.

4.1. *BSP systems.* Let T be a BSP rank-one system, $\mathcal{C} := \{T\}$ and $q_n, \sigma_{n,i}$ the parameters of a BSP construction of T , with a constant $C > 0$. For every $n \geq 0$ and $j \geq 1$, let $\sigma_0^{(n,n+j)}, \dots, \sigma_{q_n \dots q_{n+j-1}}^{(n,n+j)}$ be the spacing parameters from \mathcal{R}_n to \mathcal{R}_{n+j} , assuming that the steps for $\mathcal{R}_{n+1}, \dots, \mathcal{R}_{n+j-1}$ are skipped during the construction (we then have $\sigma_i^{(n,n+1)} = \sigma_{n,i}$ and also $\sigma_i^{(n,n+j)} = 0$ for i equal to 0 and $q_n \dots q_{n+j-1}$ by Lemma 3.6). The new cutting parameters are $q^{(n,n+j)} := q_n \dots q_{n+j-1}$ and are large enough with huge jumps j . Now, define

$$\mathcal{F}_{\mathcal{C}} := \{ (q^{(n_k, n_{k+1})}, (\sigma_0^{(n_k, n_{k+1})}, \dots, \sigma_{(q^{(n_k, n_{k+1})})}^{(n_k, n_{k+1})})) \mid 0 \leq k \leq m \mid m \geq 0, 0 = n_0 < n_1 < \dots < n_{m+1} \}.$$

Using Lemma 3.6, the new spacing parameters $\sigma_j^{(n_k, n_{k+1})}$ are bounded by C and we get

$$\sum_{1 \leq j \leq q^{(n_k, n_{k+1})}} \sigma_j^{(n_k, n_{k+1})} \leq C q^{(n_k, n_{k+1})}.$$

The set of parameters $\mathcal{F}_{\mathcal{C}}$ thus witnesses that $\{T\}$ is flexible.

4.2. *Irrational rotations.* We now consider a construction from [DEJ+23]. For every irrational number θ , Drillick *et al* give an explicit cutting-and-stacking construction of a transformation T which is the irrational rotation of angle θ when the construction yields a finite measure-preserving system.

4.2.1. *The construction in [DEJ+23].* Let θ be an irrational number and $[q_{-1}; q_0, q_1, \dots]$ its continued fraction expansion, with $q_{-1} := \lfloor \theta \rfloor$ and positive integers q_0, q_1, \dots . Let us assume that there is no integer n such that $q_k = 1$ for every $k \geq n$. We consider the sequence $(h_n)_{n \geq 0}$ defined by $h_{-1} := 0, h_0 := 1$ and $h_{k+1} := q_k h_k + h_{k-1}$ for every $k \geq 0$ (the integer h_k is the denominator of the k th convergent of the irrational number θ). Finally, for every $k \geq 0$, we set $\sigma_{k,i} = 0$ for every $i \in \{0, 1, \dots, q_k - 1\}$, and $\sigma_{k,q_k} = h_{k-1}$. Then, the sequence of cutting and stacking parameters $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{k \geq 0}$ provides a rank-one system. This system is the irrational rotation of angle θ if and only if condition (F) is satisfied, and this last condition holds if and only if the series $\sum_{k \geq 0} (1/q_k q_{k+1})$ converges (see [DEJ+23, Theorems 3.1 and 5.1]).

Remark 4.1. Equivalently, we can define rank-one systems with cutting parameters potentially equal to 1, provided that there are infinitely many cutting parameters greater than or equal to 2, but our construction of orbit equivalence is not well defined with this weaker assumption. Therefore, in the proof of Proposition 3.8 for irrational rotations, one of the main goals is to avoid cutting parameters equal to 1.

Remark 4.2. It is proven in [DEJ+23] that the set of irrational numbers θ such that the associated series $\sum_{k \geq 0} (1/q_k q_{k+1})$ converges has measure zero.

4.2.2. *Proof of Proposition 3.8 for these systems.* Let \mathcal{V} be a non-empty open subset of \mathbb{R} and

$$\mathcal{C} := \{R_\theta \mid \theta \in \mathcal{V} \cap (\mathbb{R} \setminus \mathbb{Q})\}.$$

We now prove that \mathcal{C} is a flexible class.

We first use the following basic fact from the theory of continued fractions: if A denotes the set of sequences $(q_i)_{i \geq -1}$ of integers such that q_0, q_1, \dots are positive, and if A is equipped with the induced product topology, then the map

$$(q_i)_{i \geq -1} \in A \mapsto [q_{-1}; q_0, q_1, \dots] \in \mathbb{R} \setminus \mathbb{Q}$$

is a homeomorphism (see [EW11, Lemma 3.4] for instance). We can then fix integers $n_0 \geq 0$ and $Q_{-1}, Q_0, \dots, Q_{n_0}$ (where Q_0, \dots, Q_{n_0} are positive) such that $Q_0 \dots Q_{n_0}$ is greater than or equal to 3 and the following holds: for every irrational number θ , if the first coefficients of its continued fraction expansion are $Q_{-1}, Q_0, \dots, Q_{n_0}$, then θ is in \mathcal{V} .

We write $\mathbf{Q} := (Q_0, \dots, Q_{n_0})$ and we consider the set $\tilde{\mathcal{F}}(\mathbf{Q})$ of finite sequences $(\tilde{q}_k, (\tilde{\sigma}_{k,0}, \dots, \tilde{\sigma}_{k,\tilde{q}_k}))_{0 \leq k \leq n}$ such that $n \geq n_0$ and for all $k \in \{0, \dots, n\}$,

$$\begin{aligned} \tilde{q}_k &= Q_k && \text{if } k \leq n_0, \\ \tilde{q}_k &\geq 2 && \text{if } k > n_0, \end{aligned}$$

and

$$\begin{aligned} \tilde{\sigma}_{k,i} &= 0 && \text{for } i \in \{0, \dots, \tilde{q}_k - 1\}, \\ \tilde{\sigma}_{k,\tilde{q}_k} &= \tilde{h}_{k-1} \end{aligned}$$

(where $(\tilde{h}_k)_{0 \leq k \leq n}$ is the associated height sequence and $\tilde{h}_{-1} := 0$). The finite sequences of $\tilde{\mathcal{F}}(\mathbf{Q})$ may not be finite sequences of cutting and stacking parameters in the sense of Definition 3.1, since the integers Q_0, \dots, Q_{n_0} may be equal to 1. Moreover, even if the integers Q_0, \dots, Q_{n_0} were greater than or equal to 2, we could not prove that \mathcal{C} is a flexible class with $\mathcal{F}_{\mathcal{C}} = \tilde{\mathcal{F}}(\mathbf{Q})$, since the first cutting parameters $\tilde{q}_1, \dots, \tilde{q}_{n_0}$ cannot be chosen large enough. Notice that, although we may have $\tilde{q}_k = 1$ for some $k \in \{0, 1, \dots, n_0\}$, a finite sequence $(\tilde{q}_k, (\tilde{\sigma}_{k,0}, \dots, \tilde{\sigma}_{k,\tilde{q}_k}))_{0 \leq k \leq n}$ can still define the first $(n + 1)$ steps of a cutting-and-stacking construction, and we associate to it another finite sequence $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n-n_0}$ corresponding to the cutting-and-stacking construction obtained from the previous one by skipping the steps $1, 2, \dots, n_0$. We get $h_0 = \tilde{h}_0 = 1, q_0 = Q_0 \dots Q_{n_0} \geq 3$ and for all $k \in \{1, \dots, n - n_0\}$,

$$\begin{aligned} q_k &= \tilde{q}_{n_0+k} \geq 2, \\ h_k &= \tilde{h}_{n_0+k}, \\ \sigma_{k,i} &= \tilde{\sigma}_{n_0+k,i} = 0 && \text{for } i \in \{0, \dots, q_k - 1\}, \\ \sigma_{k,q_k} &= \tilde{\sigma}_{n_0+k,\tilde{q}_{n_0+k}} = \tilde{h}_{n_0+k-1} = h_{k-1} && \text{if } k \geq 2. \end{aligned}$$

For $k = 1$, we have $\sigma_{1,q_1} = \tilde{h}_{n_0}$, where \tilde{h}_{n_0} is not equal to h_0 . Setting $C = C' := \tilde{h}_{n_0}$ (this constant only depends on Q_0, \dots, Q_{n_0}), we have $Z_1 \leq Ch_1$ and $\sigma_1 \leq C'h_0$. We immediately get the inequalities $Z_k \leq Ch_k$ and $\sigma_k \leq C'h_{k-1}$ for $k \in \{2, \dots, n - n_0\}$.

Let $\mathcal{F}(\mathbf{Q})$ be the set of finite sequences $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n-n_0}$ obtained from finite sequences $(\tilde{q}_k, (\tilde{\sigma}_{k,0}, \dots, \tilde{\sigma}_{k,q_k}))_{0 \leq k \leq n} \in \tilde{\mathcal{F}}(\mathbf{Q})$. It is now easy to check that \mathcal{C} is a flexible class, with the set of parameters $\mathcal{F}_{\mathcal{C}} := \mathcal{F}(\mathbf{Q})$ and the constants C and C' .

4.3. *Systems with a given eigenvalue.* Let θ be an irrational number and \mathcal{C} the class of rank-one systems which has $\lambda := e^{2i\pi\theta}$ as an eigenvalue. In [DV23], Danilenko and Vieprík present an explicit cutting-and-stacking construction of a system in \mathcal{C} . The parameters are chosen in the following way (see [DV23, proof of Theorem 4.1]).

4.3.1. *The construction of Danilenko and Vieprík.* For every $n \geq 1$, we fix a number $j_n \in \{1, \dots, n\}$ such that $\delta_n := |1 - \lambda^{j_n}|$ is less than $2\pi/n$. Fix $n \geq 1$, assume that $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n-1}$ has already been constructed with an auxiliary condition

$$h_n > \frac{n^4}{\delta_{n^2}}. \tag{2}$$

Danilenko and Vieprík show the existence of a sequence $(\ell_m^{(n)})_{m \geq 1}$ of positive integers less than or equal to $2\pi/\delta_{n^2}$, such that for every $m \geq 1$,

$$|1 - \lambda^{mh_n + (\ell_1^{(n)} + \dots + \ell_m^{(n)})j_n}| < \frac{2\pi}{n^2}. \tag{3}$$

Next, let q_n be an integer large enough so that the auxiliary condition (2) holds at the next step, namely

$$h_{n+1} := q_n h_n + (\ell_1^{(n)} + \dots + \ell_{q_n-1}^{(n)})j_n > \frac{(n+1)^4}{\delta_{(n+1)^2}}$$

(in [DV23], q_n is chosen as the smallest integer satisfying the property, but it is not needed, so there are infinitely many choices). Finally, the spacing parameters at this step are defined by $\sigma_{n,0} = \sigma_{n,q_n} = 0$ and $\sigma_{n,m} = \ell_m^{(n)} j_n$ for $1 \leq m \leq q_n - 1$.

With these parameters satisfying conditions (2) and (3), λ is an eigenvalue of the underlying rank-one system (see [DV23, proof of Theorem 4.1] for details).

4.3.2. *Proof of Proposition 3.8 for these systems.* Let us consider the same construction as above, but with the following auxiliary condition:

$$h_n > \max \left(\frac{n^4}{\delta_{n^2}}, \frac{(n+1)^4}{\delta_{(n+1)^2}} \right), \tag{4}$$

which is stronger than the previous auxiliary condition (2). Note that the real numbers δ_i have been fixed before setting the parameters.

The subset $\mathcal{F}_{\mathcal{C}}$ of \mathcal{P}^* is defined to be the set of finite sequences $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n}$ constructed in a recursive way. Any cutting and spacing parameter $(q_0, (\sigma_{0,0}, \dots, \sigma_{q_0,0}))$ is in $\mathcal{F}_{\mathcal{C}}$, and if $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n-1}$ is in $\mathcal{F}_{\mathcal{C}}$, then so is $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n}$ for every $(q_n, \sigma_{n,0}, \dots, \sigma_{n,q_n})$ that we can obtain at the next step, as described above, but with the stronger auxiliary condition (4).

Let $\mathbf{p} := (q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_{n \geq 0}$ be a sequence of cutting and spacing parameters. If all its projections are in \mathcal{F}_C , then \mathbf{p} provides a CSP construction with $C = 2\pi$. Indeed, we have $\sigma_{n,m} = \ell_m^{(n)} j_{n^2} \leq 2\pi n^2 / \delta_{n^2} < 2\pi h_n$. As mentioned above, conditions (2) and (3) imply that the sequence \mathbf{p} provides rank-one systems which have λ as an eigenvalue.

Finally, if $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n-1}$ is a finite sequence in \mathcal{F}_C , we can choose a large enough integer q_n so that the following holds at the next step:

$$h_{n+1} > \max \left(\frac{(n+1)^4}{\delta_{(n+1)^2}}, \frac{(n+2)^4}{\delta_{(n+2)^2}} \right)$$

(in particular, the new auxiliary condition (4) is satisfied). We use the same spacing parameters as before, namely $\sigma_{n,m} = \ell_m^{(n)} j_{n^2}$. Using $j_{n^2} \leq n^2$ and $\ell_m^{(n)} \leq (2\pi / \delta_{n^2})$, this gives

$$\sigma_n = (\ell_1^{(n)} + \dots + \ell_{q_n-1}^{(n)}) j_{n^2} \leq q_n \frac{2\pi n^2}{\delta_{n^2}} \leq q_n h_{n-1},$$

so the third point of Definition 3.7 is satisfied for $C' = 1$.

4.4. *Strongly mixing systems.* Let \mathcal{C} be the class of strongly mixing rank-one systems. We consider the construction of Ornstein in [Orn72]. The property the parameters have to satisfy at each step is given by the following lemma (in [Orn72, Lemma 3.2]), proven with a probabilistic method.

LEMMA 4.3. *Let N and K be positive integers and $\varepsilon > 0, \alpha > 0$. Then, there exist integers $m > N$ and a_1, \dots, a_m such that:*

- $|\sum_{i=j}^{j+k} a_i| \leq K$ for all $1 \leq j \leq j+k \leq m$;
- denoting by $H(\ell, k)$ the number of j such that $\sum_{i=j}^{j+k} a_i = \ell$, for $1 \leq j \leq j+k \leq m$, then $H(\ell, k) < \alpha((m-k)/K)$ for every $k < (1-\varepsilon)m$.

The set of parameters \mathcal{F}_C is defined in a recursive way, as in §4.3: any cutting and spacing parameter $(q_0, (\sigma_{0,0}, \dots, \sigma_{q_0,0}))$ is in \mathcal{F}_C , and from a finite sequence of parameters $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n-1}$ in \mathcal{F}_C , $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n}$ is also in \mathcal{F}_C if the new parameters can be written as $q_n = m$ and $\sigma_{n,i} = a_i + h_{n-1}$, where m, a_1, \dots, a_m are integers whose existence is granted by Lemma 4.3 with $N > 10^n$, $K = h_{n-1}$, $\varepsilon = 10^{n-3}$ and $\alpha = 5/4$. There are infinitely many possibilities for q_n as N can be arbitrarily large. It is shown in [Orn72] that cutting-and-stacking constructions with these parameters give strongly mixing systems, and it is clear that they are CSP with $C = 2$ and the third point of Definition 3.7 is satisfied for $C' = 2$.

5. From flexible classes to the universal odometer

The goal of this section is to prove Theorem 3.9, namely that for every $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfying $\varphi(x) = o(t^{1/3})$, any flexible class contains a rank-one system that is φ -integrably orbit equivalent to the universal odometer.

5.1. *The construction.*

5.1.1. *Overview of the construction.* We first present a natural adaptation to the case of the rank-one system of Kerr and Li’s construction of an explicit orbit equivalence between the universal odometer and any other odometers. We will then see that the quantification of the cocycles becomes more complicated due to the presence of non-zero spacing parameters.

Let $T \in \text{Aut}(X, \mu)$ be a rank-one system and consider a cutting-and-stacking construction of this transformation with the same notation $q_n, \sigma_{n,i}, \sigma_n, h_n, \mathcal{R}_n, \varepsilon_n, X_n$ as in Definition 3.2. From the sequence of Rokhlin towers \mathcal{R}_n , new towers \mathcal{R}'_n will be built as Rokhlin towers for a new system S . These towers \mathcal{R}'_n will have no spacers, that is, $\sigma'_{n,i} = 0$, so they will be partitions of X . The construction will ensure that \mathcal{R}'_n increases to the σ -algebra \mathcal{A} using the fact that it is the case for \mathcal{R}_n , so S will be an odometer. For the odometer S to be universal, we fix a sequence of prime numbers $(p_n)_{n \geq 0}$ such that every prime number appears infinitely many times, and every cutting parameter q'_n will be a multiple of p_n .

We will recursively define S on subsets increasing to X up to a null set. More precisely, if the n th tower \mathcal{R}'_n has been built and its base and its height are denoted by B'_n and h'_n , then S is provisionally defined on all the levels of the tower except the highest one and maps the i th level to the $(i + 1)$ th one. So, \mathcal{R}'_n is exactly $(B'_n, S(B'_n), \dots, S^{h'_n-1}(B'_n))$ and S is defined on $X \setminus S^{h'_n-1}(B'_n)$. To refine S , that is, to define it on a greater set, we have to build the next tower \mathcal{R}'_{n+1} and define S as for \mathcal{R}'_n . To do so and according to Definition 3.2, we have to determine a subdivision of the base B'_n into q'_n subsets $B'_{n,0}, \dots, B'_{n,q'_n-1}$. We find a function ζ_{n+1} mapping bimeasurably each $B'_{n,i}$ to $B'_{n,i+1}$ for $0 \leq i \leq q'_n - 2$. On the subset $D_{n+1} := \bigsqcup_{0 \leq i \leq q'_n-2} S^{h'_n-1}(B'_{n,i})$ of the roof $S^{h'_n-1}(B'_n)$ of \mathcal{R}'_n , S will coincide with $\zeta_{n+1}S^{-h'_n}$ and will be defined on $X \setminus S^{h'_{n+1}}(B'_{n+1}) = D_1 \sqcup \dots \sqcup D_{n+1}$, where $B'_{n+1} = B'_{n,0}$ is the base of the new Rokhlin tower \mathcal{R}'_{n+1} for S . To sum up, S is successively defined by the finite approximations obtained from the maps ζ_n . Up to conjugacy, ζ_n is exactly the addition by $(\underbrace{0, \dots, 0}_{n-1 \text{ times}}, 1, 0, 0, \dots)$ with carry over to the right

(as defined in §3.1), restricted to $[0, \dots, 0]_{n-1} \setminus [0, \dots, 0, q'_{n-1} - 1]_n$.

The construction of the maps ζ_n is by induction on $n \geq 0$. At step n , we will actually define

$$\zeta_{n+1} : B'_{n,0} \sqcup \dots \sqcup B'_{n,q'_n-2} \rightarrow B'_{n,1} \sqcup \dots \sqcup B'_{n,q'_n-1}.$$

To build ζ_{n+1} , a second induction on a parameter $m \geq n$ is required. Actually, $B'_{n,i}$ will be the disjoint union of the $B'_{n,i}(m)$ for $m \geq n$, and this inner recursion consists in choosing m bricks to define $B'_{n,i}(m)$. By definition, the m bricks will be the m levels (that is, the levels of \mathcal{R}_m) explicitly chosen to constitute $B'_{n,i}(m)$. Using powers of T , m bricks of $B'_{n,i}$ are mapped to those of $B'_{n,i+1}$ (there will be as many in $B_{n,i}$ as in $B'_{n,i+1}$) and this gives ζ_{n+1} whose orbits are included in those of T , implying immediately that the orbits of S satisfy the same property. The reverse inclusion between the orbits will be more difficult to prove and will be due to the choice of the bricks (see Remark 5.1 after the construction).

5.1.2. *The construction.* Here, T is a rank-one system in $\text{Aut}(X, \mu)$. We fix one of its cutting-and-stacking construction whose parameters are denoted as in Definition 3.2. Let $(p_n)_{n \geq 0}$ be a sequence of prime numbers such that every prime number appears infinitely many times.

In the following, we will assume that, given the cutting parameters of T , some positive integers q'_n and $t_{n,m}$ that we will introduce are well defined. In §5.3 (see Lemma 5.10), we will give conditions on the parameters of T for these quantities (and so the construction) to be well defined.

Step $n = 1$. We first build \mathcal{R}'_1 and ζ_1 by an induction over $m \geq 1$. We could denote by \mathcal{R}'_0 the trivial tower (X) with its base $B'_0 := X$. At the end of step $n = 1$, S is not yet defined on the roof of the tower \mathcal{R}'_1 , that is, on its highest level, which is a Rokhlin tower of S .

- $m = 1$: let $q'_0 > 0$ be the largest multiple of p_0 such that $q'_0 \leq q_0 - 1$. For every $0 \leq i \leq q'_0 - 1$, we define

$$B'_{0,i}(1) := T^i(B_1)$$

and

$$\zeta_1(1): \bigsqcup_{0 \leq i \leq q'_0 - 2} B'_{0,i}(1) \rightarrow \bigsqcup_{0 \leq i \leq q'_0 - 2} B'_{0,i+1}(1)$$

coinciding with T on its domain (hence, every subset $B'_{0,i}(1)$ is composed of a unique 1-brick $T^i(B_1)$).

Remark 7.1. At this step, we simply have to assume $q_0 > p_0$ for the integer q'_0 to be non-zero. However, for the well definition of other quantities at other steps, the conditions on the cutting parameters of T get increasingly more technical, which is the reason why we first assume that the parameters of T are chosen so that the positive quantities are well defined and we will then state the conditions in Lemma 5.10 (as mentioned before the beginning of the construction).

- $m > 1$: assume that the subsets $B'_{0,i}(M)$ have been built for every $1 \leq M \leq m - 1$ and $0 \leq i \leq q'_0 - 1$. Let

$$W_{1,m} := X \setminus \bigsqcup_{1 \leq M \leq m-1} \bigsqcup_{0 \leq i \leq q'_0 - 1} B'_{0,i}(M)$$

be the remaining piece of X at the end of the $(m - 1)$ th step (we could also define $W_{1,1} := X$).

Remark 7.2. Notice that m -levels are either contained in $W_{1,m}$ or disjoint from it since $X \setminus W_{1,m}$ is composed of M -levels for $1 \leq M \leq m - 1$ and the Rokhlin towers are nested. This will more generally hold true for $W_{n,m}$ with $n \geq 2$.

Let $r_{1,m}$ be the number of integers $j \in \llbracket 0, h_m - 1 \rrbracket$ such that $T^j(B_m) \subseteq W_{1,m}$, denoted by

$$0 \leq j_1^{(1,m)} < j_2^{(1,m)} < \dots < j_{r_{1,m}}^{(1,m)} < h_m.$$

Let $t_{1,m}$ be the positive integer (we assume that we can choose the cutting parameters of T for this integer to be positive, see Lemma 5.10) such that $q'_0 t_{1,m}$ is the largest multiple of q'_0 such that $q'_0 t_{1,m} < r_{1,m}$. The first $q'_0 t_{1,m}$ m -levels contained in $W_{1,m}$ are now used as m -bricks; they are split in q'_0 groups of $t_{1,m}$ m -bricks of the subsets $B'_{0,i}$ in the following way and the same will be done at steps $n > 1$ (the fact that the inequality $q'_0 t_{1,m} < r_{1,m}$ is strict and the way we make the q'_0 groups will guarantee an easy control of the cocycles, see Lemma 5.4 used for Lemmas 5.16 and 5.19). For every $0 \leq i \leq q'_0 - 1$, we define

$$B'_{0,i}(m) := \bigsqcup_{0 \leq t \leq t_{1,m}-1} T^{(j_{i+1+tq'_0}^{(1,m)})} (B_m)$$

and $\zeta_1(m)$ coinciding with $T^{(j_{i+2+ tq'_0}^{(1,m)}) - (j_{i+1+ tq'_0}^{(1,m)})}$ on $T^{(j_{i+1+ tq'_0}^{(1,m)})} (B_m)$ for every $0 \leq i \leq q'_0 - 2$ and $0 \leq t \leq t_{1,m} - 1$, so that each brick $T^{(j_{i+1+ tq'_0}^{(1,m)})} (B_m)$ is mapped on another $T^{(j_{i+2+ tq'_0}^{(1,m)})} (B_m)$. Thus, $\zeta_1(m)$ maps each $B'_{0,i}(m)$ on $B'_{0,i+1}(m)$ and this gives

$$\zeta_1(m) : \bigsqcup_{0 \leq i \leq q'_0-2} B'_{0,i}(m) \rightarrow \bigsqcup_{0 \leq i \leq q'_0-2} B'_{0,i+1}(m).$$

End of step $n = 1$. For every $0 \leq i \leq q'_0 - 1$, we define

$$B'_{0,i} := \bigsqcup_{m \geq 1} B'_{0,i}(m)$$

(the set of its m -bricks for all $m \geq 1$), $B'_1 := B'_{0,0}$ and

$$\zeta_1 : \bigsqcup_{0 \leq i \leq q'_0-2} B'_{0,i} \rightarrow \bigsqcup_{0 \leq i \leq q'_0-2} B'_{0,i+1}$$

coinciding with the maps $\zeta_1(m)$ on their respective domain (see Figure 7).

The universal odometer S we want to build is partially defined on X . More precisely, we define it on the domain $D_1 := \bigsqcup_{0 \leq i \leq q'_0-2} B'_{0,i}$ of ζ_1 so that it coincides with ζ_1 . This gives the first Rokhlin tower $\mathcal{R}'_1 := (B'_{0,0}, \dots, B'_{0,q'_0-1}) = (B'_1, S(B'_1), \dots, S^{q'_0-1}(B'_1))$. The next step will provide us with a refinement \mathcal{R}'_2 of the tower \mathcal{R}'_1 , allowing us to extend partially S on the highest level of the \mathcal{R}'_1 .

Step $n > 1$. Assume that steps $1, \dots, n - 1$ have been achieved. There are nested towers $\mathcal{R}'_1, \dots, \mathcal{R}'_{n-1}$. The k th tower \mathcal{R}'_k has $h'_k := q'_0 \dots q'_{k-1}$ levels and its base B'_k is partitioned in q'_k levels $B'_{k,0}, \dots, B'_{k,q'_k-1}$. These levels belong to \mathcal{R}'_{k+1} , whose base is $B'_{k+1} := B'_{k,0}$, with ζ_{k+1} mapping $B'_{k,i}$ to $B'_{k,i+1}$. The map S is defined on $D_1 \sqcup \dots \sqcup D_{n-1}$ using the maps $\zeta_1, \dots, \zeta_{n-1}$, where $D_1 \sqcup \dots \sqcup D_{n-1}$ corresponds to the union of all the levels of \mathcal{R}'_{n-1} except its roof.

The map S is not yet defined on the roof of \mathcal{R}'_{n-1} . By partitioning B'_{n-1} in subsets $B'_{n-1,0}, \dots, B'_{n-1,q'_{n-1}-1}$, we will define \mathcal{R}'_n which refines \mathcal{R}'_{n-1} and a function ζ_n




FIGURE 7. First step of the construction (that is, $n = 1$). In §5.4, we will define sets $E_{n,m}$ for every pair of integers (n, m) satisfying $m \geq n \geq 1$. The set $E_{1,1}$ (respectively $E_{1,2}; E_{1,3}$) is the union of the red areas (respectively red and blue areas; red, blue and green areas).

mapping $B'_{n-1,i}$ to $B'_{n-1,i+1}$. The extension of S will be defined on all the levels of \mathcal{R}'_n , except its roof (which is contained in the one of \mathcal{R}'_{n-1}). We will construct the subsets $B'_{n-1,i}$ as was done for the subsets $B'_{0,i}$, except that we only use the ‘material’ in B'_{n-1} to form the m -bricks of each $B'_{n-1,i}$. To do so, notice that the base B'_{n-1} is exactly $B'_{n-2,0}$ (the first subset in the subdivision of B'_{n-2}) which is the disjoint union of subsets of the form $B'_{n-2,0}(m)$ for $m \geq n - 1$. Moreover, for all $n - 1 \leq M \leq m$, every m -level is contained in an M -level, then we will pick the new m -bricks in $B'_{n-2,0}(n), \dots, B'_{n-2,0}(m)$. This motivates the definition of each set $W_{n,m}$ (the set of the remaining material to form m -bricks with an incremented integer m). We now discuss separately the following cases.

- $m = n$: set

$$W_{n,n} := B'_{n-2,0}(n - 1) \sqcup B'_{n-2,0}(n) \tag{5}$$

and let $r_{n,n}$ be the number of integers $j \in \llbracket 0, h_n - 1 \rrbracket$ such that $T^j(B_n) \subseteq W_{n,n}$ (note that we could have defined $r_{1,1} = q_0$), denoted by

$$0 \leq j_1^{(n,n)} < j_2^{(n,n)} < \dots < j_{r_{n,n}}^{(n,n)} < h_n.$$

Let q'_{n-1} be the largest multiple (we assume that we can choose the cutting parameters of T for this integer to be positive, see Lemma 5.10) of p_{n-1} such that $q'_{n-1} < r_{n,n}$. We then define for every $0 \leq i \leq q'_{n-1} - 1$,

$$B'_{n-1,i}(n) := T^{(j_{i+1}^{(n,n)})}(B_n),$$

meaning that among the n -levels in $W_{n,n}$, the n -bricks at step (n, n) are exactly the first q'_{n-1} ones (and set $t_{n,n} = 1$ for consistency later on). Let

$$\zeta_n(n) : \bigsqcup_{0 \leq i \leq q'_{n-1}-2} B'_{n-1,i}(n) \rightarrow \bigsqcup_{0 \leq i \leq q'_{n-1}-2} B'_{n-1,i+1}(n)$$

be the map coinciding with $T^{(j_{i+2}^{(n,n)})-(j_{i+1}^{(n,n)})}$ on each $B'_{n-1,i}(n)$, so that $B'_{n-1,i}(n)$ is mapped to $B'_{n-1,i+1}(n)$.

- $m > n$: set

$$W_{n,m} := \left(\bigsqcup_{n-1 \leq M \leq m} B'_{n-2,0}(M) \right) \setminus \left(\bigsqcup_{n \leq M \leq m-1} \bigsqcup_{0 \leq i \leq q'_{n-1}-1} B'_{n-1,i}(M) \right) \quad (6)$$

and let $r_{n,m}$ be the number of integers $j \in \llbracket 0, h_m - 1 \rrbracket$ such that $T^j(B_m) \subseteq W_{n,m}$, denoted by

$$0 \leq j_1^{(n,m)} < j_2^{(n,m)} < \dots < j_{r_{n,m}}^{(n,m)} < h_m.$$

Let $t_{n,m}$ be the positive integer (we assume that we can choose the cutting parameters of T for this integer to be positive, see Lemma 5.10) such that $q'_{n-1}t_{n,m}$ is the largest multiple of q'_{n-1} such that $q'_{n-1}t_{n,m} < r_{n,m}$. The first $q'_{n-1}t_{n,m}$ m -levels contained in $W_{n,m}$ are now used as m -bricks at step (n, m) ; they are split in q'_{n-1} groups of $t_{n,m}$ m -bricks of the subsets $B'_{n-1,i}$ in the following way. For every $0 \leq i \leq q'_{n-1} - 1$, we define

$$B'_{n-1,i}(m) := \bigsqcup_{0 \leq t \leq t_{n,m}-1} T^{(j_{i+1+tq'_{n-1}}^{(n,m)})}(B_m)$$

and $\zeta_n(m)$ coinciding with $T^{(j_{i+2+tq'_{n-1}}^{(n,m)})-(j_{i+1+tq'_{n-1}}^{(n,m)})}$ on $T^{(j_{i+1+tq'_{n-1}}^{(n,m)})}(B_m)$ for every $0 \leq i \leq q'_{n-1} - 2, 0 \leq t \leq t_{n,m} - 1$, so that each m -brick $T^{(j_{i+1+tq'_{n-1}}^{(n,m)})}(B_m)$ is mapped on another $T^{(j_{i+2+tq'_{n-1}}^{(n,m)})}(B_m)$. Thus, $\zeta_n(m)$ maps each $B'_{n-1,i}(m)$ on $B'_{n-1,i+1}(m)$ and this gives

$$\zeta_n(m) : \bigsqcup_{0 \leq i \leq q'_{n-1}-2} B'_{n-1,i}(m) \rightarrow \bigsqcup_{0 \leq i \leq q'_{n-1}-2} B'_{n-1,i+1}(m).$$

End of step n . We define for every $0 \leq i \leq q'_{n-1} - 1$,

$$B'_{n-1,i} := \bigsqcup_{m \geq n} B'_{n-1,i}(m)$$




FIGURE 8. From the first step (illustrated in Figure 7) to the second one. In B'_1 , we inductively build $B'_{1,i}(2), B'_{1,i}(2), B'_{1,i}(3), \dots$ for every $0 \leq i \leq q'_1 - 1$ (in this example, we have $q'_1 = 2$). Each set $B'_{1,i}(2)$ is composed of a unique 2-level in $B'_{0,0}(1) \sqcup B'_{0,0}(2)$ (that is, in the pale red and pale blue areas). Then each set $B'_{1,i}(3)$ is composed of 3-levels in $B'_{0,0}(1) \sqcup B'_{0,0}(2) \sqcup B'_{0,0}(3)$ (that is, in the pale red, pale blue and pale green areas) and so on. The structure that we build in $B'_1 = B'_{0,0}$ can be translated in $B'_{0,1}$ and $B'_{0,2}$ using the map ζ_1 . In §5.4, we will define sets $E_{n,m}$ for every pair of integers (n, m) satisfying $m \geq n \geq 1$. The set $E_{2,1}$ (respectively $E_{2,2}$) is the union of the areas hatched in blue (respectively in blue or green).

(the set of its m -bricks for $m \geq n$), $B'_n = B'_{n-1,0}$ and

$$\zeta_n : \bigsqcup_{0 \leq i \leq q'_{n-1}-2} B'_{n-1,i} \rightarrow \bigsqcup_{0 \leq i \leq q'_{n-1}-2} B'_{n-1,i+1}$$

coinciding with the maps $\zeta_n(m)$ on their respective domain (see Figure 8 for step $n = 2$, after the first step illustrated in Figure 7).

As the base B'_{n-1} of \mathcal{R}'_{n-1} is partitioned in $B'_{n-1,0} \sqcup \dots \sqcup B'_{n-1,q'_{n-1}-1}$, its highest level $S^{h'_{n-1}-1}(B'_{n-1})$ is partitioned in $S^{h'_{n-1}-1}(B'_{n-1,0}) \sqcup \dots \sqcup S^{h'_{n-1}-1}(B'_{n-1,q'_{n-1}-1})$. The map S is extended in the following way. On

$$D_n := S^{h'_{n-1}-1}(B'_{n-1,0}) \sqcup \dots \sqcup S^{h'_{n-1}-1}(B'_{n-1,q'_{n-1}-2}),$$

it coincides with $\zeta_n S^{-(h'_{n-1}-1)}$. So S maps $S^{h'_{n-1}-1}(B'_{n-1,i})$ on $B'_{n-1,i+1}$. This gives a Rokhlin tower \mathcal{R}'_n for S , nested in the previous one, of base $B'_n := B'_{n-1,0}$ and height $h'_n := q'_0 \dots q'_{n-1}$. Now S is defined on $(D_1 \sqcup \dots \sqcup D_{n-1}) \sqcup D_n$. The set D_n consists in the levels of \mathcal{R}'_n , except the highest one, which are contained in the highest level of \mathcal{R}'_{n-1} .

Remark 5.1. Notice that the inclusion of the S -orbits in the T -orbits is easy since S is defined from maps $\zeta_n(m)$ which are ‘piecewise powers of T ’.

The reverse inclusion will follow from the fact that we have $t_{n,n} = 1$ for every $n \geq 1$ (at step (n, n) , we form groups of only one n -level). Indeed, uniqueness implies that these chosen blocks are linked by $\zeta_n(n)$ and hence clearly by S (in contrast, an m -level, for

$m > n$, of $B'_{n-1,i}$ is mapped by $\zeta_n(m)$ to only one of the $t_{n,m}$ m -levels of $B'_{n-1,i+1}$, and not to the other). Thus, ensuring that the unique n -brick of each $B'_{n-1,i}$ is a large part of it that enables the system S to capture most of the T -orbits.

5.2. *First properties of this construction.* Recall that we consider a cutting-and-stacking construction of T with the same notation as in Definition 3.2 ($q_n, \sigma_{n,i}, \mathcal{R}_n, X_n, \varepsilon_n, \dots$), and the sequences $(h_n), (\sigma_n)$ and (Z_n) associated to the cutting and spacing parameters, and the notation $q'_n, \mathcal{R}'_n, \dots$ refer to the construction of S .

We state some important properties preparing for further results in §5.4. Many of them enable us to only take into account the combinatorics behind a cutting-and-stacking construction. We assume that all the ‘largest multiples’ (for every $n < m$, the largest multiple q'_{n-1} of p_{n-1} such that $q'_{n-1} < r_{n,n}$, and the largest multiple $q'_{n-1}t_{n,m}$ of q'_{n-1} such that $q'_{n-1}t_{n,m} < r_{n,m}$) are non-zero. In §5.3 (see Lemma 5.10), we will see how to choose the parameters for the construction to be well defined.

LEMMA 5.2. *Every tower \mathcal{R}'_n is a partition of X .*

Proof of Lemma 5.2. Let $n \geq 1$. The levels of \mathcal{R}'_n are pairwise disjoint by the definition of $(W_{n,m})_{m \geq n}$. It remains to show that \mathcal{R}'_n covers the whole space. Recall that X_n denotes the subset covered by the tower \mathcal{R}_n , and ε_n the measure of its complement.

The result holds for $n = 1$ since $\mu(W_{1,m}) \xrightarrow{m \rightarrow +\infty} 0$. Indeed, $W_{1,m+1} \cap X_m$ is the union of the m -levels which are not chosen at step $(1, m)$. By the definition of $t_{1,m}$, there are at most q'_0 . So we have $W_{1,m+1} \leq \varepsilon_m + q'_0\mu(B_m) \rightarrow 0$.

For $n > 1$, it suffices to show that the levels $B'_{n-1,0}, \dots, B'_{n-1,q'_{n-1}-1}$ of \mathcal{R}'_n form a partition of the base B'_{n-1} of \mathcal{R}'_{n-1} . We have to show that the measure of

$$\tilde{W}_{n,m} := B'_{n-1} \setminus \left(\bigsqcup_{n \leq M \leq m-1} \bigsqcup_{0 \leq i \leq q'_{n-1}-1} B'_{n-1,i}(M) \right)$$

tends to 0 as $m \rightarrow +\infty$. However, this set $\tilde{W}_{n,m}$ is the disjoint union of $\bigsqcup_{M \geq m+1} B'_{n-2,0}(M)$ and $W_{n,m}$. It is clear that

$$\mu \left(\bigsqcup_{M \geq m+1} B'_{n-2,0}(M) \right) \xrightarrow{m \rightarrow +\infty} 0$$

since μ is a finite measure. The set $W_{n,m}$ is obtained from $W_{n,m-1}$ by adding $B'_{n-2,0}(m)$ and removing $q'_{n-1}t_{n,m-1}$ $(m - 1)$ -levels. Thus, we have $\mu(W_{n,m}) \xrightarrow{m \rightarrow +\infty} 0$ by the definition of $(t_{n,m})_{m \geq n}$. Hence, we have $\mu(\tilde{W}_{n,m}) \xrightarrow{m \rightarrow +\infty} 0$ and we are done. □

As a consequence, if $(\mathcal{R}'_n)_n$ increases to the σ -algebra \mathcal{A} (this will be proved in Corollary 5.14), then S is a rank-one system without spacer, so this is an odometer.

LEMMA 5.3. *Let $n \geq 1$. On the base B'_n of the n th S -Rokhlin tower \mathcal{R}'_n , S is defined as follows. For every $0 \leq i \leq h'_n - 1$, we have*

$$S^i = \zeta_1^{i_0} \dots \zeta_n^{i_{n-1}} \quad \text{on } B'_n$$

with $i_0 \in \llbracket 0, q'_0 - 1 \rrbracket, \dots, i_{n-1} \in \llbracket 0, q'_{n-1} - 1 \rrbracket$ such that

$$i = \sum_{\ell=0}^{n-1} q'_0 \cdots q'_{\ell-1} i_\ell = \sum_{\ell=0}^{n-1} h'_\ell i_\ell.$$

Proof of Lemma 5.3. By induction over $n \geq 1$. It is clear for $n = 1$ since S coincides with ζ_1 on the levels of \mathcal{R}'_1 except its roof. Assume that the result holds for $n \geq 1$. The tower \mathcal{R}'_n is divided in q'_n subcolumns whose levels are exactly those of \mathcal{R}'_{n+1} , and the i_n th subcolumn ($0 \leq i_n \leq h'_n - 1$) is the S -Rokhlin tower of height h'_n and base B'_{n,i_n} . Let $0 \leq i \leq h'_{n+1} - 1$. By the equality $B'_{n+1} = B'_{n,0}$ and by the definition of S from ζ_{n+1} (at the end of step $n + 1$ of the construction), $S^i = S^j \zeta_{n+1}^{i_n}$ on B'_{n+1} for non-negative integers i_n and j such that $i = i_n h'_n + j$ and $j < h'_n$. The set $\zeta_{n+1}^{i_n}(B'_{n,0})$ is equal to B'_{n,i_n} , so this is a subset of B'_n ; hence the result by the induction hypothesis. \square

Therefore, the subset D_n defined in the construction can be written as follows:

$$\begin{aligned} D_n &= \zeta_1^{q'_0-1} \cdots \zeta_{n-1}^{q'_{n-2}-1} \left(\bigsqcup_{0 \leq i_n \leq q'_{n-1}-2} B'_{n-1,i_n} \right) \\ &= \zeta_1^{q'_0-1} \cdots \zeta_{n-1}^{q'_{n-2}-1} \left(\bigsqcup_{0 \leq i_n \leq q'_{n-1}-2} \zeta_n^{i_n}(B'_{n-1,0}) \right) \end{aligned} \tag{7}$$

and S coincides with $\zeta_n \zeta_{n-1}^{-(q'_{n-2}-1)} \cdots \zeta_1^{-(q'_0-1)}$ on D_n .

By the cocycle of $\zeta_n(m)$, we mean the integer-valued map defined on the domain of $\zeta_n(m)$ and which maps x to the unique integer k satisfying $\zeta_n(m)x = T^k x$.

LEMMA 5.4. *The cocycle of $\zeta_n(m)$ is positive and bounded above by $h_{m-1} + Z_{m-1}$.*

Proof of Lemma 5.4. By definition, for fixed integers $0 \leq i \leq q'_{n-1} - 2$ and $0 \leq t \leq t_{n,m} - 1$, the cocycle on $\mathbb{B} := T^{(j_{i+1+tq'_{n-1}})^{(n,m)}}(B_m)$ takes the value $\Delta j := j_{i+2+tq'_{n-1}}^{(n,m)} - j_{i+1+tq'_{n-1}}^{(n,m)}$. Let us recall that the integers

$$0 \leq j_1^{(n,m)} < j_2^{(n,m)} < \cdots < j_{r_{n,m}}^{(n,m)} < h_m$$

are the sets of indices $j \in \llbracket 0, h_m - 1 \rrbracket$ such that $T^j(B_m) \subseteq W_{n,m}$. Thus, Δj is obviously positive. Let us fix an $(m - 1)$ -level \mathbb{B}^* which is not chosen at step $(n, m - 1)$, so it is contained in $W_{n,m}$. If m is equal to n , we can choose $\mathbb{B}^* = B'_{n-2,0}(n - 1)$. For $m > n$, the existence of \mathbb{B}^* is granted by the fact that we have $q'_{n-1} t_{n,m-1} < r_{n,m-1}$. We write it as $\mathbb{B}^* = T^{k_0}(B_{m-1})$, where k_0 is an integer in $\llbracket 0, h_{m-1} - 1 \rrbracket$.

By definition, Δj is the least positive integer j such that $T^j(\mathbb{B})$ is in $W_{n,m}$. Moreover, the m -levels of \mathbb{B}^* are in $W_{n,m}$. Therefore, the consecutive m -levels $T(\mathbb{B}), \dots, T^{\Delta j-1}(\mathbb{B})$ are not in \mathbb{B}^* .

First case. In the tower \mathcal{R}_m , assume that the m -levels $T(\mathbb{B}), \dots, T^{\Delta j-1}(\mathbb{B})$ are before $T^{k_0}(B_{m-1,0})$, that is, before the first m -level of \mathbb{B}^* . Therefore, the enumeration $\mathbb{B}, \dots, T^{\Delta j}(\mathbb{B})$ is included in the enumeration

$$\Sigma_{m-1,0,1}, \dots, \Sigma_{m-1,0,\sigma_{m-1,0}}, B_{m-1,0}, \dots, T^{k_0}(B_{m-1,0}),$$

implying that $\Delta j \leq \sigma_{m-1,0} + k_0 \leq Z_{m-1} + h_{m-1}$.

Second case. Now assume that $T(\mathbb{B}), \dots, T^{\Delta j-1}(\mathbb{B})$ are after $T^{k_0}(B_{m-1,q_{m-1}-1})$, that is, after the last m -level of \mathbb{B}^* . Therefore, the enumeration $\mathbb{B}, \dots, T^{\Delta j}(\mathbb{B})$ is included in the enumeration

$$T^{k_0}(B_{m-1,q_{m-1}-1}), \dots, T^{h_{m-1}-1}(B_{m-1,q_{m-1}-1}), \Sigma_{m-1,q_{m-1},1}, \dots, \Sigma_{m-1,q_{m-1},\sigma_{m-1,q_{m-1}}},$$

and we get $\Delta j \leq (h_{m-1} - k_0 - 1) + \sigma_{m-1,q_{m-1}} \leq h_{m-1} + Z_{m-1}$.

Third case. Finally, if $T(\mathbb{B}), \dots, T^{\Delta j-1}(\mathbb{B})$ are between $T^{k_0}(B_{m-1,i})$ and $T^{k_0}(B_{m-1,i+1})$ for some $0 \leq i \leq q_{m-1} - 2$, that is, between two consecutive m -levels of \mathbb{B}^* , then the enumeration $\mathbb{B}, \dots, T^{\Delta j}(\mathbb{B})$ is included in the enumeration

$$T^{k_0}(B_{m-1,i}), \dots, T^{h_{m-1}-1}(B_{m-1,i}), \Sigma_{m-1,i,1}, \dots, \Sigma_{m-1,i,\sigma_{m-1,i}}, \\ B_{m-1,i+1}, \dots, T^{k_0}(B_{m-1,i+1}),$$

which gives $\Delta j \leq (h_{m-1} - k_0 - 1) + \sigma_{m-1,i} + (k_0 + 1) \leq h_{m-1} + Z_{m-1}$. □

LEMMA 5.5. *An m -brick at step n is included in an M -brick at step $n - 1$ for some $n - 1 \leq M \leq m$.*

Proof of Lemma 5.5. This follows directly from the definition of $W_{n,m}$ in the construction (see §5.1). Indeed, the ‘ (M) ’ in ‘ $B'_{n-2,0}(M)$ ’ means that we only consider the M -bricks, at step $n - 1$, composing $B'_{n-2,0}$. □

We now present a combinatorial description of the construction.

LEMMA 5.6. *The quantities $r_{n,m}, q_n, q'_n, t_{n,m}, \sigma_n$ satisfy the following recurrence relation:*

$$\begin{aligned} & t_{0,1} := 0; \\ & \text{for } m \geq 2, \quad t_{0,m} := \sigma_{m-1}; \\ & \text{for } m = n \geq 1, \quad \begin{cases} r_{n,n} = q_{n-1} + t_{n-1,n}, \\ q'_{n-1} = \left\lfloor \frac{r_{n,n} - 1}{p_{n-1}} \right\rfloor p_{n-1}, \\ t_{n,n} = 1; \end{cases} \\ & \text{for } m > n \geq 1, \quad \begin{cases} r_{n,m} = q_{m-1}(r_{n,m-1} - q'_{n-1}t_{n,m-1}) + t_{n-1,m}, \\ t_{n,m} = \left\lfloor \frac{r_{n,m} - 1}{q'_{n-1}} \right\rfloor. \end{cases} \end{aligned}$$

During the construction, some integers have been defined for consistency ($r_{1,1} := q_0, t_{n,n} := 1$). Note that in this lemma, we also define the integers $t_{n,m}$ for $n = 0$. This enables us to extend the relations

$$r_{n,n} = q_{n-1} + t_{n-1,n} \quad \text{and} \quad r_{n,m} = q_{m-1}(r_{n,m-1} - q'_{n-1}t_{n,m-1}) + t_{n-1,m}$$

for $n = 1$.

Proof of Lemma 5.6. Case $n = 1$. For $m = 1$, the $r_{1,1}$ 1-levels potentially chosen to be 1-bricks are exactly the levels of \mathcal{R}_1 , so we have $r_{1,1} = q_0 + t_{0,1}$ since $t_{0,1} := 0$. We choose q'_0 of them, where q'_0 is the largest multiple of p_0 such that $q'_0 < r_{1,1}$, so q'_0 is equal to $\lfloor (r_{1,1} - 1)/p_0 \rfloor p_0$. Finally, q'_0 is obviously equal to $q'_0 t_{1,1}$ since $t_{1,1} := 1$. For $m > 1$, there are $r_{n,m}$ m -levels in $W_{1,m}$: some of them are in the $r_{1,m-1} - q'_0 t_{1,m-1}$ $(m - 1)$ -levels which are not chosen at step $(1, m - 1)$ and the other are the spacers from \mathcal{R}_{m-1} to \mathcal{R}_m . So, we have

$$r_{1,m} = q_{m-1}(r_{1,m-1} - q'_0 t_{1,m-1}) + \sigma_{m-1}$$

and we set $t_{0,m} := \sigma_{m-1}$. We choose $q'_0 t_{1,m}$ of them as m -bricks, where $q'_0 t_{1,m}$ is the largest multiple of q'_0 such that $q'_0 t_{1,m} < r_{1,m}$, that is, $t_{1,m} := \lfloor (r_{1,m} - 1)/q'_0 \rfloor$.

Case $n > 1$. For $m = n$, there are $r_{n,n}$ n -levels in $W_{n,n} = B'_{n-2,0}(n - 1) \sqcup B'_{n-2,0}(n)$. First, since we have $t_{n-1,n-1} = 1$, the set $B'_{n-2,0}(n - 1)$ is an $(n - 1)$ -brick at step $n - 1$ and it contains q_{n-1} n -levels. Second, $B'_{n-2,0}(n)$ is the union of $t_{n-1,n}$ n -bricks. Hence, we have $r_{n,n} = q_{n-1} + t_{n-1,n}$. By definition, q'_{n-1} is equal to $\lfloor (r_{n,n} - 1)/p_{n-1} \rfloor p_{n-1}$ and obviously to $q'_{n-1} t_{n,n}$ with $t_{n,n} := 1$. For $m > n$, there are $r_{n,m}$ m -levels in $W_{n,m}$. This set is composed of

$$\left(\bigsqcup_{n-1 \leq M \leq m-1} B'_{n-2,0}(M) \right) \setminus \left(\bigsqcup_{n \leq M \leq m-1} \bigsqcup_{0 \leq i \leq q'_{n-1}-1} B'_{n-1,i}(M) \right)$$

and

$$B'_{n-2,0}(m).$$

The first one is the union of the $r_{n,m-1} - q'_{n-1} t_{n,m-1}$ $(m - 1)$ -levels which are not chosen at step $(n, m - 1)$, and the second one is built at step $(n - 1, m)$ from its $t_{n-1,m}$ m -bricks. So, we have

$$r_{n,m} = q_{m-1}(r_{n,m-1} - q'_{n-1} t_{n,m-1}) + t_{n-1,m}.$$

We choose $q'_{n-1} t_{n,m}$ of these m -levels as m -bricks at this step, where $q'_{n-1} t_{n,m}$ is the largest multiple of q'_{n-1} such that $q'_{n-1} t_{n,m} < r_{n,m}$, that is, $t_{n,m} := \lfloor (r_{n,m} - 1)/q'_{n-1} \rfloor$. \square

It will be more convenient to use the following slight modification of Lemma 5.6:

$$\begin{aligned} & t_{0,1} = 0; \\ \text{for } m \geq 2, & \quad t_{0,m} = \sigma_{m-1}; \\ \text{for } m = n \geq 1, & \quad \begin{cases} r_{n,n} = q_{n-1} + t_{n-1,n}, \\ q'_{n-1} \leq r_{n,n} - 1, \\ t_{n,n} = 1; \end{cases} \\ \text{for } m > n \geq 1, & \quad \begin{cases} r_{n,m} \leq q_{m-1} q'_{n-1} + t_{n-1,m}, \\ t_{n,m} \leq \frac{r_{n,m} - 1}{q'_{n-1}}. \end{cases} \end{aligned} \tag{8}$$

This is a consequence of the inequalities $\lfloor x \rfloor \leq x$ and $r_{n,m-1} - q'_{n-1}t_{n,m-1} \leq q'_{n-1}$ (by the definition of $t_{n,m-1}$).

As the strategy will be to recursively choose large enough cutting parameters q_n for T , we would like to understand the asymptotic behaviour of q'_n as q_n increases. Then, the goal is to find bounds for q'_n/q_n .

LEMMA 5.7. *For every $n \geq 0$, we have*

$$q'_n \geq q_n - (1 + p_n).$$

Proof of Lemma 5.7. Using the equalities $q'_n = \lfloor r_{n+1,n+1} - 1/p_n \rfloor p_n$ and $r_{n+1,n+1} = q_n + t_{n,n+1}$ in Lemma 5.6, where the integer $t_{n,n+1}$ is non-negative, we get

$$q'_n \geq \left(\frac{r_{n+1,n+1} - 1}{p_n} - 1 \right) p_n \geq q_n - 1 - p_n$$

and we are done. □

We have found a lower bound for q'_n/q_n (up to some additional term $-(1 + p_n)$). Let us find an upper bound.

LEMMA 5.8. *For every $n \geq 1$, we have*

$$q'_n \leq 3q_n + \frac{\sigma_n}{q'_0 \cdots q'_{n-1}}.$$

With an asymptotic control on σ_n , using flexible classes, we will be able to get $q'_n \leq 4q_n$ (see Lemma 5.12).

Proof of Lemma 5.8. By induction over $i \in \llbracket 0, n - 1 \rrbracket$ (with $n \geq 1$) and using equation (8), we show that

$$q'_n \leq q_n \left(2 + \sum_{j=1}^i \prod_{k=1}^j \frac{1}{q'_{n-k}} \right) + t_{n-1-i,n+1} \prod_{k=1}^{i+1} \frac{1}{q'_{n-k}}.$$

For $i = 0$, we have $q'_n < r_{n+1,n+1} = q_n + t_{n,n+1}$ and

$$t_{n,n+1} \leq \frac{r_{n,n+1} - 1}{q'_{n-1}} \leq \frac{1}{q'_{n-1}} (q_n q'_{n-1} + t_{n-1,n+1}) = q_n + \frac{t_{n-1,n+1}}{q'_{n-1}},$$

so we get $q'_n \leq 2q_n + (t_{n-1,n+1}/q'_{n-1})$. For $0 \leq i \leq n - 2$, we have

$$t_{n-1-i,n+1} \leq \frac{r_{n-1-i,n+1} - 1}{q'_{n-2-i}} \leq \frac{1}{q'_{n-2-i}} (q_n q'_{n-2-i} + t_{n-2-i,n+1}) = q_n + \frac{t_{n-2-i,n+1}}{q'_{n-2-i}}.$$

If the result holds true for i , we get

$$\begin{aligned} q'_n &\leq q_n \left(2 + \sum_{j=1}^i \prod_{k=1}^j \frac{1}{q'_{n-k}} \right) + \left(q_n + \frac{t_{n-2-i,n+1}}{q'_{n-2-i}} \right) \prod_{k=1}^{i+1} \frac{1}{q'_{n-k}} \\ &= q_n \left(2 + \sum_{j=1}^{i+1} \prod_{k=1}^j \frac{1}{q'_{n-k}} \right) + t_{n-1-(i+1),n} \prod_{k=1}^{i+2} \frac{1}{q'_{n-k}}, \end{aligned}$$

so the result is also true for $i + 1$.

Taking $i = n - 1$, this gives the lemma since $q'_\ell \geq 2$ for every integer $\ell \geq 1$. □

5.3. *Towards flexible classes.* We now explain why flexible classes fit in this construction.

First, a condition for the construction to be well defined needs an inductive choice of the cutting parameters $(q_n)_{n \geq 0}$ of T (see Lemma 5.9). Second, a control on the spacing parameters will imply useful asymptotic controls for the quantification of the cocycles (see Lemma 5.12). Note that in the proof of Theorem 3.9 (see §5.5), we will need other estimates to quantify the cocycles. It will be possible, again using the definition of a flexible class, to inductively build large enough cutting parameters to have these estimates.

If the parameters are chosen according to a set $\mathcal{F}_C \subseteq \mathcal{P}^*$ associated to a flexible class \mathcal{C} , the underlying rank-one system has the desired property, that is, it is in \mathcal{C} , and is orbit equivalent to the universal odometer, with some quantification guaranteed by the control of the spacing parameters and by the fact that the cutting parameters q_n have been recursively chosen and large enough.

LEMMA 5.9. *Let T be a rank-one system with cutting and spacing parameters*

$$(q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_{n \geq 0}$$

such that the construction in §5.1 is well defined. Then, for every $n \in \mathbb{N}$, q'_n only depends on $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n}$.

Proof of Lemma 5.9. This directly follows from Lemma 5.6. □

Then, the main novelty in this paper is to build the rank-one system T while we are building the universal odometer S . Once (q'_0, \dots, q'_n) has been built from $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n}$, we are free to choose $(q_{n+1}, (\sigma_{n+1,0}, \dots, \sigma_{n+1,q_{n+1}}))$ for the definition of q'_{n+1} . The recursive definition of the cutting parameters is one of the main ideas behind the definition of a flexible class, and it allows to find cutting parameters satisfying some assumptions, for example, the assumptions of the following lemma.

LEMMA 5.10. *Assume that for every $n \in \mathbb{N}$,*

$$q_n > \max(p_n, q'_0, \dots, q'_{n-1}). \tag{9}$$

Then, the construction is well defined, that is, all the ‘largest multiples’ are non-zero (that is, the largest multiple q'_{n-1} of p_{n-1} such that $q'_{n-1} < r_{n,n}$, and the largest multiple $q'_{n-1} t_{n,m}$ of q'_{n-1} such that $q'_{n-1} t_{n,m} < r_{n,m}$).

Remark 5.11. Without loss of generality, we can assume that p_0 is equal to 2. Therefore, the assumption of Lemma 5.10 for $n = 0$ requires that q_0 is greater than 2, which explains the second item of Definition 3.7.

Proof of Lemma 5.10. First, let us prove this result at step $n = 1$ of the outer recursion. At step $m = 1$ of the inner recursion, q_0 is greater than p_0 , so q'_0 (the largest multiple of p_0 such that $q'_0 \leq q_0 - 1$) is positive. For a step $m > 1$, notice that there exists an $(m - 1)$ -level which is not chosen at the previous step (as we have $r_{1,m-1}$ $(m - 1)$ -levels in $W_{1,m-1}$ and we choose $q'_0 t_{1,m-1}$ of them, with $q'_0 t_{1,m-1} < r_{1,m-1}$) so its q_{m-1} m -levels are in $W_{1,m}$ and this gives $r_{1,m} \geq q_{m-1}$. Therefore, we have $r_{1,m} > q'_0$ and $t_{1,m}$ is non-zero.

Now consider a step $n > 1$ of the outer recursion. For $m = n$, $B'_{n-2,0}(n - 1)$ is an $(n - 1)$ -level in $W_{n,n}$, so we have $r_{n,n} \geq q_{n-1} > p_{n-1}$, and hence the positivity of q'_{n-1} . For $m > n$, we have $r_{n,m} \geq q_{m-1}$ (same argument as for $n = 1$), this implies $r_{n,m} > q'_{n-1}$ and $t_{n,m}$ is positive. □

The next lemma refines the estimate given by Lemma 5.8, with assumptions which will be satisfied in the context of flexible classes.

LEMMA 5.12. *Let $(q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_{n \geq 0}$ be the parameters of a CSP construction of T with associated constant $C > 0$. Assume that there exists a constant $C' > 0$ such that*

$$\text{for all } n \geq 1, \quad \sigma_n \leq C' q_n h_{n-1} \quad \text{and} \quad q_n - (1 + p_n) \geq C' h_n$$

(for instance, if the third point of Definition 3.7 holds and if, given $(q_k, (\sigma_{0,k}, \dots, \sigma_{q_k,k}))_{0 \leq k \leq n-1}$, q_n is chosen large enough). Then, we get the following bound:

$$\text{for all } n \in \mathbb{N}, \quad \frac{q'_n}{q_n} \leq 4.$$

Proof of Lemma 5.12. For $n = 0$, this is a consequence of the inequality $q'_0 \leq q_0 - 1$. Now let us prove the result for $n \geq 1$. Using Lemma 5.8, it suffices to get

$$\text{for all } n \geq 1, \quad \frac{\sigma_n}{q'_0 \cdots q'_{n-1}} \leq q_n.$$

However, we have

$$\sigma_n \leq C' q_n h_{n-1} \leq q_n (q_{n-1} - (1 + p_{n-1})),$$

and the right-hand side is bounded above by $q_n q'_{n-1}$ (by Lemma 5.7), so the result follows from the inequality $q'_{n-1} \leq q'_0 \cdots q'_{n-1}$. □

5.4. *Equality of the orbits, universal odometer and quantitative control of the cocycles.* Recall the notation for the construction of T by cutting and stacking, where $(q_n)_n$ and $(\sigma_{n,i})_{n,i}$ are respectively the cutting and spacing parameters. The tower \mathcal{R}_n is the n th T -Rokhlin tower, its height is h_n , it covers the subset X_n of X , ε_n is the measure of its complement, Z_n is the maximum of the spacing parameters over the first n steps and M_0 is the measure of the unique 0-level B_0 .

We use similar notation q'_n, h'_n and \mathcal{R}'_n for S . We also set

$$H'_n := h'_1 + \dots + h'_n$$

for all $n \geq 1$, and $H'_0 := 0$.

The construction is assumed to be well defined, considering a cutting-and-stacking definition of T with parameters satisfying the criterion (9) (see Lemma 5.10). Since S is piecewise given by powers of T , the S -orbits are included in the T -orbits. It remains to show the reverse inclusion, to prove that $(\mathcal{R}'_n)_{n \geq 0}$ is increasing to the σ -algebra \mathcal{A} and to quantify the cocycles.

As in [KL24], we set

$$E_{n,m} := \bigsqcup_{i=0}^{h'_n-1} S^i \left(\bigsqcup_{n \leq M \leq m} B'_{n-1,0}(M) \right) = \bigsqcup_{\substack{0 \leq i_0 \leq q'_0-1 \\ \vdots \\ 0 \leq i_{n-1} \leq q'_{n-1}-1}} \zeta_1^{i_0} \dots \zeta_n^{i_{n-1}} \left(\bigsqcup_{n \leq M \leq m} B'_{n-1,0}(M) \right)$$

and

$$K_n := \bigsqcup_{\substack{0 < i \leq h_n-1 \\ T^{i-1}(B_n) \sqcup T^i(B_n) \subseteq E_{n,n}}} T^i(B_n).$$

Since $B'_{n-1,0}$ is exactly the base B'_n of \mathcal{R}'_n , the subsets $S^i(B'_{n-1,0})$, for $0 \leq i \leq h'_n - 1$, are exactly the levels of \mathcal{R}'_n which is a partition of X . So the motivation behind the definition of $E_{n,m}$ is first to approximate $B'_{n-1,0}$ by its M -bricks for $n \leq M \leq m$, and then the set $E_{n,m}$ is actually the union of the M -bricks, for $n \leq M \leq m$, of step n of the outer recursion, and their translates by S in the other levels in \mathcal{R}'_n (the sets $E_{1,1}, E_{1,2}, E_{1,3}, E_{2,1}$ and $E_{2,2}$ are illustrated in Figures 7 and 8). We get a better approximation of X as m increases and notice that $E_{n,m}$ is a subset of X_m since every M -brick, for $n \leq M \leq m$, is a union of m -levels. Finally, the sets K_n , for $n \geq 1$, are introduced to show that the system S captures the T -orbits (recall Remark 5.1).

LEMMA 5.13. *The following holds:*

$$\mu(X_m \setminus E_{n,m}) \leq \begin{cases} \frac{H'_n}{h_m} & \text{for } n < m, \\ \frac{H'_{n-1} + p_{n-1}h'_{n-1}}{h_n} & \text{for } n = m. \end{cases}$$

Proof of Lemma 5.13. We prove the inclusions

$$E_{n,m} \subset E_{n-1,m} \subset \dots \subset E_{2,m} \subset E_{1,m} \subset X_m,$$

and we bound the measures of $X_m \setminus E_{1,m}$ and each set $E_{k,m} \setminus E_{k-1,m}$. The result follows from the decomposition

$$X_m \setminus E_{n,m} = (X_m \setminus E_{1,m}) \sqcup \bigsqcup_{2 \leq k \leq n} (E_{k-1,m} \setminus E_{k,m}) \tag{10}$$

and σ -additivity of μ .

The set $E_{1,m}$ is composed of m -levels, so it is contained in X_m . If $m = 1$, then $X_m \setminus E_{1,m}$ is the disjoint union of $r_{1,1} - q'_0$ 1-levels (see step (1, 1) of the construction). If $m > 1$, then $X_m \setminus E_{1,m}$ is the disjoint union of $r_{1,m} - q'_0 t_{1,m}$ m -levels (see step (1, m) of the construction). By definition of q'_0 (if $m = 1$) or $t_{1,m}$ (if $m > 1$), we thus have

$$\mu(X_m \setminus E_{1,m}) \leq \begin{cases} \frac{p_0}{h_m} & \text{if } m = 1, \\ \frac{h'_1}{h_m} & \text{if } m > 1 \end{cases}$$

(recall that $h'_1 = q'_0$).

Let $k \in \llbracket 2, n \rrbracket$. The function ζ_k has been built to map each M -brick ($M \geq k$) at step k to another. However, such a brick is contained in an M' -brick ($k - 1 \leq M' \leq M$) from the previous step $k - 1$ (see Lemma 5.5). We then have

$$\bigsqcup_{0 \leq i_{k-1} \leq q'_{k-1}-1} \zeta_k^{i_{k-1}} \left(\bigsqcup_{k \leq M \leq m} B'_{k-1,0}(M) \right) \subseteq \bigsqcup_{k-1 \leq M \leq m} B'_{k-2,0}(M).$$

Applying $\zeta_1^{i_0} \dots \zeta_{k-1}^{i_{k-2}}$ and considering the union over i_0, \dots, i_{k-2} , we get the inclusion $E_{k,m} \subseteq E_{k-1,m}$ and the equality

$$\begin{aligned} & E_{k-1,m} \setminus E_{k,m} \\ &= \bigsqcup_{\substack{0 \leq i_0 \leq q'_0-1 \\ \vdots \\ 0 \leq i_{k-2} \leq q'_{k-2}-1}} \zeta_1^{i_0} \dots \zeta_{k-1}^{i_{k-2}} \left(\underbrace{\left(\bigsqcup_{k-1 \leq M \leq m} B'_{k-2,0}(M) \right) \setminus \left(\bigsqcup_{k \leq M \leq m} \bigsqcup_{0 \leq i_{k-1} \leq q'_{k-1}-1} B'_{k-1,i_k}(M) \right)}_{=: [*]} \right). \end{aligned}$$

So the measure of $E_{k-1,m} \setminus E_{k,m}$ is $q'_0 \dots q'_{k-2} \mu([*]) = h'_{k-1} \mu([*])$ by T -invariance. The set $[*]$ is obtained from $W_{k,m}$ (see equations (5) and (6)) by removing the m -bricks that have been chosen at step (k, m) . If $m = k$, then $[*]$ is the disjoint union of $r_{k,k} - q'_{k-1}$ m -levels (see step (k, k) of the construction). If $m > k$, then $[*]$ is the disjoint union of $r_{k,m} - q'_{k-1} t_{k,m}$ m -levels (see step (k, m) of the construction). By definition of q'_{k-1} (if $m = k$) or $t_{k,m}$ (if $m > k$), we thus have

$$\mu([*]) \leq \begin{cases} \frac{p_{k-1}}{h_m} & \text{if } m = k, \\ \frac{q'_{k-1}}{h_m} & \text{if } m > k, \end{cases}$$

and

$$\mu(E_{k-1,m} \setminus E_{k,m}) \leq \begin{cases} \frac{h'_{k-1} p_{k-1}}{h_m} & \text{if } m = k, \\ \frac{h'_k}{h_m} & \text{if } m > k. \end{cases}$$

Using equation (10) and σ -additivity of μ , we get the following inequalities. If $m > n$, we get

$$\mu(X_m \setminus E_{n,m}) = \mu(X_m \setminus E_{1,m}) + \sum_{2 \leq k \leq n} \mu(E_{k-1,m} \setminus E_{k,m}) \leq \sum_{1 \leq k \leq n} \frac{h'_k}{h_m} = \frac{H'_n}{h_m}.$$

If $m = n$, we get

$$\begin{aligned} \mu(X_m \setminus E_{n,m}) &= (\mu(X_m \setminus E_{1,m}) + \sum_{2 \leq k \leq m-1} \mu(E_{k-1,m} \setminus E_{k,m})) + \mu(E_{n-1,n} \setminus E_{n,n}) \\ &\leq \sum_{1 \leq k \leq m-1} \frac{h'_k}{h_m} + \frac{p_{n-1}h'_{n-1}}{h_n} \\ &= \frac{H'_{n-1}}{h_n} + \frac{p_{n-1}h'_{n-1}}{h_m} \end{aligned}$$

and we are done. □

The quantity $H'_{n-1} + p_{n-1}h'_{n-1}$ only depends on q'_1, \dots, q'_{n-2} which only depend on $(q_i, (\sigma_{i,j})_{0 \leq j \leq q_i})_{0 \leq i \leq n-2}$ (see Lemma 5.9), and h_n is larger than $q_1 \dots q_{n-1}/M_0$ with q_{n-1} appearing at step $n - 1$. Then, the strategy will be to recursively choose the cutting parameters q_{n-1} so that

$$\frac{H'_{n-1} + p_{n-1}h'_{n-1}}{h_n} \xrightarrow{n \rightarrow +\infty} 0. \tag{11}$$

As $\mu(X_n) \xrightarrow{n \rightarrow +\infty} 1$, this gives $\mu(E_{n,n}) \xrightarrow{n \rightarrow +\infty} 1$ by Lemma 5.13.

COROLLARY 5.14. *If $\mu(E_{n,n}) \xrightarrow{n \rightarrow +\infty} 1$, then S is the universal odometer.*

Proof of Corollary 5.14. By the definition of q'_n at step (n, n) and by choice of the sequence (p_n) , every prime number appears infinitely many times as a prime factor among the integers q'_0, q'_1, q'_2, \dots . If S is an odometer, then it is clearly universal. It remains to show that $(\mathcal{R}'_n)_{n \in \mathbb{N}}$ increases to the σ -algebra \mathcal{A} . Then, S is a rank-one system with zero spacing parameters by Lemma 5.2, so this is an odometer.

Consider a subsequence $(n_k)_{k \geq 0}$ such that the series $\sum_{k \geq 0} \mu((E_{n_k, n_k})^c)$ is convergent. By the Borel–Cantelli lemma, the set $X_0 := \bigcup_{j \geq 0} \bigcap_{k \geq j} E_{n_k, n_k}$ is of full measure. Let $x, y \in X_0$. Assume that they belong to the same level of \mathcal{R}'_n for every n larger than some threshold N_0 . The goal is to show that x and y are equal, so that $(\mathcal{R}'_n)_{n \in \mathbb{N}}$ separates the points of a set of full measure and hence it increases to \mathcal{A} .

By the definition of X_0 , there exists an infinite subset I of \mathbb{N} , bounded below by N_0 , such that $E_{n,n}$ contains x and y for every $n \in I$. Let us fix an integer $n \in I$. By the definition of $E_{n,n}$, x is in some $S^i(B'_{n-1,0}(n))$ and y in some $S^j(B'_{n-1,0}(n))$, for $0 \leq i, j \leq q'_0 \dots q'_{n-1} - 1$. However, x and y are in the same level of \mathcal{R}'_n ; furthermore, $S^i(B'_{n-1,0}(n))$ is included in the level $S^i(B'_n)$ and $S^j(B'_{n-1,0}(n))$ in the level $S^j(B'_n)$, so we have $i = j$. Moreover, since we have $t_{n,n} = 1$, all the sets $S^k(B'_{n-1,0}(n))$ are n -levels, that is, levels of the n th T -Rokhlin tower \mathcal{R}_n , so x and y are in the same n -level. This holds

for every $n \in I$, so for infinitely many n . Moreover, $(\mathcal{R}_n)_{n \in \mathbb{N}}$ separates the points up to a null set, since T is rank-one; hence the result. \square

LEMMA 5.15. For every $n \in \mathbb{N}$, we have

$$\mu(K_n) \geq \mu(X_n) - \mu(B_n) - 2\mu(X_n \setminus E_{n,n}).$$

Moreover, $\mu(K_n) \xrightarrow{n \rightarrow +\infty} 1$ if $\mu(E_{n,n}) \xrightarrow{n \rightarrow +\infty} 1$.

Proof of Lemma 5.15. The set K_n is equal to $(E_{n,n} \setminus B_n) \setminus T(X_n \setminus E_{n,n})$, so we get

$$\begin{aligned} \mu(K_n) &\geq \mu(E_{n,n} \setminus B_n) - \mu(T(X_n \setminus E_{n,n})) \\ &\geq \mu(E_{n,n}) - \mu(B_n) - \mu(X_n \setminus E_{n,n}) \\ &= \mu(X_n) - \mu(B_n) - 2\mu(X_n \setminus E_{n,n}). \end{aligned}$$

The second result follows from the fact that $\mu(X_n) \xrightarrow{n \rightarrow +\infty} 1$ and $\mu(B_n) \xrightarrow{n \rightarrow +\infty} 0$. \square

LEMMA 5.16. For every $x \in K_n$, there exists $k \in \mathbb{Z}$ such that

$$|k| \leq 4(h_{n-1} + Z_{n-1})(h'_{n-1})^2$$

and $T^{-1}x = S^kx$.

Proof of Lemma 5.16. Let $x \in K_n$. By the definition of K_n , the points x and $T^{-1}x$ are in $E_{n,n}$ and there exists $1 \leq i \leq h_n - 1$ such that $x \in T^i(B_n)$. Writing $E_{n,n}$ as

$$E_{n,n} = \bigsqcup_{\substack{0 \leq i \leq h'_{n-1}-1 \\ 0 \leq i_{n-1} \leq q'_{n-1}-1}} S^i \zeta_n^{i_{n-1}}(B'_{n-1,0}(n)) = \bigsqcup_{\substack{0 \leq i \leq h'_{n-1}-1 \\ 0 \leq i_{n-1} \leq q'_{n-1}-1}} S^i(B'_{n-1,i_{n-1}}(n)),$$

it is clear that there exist $0 \leq k_0, k_1 \leq h'_{n-1} - 1$ such that $y := S^{-k_0}x$ and $z := S^{-k_1}T^{-1}x$ are in $\bigsqcup_{0 \leq i_{n-1} \leq q'_{n-1}-1} B'_{n-1,i_{n-1}}(n)$.

We first show that we can write $y = \zeta_n^{k_2}z$ for some k_2 , using the fact that ζ_n connects the n -bricks of step (n, n) of the construction (since $t_{n,n} = 1$). Second, ζ_n can be written as a power of S and the equality $y = S^{k_3}z$ holds for some k_3 that we will be able to bound by Lemma 5.4. Finally, the result follows from the bound for each integer k_0, k_1, k_3 .

Step 1: Finding k_2 such that $y = \zeta_n^{k_2}z$. Using Lemma 5.3, we can write

$$x = \zeta_1^{i_0} \dots \zeta_{n-1}^{i_{n-2}}y \quad \text{and} \quad T^{-1}x = \zeta_1^{j_0} \dots \zeta_{n-1}^{j_{n-2}}z$$

for some integers $0 \leq i_0, j_0 \leq q'_0 - 1, \dots, 0 \leq i_{n-2}, j_{n-2} \leq q'_{n-2} - 1$, and there exist $0 \leq i_{n-1}, j_{n-1} \leq q'_{n-1} - 1$ such that

$$y \in B'_{n-1,i_{n-1}}(n) \quad \text{and} \quad z \in B'_{n-1,j_{n-1}}(n).$$

More precisely, by Lemma 5.5, and the fact that y and z are in n -bricks at step (n, n) , we have

$$x = \zeta_1(M_1)^{i_0} \dots \zeta_{n-1}(M_{n-1})^{i_{n-2}}y \quad \text{and} \quad T^{-1}x = \zeta_1(L_1)^{j_0} \dots \zeta_{n-1}(L_{n-1})^{j_{n-2}}z$$

with $k \leq L_k, M_k \leq n$ for every $1 \leq k \leq n - 1$. By construction, T and the maps $\zeta_k(m)$, for $1 \leq k \leq n - 1$ and $k \leq m \leq n$, satisfy the following property: for every n -level $T^k(B_n)$, with $0 \leq k \leq h_n - 1$, contained in the domain of the map, if it is mapped to another n -level $T^{k+\ell}(B_n)$, with $0 \leq k + \ell \leq h_n - 1$, then the application coincides with T^ℓ on $T^k(B_n)$. In other words, it consists in going up or down $|\ell|$ floors in the tower \mathcal{R}_n , without going above its roof or below its base. Therefore, from $B'_{n-1,i_{n-1}}(n)$ to $B'_{n-1,j_{n-1}}(n)$, the map

$$\tilde{S} := (\zeta_1(L_1)^{j_0} \dots \zeta_{n-1}(L_{n-1})^{j_{n-2}})^{-1} T^{-1} \zeta_1(M_1)^{i_0} \dots \zeta_{n-1}(M_{n-1})^{i_{n-2}}$$

consists in successively going up or down in the tower, so this is a power of T given by the difference between the floor of $B'_{n-1,i_{n-1}}(n)$ and that of $B'_{n-1,j_{n-1}}(n)$. The map $\zeta_n^{j_{n-1}-i_{n-1}}$ also satisfies this property, and thus $\zeta_n^{j_{n-1}-i_{n-1}}$ and \tilde{S} coincide on $B'_{n-1,i_{n-1}}(n)$ and $y = \zeta_n^{k_2} z$ with $k_2 := j_{n-1} - i_{n-1}$.

Step 2: Finding k_3 such that $y = S^{k_3} z$. Using Lemma 5.3 and the equality $\zeta_n^i(B'_n) = B'_{n-1,i}$, we have $S^{h'_{n-1}(j_{n-1}-i_{n-1})} y = z$, we set $k_3 := h'_{n-1}(j_{n-1} - i_{n-1})$ and it remains to find a bound for $j_{n-1} - i_{n-1}$. We need to get more information on the power of T , denoted by T^ℓ , which coincides with \tilde{S} on $B'_{n-1,i_{n-1}}(n)$. By Lemma 5.4 and the definition of \tilde{S} , we get

$$\begin{aligned} |\ell| &\leq (h_{n-1} + Z_{n-1})(i_0 + \dots + i_{n-2}) + 1 + (h_{n-1} + Z_{n-1})(j_0 + \dots + j_{n-2}) \\ &\leq 2(h_{n-1} + Z_{n-1})(q'_0 + \dots + q'_{n-2}) + 1 \\ &\leq 3(h_{n-1} + Z_{n-1})(q'_0 + \dots + q'_{n-2}), \end{aligned}$$

where ‘+1’ comes from ‘ T^{-1} ’ in the expression of \tilde{S} and has been bounded by $(h_{n-1} + Z_{n-1})(q'_0 + \dots + q'_{n-2})$. The sum $q'_0 + \dots + q'_{n-2}$ is less than the product $q'_0 \dots q'_{n-2} = h'_{n-1}$, which gives

$$|\ell| \leq 3(h_{n-1} + Z_{n-1})h'_{n-1}.$$

Since ζ_n has a positive cocycle (by Lemma 5.4), the equality $\zeta_n^{(j_{n-1}-i_{n-1})} = T^\ell$ implies $|\ell| \geq |j_{n-1} - i_{n-1}|$. Therefore, we find the bound

$$|k_3| \leq 3(h_{n-1} + Z_{n-1})(h'_{n-1})^2.$$

Step 3: Bounding the integer k such that $T^{-1}x = S^k x$. By the definition of k_0, k_1 and k_3 , $T^{-1}x$ is equal to $S^k x$ with $k := k_1 - k_3 - k_0$ which is thus bounded as follows:

$$\begin{aligned} |k| &\leq |k_0| + |k_1| + |k_3| \\ &\leq 2(h'_{n-1} - 1) + 3(h_{n-1} + Z_{n-1})(h'_{n-1})^2 \\ &\leq 4(h_{n-1} + Z_{n-1})(h'_{n-1})^2, \end{aligned}$$

and hence the result. □

COROLLARY 5.17. *If $\mu(E_{n,n}) \xrightarrow{n \rightarrow +\infty} 1$, then T and S have the same orbits.*

Proof of Corollary 5.17. It is clear that the S -orbits are contained in the T -orbits. By Lemma 5.15, $\bigcup_{n \in \mathbb{N}} K_n$ is of full measure, so the reverse inclusion follows from Lemma 5.16. \square

Remark 5.18. Corollary 5.17 holds for every rank-one system T . Indeed, skipping steps in the cutting-and-stacking process of T recursively increases the cutting parameters q_n , it enables us to get criteria (9) and (11) (the first one implies that the construction in §5.1 is well defined, the second one that $\mu(E_{n,n}) \rightarrow 1$).

However, the quantification of the cocycles will not necessarily hold for all the rank-one systems, since we will need to control the quantities Z_n depending on the spacing parameters (see §5.5).

Note that by Dye’s theorem, it was already known that every rank-one system is orbit equivalent to the universal odometer, but the proof of this theorem does not provide an explicit orbit equivalence, and thus prevents us from quantifying the cocycles.

Now the goal is to control the cocycle c_S . The equalities in equation (7) in §5.2 and the decomposition of $B_{n-1,i}$ in bricks motivate the following definition:

$$\begin{aligned} \text{for all } m \geq n \geq 1, \quad D_n(m) &:= \zeta_1^{q'_0-1} \cdots \zeta_{n-1}^{q'_{n-2}-1} \left(\bigsqcup_{0 \leq i_n \leq q'_{n-1}-2} B'_{n-1,i_n}(m) \right) \\ &= \zeta_1^{q'_0-1} \cdots \zeta_{n-1}^{q'_{n-2}-1} \left(\bigsqcup_{0 \leq i_n \leq q'_{n-1}-2} \zeta_n^{i_n}(m)(B'_{n-1,0}) \right). \end{aligned} \quad (12)$$

It is the union of all the translates of the m -bricks at step (n, m) composing D_n . Note that S coincides with $\zeta_n(m)\zeta_{n-1}^{-(q'_{n-2}-1)} \cdots \zeta_1^{-(q'_0-1)}$ on $D_n(m)$ (since it coincides with $\zeta_n \zeta_{n-1}^{-(q'_{n-2}-1)} \cdots \zeta_1^{-(q'_0-1)}$ and ζ_n coincides with $\zeta_n(m)$ on the m -bricks at step n). The partition of D_n into such subsets $D_n(m)$, for $m \geq n$, gives a fine control of the cocycle c_S .

LEMMA 5.19. For $1 \leq n < m$, $D_n(m)$ is contained in $X_m \setminus E_{n,m-1}$ and we have

$$\mu(D_n(m)) \leq \begin{cases} \varepsilon_{m-1} - \varepsilon_m + \frac{H'_n}{h_{m-1}} & \text{if } m > n + 1, \\ \varepsilon_{m-1} - \varepsilon_m + \frac{H'_{n-1} + p_{n-1}h'_{n-1}}{h_n} & \text{if } m = n + 1. \end{cases}$$

For all $n \geq 1$, we have

$$\mu(D_n(n)) \leq \frac{q'_{n-1}}{h_n}.$$

Moreover, for every $x \in D_n(m)$,

$$|c_S(x)| \leq (h_{m-1} + Z_{m-1})h'_{n-1}.$$

Proof of Lemma 5.19. For $1 \leq n < m$, $D_n(m)$ is composed of translates of the m -bricks used at step (n, m) , so it is disjoint from the translates of the M -bricks used at step (n, M) for $n \leq M \leq m - 1$; hence the inclusion $D_{n,m} \subseteq X_m \setminus E_{n,m-1}$. The bound for $\mu(D_n(m))$

follows from the decomposition $X_m \setminus E_{n,m-1} = (X_m \setminus X_{m-1}) \sqcup (X_{m-1} \setminus E_{n,m-1})$ and Lemma 5.13.

For $n \geq 1$, by the definition of $D_n(n)$ and the ζ_i -invariance of the measure, we get

$$\mu(D_n(n)) = (q'_{n-1} - 1)\mu(B'_{n-1,0}(n)) \leq q'_{n-1}\mu(B'_{n-1,0}(n));$$

hence the result, since $B'_{n-1,0}(n)$ is an n -level, so it has measure less than $1/h_n$.

For the cocycle c_S , we first decompose $D_n(m)$ in the following way:

$$D_n(m) = \bigsqcup_{\ell} \underbrace{\zeta_1^{q'_0-1} \cdots \zeta_{n-1}^{q'_{n-2}-1}}_{=:D^\ell}(\beta_\ell),$$

where $(\beta_\ell)_\ell$ is the family of m -bricks, at step (n, m) , which constitute the subset $\bigsqcup_{0 \leq i_n \leq q'_{n-1}-2} B'_{n-1,i_n}(m)$. For a fixed ℓ , by Lemma 5.5, there exist $1 \leq L_1 \leq m, \dots, n-1 \leq L_{n-1} \leq m$ such that

$$D^\ell = \zeta_1^{q'_0-1}(L_1) \cdots \zeta_{n-1}^{q'_{n-2}-1}(L_{n-1})(\beta_\ell)$$

and, on this subset, S coincides with $\zeta_n(m)\zeta_{n-1}^{-(q'_{n-2}-1)}(L_{n-1}) \cdots \zeta_1^{-(q'_0-1)}(L_1)$. Then, using Lemma 5.4, we get

$$\begin{aligned} |(c_S)_{|D^\ell}| &\leq (h_{m-1} + Z_{m-1})((q'_0 - 1) + \cdots + (q'_{n-2} - 1) + 1) \\ &\leq (h_{m-1} + Z_{m-1})q'_0 \cdots q'_{n-2} \\ &= (h_{m-1} + Z_{m-1})h'_{n-1}, \end{aligned}$$

and hence the result. □

5.5. *Proof of Theorem 3.9.* Let T be a rank-one system whose parameters satisfy criteria (9) and (11). The first one ensures that the construction is well defined (Lemma 5.10), the second one implies $\mu(E_{n,n}) \rightarrow 1$ (Lemma 5.13), so we have an orbit equivalence between T and S (Lemma 5.17). We can then define the cocycles $c_T, c_S: X \rightarrow \mathbb{Z}$ by

$$\text{for all } x \in X, \quad Tx = S^{c_T(x)}x \quad \text{and} \quad Sx = T^{c_S(x)}x.$$

In Lemmas 5.16 and 5.19, we obtained bounds for the cocycles on precise subsets covering $X: (K_n)_n$ for $c_T, (D_n(m))_{n,m}$ for c_S . This will provide a bound for the φ -integral of each cocycle. However, first, we need to change φ via the following lemma inspired by [CJLMT23, Lemma 2.12]. Without loss of generality, φ has the properties given by the lemma and this will simplify the bound for each φ -integral.

LEMMA 5.20. *Let $0 < \alpha \leq 1$ and $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ satisfying $\varphi(t) = o(t^\alpha)$. Then, there exists $\Phi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with the following properties:*

- Φ is increasing;
- Φ is subadditive: for all $t, s \in \mathbb{R}_+, \Phi(t + s) \leq \Phi(t) + \Phi(s)$;
- $\Phi(t) = o(t^\alpha)$;
- $\varphi(t) = O(\Phi(t))$.

Proof of Lemma 5.20. Set

$$\begin{aligned} \theta: \mathbb{R}_+^* &\rightarrow \mathbb{R}_+ \\ t &\mapsto \min\left(1, \sup_{s \geq t} \frac{\varphi(s) + 1}{s}\right) \end{aligned}$$

and

$$\begin{aligned} \Phi: \mathbb{R}_+ &\rightarrow \mathbb{R}_+ \\ t &\mapsto \int_0^t \theta(s) \, ds. \end{aligned}$$

The map θ is positive-valued and non-increasing, so Φ is an increasing and subadditive function satisfying $\Phi(t) \geq t\theta(t)$ for every $t \in \mathbb{R}_+$. The assumption $\varphi(t) = o(t^\alpha)$ implies that $\theta(t) = \sup_{s \geq t} (\varphi(s) + 1)/s$ for $t > 0$ large enough, so we have

$$\Phi(t) \geq t\theta(t) \geq t \sup_{s \geq t} \frac{\varphi(s) + 1}{s} \geq \varphi(t).$$

Finally, for a fixed $\varepsilon > 0$, there exists $t_0 > 0$ such that $\varphi(s) \leq \varepsilon s^\alpha$ for every $s \geq t_0$. For every $t \geq t_0$, this gives

$$\sup_{s \geq t} \frac{\varphi(s) + 1}{s} \leq \sup_{s \geq t} \left(\frac{\varepsilon}{s^{1-\alpha}} + \frac{1}{s}\right) = \frac{\varepsilon}{t^{1-\alpha}} + \frac{1}{t}$$

and for every $t \geq t_0$, we have

$$\int_{t_0}^t \theta(s) \, ds \leq \int_{t_0}^t \left(\frac{\varepsilon}{s^{1-\alpha}} + \frac{1}{s}\right) \, ds = \frac{\varepsilon}{\alpha} t^\alpha + \ln t - \frac{\varepsilon}{\alpha} t_0^\alpha - \ln t_0,$$

and hence $\Phi(t) = o(t^\alpha)$. □

LEMMA 5.21. *Assume that criteria (9) (in Lemma 5.10) and (11) (after Lemma 5.13) are satisfied. Let $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be an increasing and subadditive map. Then, setting*

$$\begin{aligned} \Delta(n) &:= (1 + 2(H'_n + p_n h'_n))(h'_n)^2 \left(\frac{\varphi(h_{n+1}^3)}{h_{n+1}} + \frac{\varphi(Z_{n+1} h_{n+1}^2)}{h_{n+1}}\right), \\ \Delta_\varepsilon(n) &:= \varepsilon_{n+1} (h'_n)^2 (\varphi(h_{n+1}^3) + \varphi(Z_{n+1} h_{n+1}^2)), \end{aligned}$$

we have the following bound:

$$\int_X \varphi(|c_T(x)|) \, d\mu \leq \varphi(4(h_0 + Z_0)(h'_0)^2) + 4 \sum_{n=0}^{+\infty} \Delta(n) + 4 \sum_{n=0}^{+\infty} \Delta_\varepsilon(n). \tag{13}$$

Proof of Lemma 5.21. Motivated by Lemma 5.16, we will rather quantify the cocycle $c_{T^{-1}}$ defined on X (up to a null set) by

$$T^{-1}x = S^{c_{T^{-1}}(x)}x.$$

It is equivalent to quantifying c_T since we have

$$\text{for all } x \in X, \quad c_{T^{-1}}(x) = -c_T(T^{-1}x)$$

and μ is T -invariant.

Let $(K'_n)_{n>0}$ be the partition of X inductively defined by

$$\begin{cases} K'_1 := K_1, \\ \text{for all } n > 0, K'_{n+1} := K_{n+1} \setminus (K_1 \cup \dots \cup K_n). \end{cases}$$

The subsets K'_n are pairwise disjoint and cover the whole space since we have

$$K'_1 \cup \dots \cup K'_n = K_1 \cup \dots \cup K_n$$

and $\mu(K_n) \rightarrow 1$ (using Lemma 5.15). By the fact that K_n is included in X_n , and by Lemmas 5.15 and 5.13, we have

$$\begin{aligned} \mu(K'_{n+1}) &\leq \mu(X \setminus K_n) \\ &= \mu(X \setminus X_n) + \mu(X_n \setminus K_n) \\ &\leq \varepsilon_n + \mu(B_n) + 2\mu(X_n \setminus E_{n,n}) \\ &\leq \varepsilon_n + \frac{1 + 2(H'_{n-1} + p_{n-1}h'_{n-1})}{h_n}. \end{aligned}$$

Since K'_{n+1} is contained in K_{n+1} , Lemma 5.16 implies

$$\text{for all } x \in K'_{n+1}, \quad |c_{T^{-1}}(x)| \leq 4(h_n + Z_n)(h'_n)^2.$$

We then get

$$\begin{aligned} \int_X \varphi(|c_T(x)|) d\mu &= \int_X \varphi(|c_{T^{-1}}(x)|) d\mu \\ &= \sum_{n=0}^{+\infty} \int_{K'_{n+1}} \varphi(|c_{T^{-1}}(x)|) d\mu \\ &\leq \sum_{n=0}^{+\infty} \mu(K'_{n+1})\varphi(4(h_n + Z_n)(h'_n)^2) \\ &\leq \varphi(4(h_0 + Z_0)(h'_0)^2) \\ &\quad + \sum_{n=1}^{+\infty} \left(\varepsilon_n + \frac{1 + 2(H'_{n-1} + p_{n-1}h'_{n-1})}{h_n} \right) \varphi(4(h_n + Z_n)(h'_n)^2). \end{aligned}$$

Now we use the assumptions on φ to simplify the previous bound. We have $h'_n = h'_{n-1}q'_{n-1} \leq h'_{n-1}h_n$ (by construction, we have $q'_{n-1} \leq r_{n,n} \leq h_n$). By monotonicity and subadditivity, this yields

$$\begin{aligned} \varphi(4(h_n + Z_n)(h'_n)^2) &\leq \varphi(4(h_n + Z_n)(h'_{n-1}h_n)^2) \\ &\leq 4(h'_{n-1})^2(\varphi(h_n^3) + \varphi(Z_nh_n^2)) \end{aligned}$$

and we get the bound (13). □

LEMMA 5.22. Assume that criteria (9) (in Lemma 5.10) and (11) (after Lemma 5.13) are satisfied and that the following holds:

$$\text{for all } n \geq 0, \quad \frac{q'_n}{q_n} \leq 4$$

(this is an assumption that we will be able to get by Lemma 5.12, using flexible classes).

Let $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be an increasing and subadditive map. Then, setting

$$\begin{aligned} \Gamma_1(n) &:= 4h'_n \left(\frac{\varphi(h_{n+1}^2)}{h_{n+1}} + \frac{\varphi(Z_{n+1}h_{n+1})}{h_{n+1}} \right), \\ \Gamma_2(n) &:= (H'_n + p_n h'_n) h'_n \left(\frac{\varphi(h_{n+1})}{h_{n+1}} + \frac{\varphi(Z_{n+1})}{h_{n+1}} \right), \\ \Gamma_3(n, m) &:= H'_n h'_{n-1} \left(\frac{\varphi(h_m)}{h_m} + \frac{\varphi(Z_m)}{h_m} \right), \\ \Gamma_\varepsilon(n, m) &:= \varepsilon_m h'_n (\varphi(h_m) + \varphi(Z_m)), \end{aligned}$$

we have the following bound:

$$\begin{aligned} \int_X \varphi(|c_S|) d\mu &\leq \mu(D_1(1))\varphi((h_0 + Z_0)h'_0) \\ &\quad + \sum_{n \geq 0} \Gamma_1(n) + \sum_{n \geq 0} \Gamma_2(n) + \sum_{n \geq 1} \sum_{m \geq n+1} \Gamma_3(n, m) \\ &\quad + \sum_{n \geq 0} \sum_{m \geq n+1} \Gamma_\varepsilon(n, m). \end{aligned} \tag{14}$$

Proof of Lemma 5.22. By Lemma 5.19, for each subset $D_n(m)$, we found a bound for the cycle c_S on it, we then get

$$\begin{aligned} \int_X \varphi(|c_S|) d\mu &= \sum_{m \geq n \geq 1} \int_{D_n(m)} \varphi(|c_S|) d\mu \\ &\leq \sum_{m \geq n \geq 1} \mu(D_n(m))\varphi((h_{m-1} + Z_{m-1})h'_{n-1}) \\ &\leq \mu(D_1(1))\varphi((h_0 + Z_0)h'_0) \\ &\quad + \sum_{n \geq 2} \gamma_1(n) + \sum_{n \geq 1} \gamma_2(n) + \sum_{n \geq 1} \sum_{m \geq n+2} \gamma_3(n, m), \end{aligned}$$

where

$$\begin{aligned} \gamma_1(n) &:= \mu(D_n(n))\varphi((h_{n-1} + Z_{n-1})h'_{n-1}), \\ \gamma_2(n) &:= \mu(D_n(n+1))\varphi((h_n + Z_n)h'_{n-1}), \\ \gamma_3(n, m) &:= \mu(D_n(m))\varphi((h_{m-1} + Z_{m-1})h'_{n-1}). \end{aligned}$$

Lemma 5.19 also yields a bound for the measure of each set $D_n(m)$, which implies

$$\begin{aligned} \gamma_1(n) &\leq \frac{q'_{n-1}}{h_n} \varphi((h_{n-1} + Z_{n-1})h'_{n-1}), \\ \gamma_2(n) &\leq \left(\varepsilon_n + \frac{H'_{n-1} + p_{n-1}h'_{n-1}}{h_n} \right) \varphi((h_n + Z_n)h'_{n-1}), \\ \gamma_3(n, m) &\leq \left(\varepsilon_{m-1} + \frac{H'_n}{h_{m-1}} \right) \varphi((h_{m-1} + Z_{m-1})h'_{n-1}). \end{aligned}$$

For all $n \geq 2$, note that we have

$$\begin{aligned} \varphi((h_{n-1} + Z_{n-1})h'_{n-1}) &\leq \varphi((h_{n-1} + Z_{n-1})h'_{n-2}h_{n-1}) \\ &\leq h'_{n-2}(\varphi(h_{n-1}^2) + \varphi(Z_{n-1}h_{n-1})) \end{aligned}$$

and

$$\frac{q'_{n-1}}{h_n} \leq \frac{q'_{n-1}}{h_{n-1}q_{n-1}} \leq \frac{4}{h_{n-1}},$$

so we get

$$\gamma_1(n) \leq 4h'_{n-2} \left(\frac{\varphi(h_{n-1}^2)}{h_{n-1}} + \frac{\varphi(Z_{n-1}h_{n-1})}{h_{n-1}} \right) = \Gamma_1(n - 2).$$

For $\gamma_2(n)$ and $\gamma_3(n, m)$, note that we have

$$\text{for all } n \geq 1, \text{ for all } m \geq n + 1, \quad \varphi((h_{m-1} + Z_{m-1})h'_{n-1}) \leq h'_{n-1}(\varphi(h_{m-1}) + \varphi(Z_{m-1})),$$

so we get

$$\begin{aligned} \gamma_2(n) &\leq \left(\varepsilon_n + \frac{H'_{n-1} + p_{n-1}h'_{n-1}}{h_n} \right) h'_{n-1}(\varphi(h_n) + \varphi(Z_n)) \\ &= \varepsilon_n h'_{n-1}(\varphi(h_n) + \varphi(Z_n)) + (H'_{n-1} + p_{n-1}h'_{n-1})h'_{n-1} \left(\frac{\varphi(h_n)}{h_n} + \frac{\varphi(Z_n)}{h_n} \right) \\ &= \Gamma_\varepsilon(n - 1, n) + \Gamma_2(n - 1) \end{aligned}$$

and

$$\begin{aligned} \gamma_3(n, m) &\leq (\varepsilon_{m-1} + \frac{H'_n}{h_{m-1}})h'_{n-1}(\varphi(h_{m-1}) + \varphi(Z_{m-1})) \\ &= \varepsilon_{m-1}h'_{n-1}(\varphi(h_{m-1}) + \varphi(Z_{m-1})) + H'_n h'_{n-1} \left(\frac{\varphi(h_{m-1})}{h_{m-1}} + \frac{\varphi(Z_{m-1})}{h_{m-1}} \right). \\ &= \Gamma_\varepsilon(n - 1, m - 1) + \Gamma_3(n, m - 1) \end{aligned}$$

The bound (14) now follows immediately. □

Proof of Theorem 3.9. Let \mathcal{C} be a flexible class and $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ a map satisfying $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$. If $\Phi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is another map satisfying $\varphi(t) = O(\Phi(t))$, then Φ -integrability implies φ -integrability. Therefore, without loss of generality, we assume that φ satisfies the assumptions of Lemma 5.20, that is, φ is increasing and subadditive.

Using the definition of a flexible class, we will build T with large enough and inductively chosen cutting parameters q_n . Let $\mathcal{F}_{\mathcal{C}}$ be an associated set of parameters, and fix the associated constants C and C' given in Definition 3.7. First, choose any cutting and spacing parameter $(q_0, (\sigma_{0,0}, \dots, \sigma_{0,q_0}))$ in $\mathcal{F}_{\mathcal{C}}$ such that $q_0 \geq 3$. Without loss of generality, we assume $p_0 = 2$ and we get $q_0 > p_0$, as required in the assumption of Lemma 5.10 for $n = 0$. For a fixed $n \geq 1$, assume that $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n-1}$ has already been determined in $\mathcal{F}_{\mathcal{C}}$, this immediately gives q'_0, \dots, q'_{n-1} (see Lemma 5.9). The goal is

to find the next parameters with q_n large enough. Consider $\kappa_n > 0$ such that for every $t \geq h_n \kappa_n$, the following hold:

$$\kappa_n > \max(p_n, q'_0, \dots, q'_{n-1}); \tag{15}$$

$$\frac{H'_n + p_n h'_n}{t} \leq \frac{1}{n}. \tag{16}$$

The assumption $\varphi(t) = o(t^{1/3})$ also implies the following inequations for a large enough κ_n :

$$(1 + 2(H'_n + p_n h'_n))(h'_n)^2 \left(\frac{\varphi(t^3)}{t} + \frac{\varphi(Ct^3)}{t} \right) \leq \frac{1}{2^n}; \tag{17}$$

$$(h'_n)^2(\varphi(t^3) + \varphi(Ct^3)) \leq \frac{t}{2^n q_0 \dots q_{n-1}}; \tag{18}$$

$$4h'_n \left(\frac{\varphi(t^2)}{t} + \frac{\varphi(Ct^2)}{t} \right) \leq \frac{1}{2^n}; \tag{19}$$

$$(H'_n + p_n h'_n)h'_n \left(\frac{\varphi(t)}{t} + \frac{\varphi(Ct)}{t} \right) \leq \frac{1}{2^{n+1}}; \tag{20}$$

$$\text{for all } 1 \leq \ell \leq n, \quad H'_\ell h'_{\ell-1} \left(\frac{\varphi(t)}{t} + \frac{\varphi(Ct)}{t} \right) \leq \frac{1}{2^{n+2}}; \tag{21}$$

$$\text{for all } 0 \leq \ell \leq n, \quad h'_\ell(\varphi(t) + \varphi(Ct)) \leq \frac{t}{2^n q_0 \dots q_{n-1}}, \tag{22}$$

for every $t \geq h_n \kappa_n$. With inequations (17), (18), (19), (20), (21) and (22), we will respectively find bounds for the quantities $\Delta(n)$, $\Delta_\varepsilon(n)$, $\Gamma_1(n)$, $\Gamma_2(n)$, $\Gamma_3(n, m)$ and $\Gamma_\varepsilon(n, m)$ (see Lemmas 5.21 and 5.22).

We then set a new cutting parameter $q_n \geq \kappa_n$ large enough with associated spacing parameters $\sigma_{n,0}, \dots, \sigma_{n,q_n}$ so that $(q_k, (\sigma_{k,0}, \dots, \sigma_{k,q_k}))_{0 \leq k \leq n} \in \mathcal{F}_C$, $\sigma_n \leq C' q_n h_{n-1}$ and the following additional assumptions are satisfied:

$$q_n - (1 + p_n) \geq C' h_n \tag{23}$$

and

$$\text{for all } 0 \leq k \leq n - 1, \quad q_n \geq C' 2^{n-k} q_k. \tag{24}$$

Let (h_n) , (σ_n) and (Z_n) be the sequences associated to $\mathbf{p} := (q_n, (\sigma_{n,0}, \dots, \sigma_{n,q_n}))_{n \geq 0} \in \mathcal{P}^{\mathbb{N}}$ (as described in Definition 3.1), (h'_n) the height sequence of the cutting sequence $(q'_n)_{n \geq 0}$ for the universal odometer that we build. We first check that the underlying system is finite measure preserving, that is, condition (F) in Definition 3.2 is satisfied. However, we have

$$\frac{\sigma_n}{h_{n+1}} \leq \frac{C' q_n h_{n-1}}{q_n q_{n-1} h_{n-1}} = \frac{C'}{q_{n-1}},$$

so the summability easily follows from inequality (24). The underlying system preserves a probability measure, so it is rank-one. Moreover it belongs to \mathcal{C} by the definition of a flexible class.

Inequality (15) ensures that criterion (9) holds and that the construction in §5.1 is well defined (see Lemma 5.10). Using $h_{n+1} \geq h_n q_n$, the limit in equation (11) is a consequence of inequality (16) and implies $\mu(E_{n,n}) \rightarrow 1$. Inequality (23) implies

$$\text{for all } n \in \mathbb{N}, \quad \frac{q'_n}{q_n} \leq 4$$

(see Lemma 5.12).

Then, Lemmas 5.21 and 5.22 imply that the bounds in equation (13) for the φ -integral of c_T and equation (14) for the φ -integral of c_S hold. It remains to prove that these bounds are finite, namely that the series

$$\begin{aligned} & \sum_{n \geq 0} \Delta(n), \sum_{n \geq 0} \Delta_\varepsilon(n), \sum_{n \geq 0} \Gamma_1(n), \sum_{n \geq 0} \Gamma_2(n), \\ & \sum_{n \geq 1} \sum_{m \geq n+1} \Gamma_3(n, m) \quad \text{and} \quad \sum_{n \geq 0} \sum_{m \geq n+1} \Gamma_\varepsilon(n, m) \end{aligned}$$

converge.

Using the monotonicity of φ and the inequalities $Z_{n+1} \leq Ch_{n+1}$ and (17) for $t = h_{n+1}$ (which is greater or equal to $h_n \kappa_n$), we get $\Delta(n) \leq (1/2^n)$, so the series $\sum_{n \geq 0} \Delta(n)$ converges. It is also straightforward to see that the series $\sum_{n \geq 0} \Gamma_1(n)$ and $\sum_{n \geq 0} \Gamma_2(n)$ are convergent, using inequalities (19) and (20). Inequality (21) implies $\Gamma_3(n, m) \leq (1/2^{m+1})$, so we get

$$\sum_{m \geq n+1} \Gamma_3(n, m) \leq \frac{1}{2^{n+1}}$$

for every $n \geq 0$, and the series $\sum_{n \geq 1} \sum_{m \geq n+1} \Gamma_3(n, m)$ converges.

For the other series $\sum_{n \geq 0} \Delta_\varepsilon(n)$ and $\sum_{n \geq 0} \sum_{m \geq n+1} \Gamma_\varepsilon(n, m)$, we have to control the sequence (ε_n) (recall that $\varepsilon_n := \mu((X_n)^c)$). Denote by M_0 the measure of B_0 (the unique level of the T -Rokhlin tower \mathcal{R}_0). For every $n \geq 1$, we have

$$\varepsilon_n = \sum_{k \geq n} \frac{M_0}{q_0 \dots q_k} \sigma_k \leq \sum_{k \geq n} \frac{M_0 C' h_{k-1}}{q_0 \dots q_{k-1}} \leq \sum_{k \geq n} \frac{C'}{q_{k-1}} \leq \frac{1}{q_{n-1}} \sum_{k \geq n} \frac{1}{2^{k-n}} \leq \frac{2}{q_{n-1}},$$

using Lemma 3.4 and inequation (24).

Given $n \geq 0$, inequation (22) and Lemma 3.4 imply

$$(h'_n)^2(\varphi(h_{n+1}^3) + \varphi(Z_{n+1}h_{n+1}^2)) \leq \frac{h_{n+1}}{2^n q_0 \dots q_{n-1}} \leq \frac{q_n}{2^n M_0}.$$

Combining this with the inequality $\varepsilon_{n+1} \leq 2/q_n$, we then get

$$\Delta_\varepsilon(n) = \varepsilon_{n+1}(h'_n)^2(\varphi(h_{n+1}^3) + \varphi(Z_{n+1}h_{n+1}^2)) \leq \frac{1}{2^{n-1} M_0},$$

so the series $\sum_{n \geq 0} \Delta_\varepsilon(n)$ converges.

For fixed integers $n \geq 0$ and $m \geq n + 1$, inequation (22) and Lemma 3.4 imply

$$h'_n(\varphi(h_m) + \varphi(Z_m)) \leq \frac{h_m}{2^{m-1}q_0 \dots q_{m-2}} \leq \frac{q_{m-1}}{2^{m-1}M_0}.$$

Combining this with the inequality $\varepsilon_m \leq 2/q_{m-1}$, we then get

$$\Gamma_\varepsilon(n, m) = \varepsilon_m h'_n(\varphi(h_m) + \varphi(Z_m)) \leq \frac{1}{2^{m-2}M_0}.$$

This gives

$$\sum_{m \geq n+1} \Gamma_\varepsilon(n, m) \leq \frac{1}{2^{n-2}M_0}$$

for every $n \geq 0$, so the series $\sum_{n \geq 0} \sum_{m \geq n+1} \Gamma_\varepsilon(n, m)$ converges.

Therefore, the cocycles c_T and c_S are φ -integrable as wanted, which concludes the proof. □

Remark 5.23. For φ -integrability of c_S , we only need to control quantities of the form $\varphi(u^2)/u$ and $\varphi(u)/u$ ($\varphi(u^3)/u$ does not appear). Therefore, Theorem 3.9 can be stated with a stronger quantification on the cocycle c_S , namely ψ -integrability with $\psi(t) = o(t^{1/2})$ (it suffices to replace $t^{1/3}$ by $t^{1/2}$ in inequation (22)).

We are now able to prove Theorem D.

Proof of Theorem D. Let $\varphi: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a map satisfying $\varphi(t) \underset{t \rightarrow +\infty}{=} o(t^{1/3})$. By Lemma 5.20, we may and do assume that φ is increasing and subadditive.

Given a flexible class \mathcal{C} , an associated set of parameters $\mathcal{F}_{\mathcal{C}}$ and constants C and C' , the last proof shows that we can choose the parameters in the following way. First, we choose any cutting and spacing parameter $(q_0, (\sigma_{0,0}, \dots, \sigma_{0,q_0}))$ in $\mathcal{F}_{\mathcal{C}}$, with $q_0 \geq 3$. Then, if $\mathbf{p}_n := (q_k, (\sigma_{k,0}, \dots, \sigma_{0,q_k}))_{0 \leq k \leq n-1}$ has been set, there exists a constant depending on $\varphi, \mathcal{F}_{\mathcal{C}}, C, C'$ and \mathbf{p}_n , denoted by $K_\varphi(\mathcal{F}_{\mathcal{C}}, C, C', \mathbf{p}_n)$, such that conditions (15), (16), (17), (18), (19), (20), (21), (22), (23) and (24) hold for every $q_n \geq K_\varphi(\mathcal{F}_{\mathcal{C}}, C, C', \mathbf{p}_n)$, and it remains to find such an integer q_n and spacing parameters $\sigma_{n,0}, \dots, \sigma_{n,q_n}$ such that $\mathbf{p}_{n+1} := (q_k, (\sigma_{k,0}, \dots, \sigma_{0,q_k}))_{0 \leq k \leq n}$ is in $\mathcal{F}_{\mathcal{C}}$ and the inequality $\sigma_n \leq C'q_n h_{n-1}$ holds.

Let $\mathbf{Q} := (Q_{-1}, \dots, Q_{n_0})$ be a sequence of integers, where n_0, Q_0, \dots, Q_{n_0} are positive and $Q_0 \dots Q_{n_0} \geq 3$, and let us consider the set of parameters $\mathcal{F}(\mathbf{Q})$ built in §4.2, and the associated constants $C_{\mathbf{Q}}$ and $C'_{\mathbf{Q}}$. In this case, the spacing parameters $\sigma_{k,i}$ at step k are equal to 0 or h_{k-1} , so they are determined by the previous cutting parameters. Moreover, the first cutting parameter q_0 is equal to $Q_0 \dots Q_{n_0}$. Therefore, for every finite sequence $\mathbf{p}_n := (q_k, (\sigma_{k,0}, \dots, \sigma_{0,q_k}))_{0 \leq k \leq n-1}$ in $\mathcal{F}(\mathbf{Q})$, we write $K_\varphi(\mathbf{Q}, q_1, \dots, q_{n-1})$ instead of $K_\varphi(\mathcal{F}(\mathbf{Q}), C_{\mathbf{Q}}, C'_{\mathbf{Q}}, \mathbf{p}_n)$.

Recall that A denotes the set of sequences $(q_i)_{i \geq -1}$ of integers such that q_0, q_1, \dots are positive. To every sequence $\boldsymbol{\varepsilon} = (\varepsilon_i)_{i \geq 0} \in \{0, 1\}^{\mathbb{N}}$, we associate a sequence $q(\boldsymbol{\varepsilon}) \in A$ inductively defined by

$$q(\boldsymbol{\varepsilon})_0 = q_0,$$

$$\text{for all } i \geq 0, \quad q(\boldsymbol{\varepsilon})_{i+1} = K_\varphi(\mathbf{Q}, q(\boldsymbol{\varepsilon})_1, \dots, q(\boldsymbol{\varepsilon})_i) + \varepsilon_i.$$

Every sequence $\boldsymbol{\varepsilon} = (\varepsilon_i)_{i \geq 0} \in \{0, 1\}^{\mathbb{N}}$ provides a sequence of parameters in $\mathcal{F}(\boldsymbol{Q})$, whose cutting parameters are $q(\boldsymbol{\varepsilon})_0, q(\boldsymbol{\varepsilon})_1, \dots$, and which gives rise to the irrational rotation of angle $\theta(\boldsymbol{\varepsilon}) := [Q_{-1}, \dots, Q_{n_0}, q(\boldsymbol{\varepsilon})_1, q(\boldsymbol{\varepsilon})_2, \dots]$.

Let us now consider a non-empty open subset \mathcal{V} of \mathbb{R} and a finite sequence \boldsymbol{Q} so that $\theta(\boldsymbol{\varepsilon})$ is in \mathcal{V} for every $\boldsymbol{\varepsilon} \in \{0, 1\}^{\mathbb{N}}$. We get that the set of irrational numbers θ in \mathcal{V} such that the irrational rotation of angle θ is φ -integrably orbit equivalent to the universal odometer contains the set $\{q(\boldsymbol{\varepsilon}) \mid \boldsymbol{\varepsilon} \in \{0, 1\}^{\mathbb{N}}\}$, so it is uncountable using the facts that the map $\boldsymbol{\varepsilon} \in \{0, 1\}^{\mathbb{N}} \mapsto q(\boldsymbol{\varepsilon}) \in A$ is injective and the continued fraction expansion is unique for every irrational number. \square

Acknowledgements. I am very grateful to my advisors François Le Maître and Romain Tessera for their support, fruitful discussions and valuable advice. I also wish to thank Mathieu Da Silva, Victor Dubach and Fabien Hoareau for their useful comments on the paper. Finally, I am very grateful to the referee for their careful reading, for many corrections and for pointing out mistakes in the proof of an earlier version of Proposition 1.2.

REFERENCES

- [Aus16] T. Austin. Behaviour of entropy under bounded and integrable orbit equivalence. *Geom. Funct. Anal.* **26**(6) (2016), 1483–1525.
- [Bel69] R. M. Belinskaya. Partitions of Lebesgue space in trajectories defined by ergodic automorphisms. *Funct. Anal. Appl.* **2**(3) (1969), 190–199.
- [Cha69] R. V. Chacon. Weakly mixing transformations which are not strongly mixing. *Proc. Amer. Math. Soc.* **22**(3) (1969), 559–562.
- [CJLMT23] A. Carderi, M. Joseph, F. Le Maître and R. Tessera. Belinskaya’s theorem is optimal. *Fund. Math.* **263**(1) (2023), 51–90.
- [DEJ+23] H. Drillick, A. Espinosa-Dominguez, J. N. Jones-Baro, J. Leng, Y. Mandelshtam and C. E. Silva. Non-rigid rank-one infinite measures on the circle. *Dyn. Syst.* **38**(2) (2023), 275–300.
- [Dow11] T. Downarowicz. *Entropy in Dynamical Systems (New Mathematical Monographs, 18)*. Cambridge University Press, Cambridge, 2011.
- [DV23] A. I. Danilenko and M. I. Vieprik. Explicit rank-1 constructions for irrational rotations. *Studia Math.* **270**(2) (2023), 121–144.
- [Dye59] H. A. Dye. On groups of measure preserving transformations. I. *Amer. J. Math.* **81**(1) (1959), 119–159.
- [EW11] M. Einsiedler and T. Ward. *Ergodic Theory: With a View Towards Number Theory*. Springer, London, 2011.
- [Fer97] S. Ferenczi. Systems of finite rank. *Colloq. Math.* **73**(1) (1997), 35–65.
- [FRW11] M. Foreman, D. Rudolph and B. Weiss. The conjugacy problem in ergodic theory. *Ann. of Math.* (2) **173**(3) (2011), 1529–1586.
- [GZ19] S. Gao and C. Ziegler. Topological mixing properties of rank-one subshifts. *Trans. London Math. Soc.* **6**(1) (2019), 1–21.
- [HVN42] P. R. Halmos and J. Von Neumann. Operator methods in classical mechanics. II. *Ann. of Math.* (2) **43**(2) (1942), 332–350.
- [Jun76] A. Del Junco. Transformations with discrete spectrum are stacking transformations. *Canad. J. Math.* **28**(4) (1976), 836–839.
- [KL17] D. Kerr and H. Li. *Ergodic Theory: Independence and Dichotomies (Springer Monographs in Mathematics)*. Springer International Publishing, Cham, 2017.
- [KL21] D. Kerr and H. Li. Entropy, Shannon orbit equivalence, and sparse connectivity. *Math. Ann.* **380**(3) (2021), 1497–1562.
- [KL24] D. Kerr and H. Li. Entropy, virtual Abelianness and Shannon orbit equivalence. *Ergod. Th. & Dynam. Sys.*. Published online 10 April 2024.

- [Orn70] D. Ornstein. Bernoulli shifts with the same entropy are isomorphic. *Adv. Math.* **4**(3) (1970), 337–352.
- [Orn72] D. S. Ornstein. On the root problem in ergodic theory. *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, Volume 2: Probability Theory*. Vol. 6.2. Ed. L. M. Le Cam, J. Neyman and E. L. Scott. University of California Press, Berkeley, CA, 1972, pp. 347–357.
- [ORW82] D. S. Ornstein, D. J. Rudolph and B. Weiss. Equivalence of measure preserving transformations. *Mem. Amer. Math. Soc.* **37**(262) (1982), xii+116 pp.
- [OW87] D. S. Ornstein and B. Weiss. Entropy and isomorphism theorems for actions of amenable groups. *J. Anal. Math.* **48**(1) (1987), 1–141.
- [VO16] M. Viana and K. Oliveira. *Foundations of Ergodic Theory (Cambridge Studies in Advanced Mathematics, 151)*. Cambridge University Press, Cambridge, 2016.