



Complex linear differential equations with solutions in weighted Dirichlet spaces and derivative Hardy spaces

Qingze Lin and Huayou Xie*

Abstract. In this paper, by the use of n -th derivative characterization, we obtain several some sufficient conditions for all solutions of the complex linear differential equation

$$f^{(n)} + A_{n-1}(z)f^{(n-1)} + \cdots + A_1(z)f' + A_0(z)f = A_n(z)$$

to lie in weighted Dirichlet spaces and derivative Hardy spaces, respectively, where $A_i(z)$ ($i = 0, 1, \dots, n$) are analytic functions defined in the unit disc. This work continues the lines of the investigations by Heittokangas, et al. for growth estimates about the solutions of the above equation.

1 Introduction

Denote by \mathbb{D} the open unit disc in the complex plane and by $\partial\mathbb{D} = \{z : |z| = 1\}$ the unit circle. Define $H(\mathbb{D})$ as the space of all analytic functions on \mathbb{D} .

In 1982, the complex second-order equation

$$f'' + A(z)f = 0,$$

where $A(z) \in H(\mathbb{D})$, was investigated by Pommerenke [37]. By means of Carleson measures, he showed some sufficient conditions on the analytic function $A(z)$ such that all solutions of the above equation lie in Hardy spaces. Later on, complex linear differential equations of second and even higher orders on the unit disc attracted the attention of many scholars. In 2000, Heittokangas [19] investigated the growth of the solutions of the equation

$$f^{(n)} + A_{n-1}(z)f^{(n-1)} + \cdots + A_1(z)f' + A_0(z)f = A_n(z) \quad (1.1)$$

where $n \geq 2$. He wished to find some sufficient conditions for the coefficients $A_i(z)$ ($i = 0, 1, \dots, n$) such that all solutions of the equation (1.1) lie in some function spaces (i.e., weighted Hardy spaces, Bloch type spaces and general function spaces $F(p, q, s)$). In [25], two sufficient conditions for all solutions of equation (1.1) to lie in growth spaces H_α^∞ were presented by Huusko et al. Recall that for $0 < \alpha < \infty$, H_α^∞ is the space consisting of all functions $f \in H(\mathbb{D})$ such that

$$\|f\|_{H_\alpha^\infty} := \sup_{z \in \mathbb{D}} |f(z)|(1 - |z|^2)^\alpha < \infty.$$

2020 Mathematics Subject Classification: 34M10; 30H05; 31C25.

Keywords: complex linear differential equation, weighted Dirichlet spaces, derivative Hardy spaces..

*Corresponding author

Recently, equation (1.1) had been extensively studied in some other function spaces, such as weighted Fock spaces [24], Morrey spaces [39] and Dirichlet-Morrey spaces [40]. In particular, for the case of $A_n(z) = 0$, a lot of works had been done by a number of researchers. For instance, by means of sharp estimates of logarithmic derivatives, Gundersen-Steinbart-Wang [17] showed that every solution f of the linear differential equation

$$f^{(n)} + p_{n-1}(z)f^{(n-1)} + \cdots + p_1(z)f' + p_0(z)f = 0$$

where $p_i(z)$ are polynomials and $p_0 \neq 0$, is entire of finite rational order. The related problems in the unit disc were later considered by Chyzhykov et al. [5]. Indeed, they investigated the impact of the increasing in coefficients on the growth of solutions of the equation

$$f^{(n)} + A_{n-1}(z)f^{(n-1)} + \cdots + A_1(z)f' + A_0(z)f = 0. \quad (1.2)$$

Later on, Korhonen and Rättyä [26] continued the work of Chyzhykov et al. to show a precise estimate for the growth of solutions of the equation (1.2). In [21], Heittokangas et al. studied the growth relation between the coefficients and the solutions of the equation (1.2) in weighted Bergman or Hardy spaces. Indeed, they show that when the coefficients in the equation (1.2) belong to weighted Bergman or Hardy spaces, then all solutions are of some finite orders of growth, measured according to the Nevanlinna characteristic. For more related results, we refer the readers to [4, 15, 20, 27, 28, 42].

In this paper, motivated by the above works, we are interested in studying the sufficient conditions for all solutions of the equation (1.1) to lie in derivative Hardy spaces and weighted Dirichlet spaces. Now, let's recall their definitions.

Definition 1.1 Let $0 < p < \infty$. The derivative Hardy space S^p is a proper subspace of Hardy space H^2 and consists of $f \in H(\mathbb{D})$ satisfying

$$\|f\|_{S^p} := (|f(0)|^p + \|f'\|_{H^p})^{1/p} = \left(|f(0)|^p + \sup_{0 < r < 1} \frac{1}{2\pi} \int_0^{2\pi} |f'(re^{i\theta})|^p d\theta \right)^{1/p} < \infty.$$

In particular, for any $f(z) = \sum_{n=0}^{\infty} a_n z^n \in H(\mathbb{D})$, it holds that $\|f\|_{S^2}^2 = |f(0)|^2 + \sum_{n=1}^{\infty} n^2 |a_n|^2$. In 1978, Roan [38] started on the investigations for the boundedness of composition operators on derivative Hardy spaces. Related problems were also investigated by MacCluer [35]. After their works, Contreras and Hernández-Díaz [6] made a systematic work on the boundedness, compactness, complete continuity and weak compactness of weighted composition operators on derivative Hardy spaces. Recently, Lin, et al. [32] showed the boundedness of Volterra type operators on derivative Hardy spaces. Other intriguing issues about derivative Hardy spaces have been studied, including linear isometries [36], invariant subspace problems [7, 8, 29, 32], order boundedness of weighted composition operators [31, 33] and so forth. For more results about the derivative Hardy spaces, we refer the readers to [2, 9, 16, 18, 22, 23] and the references therein.

For $0 < p < \infty$ and $-1 < \alpha < \infty$, the weighted Bergman space A_α^p consists of all $f \in H(\mathbb{D})$ such that

$$\|f\|_{A_\alpha^p}^p := \int_{\mathbb{D}} |f(w)|^p dA_\alpha(w) < \infty$$

where $dA(w) = (1/\pi)dxdy$ is the normalized Lebesgue area measure on \mathbb{D} and $dA_\alpha(w) = (1 + \alpha)(1 - |w|^2)^\alpha dA(w)$ is the weighted Lebesgue measure (See [10] or [43] for more information about Bergman spaces).

Definition 1.2 For $0 < p < \infty$ and $-1 < \alpha < \infty$, the weighted Dirichlet space \mathcal{D}_α^p consists of all $f \in H(\mathbb{D})$ such that

$$\|f\|_{\mathcal{D}_\alpha^p} := \left(|f(0)|^p + \int_{\mathbb{D}} |f'(w)|^p dA_\alpha(w) \right)^{1/p} < \infty.$$

It is obvious that for any $f \in H(\mathbb{D})$, $f \in \mathcal{D}_\alpha^p$ if and only if $f' \in A_\alpha^p$. When $p < \alpha + 1$, the weighted Dirichlet space \mathcal{D}_α^p coincides with the weighted Bergman space $A_{\alpha-p}^p$ with equivalent norms. If $p > \alpha + 2$, the weighted Dirichlet space \mathcal{D}_α^p is contained in the essentially bounded space H^∞ (see [41, Theorem 4.2]).

In 1999, Wu [41] gave the Carleson measure characterization for the weighted Dirichlet space \mathcal{D}_α^p when $p \geq \alpha + 1$. In addition, he provided a sufficient and necessary condition, in terms of Carleson measures, for boundedness of multiplication operators on such weighted Dirichlet spaces. Related studies also appeared in the work of Arcozzi-Rochberg-Sawyer [3]. Continuing their researches, Girela and Peláez [12] obtained complete characterizations, in terms of Carleson measures, of conditions that enable the weighted Dirichlet spaces \mathcal{D}_α^p to be embedded into the Lebesgue spaces $L^q(d\mu)$ for $q > p > 0$, where $d\mu$ is a positive Borel measure on \mathbb{D} . Later, the characterizations of boundedness and compactness of multiplication operators and some integration operators on weighted Dirichlet spaces were obtained in [11, 30]. See [13] and [14] for related studies about weighted Dirichlet spaces.

Now, we state our main results. Theorem 1.1 and 1.2 show two sufficient conditions for all solutions of the equation (1.1) to lie in weighted Dirichlet spaces. Theorems 1.3 and 1.4 show two sufficient conditions for all solutions of the equation (1.1) to lie in derivative Hardy spaces.

Theorem 1.1 Let $0 < p < \infty$ and $\alpha > -1$. Let n be a positive integer and $A_i \in H(\mathbb{D})$, $i = 0, 1, \dots, n$. Assume that the following statements hold:

$$C_1 := \int_{\mathbb{D}} \left| \int_0^z \int_0^{\xi_1} \dots \int_0^{\xi_{n-2}} A_n(\xi_{n-1}) d\xi_{n-1} \dots d\xi_1 \right|^p dA_\alpha(z) < \infty,$$

$$C_2 := \int_{\mathbb{D}} \left(\sum_{m=1}^{n-1} \int_0^z \int_0^{\xi_1} \dots \int_0^{\xi_{m-1}} \frac{\left| \sum_{k=1}^m A_{n-k}^{(m-k)}(\xi_m) \right|}{(1 - |\xi_m|^2)^{\frac{\alpha+2}{p}}} d\xi_m \dots d\xi_1 \right)^p dA_\alpha(z)$$

and

$$C_3 := \begin{cases} \int_{\mathbb{D}} \left(\int_0^z \int_0^{\xi_1} \dots \int_0^{\xi_{n-2}} \frac{|A_0(\xi_{n-1})|}{(1-|\xi_{n-1}|^2)^{\frac{\alpha+2-p}{p}}} d\xi_{n-1} \dots d\xi_1 \right)^p dA_\alpha(z), & \text{if } p < \alpha + 2; \\ \int_{\mathbb{D}} \left(\int_0^z \int_0^{\xi_1} \dots \int_0^{\xi_{n-2}} |A_0(\xi_{n-1})| \left(\log \frac{2}{1-|\xi_{n-1}|^2} \right)^{\frac{p-1}{p}} d\xi_{n-1} \dots d\xi_1 \right)^p dA_\alpha(z), & \text{if } p = \alpha + 2; \\ \int_{\mathbb{D}} \left(\int_0^z \int_0^{\xi_1} \dots \int_0^{\xi_{n-2}} |A_0(\xi_{n-1})| d\xi_{n-1} \dots d\xi_1 \right)^p dA_\alpha(z), & \text{if } p > \alpha + 2. \end{cases}$$

are two positive constants satisfying $C_2 + C_3 < 1/d$, where $d > 0$ is sufficiently large and is only related to n, p . Then all solutions of the equation (1.1) lie in \mathcal{D}_α^p .

Theorem 1.2 Let $0 < p < \infty$ and $\alpha > -1$. Let n be a positive integer and $A_i \in H(\mathbb{D})$, $i = 0, 1, \dots, n$. Assume that the following statements hold:

$$K_1 := \int_{\mathbb{D}} |A_n(z)|^p (1 - |z|^2)^{pn-p} dA_\alpha(z) < \infty,$$

$$K_2 := \begin{cases} \int_{\mathbb{D}} |A_0(z)|^p (1 - |z|^2)^{pn-\alpha-2} dA_\alpha(z), & \text{if } p < \alpha + 2; \\ \int_{\mathbb{D}} |A_0(z)|^p (1 - |z|^2)^{pn-p} \left(\log \frac{2}{1-|z|^2} \right)^{p-1} dA_\alpha(z), & \text{if } p = \alpha + 2; \\ \int_{\mathbb{D}} |A_0(z)|^p (1 - |z|^2)^{pn-p} dA_\alpha(z), & \text{if } p > \alpha + 2, \end{cases}$$

and

$$K_3 := \sum_{i=1}^{n-1} \|A_i\|_{H_{n-i}^\infty}^p$$

are two positive constants satisfying $K_2 + K_3 < 1/c$, where $c > 0$ is sufficiently large and is only related to n, p . Then all solutions of the equation (1.1) lie in \mathcal{D}_α^p .

Theorem 1.3 Let $0 < p < \infty$. Let n be a positive integer and $A_i \in H(\mathbb{D})$, $i = 0, 1, \dots, n$. Assume that the following statements hold:

$$C_1 := \sup_{0 < r < 1} \frac{1}{2\pi} \int_0^{2\pi} \left| \int_0^{re^{i\theta}} \int_0^{\xi_1} \dots \int_0^{\xi_{n-2}} A_n(\xi_{n-1}) d\xi_{n-1} \dots d\xi_1 \right|^p d\theta < \infty,$$

$$C_2 := \sup_{0 < r < 1} \frac{1}{2\pi} \int_0^{2\pi} \left(\sum_{m=1}^{n-1} \int_0^{re^{i\theta}} \int_0^{\xi_1} \dots \int_0^{\xi_{m-1}} \frac{|\sum_{k=1}^m A_{n-k}(\xi_m)|}{(1-|\xi_m|^2)^{1/p}} d\xi_m \dots d\xi_1 \right)^p d\theta$$

and

$$C_3 := \begin{cases} \sup_{0 < r < 1} \frac{1}{2\pi} \int_0^{2\pi} \left(\int_0^{re^{i\theta}} \int_0^{\xi_1} \dots \int_0^{\xi_{n-2}} \frac{|A_0(\xi_{n-1})|}{(1-|\xi_{n-1}|^2)^{\frac{1-p}{p}}} d\xi_{n-1} \dots d\xi_1 \right)^p d\theta, & \text{if } 0 < p < 1; \\ \sup_{0 < r < 1} \frac{1}{2\pi} \int_0^{2\pi} \left(\int_0^{re^{i\theta}} \int_0^{\xi_1} \dots \int_0^{\xi_{n-2}} |A_0(\xi_{n-1})|^p d\xi_{n-1} \dots d\xi_1 \right) d\theta, & \text{if } 1 \leq p < \infty. \end{cases}$$

are two positive constants satisfying $C_2 + C_3 < 1/d$, where $d > 0$ is sufficiently large and is only related to n, p . Then all solutions of the equation (1.1) lie in S^p .

Theorem 1.4 Let $0 < p < \infty$. Let n be a positive integer and $A_i \in H(\mathbb{D}), i = 0, 1, \dots, n$. Assume that the following statements hold:

$$K_1 := \int_{\partial\mathbb{D}} \left(\int_{S(\xi)} |A_n(z)|^2 (1 - |z|^2)^{2n-4} dA(z) \right)^{p/2} dm(\xi) < \infty,$$

$$K_2 := \begin{cases} \int_{\partial\mathbb{D}} \left(\int_{S(\xi)} |A_0(z)|^2 (1 - |z|^2)^{2(n-1-\frac{1}{p})} dA(z) \right)^{p/2} dm(\xi), & \text{if } 0 < p < 1; \\ \int_{\partial\mathbb{D}} \left(\int_{S(\xi)} |A_0(z)|^2 (1 - |z|^2)^{2n-4} dA(z) \right)^{p/2} dm(\xi), & \text{if } 1 \leq p < \infty, \end{cases}$$

and

$$K_3 := \sum_{i=1}^{n-1} \min \left\{ \|A_i\|_{H_{n-i}^\infty}^p, \int_{\partial\mathbb{D}} \left(\int_{S(\xi)} |A_i(z)|^2 (1 - |z|^2)^{2(n-j-1-\frac{1}{p})} dA(z) \right)^{p/2} dm(\xi) \right\}$$

are two positive constants satisfying $K_2 + K_3 < 1/c$, where $c > 0$ is sufficiently large and is only related to n, p . Then all solutions of the equation (1.1) lie in S^p .

The structure of this paper is organized as follows.

In Section 2, we collect some preliminary lemmas that will be used throughout the paper. In Section 3, we prove our main results.

Throughout this paper, for any two positive functions $f(x)$ and $g(x)$, we write $f \lesssim g$ if $f \leq cg$ holds, where c is a positive constant independent of the variable x . We write $f \approx g$ whenever $f \lesssim g \lesssim f$. Moreover, the value of “ c ” may vary from line to line but will remain independent of the main variables.

2 Preliminaries

First, we need the following two lemmas, which will be used frequently later.

Lemma 2.1 [9] Suppose that N is a positive integer and $b_n \geq 0$ for $n = 1, 2, \dots, N$. Then

$$\left(\sum_{n=1}^N b_n \right)^p \leq \sum_{n=1}^N b_n^p, \quad 0 < p \leq 1$$

and

$$\left(\sum_{n=1}^N b_n \right)^p \leq N^{p-1} \left(\sum_{n=1}^N b_n^p \right), \quad 1 \leq p < \infty.$$

Lemma 2.2 [43, Theorem 4.28] Suppose $p > 0, \alpha > -1, n \geq 0$, and $f \in H(\mathbb{D})$. Then

$$\|f\|_{A_\alpha^p} \approx \sum_{i=0}^{n-1} |f^{(i)}(0)|^p + \int_{\mathbb{D}} |f^{(n)}(z)|^p (1 - |z|^2)^{n p} dA_\alpha(z).$$

Moreover, we need to use the following equivalent norms of H^p :

Lemma 2.3 ([1, p. 125]) Suppose $0 < p < \infty$ and $f \in H^p$. Then

$$\|f\|_{H^p}^p \approx \sum_{i=0}^{n-1} |f^{(i)}(0)|^p + \int_{\partial\mathbb{D}} \left(\int_{S(\xi)} |f^{(n)}(z)|^2 (1-|z|^2)^{2n-2} dA(z) \right)^{p/2} dm(\xi).$$

Next, the growth estimates of elements in S^p and \mathcal{D}_α^p are given in the following two lemmas, respectively.

Lemma 2.4 (see [32, 33]). If $1 \leq p < \infty$, then for any $f \in S^p$, it holds that $|f(z)| \leq \pi \|f\|_{S^p}$, $z \in \mathbb{D}$. If $0 < p < 1$, then for any $f \in S^p$,

$$|f(z)| \lesssim \frac{\|f\|_{S^p}}{(1-|z|^2)^{1/p-1}}, \quad \forall z \in \mathbb{D}.$$

Lemma 2.5 [41] Let $0 < p < \infty$ and $\alpha > -1$. If $f \in \mathcal{D}_\alpha^p$, then

- (1) $|f(z)| \lesssim \frac{\|f\|_{\mathcal{D}_\alpha^p}}{(1-|z|^2)^{\frac{\alpha+2-p}{p}}}$, whenever $p < \alpha + 2$;
- (2) $|f(z)| \lesssim \left(\log \frac{2}{1-|z|^2} \right)^{\frac{p-1}{p}} \|f\|_{\mathcal{D}_\alpha^p}$, whenever $p = \alpha + 2$;
- (3) $|f(z)| \leq \|f\|_{\mathcal{D}_\alpha^p}$, whenever $p > \alpha + 2$.

In addition, we have to use the following growth estimates for the n -th order derivative function.

Lemma 2.6 Let $0 < p < \infty, \alpha > -1$ and n be a positive integer. If $f \in \mathcal{D}_\alpha^p$, then

$$|f^{(n)}(z)| \lesssim \frac{\|f\|_{\mathcal{D}_\alpha^p}}{(1-|z|^2)^{\frac{2+\alpha}{p}+n-1}}, \quad z \in \mathbb{D}.$$

Proof For any $z \in \mathbb{D}$, by [34, Lemma 2.1], we know

$$|f^{(n-1)}(z)|^p \lesssim \frac{\int_{D_r(z)} |f(w)|^p dA_\alpha(w)}{(1-|z|^2)^{2+\alpha+(n-1)p}} \lesssim \frac{\int_{\mathbb{D}} |f(w)|^p dA_\alpha(w)}{(1-|z|^2)^{2+\alpha+(n-1)p}}.$$

Letting $f = g'$ yields

$$|g^{(n)}(z)|^p \lesssim \frac{\int_{\mathbb{D}} |g'(w)|^p dA_\alpha(w)}{(1-|z|^2)^{2+\alpha+(n-1)p}},$$

which is the desired result. ■

By Lemma 2.3, it is not difficult to obtain the following result.

Lemma 2.7 Let $0 < p < \infty$ and n be a non-negative integer. If $f \in S^p$, then

$$|f^{(n)}(z)| \lesssim \frac{\|f\|_{S^p}}{(1-|z|^2)^{1/p+n-1}}, \quad z \in \mathbb{D}.$$

3 The proof of the main results

Proof of Theorem 1.1: Assume that f is a solution of equation (1.1), then

$$f_r^{(n)}(z) + \sum_{j=0}^{n-1} B_j(z) f_r^{(j)}(z) = B_n(z), \quad z \in \mathbb{D}, \tag{3.1}$$

where $f_r(z) = f(rz)$, $B_j(z) = B_j(z, r) = r^{n-j} A_j(rz)$, $j = 0, \dots, n - 1$, $B_n(z) = r^n A_n(rz)$, $0 \leq r < 1$.

By using the equation

$$f(z) = \int_0^z f'(\xi) d\xi + f(0)$$

$n - 1$ times, we get

$$\begin{aligned} f_r'(z) &= \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} f_r^{(n)}(\xi_{n-1}) d\xi_{n-1} \cdots d\xi_1 + \sum_{j=0}^{n-2} \frac{f_r^{(j+1)}(0)}{j!} z^j \\ &= \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} \left(B_n(\xi_{n-1}) - \sum_{j=0}^{n-1} B_j(\xi_{n-1}) f_r^{(j)}(\xi_{n-1}) \right) d\xi_{n-1} \cdots d\xi_1 + \sum_{j=0}^{n-2} \frac{f_r^{(j+1)}(0)}{j!} z^j \end{aligned}$$

Combining this with Lemma 2.1, we obtain

$$\begin{aligned} \|f_r\|_{\mathcal{D}_\alpha^p}^p &= \int_{\mathbb{D}} |f_r'(z)|^p dA_\alpha(z) + |f_r(0)|^p \\ &= \int_{\mathbb{D}} \left| \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} \left(B_n(\xi_{n-1}) - \sum_{j=0}^{n-1} B_j(\xi_{n-1}) f_r^{(j)}(\xi_{n-1}) \right) d\xi_{n-1} \cdots d\xi_1 \right. \\ &\quad \left. + \sum_{j=0}^{n-2} \frac{f_r^{(j+1)}(0)}{j!} z^j \right|^p dA_\alpha(z) + |f_r(0)|^p \\ &\lesssim \int_{\mathbb{D}} \left| \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} B_n(\xi_{n-1}) d\xi_{n-1} \cdots d\xi_1 \right|^p dA_\alpha(z) \\ &\quad + \int_{\mathbb{D}} \left| \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} \sum_{j=1}^{n-1} B_j(\xi_{n-1}) f_r^{(j)}(\xi_{n-1}) d\xi_{n-1} \cdots d\xi_1 \right|^p dA_\alpha(z) \\ &\quad + \int_{\mathbb{D}} \left| \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} B_0(\xi_{n-1}) f_r(\xi_{n-1}) d\xi_{n-1} \cdots d\xi_1 \right|^p dA_\alpha(z) \\ &\quad + \int_{\mathbb{D}} \left| \sum_{j=0}^{n-2} \frac{f_r^{(j+1)}(0)}{j!} z^j \right|^p dA_\alpha(z) + |f_r(0)|^p \\ &= I_1 + I_2 + I_3 + I_4 + I_5. \end{aligned}$$

Let us first discuss I_2 . From [25, Lemma 12], we have

$$I_2 = \int_{\mathbb{D}} \left| \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} \sum_{j=1}^{n-1} \sum_{i=0}^{j-1} (-1)^i \binom{j-1}{i} (f'_r \cdot B_j^{(i)})^{(j-i-1)}(\xi_{n-1}) d\xi_{n-1} \cdots d\xi_1 \right|^p dA_\alpha(z)$$

As

$$\begin{aligned} & \int_0^{\xi_{n-1-(j-i-1)}} \int_0^{\xi_{n-(j-i-1)}} \cdots \int_0^{\xi_{n-2}} (f'_r \cdot B_j^{(i)})^{(j-i-1)}(\xi_{n-1}) d\xi_{n-1} \cdots d_{n-(j-i-1)} \\ &= \int_0^{\xi_{n-1-(j-i-1)}} \int_0^{\xi_{n-(j-i-1)}} \cdots \int_0^{\xi_{n-3}} \left[(f'_r \cdot B_j^{(i)})^{(j-i-2)}(\xi_{n-2}) - (f'_r \cdot B_j^{(i)})^{(j-i-2)}(0) \right] d\xi_{n-2} \cdots \\ &= B_j^{(i)}(\xi_{n-1-(j-i-1)}) f'_r(\xi_{n-1-(j-i-1)}) - \sum_{t=0}^{j-i-2} \frac{(B_j^{(i)} f'_r)^{(t)}(0)}{t!} \xi_{n-1-(j-i-1)}^t, \end{aligned}$$

then

$$\begin{aligned} I_2 &= \int_{\mathbb{D}} \left| \sum_{j=1}^{n-1} \sum_{i=0}^{j-1} (-1)^i \binom{j-1}{i} \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2-(j-i-1)}} \left[B_j^{(i)}(\xi_{n-1-(j-i-1)}) f'_r(\xi_{n-1-(j-i-1)}) \right. \right. \\ &\quad \left. \left. - \sum_{t=0}^{j-i-2} \frac{(B_j^{(i)} f'_r)^{(t)}(0)}{t!} \xi_{n-1-(j-i-1)}^t \right] d\xi_{n-1-(j-i-1)} \cdots d\xi_1 \right|^p dA_\alpha(z). \end{aligned}$$

We relabel the indices as follows:

$$j = n - k, \quad i = m - k.$$

It follows that

$$\begin{aligned} I_2 &= \int_{\mathbb{D}} \left| \sum_{m=1}^{n-1} \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{m-1}} \left[\sum_{k=1}^m (-1)^{m-k} \binom{n-k-1}{m-k} B_{n-k}^{(m-k)}(\xi_m) \right] f'_r(\xi_m) d\xi_m \cdots d\xi_1 \right. \\ &\quad \left. - \sum_{j=1}^{n-1} \sum_{i=0}^{j-1} \sum_{t=0}^{j-i-2} (-1)^i \binom{j-1}{i} \frac{(B_j^{(i)} f'_r)^{(t)}(0)}{(n-j+i+t)!} z^{n-j+i+t} \right|^p dA_\alpha(z) \\ &\lesssim \int_{\mathbb{D}} \left| \sum_{m=1}^{n-1} \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{m-1}} \left[\sum_{k=1}^m B_{n-k}^{(m-k)}(\xi_m) \right] f'_r(\xi_m) d\xi_m \cdots d\xi_1 \right|^p dA_\alpha(z) \\ &\quad + \int_{\mathbb{D}} \left| \sum_{j=1}^{n-1} \sum_{i=0}^{j-1} \sum_{t=0}^{j-i-2} \frac{(B_j^{(i)} f'_r)^{(t)}(0)}{(n-j+i+t)!} z^{n-j+i+t} \right|^p dA_\alpha(z) \\ &= I_{21} + I_{22}. \end{aligned}$$

By Lemma 2.6, we have

$$\begin{aligned}
 I_{21} &\leq \int_{\mathbb{D}} \left(\sum_{m=1}^{n-1} \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{m-1}} \left| \sum_{k=1}^m B_{n-k}^{(m-k)}(\xi_m) \right| |f_r'(\xi_m)| d\xi_m \cdots d\xi_1 \right)^p dA_\alpha(z) \\
 &\lesssim \|f_r\|_{\mathcal{D}_\alpha^p}^p \int_{\mathbb{D}} \left(\sum_{m=1}^{n-1} \int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{m-1}} \frac{\left| \sum_{k=1}^m B_{n-k}^{(m-k)}(\xi_m) \right|}{(1-|\xi_m|^2)^{\frac{\alpha+2}{p}}} d\xi_m \cdots d\xi_1 \right)^p dA_\alpha(z) \\
 &\lesssim \|f_r\|_{\mathcal{D}_\alpha^p}^p C_2.
 \end{aligned}$$

On the other hand, we have

$$I_4 + I_{22} = \sup_{z \in \mathbb{D}} \left(\left| \sum_{j=0}^{n-2} \frac{f_r^{(j+1)}(0)}{j!} z^j \right|^p + \left| \sum_{j=1}^{n-1} \sum_{i=0}^{j-1} \sum_{t=0}^{j-i-2} \frac{B_j^{(i)} f_r^{(t)}(0)}{(n-j+i+t)!} z^{n-j+i+t} \right|^p \right) \leq C_f < \infty.$$

Using Lemma 2.5 again, we have

$$I_3 \lesssim \int_{\mathbb{D}} \left(\int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} |B_0(\xi_{n-1})| |f_r(\xi_{n-1})| d\xi_{n-1} \cdots d\xi_1 \right)^p dA_\alpha(z) \lesssim \|f_r\|_{\mathcal{D}_\alpha^p}^p C_3,$$

where

$$C_3 = \begin{cases} \int_{\mathbb{D}} \left(\int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} \frac{|B_0(\xi_{n-1})|}{(1-|\xi_{n-1}|^2)^{\frac{\alpha+2-p}{p}}} d\xi_{n-1} \cdots d\xi_1 \right)^p dA_\alpha(z), & \text{if } p < \alpha + 2; \\ \int_{\mathbb{D}} \left(\int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} |B_0(\xi_{n-1})| \left(\log \frac{2}{1-|\xi_{n-1}|^2} \right)^{\frac{p-1}{p}} d\xi_{n-1} \cdots d\xi_1 \right)^p dA_\alpha(z), & \text{if } p = \alpha + 2; \\ \int_{\mathbb{D}} \left(\int_0^z \int_0^{\xi_1} \cdots \int_0^{\xi_{n-2}} |B_0(\xi_{n-1})| d\xi_{n-1} \cdots d\xi_1 \right)^p dA_\alpha(z) & \text{if } p > \alpha + 2. \end{cases}$$

Consequently,

$$\|f_r\|_{\mathcal{D}_\alpha^p}^p \lesssim \frac{C_1 + I_{22} + I_4 + I_5}{1 - d(C_2 + C_3)} < \infty,$$

which gives that $f \in \mathcal{D}_\alpha^p$ as $r \rightarrow 1^-$. This completes the proof. □

Proof of Theorem 1.2: Assume that f is a solution of equation (1.1), then we have

$$f_r^{(n)}(z) + \sum_{j=0}^{n-1} B_j(z) f_r^{(j)}(z) = B_n(z), \quad z \in \mathbb{D},$$

where $f_r(z) = f(rz)$, $B_j(z) = B_j(z, r) = r^{n-j} A_j(rz)$, $j = 0, \dots, n-1$, $B_n(z) = r^n A_n(rz)$, $0 \leq r < 1$.

By Lemma 2.1 and Lemma 2.2, we have

$$\begin{aligned} \|f_r\|_{\mathcal{D}_\alpha^p}^p &\approx \int_{\mathbb{D}} |f_r^{(n)}(z)|^p (1 - |z|^2)^{pn-p} dA_\alpha(z) + \sum_{k=0}^{n-1} |f_r^{(k)}(0)|^p \\ &= \int_{\mathbb{D}} \left| B_n(z) - \sum_{j=0}^{n-1} B_j(z) f_r^{(j)}(z) \right|^p (1 - |z|^2)^{pn-p} dA_\alpha(z) + \sum_{k=0}^{n-1} |f_r^{(k)}(0)|^p \\ &\leq \int_{\mathbb{D}} \left(|B_n(z)| + |B_0(z) f_r(z)| + \sum_{j=1}^{n-1} |B_j(z) f_r^{(j)}(z)| \right)^p (1 - |z|^2)^{pn-p} dA_\alpha(z) + \sum_{k=0}^{n-1} |f_r^{(k)}(0)|^p \\ &\lesssim \int_{\mathbb{D}} |B_n(z)|^p (1 - |z|^2)^{pn-p} dA_\alpha(z) + \int_{\mathbb{D}} |B_0(z) f_r(z)|^p (1 - |z|^2)^{pn-p} dA_\alpha(z) \\ &\quad + \sum_{j=1}^{n-1} \int_{\mathbb{D}} |B_j(z) f_r^{(j)}(z)|^p (1 - |z|^2)^{pn-p} dA_\alpha(z) + \sum_{k=0}^{n-1} |f_r^{(k)}(0)|^p \\ &= I_1 + I_2 + I_3 + I_4. \end{aligned}$$

Applying Lemma 2.5 leads to

$$I_2 \lesssim \begin{cases} \|f_r\|_{\mathcal{D}_\alpha^p}^p \int_{\mathbb{D}} |B_0(z)|^p (1 - |z|^2)^{pn-\alpha-2} dA_\alpha(z), & \text{if } p < \alpha + 2; \\ \|f_r\|_{\mathcal{D}_\alpha^p}^p \int_{\mathbb{D}} |B_0(z)|^p (1 - |z|^2)^{pn-p} \left(\log \frac{2}{1-|z|^2} \right)^{p-1} dA_\alpha(z), & \text{if } p = \alpha + 2; \\ \|f_r\|_{\mathcal{D}_\alpha^p}^p \int_{\mathbb{D}} |B_0(z)|^p (1 - |z|^2)^{pn-p} dA_\alpha(z) & \text{if } p > \alpha + 2. \end{cases}$$

On the other hand,

$$\begin{aligned} I_3 &\leq \sum_{j=1}^{n-1} \left(\sup_{z \in \mathbb{D}} |B_j(z)|^p (1 - |z|^2)^{p(n-j)} \right) \int_{\mathbb{D}} |f_r^{(j)}(z)|^p (1 - |z|^2)^{pj-p} dA_\alpha(z) \\ &\leq \sum_{j=1}^{n-1} \|B_j\|_{H_{n-j}^\infty}^p \|f_r\|_{\mathcal{D}_\alpha^p}^p. \end{aligned}$$

Consequently, $I_3 \lesssim K_3 \|f_r\|_{\mathcal{D}_\alpha^p}^p$. It follows from the assumption that

$$\|f_r\|_{\mathcal{D}_\alpha^p}^p \lesssim \frac{K_1 + I_4}{1 - c(K_2 + K_3)} < \infty,$$

for $0 \leq r < 1$. Letting $r \rightarrow 1^-$ gives that $f \in \mathcal{D}_\alpha^p$. This completes the proof. □

Remark 3.1 Although the sufficient condition in Theorem 1.1 and 1.2 is more complicated, in fact, we can illustrate the feasibility of that sufficient condition by some examples. Let us consider the complex second-order equation

$$f'' + A(z)f = 0.$$

Fix a constant-valued function $A(z) = k (k \in \mathbb{R})$ satisfying the condition

$$\int_{\mathbb{D}} |A_0(z)|^p (1 - |z|^2)^p dA_\alpha(z) < \frac{1}{c}.$$

If $k < 0$, we can easily find that the solution to the equation $f'' + kf = 0$ has a solution base $\{f_1, f_2\}$, where

$$f_1(z) = e^{\sqrt{-k}z} \text{ and } f_2(z) = e^{-\sqrt{-k}z}.$$

If $k > 0$, then the equation $f'' + kf = 0$ has a solution base $\{f_1, f_2\}$, where

$$f_3(z) = \cos(kz) \text{ and } f_4(z) = \sin(kz).$$

It can be seen that f_1, f_2, f_3 and f_4 belongs to the weighted Dirichlet spaces $\mathcal{D}_\alpha^p (p > \alpha + 2)$.

On the other hand, if we consider the equation $f^{(n)} = 0$, then it is easy to know that all solutions of this equation are

$$f(z) = a_0 + a_1z + a_2z^2 + a_3z^3 + \dots + a_{n-1}z^{n-1}.$$

Obviously, $f \in \mathcal{D}_\alpha^p$.

Proof of Theorem 1.3: The proof can be accomplished by using Lemma 2.1, Lemma 2.4 and Lemma 2.7, and the similar proof of Theorem 1.1. □

Proof of Theorem 1.4: The proof can be accomplished by using Lemma 2.3, Lemma 2.4 and Lemma 2.7, and the similar proof of Theorem 1.2. □

The following two corollaries provide some stronger sufficient conditions making the solutions of the equation 1.1 lie in weighted Dirichlet spaces.

The first one is a variant of Theorem 1.1.

Corollary 3.2 Let $0 < p < \infty$ and $\alpha > -1$. Let n be a positive integer and $A_i \in H(\mathbb{D})$, $i = 0, 1, \dots, n$. Assume that the following statements hold:

$$C_1 := \int_{\mathbb{D}} \left| \int_0^z \int_0^{\xi_1} \dots \int_0^{\xi_{n-2}} A_n(\xi_{n-1}) d\xi_{n-1} \dots d\xi_1 \right|^p dA_\alpha(z) < \infty,$$

$$C_2 := \int_{\mathbb{D}} \left(\sum_{m=1}^{n-1} \int_0^z \int_0^{\xi_1} \dots \int_0^{\xi_{m-1}} \frac{\left| \sum_{k=1}^m A_{n-k}^{(m-k)}(\xi_m) \right|}{(1 - |\xi_m|^2)^{\frac{\alpha+2}{p}}} d\xi_m \dots d\xi_1 \right)^p dA_\alpha(z),$$

and

$$O_3 := \int_{\mathbb{D}} \left(\int_0^z \int_0^{\xi_1} \dots \int_0^{\xi_{n-2}} \frac{|A_0(\xi_{n-1})|}{(1 - |\xi_{n-1}|^2)^{\frac{\alpha+2+p}{p}}} d\xi_{n-1} \dots d\xi_1 \right)^p dA_\alpha(z)$$

are two positive constants satisfying $C_2 + O_3 < 1/d$, where $d > 0$ is sufficiently large and is only related to n, p and the growth of functions in the weighted Dirichlet spaces. Then all solutions of the equation (1.1) lie in \mathcal{D}_α^p .

The second one is a variant of Theorem 1.2.

Corollary 3.3 Let $0 < p < \infty$ and $\alpha > -1$. Let n be a positive integer and $A_i \in H(\mathbb{D})$, $i = 0, 1, \dots, n$. Assume that the following statements hold:

$$K_1 := \int_{\mathbb{D}} |A_n(z)|^p (1 - |z|^2)^{pn-p} dA_{\alpha}(z) < \infty,$$

$$Q_2 := \int_{\mathbb{D}} |A_0(z)|^p (1 - |z|^2)^{pn-p-\alpha-2} dA_{\alpha}(z),$$

and

$$Q_3 := \sum_{i=1}^{n-1} \int_{\mathbb{D}} |A_i(z)|^p (1 - |z|^2)^{p(n-i)-(2+\alpha)} dA_{\alpha}(z)$$

are two positive constants satisfying $Q_2 + Q_3 < 1/c$, where $c > 0$ is sufficiently large and is only related to n, p . Then all solutions of the equation (1.1) lie in \mathcal{D}_{α}^p .

Remark 3.4 Actually, in the proof of Theorem 1.2, by Lemma 2.6, we have

$$I_3 \lesssim \|f_r\|_{\mathcal{D}_{\alpha}^p}^p \sum_{j=1}^{n-1} \int_{\mathbb{D}} |B_j(z)|^p (1 - |z|^2)^{p(n-j)-(2+\alpha)} dA_{\alpha}(z) \leq Q_3 \|f_r\|_{\mathcal{D}_{\alpha}^p}^p.$$

It is worth noting that the condition Q_3 in the above corollary is stronger than the condition K_3 in Theorem 1.2, which can be obtained by applying [43, Proposition 4.13]

$$|f(z)|(1 - |z|^2)^{p(n-i)} \lesssim \int_{\mathbb{D}} |f(w)|^p (1 - |w|^2)^{p(n-j)-(2+\alpha)} dA_{\alpha}(w)$$

for any $f \in H(\mathbb{D})$.

Acknowledgments. The authors are grateful to the reviewer for his (or her) helpful comments. Qingze Lin is supported by STU Scientific Research Initiation Grant (No. NTF24015T).

References

- [1] P. Ahern, J. Bruna, *Maximal and area integral characterizations of Hardy-Sobolev spaces in the unit ball of C^n* , Rev. Mat. Iberoamericana 4 (1988), no. 1, 123–153.
- [2] R. F. Allen, K. C. Heller, M. A. Pons, *Multiplication operators on $S^2(\mathbb{D})$* , Acta Sci. Math. (Szeged) 81 (2015), no. 3-4, 575–587.
- [3] N. Arcozzi, R. Rochberg, E. Sawyer, *Carleson measures for analytic Besov spaces*, Rev. Mat. Iberoamericana 18 (2002), no. 2, 443–510.
- [4] D. Benbourenane, L. R. Sons, *On global solutions of complex differential equations in the unit disk*, Complex Var. Theory Appl. 49 (2004), no. 13, 913–925.
- [5] I. Chyzykhov, G. Gundersen, J. Heittokangas, *Linear differential equations and logarithmic derivative estimates*, Proc. London Math. Soc. (3) 86 (2003), no. 3, 735–754.
- [6] M. D. Contreras, A. G. Hernández-Díaz, *Weighted composition operators on spaces of functions with derivative in a Hardy space*, J. Operator Theory 52 (2004), no. 1, 173–184.
- [7] Ž. Čučković, B. Paudyal, *Invariant subspaces of the shift plus complex Volterra operator*, J. Math. Anal. Appl. 426 (2015), no. 2, 1174–1181.
- [8] Ž. Čučković, B. Paudyal, *The lattices of invariant subspaces of a class of operators on the Hardy space*, Arch. Math. (Basel) 110 (2018), no. 5, 477–486.

- [9] P. L. Duren, *Theory of H^p spaces*, Pure and Applied Mathematics, Vol. 38. Academic Press, New York-London, 1970.
- [10] P. L. Duren, A. Schuster, *Bergman spaces*, Mathematical Surveys and Monographs, 100. American Mathematical Society, Providence, RI, 2004.
- [11] P. Galanopoulos, D. Girela, J. Á. Peláez, *Multipliers and integration operators on Dirichlet spaces*, Trans. Amer. Math. Soc. **363** (2011), no. 4, 1855–1886.
- [12] D. Girela, J. Á. Peláez, *Carleson measures, multipliers and integration operators for spaces of Dirichlet type*, J. Funct. Anal. **241** (2006), no. 1, 334–358.
- [13] D. Girela, J. Á. Peláez, *Carleson measures for spaces of Dirichlet type*, Integral Equations Operator Theory **55** (2006), no. 3, 415–427.
- [14] D. Girela, J. Á. Peláez, *Growth properties and sequences of zeros of analytic functions in spaces of Dirichlet type*, J. Aust. Math. Soc. **80** (2006), no. 3, 397–418.
- [15] J. Gröhn, J. Huusko, J. Rättyä, *Linear differential equations with slowly growing solutions*, Trans. Amer. Math. Soc. **370** (2018), no. 10, 7201–7227.
- [16] C. Gu, S. Luo, *Composition and multiplication operators on the derivative Hardy space $S^2(\mathbb{D})$* , Complex Var. Elliptic Equ. **63** (2018), no. 5, 599–624.
- [17] G. G. Gundersen, E. M. Steinbart, S. Wang, *The possible orders of solutions of linear differential equations with polynomial coefficients*, Trans. Amer. Math. Soc. **350** (1998), no. 3, 1225–1247.
- [18] A. Gupta, B. Gupta, *On k -composition and k -Hankel composition operators on the derivative Hardy space*, Banach J. Math. Anal. **14** (2020), no. 4, 1602–1629.
- [19] J. Heittokangas, *On complex differential equations in the unit disc*, Dissertation, University of Joensuu, Joensuu, 2000. Ann. Acad. Sci. Fenn. Math. Diss. No. 122 (2000), 54 pp.
- [20] J. Heittokangas, R. Korhonen, J. Rättyä, *Growth estimates for solutions of linear complex differential equations*, Ann. Acad. Sci. Fenn. Math. **29** (2004), no. 1, 233–246.
- [21] J. Heittokangas, R. Korhonen, J. Rättyä, *Linear differential equations with coefficients in weighted Bergman and Hardy spaces*, Trans. Amer. Math. Soc. **360** (2008), no. 2, 1035–1055.
- [22] K. Heller, *Composition Operators on $S^2(\mathbb{D})$* , Thesis (Ph.D.)—University of Virginia. 2010. 143 pp.
- [23] K. Heller, *Adjoint of linear fractional composition operators on $S^2(\mathbb{D})$* , J. Math. Anal. Appl. **394** (2012), no. 2, 724–737.
- [24] G. Hu, J. Huusko, J. Long, Y. Sun, *Linear differential equations with solutions lying in weighted Fock spaces*, Complex Var. Elliptic Equ. **66** (2021), no. 2, 194–208.
- [25] J. Huusko, T. Korhonen, A. Reijonen, *Linear differential equations with solutions in the growth space H_{∞}^{∞}* , Ann. Acad. Sci. Fenn. Math. **41** (2016), no. 1, 399–416.
- [26] R. Korhonen, J. Rättyä, *Linear differential equations in the unit disc with analytic solutions of finite order*, Proc. Amer. Math. Soc. **135** (2007), no. 5, 1355–1363.
- [27] H. Li, S. Li, *Nonlinear differential equation and analytic function spaces*, Complex Var. Elliptic Equ. **63** (2018), no. 1, 136–149.
- [28] H. Li, H. Wulan, *Linear differential equations with solutions in the Q_K spaces*, J. Math. Anal. Appl. **375** (2011), no. 2, 478–489.
- [29] Q. Lin, *The invariant subspaces of the shift plus integer multiple of the Volterra operator on Hardy spaces*, Arch. Math. (Basel) **111** (2018), no. 5, 513–522.
- [30] Q. Lin, *Volterra type operators on weighted Dirichlet spaces*, Chinese Ann. Math. Ser. B **42** (2021), no. 4, 601–612.
- [31] Q. Lin, *Order boundedness of weighted composition operators between two classes of function spaces (Chinese)*, Acta Math. Sinica (Chinese Ser.) **65** (2022), no. 2, 317–324.
- [32] Q. Lin, J. Liu, Y. Wu, *Volterra type operators on $S^p(\mathbb{D})$ spaces*, J. Math. Anal. Appl. **461** (2018), no. 2, 1100–1114.
- [33] Q. Lin, J. Liu, Y. Wu, *Order boundedness of weighted composition operators on weighted Dirichlet spaces and derivative Hardy spaces*, Bull. Belg. Math. Soc. Simon Stevin **27** (2020), no. 4, 627–637.
- [34] D. H. Luecking, *Forward and reverse Carleson inequalities for functions in Bergman spaces and their derivatives*, Amer. J. Math. **107** (1985), no. 1, 85–111.
- [35] B. D. MacCluer, *Composition operators on S^p* , Houston J. Math. **13** (1987), no. 2, 245–254.
- [36] W. P. Novinger, D. M. Oberlin, *Linear isometries of some normed spaces of analytic functions*, Canad. J. Math. **37** (1985), no. 1, 62–74.

- [37] Ch. Pommerenke, *On the mean growth of the solutions of complex linear differential equations in the disk*, Complex Variables Theory Appl. **1** (1982/83), no. 1, 23–38.
- [38] R. C. Roan, *Composition operators on the space of functions with HP -derivative*, Houston J. Math. **4** (1978), no. 3, 423–438.
- [39] Y. Sun, J. Liu, G. Hu, *Complex linear differential equations with solutions in the H_K^2 spaces*, Complex Var. Elliptic Equ. **67** (2022), no. 11, 2577–2588.
- [40] Y. Sun, B. Liu, J. L. Liu, *Complex linear differential equations with solutions in Dirichlet-Morrey spaces*, Anal. Math. **49** (2023), no. 1, 295–306.
- [41] Z. Wu, *Carleson measures and multipliers for Dirichlet spaces*, J. Funct. Anal. **169** (1999), no. 1, 148–163.
- [42] L. Xiao, *Differential equations with solutions lying in the $F(p, q, s)$ space*, Complex Var. Elliptic Equ. **63** (2018), no. 1, 116–135.
- [43] K. Zhu, *Operator Theory in Function Spaces*, Second edition. Mathematical Surveys and Monographs, 138. Amer. Math. Soc, Providence. 2007.

Department of Mathematics, Shantou University, Shantou, China, 515063
e-mail: gdlqz@e.gzhu.edu.cn.

Department of Mathematics, Sun Yat-sen University, Guangzhou, China, 510275
e-mail: xiehy33@mail2.sysu.edu.cn.