

PERTURBED BILLIARD SYSTEMS II BERNOULLI PROPERTIES

IZUMI KUBO AND HIROSHI MURATA*)

§1. Introduction

One of the authors has shown the ergodicity of the perturbed billiard system which can describe the motion of a particle in a potential field of a special type [5], [6]. Since then, some development has been made, and we are now able to show the Bernoulli property of the system in this article. We hope, the result gives a new progress in statistical mechanics. Our method of the proof is inspired by the idea of D. S. Ornstein and B. Weiss [9], which has been used by G. Gallavotti and D. S. Ornstein [3] for a Sinai billiard system.

A perturbed billiard transformation will be prescribed in §3. Roughly speaking, it is an automorphism T_* of two dimensional measure space (M, ν) which can be expressed as the product of T_1 and T , where T_1 is a ν -preserving C^2 -diffeomorphism of M and where T is a Sinai billiard transformation. Such an automorphism T_* appears in a dynamical system of a particle moving in a potential field which is a composition of several finite range potentials (see [5], [6]). In order to discuss such a perturbed billiard system we need three assumptions (H-1)~(H-3), which specify the diffeomorphism T_1 . Under these assumptions, the perturbation of T by T_1 is not so much. Details of them will be found in §3.

Our main results are the following:

THEOREM 2. *Under the assumptions (H-1)~(H-3), partitions $\alpha^{(c)}$ and $\alpha^{(e)}$ are weak Bernoulli generators for T_* . Thus T_* is isomorphic to a Bernoulli shift.*

Here $\alpha^{(c)}$ and $\alpha^{(e)}$ are partitions of M whose elements are connected

Received June 7, 1975.

Revised December 4, 1979.

*) This author was partially supported by Yukawa Fellowship.

components such that the restrictions of T_*^{-1} and T_* on them are continuous, respectively.

THEOREM 3. *Under the assumptions (H-1)~(H-3), every countable partition α is weakly Bernoullian for T_* , whenever $\log d(x; \partial\alpha)$ is integrable.*

Here $d(x; \partial\alpha)$ is the distance between a point x and the union $\partial\alpha$ of the boundaries of all elements of α .

THEOREM 5. *Under the assumptions (H-1)~(H-3) and (f-1)~(f-3), if $\{S_t^f\}$ is a K -system and α is a finite partition such that $\log \tilde{d}(w; \partial\alpha)$ is integrable, then α is very weakly Bernoullian for $\{S_t^f\}$ ($t \neq 0$). Furthermore $\{S_t^f\}$ is a Bernoulli flow.*

As stated in Corollary 5.3, $\{S_t^f\}$ is a K -system if it does not have any point spectrum. With this result, we have a stronger assertion Corollary 6.2. Here $\{S_t^f\}$ is a flow of Kakutani-Ambrose type whose basic transformation and ceiling function are T_* and $f(x)$, respectively. There we assume the conditions (f-1)~(f-3), which are prescribed in § 5, so as $f(x)$ to be regular. Actually, if $f(x)$ is positive and smooth on M , then they are obviously satisfied. Our formulation is complicated, but necessary in order to apply to the case of dynamical system on a potential field as described above.

In section 2 some lemmas to make easier checking weak Bernoulli property will be given. In section 3 some fundamental results of the perturbed billiard transformation T_* , which have been shown in [5], will be summarized. In section 4 the proofs of Theorem 2 and Theorem 3 will be shown appealing lemmas in § 2. The most complicated parts of the proofs are in the estimations of densities of measures related to transversal fibres of T_* . In section 5 we will discuss on the construction of transversal fibres and the K -properties of the flow $\{S_t^f\}$. In section 6 the proof of Theorem 5 will be shown by using properties of the transversal fibres.

Lastly we remark that the same results can be obtained for more general T_1 as discussed in [6], since the properties stated in § 3 are also true for the general case.

§ 2. Weak and very weak Bernoulli partitions

Let (M, ν) be a Lebesgue space with total mass $\nu(M) = 1$, and let T

be a bimeasurable measure preserving transformation on M . D. S. Ornstein gave the following definitions:

DEFINITION 2.1. A countable partition α of M is said to be *weakly Bernoullian* for T , [if for any $\varepsilon > 0$ there exists $N \geq 0$ such that for all $N'' \geq N' \geq N$, all $n \geq 0$, and ε -a.e. B in $\bigvee_{i=N'}^{N''} T^{-i}\alpha$

$$(2.1) \quad \sum_{\substack{A \in \bigvee_{i=0}^n T^i \alpha}} |\nu(A) - \nu(A|B)| < \varepsilon .$$

Here “ ε -a.e. B in ξ ” means “except element B of the partition ξ which is included in a set of measure ε ”. For two countable partitions $\alpha = \{A_i\}$ and $\beta = \{B_i\}$ of M , define the usual metric by

$$d(\alpha, \beta) \equiv \sum_i \nu(A_i \ominus B_i) ,$$

where $A \ominus B$ denotes the symmetric difference of the sets A and B . For given two sequences of partitions $\{\alpha_j\}_1^n$ and $\{\beta_j\}_1^n$ write

$$\{\alpha_j\}_1^n \sim \{\beta_j\}_1^n$$

if for all $k_j, 1 \leq j \leq n$,

$$\mu_X \left(\bigcap_1^n A_{k_j}^{(j)} \right) = \mu_Y \left(\bigcap_1^n B_{k_j}^{(j)} \right) ,$$

where $\alpha_j = \{A_1^{(j)}, \dots, A_{a^{(j)}}^{(j)}\}$ are partitions on (X, μ_X) and $\beta_j = \{B_1^{(j)}, \dots, B_{b^{(j)}}^{(j)}\}$ are partitions on (Y, μ_Y) . Further define the metric \bar{d} by

$$\bar{d}(\{\alpha_j\}_1^n, \{\beta_j\}_1^n) = \inf_{\{\bar{\alpha}_j\}, \{\bar{\beta}_j\}} \frac{1}{n} \sum_1^n d(\bar{\alpha}_j, \bar{\beta}_j) ,$$

where $\{\bar{\alpha}_j\}$ and $\{\bar{\beta}_j\}$ run over all pairs of partitions on the same space such that $\{\alpha_j\}_1^n \sim \{\bar{\alpha}_j\}_1^n$ and $\{\beta_j\}_1^n \sim \{\bar{\beta}_j\}_1^n$. Let α be a partition and E be a subset of M . Then the normalized measure $\nu_E(A) = \nu(A \cap E)/\nu(E)$ will be associated to $\alpha|_E$.

DEFINITION 2.2. A finite partition α is said to be *very weakly Bernoullian* for T , if for every $\varepsilon > 0$ there exists $N \geq 0$ such that for all $N'' \geq N' \geq N$, all $n \geq 0$ and ε -a.e. B in $\bigvee_{i=N'}^{N''} T^{-i}\alpha$,

$$(2.2) \quad \bar{d}(\{T^i \alpha\}_1^n, \{T^i \alpha|B\}_1^n) \leq \varepsilon .$$

D. S. Ornstein and others [2], [4], [7], [8] have shown the following theorem:

THEOREM A. *If one of the following conditions is satisfied, then T is isomorphic to a Bernoulli shift.*

- (i) *There is a weak Bernoulli generator for T .*
- (ii) *There is a sequence of weak Bernoulli partitions α_n for T such that $\bigvee_{i=-\infty}^{\infty} T^i \alpha_n \uparrow \epsilon^*$ as $n \rightarrow \infty$.*
- (iii) *There is a sequence of very weak Bernoulli partitions α_n for T such that $\bigvee_{i=-\infty}^{\infty} T^i \alpha_n \uparrow \epsilon$ as $n \rightarrow \infty$.*

In order to apply this theorem to a perturbed billiard system, it is convenient to prepare the following lemmas.

LEMMA 2.3. (i) *If for any $\epsilon > 0$ and $\delta > 0$, there exist a natural number N and a finite family \mathcal{F} of disjoint subsets of M with $\sum_{F \in \mathcal{F}} \nu(F) \geq 1 - \delta$ such that for all $N'' \geq N' \geq N$, all $n \geq 0$ and ϵ -a.e. B in $\bigvee_{i=N'}^{N''} T^{-i} \alpha$,*

$$(2.3) \quad \sum_{\substack{A \in \bigvee_{i=0}^n T^i \alpha \\ i=0}} |\nu(A|F) - \nu(A|F \cap B)| < \delta \quad \text{for any } F \text{ in } \mathcal{F},$$

$$(2.4) \quad \sum_{F \in \mathcal{F}} |\nu(F) - \nu(F|B)| < \delta,$$

then the partition α is weakly Bernoullian for T .

(ii) *In (i), the condition (2.4) is unnecessary if T is a K -system and the entropy of α is finite.*

(iii) *In (i), (2.3) is fulfilled if for all A in $\bigvee_{i=0}^n T^i \alpha$, all F in \mathcal{F} and ϵ -a.e. B in $\bigvee_{i=N'}^{N''} T^{-i} \alpha$,*

$$(2.5) \quad \left| \frac{\nu(A \cap B \cap F)\nu(F)}{\nu(A \cap F)\nu(B \cap F)} - 1 \right| < \delta.$$

Proof. Put $F_0 = M - \bigcup_{F \in \mathcal{F}} F$, then one has

$$\begin{aligned} \nu(F_0|B) &\leq 1 - \sum_F \nu(F|B) \leq 1 - (1 - \delta) \sum_F \nu(F) \\ &\leq 1 - (1 - \delta)^2 \end{aligned}$$

by (2.4) and by $\sum \nu(F) \geq 1 - \delta$. From (2.3) the estimate

$$\begin{aligned} &\sum_F \sum_A |\nu(A \cap F) - \nu(A \cap F|B)| \\ &\leq \sum_{A, F} |\nu(A|F)\nu(F) - \nu(A|B \cap F)\nu(F|B)| \end{aligned}$$

^{*}) The symbol ϵ denotes the partition into the individual points.

$$\begin{aligned} &\leq \sum_{A,F} |\nu(A|F) - \nu(A|B \cap F)| \nu(F) + \sum_{A,F} \nu(A|B \cap F) |\nu(F) - \nu(F|B)| \\ &\leq 2\delta \end{aligned}$$

is obtained. Hence for 2ε -a.e. B in $\bigvee_{i=N'}^{N''} T^{-i}\alpha$,

$$\begin{aligned} &\sum_A |\nu(A) - \nu(A|B)| \\ &\leq \sum_{F,A} |\nu(A \cap F) - \nu(A \cap F|B)| + \sum_A \{\nu(A \cap F_0) + \nu(A \cap F_0|B)\} \\ &\leq 2\delta + \delta + 1 - (1 - \delta)^2 \leq 5\delta. \end{aligned}$$

Thus (i) is proved. If T is a K -system, then for the given \mathcal{F} there exists N such that for all $N'' \geq N' \geq N$, all F in \mathcal{F} and ε -a.e. B in $\bigvee_{i=N'}^{N''} T^{-i}\alpha$,

$$|\nu(F) - \nu(F|B)| \leq \delta\nu(F).$$

It is easily seen that (2.5) implies (2.3).

Q.E.D.

A mapping ϕ from X to Y is called ε -measure preserving if there exists a subset E of X with $\mu_x(E) \leq \varepsilon$ such that for all $A \subset X - E$

$$\left| \frac{\mu_y(\phi A)}{\mu_x(A)} - 1 \right| < \varepsilon.$$

Let $e(n)$ be the function on ordinal numbers defined by $e(0) = 0$, $e(n) = 1$ for $n \neq 0$. For a given partition $\alpha = \{A_j\}$, the name function of α is defined by $\ell(x) \equiv j$ if x is in A_j . The following lemma is due to D. S. Ornstein and B. Weiss [9].

LEMMA 2.4. Let $\{\alpha_i\}_1^n$ be partitions of X with name functions $\ell_i(x)$, and $\{\beta_i\}_1^n$ be partitions of Y with name functions $m_i(y)$. If there is an ε -measure preserving mapping ϕ from X to Y such that

$$\frac{1}{n} \sum_1^n e(\ell_i(x) - m_i(\phi x)) \leq \varepsilon$$

holds for x in $X - E$ with $\mu_x(E) \leq \varepsilon$, then

$$\bar{d}(\{\alpha_i\}_1^n, \{\beta_i\}_1^n) \leq 16\varepsilon.$$

The lemma is easily proved, but it is useful to check (2.2) for a suitable partition α (cf. § 6).

§ 3. Perturbed billiard systems

In the previous article by one of the authors [5], a perturbed billiard system was defined as follows. Let \bar{Q}_ι , $\iota = 1, 2, \dots, I$ be disjoint strictly

convex domains in a 2-dimensional torus T whose boundaries ∂Q_i are closed curves of C^3 -class. Put $Q = T - \bigcup_i \bar{Q}_i$ and $\partial Q = \bigcup_i \partial Q_i$ and put $M_0 = \{(q, p); q \in Q, p = (\cos \omega, \sin \omega), 0 \leq \omega < 2\pi\}$. The flow $\{S_t\}$ on M_0 which describes the motion of a particle moving around in Q with unit speed and with elastic collision at ∂Q is called a Sinai billiard system in Q [11], [12]; the particle moves along straight lines in the interior of Q with speed one, and is reflected at ∂Q according to the law "the angle of reflection is equal to the angle of incidence". Denote by M the set of all unit incident vectors at ∂Q . Then every element $x = (q, p)$ of M can be represented by coordinates (ι, r, φ) , where ι is the number of ∂Q_i containing q , r is the arclength between q and a fixed origin in ∂Q_i , measured along ∂Q_i clockwise, and φ is the angle between p and the inward normal of ∂Q_i at q . For x in M_0 , put

$$(3.1) \quad \begin{aligned} v(x) &= \inf \{t \geq 0; S_t x \text{ collides with } \partial Q\} \\ \tau(x) &= \sup \{t < 0; S_t x \text{ collides with } \partial Q\}. \end{aligned}$$

Then almost every point x in M_0 (with respect to the measure $dq d\omega$) is parametrized by (ι, r, φ, v) , where $v = v(x)$ and (ι, r, φ) represents the point $S_v x$ in M . One can define a transformation T of M , which is called a *Sinai billiard transformation*, by

$$(3.2) \quad Tx = S_{\tau(x)-v}x \quad \text{for } x \text{ in } M.$$

Then $\{S_{-t}\}$ is a *Kakutani-Ambrose flow* with the basic space M , the basic transformation T and the ceiling function $-\tau(x)$. The invariant measure μ of $\{S_t\}$ determined by Liouville's theorem is expressed in the form

$$(3.3) \quad d\mu = -\mu_0 \cos \varphi d\varphi dr dv d\iota$$

and the corresponding invariant measure of T is expressed in the form

$$(3.4) \quad d\nu = -\nu_0 \cos \varphi d\varphi dr d\iota$$

with $\mu_0 = (2\pi |Q|)^{-1}$ and $\nu_0 = (2|\partial Q|)^{-1}$, where $|Q|$ is the volume of Q and $|\partial Q|$ is the total arclength of the boundary ∂Q .

DEFINITION 3.1. A transformation T_* of M is called a *perturbed billiard transformation* if T_* is expressed in the form

$$(3.5) \quad T_* = T_1 T,$$

where T is the Sinai billiard transformation and T_1 is a C^2 -diffeomorphism

of M which preserves the measure ν and each $M^{(\iota)} \equiv \{(\iota, r, \varphi); (\iota, r) \in \partial Q_i\}$.

In [5], a special class of perturbed billiard transformations has been investigated*):

(H-1) $T_1(\iota, r, \varphi) = (\iota, r - H(\iota, \varphi), \varphi)$, where $H(\iota, \varphi)$ is a function of C^2 -class and satisfies $H(\iota, (\pi/2)) = H(\iota, (3/2)\pi) = 0$,

(H-2) every \bar{Q}_i are disjoint strictly convex domains whose boundaries are curves of C^3 -class,

(H-3) $\min_{\iota, \varphi} \{h(\iota, \varphi) + [\max_r k(\iota, r) + (\min_{\iota, r, \varphi'} |\tau(\iota, r, \varphi')|)^{-1}]^{-1}\} > 0$, where $h(\iota, \varphi) \equiv dH(\iota, \varphi)/d\varphi$ and $k(\iota, r)$ is the curvature of ∂Q_i at (ι, r) .

Under the above three assumptions the ergodicity and the K -property of the perturbed billiard transformation T_* were shown in [5]. In order to describe the results, it is necessary to introduce notation and terminology. A connected curve $\gamma; \varphi = \psi(r)$ in $M^{(\iota)}$ is called K -increasing if for $r \neq r'$

$$(3.6) \quad k_{\min} \leq \frac{\psi(r) - \psi(r')}{r - r'} \leq K_{\max}(\iota)$$

holds, where

$$k_{\min} \equiv \min_{\iota, r} k(\iota, r) \quad \text{and} \quad K_{\max}(\iota) \equiv \max_r k(\iota, r) + \left(\min_{r, \varphi} |\tau(T_*^{-1}(\iota, r, \varphi))| \right)^{-1}.$$

A connected curve $\gamma; \varphi = \psi(r)$ in $M^{(\iota)}$ is called K -decreasing, if for $r \neq r'$

$$(3.7) \quad K_{\min} \leq -\frac{\psi(r) - \psi(r')}{r - r'} \leq K_{\max}$$

holds, where $K_{\min} \equiv [\max_{\iota, \varphi} h(\iota, \varphi) + k_{\min}^{-1}]^{-1}$ and $K_{\max} \equiv \max_{\iota} [\min_{\varphi} h(\iota, \varphi) + K_{\max}(\iota)^{-1}]^{-1}$. Put $S \equiv \{(\iota, r, \varphi) \in M; \varphi = \pi/2 \text{ or } 3\pi/2\}$. Then $T_*^{-1}S$ (resp. T_*S) is called the curves of discontinuity of T_* (resp. T_*^{-1}). The image $T_*^{-1}S$ (resp. T_*S) consists of a countable number of K -increasing (resp. K -decreasing) curves. The curves of $T_*^{-1}S$ (resp. T_*S) decompose M into connected components and define the partition $\alpha^{(e)}$ (resp. $\alpha^{(c)}$) into the components $\{X_j^{(e)}\}$ (resp. $\{X_j^{(c)}\}$). Then T_* (resp. T_*^{-1}) is continuous in the interior of each component and belongs to C^2 -class. If T_* (resp. T_*^{-1}) is continuous on a connected K -decreasing (resp. K -increasing) curve γ , then so is the image of γ .

For a point $x = (\iota, r, \varphi)$ in M , put $x_i = (\iota_i, r_i, \varphi_i) \equiv T_*^{-1}(\iota, r, \varphi)$, $(\iota_i, r'_i, \varphi'_i) \equiv T_*^{-1}x_i$, $k_i \equiv k(\iota_i, r_i)$, $k'_i \equiv k(\iota_i, r'_i)$, $h_i \equiv h(\iota_i, \varphi_i)$ and $\tau_i \equiv \tau(\iota_i, r_i, \varphi_i)$. Define

*) A more general case was discussed in [6].

functions $b_n(x; t)$, $-\infty < n < \infty$, of (x, t) in $M \times (-\infty, \infty)$ by

$$\begin{aligned}
 b_1(x; t) &\equiv \frac{(\cos \varphi + k'\tau_1)(h + t) + \tau_1}{(k_1 \cos \varphi + k' \cos \varphi_1 + k_1 k' \tau_1)(h + t) + \cos \varphi_1 + k_1 \tau_1} \\
 b_{n+1}(x; t) &\equiv b_1(T_*^{-n}x; b_n(x; t)) \quad n \geq 1 \\
 b_0(x; t) &\equiv t \\
 b_{-1}(x; t) &\equiv -h - \frac{(\cos \varphi + kt)t - \tau}{(k \cos \varphi_{-1} + k'_{-1} \cos \varphi + k k'_{-1} \tau)t - (\cos \varphi_{-1} + k'_{-1} \tau)} \\
 b_{-n-1}(x; t) &\equiv b_{-1}(T_*^n x; b_{-n}(x; t)) \quad n \geq 1.
 \end{aligned}$$

Suppose that γ and $T_*^{-n}\gamma$ are given by the equations $r = u(\varphi)$ and $r_n = u_n(\varphi_n)$, respectively, with $T_*^{-n}(\iota, u(\varphi), \varphi) = (\iota_n, u_n(\varphi_n), \varphi_n)$. Then the formula

$$\frac{du_n(\varphi_n)}{d\varphi_n} = b_n\left(\iota, r, \varphi; \frac{du(\varphi)}{d\varphi}\right)$$

holds for all n . Further one can see that for (x, t) in $M \times [1/K_{\min}, \infty)$ $b_n(T_*^n x; t)$ converges to a positive function $1/\chi^{(e)}(x)$ and that $b_{-n}(T_*^{-n} x; -t)$ converges to a negative function $1/\chi^{(c)}(x)$. The function $\chi^{(c)}$ (resp. $\chi^{(e)}$) is continuous at x not in $\bigcup_{n=0}^{\infty} T_*^n S$ (resp. $\bigcup_{n=0}^{\infty} T_*^{-n} S$) and satisfies

$$K_{\min} \leq -\chi^{(c)}(x) \leq K_{\max} \quad (\text{resp. } k_{\min} \leq \chi^{(e)}(x) \leq K_{\max}(\iota)).$$

THEOREM 1. (i) $\alpha^{(c)}$ and $\alpha^{(e)} = T_*^{-1}\alpha^{(c)}$ are generators for T_* with the same finite entropy.

(ii) Almost every element of $\zeta^{(c)} \equiv \bigvee_{i=0}^{\infty} T_*^i \alpha^{(c)}$ is a connected K -decreasing curve whose gradient* at x is equal to $\chi^{(c)}(x)$. Alternatively, almost every element of $\zeta^{(e)} \equiv \bigvee_{i=0}^{\infty} T_*^{-i} \alpha^{(e)}$ is a connected K -increasing curve whose gradient at x is equal to $\chi^{(e)}(x)$.

(iii) T_* is a K -system. Actually, the partition $\zeta^{(c)}$ and the partition $\zeta^{(e)}$ satisfy the following conditions:

$$\begin{aligned}
 T_*^{-1}\zeta^{(c)} &> \zeta^{(c)}, \quad T_*\zeta^{(e)} > \zeta^{(e)}, \\
 \bigvee_i T_*^i \zeta^{(c)} &= \bigvee_i T_*^i \zeta^{(e)} = \epsilon \\
 \bigwedge_i T_*^i \zeta^{(c)} &= \bigwedge_i T_*^i \zeta^{(e)} = \text{the trivial partition}.
 \end{aligned}$$

By the theorem, in order to show the Bernoulli property of T_* , it is enough to give a family \mathcal{F} which satisfies the condition (2.5) in Lemma 2.3. For this purpose, it is necessary to investigate the structure of the

* When a curve γ is given by the equation $\varphi = \psi(r)$, the gradient of γ at $x = (r, \varphi)$ is $d\psi/dr$.

measure μ in connection with the partitions $\zeta^{(e)}$ and $\zeta^{(e)}$. Denote by $\gamma^{(e)}(x)$ the curve which is the element of $\zeta^{(e)}$ involving x . Alternatively, denote by $\gamma^{(e)}(x)$ the curve which is the element of $\zeta^{(e)}$ including x . For two decreasing curves γ and γ' , define the canonical mapping $\Psi_{\gamma,\gamma'}^{(e)}$ by

$$(3.8) \quad \begin{aligned} \Psi_{\gamma,\gamma'}^{(e)}: \gamma &\rightarrow \gamma' \\ \Psi_{\gamma,\gamma'}^{(e)}: x &\mapsto \gamma^{(e)}(x) \cap \gamma' . \end{aligned}$$

Let σ_γ and $\sigma_{\gamma'}$ be the measures on γ and γ' respectively defined as follows; for $\bar{\gamma}$ in γ and $\bar{\gamma}'$ in γ'

$$\sigma_\gamma(\bar{\gamma}) = \int_{\bar{\gamma}} |d\varphi| \quad \text{and} \quad \sigma_{\gamma'}(\bar{\gamma}') = \int_{\bar{\gamma}'} |d\varphi| .$$

Define the measure $\Psi_{\gamma,\gamma'}^{(e)}\sigma_{\gamma'}$ by

$$\Psi_{\gamma,\gamma'}^{(e)}\sigma_{\gamma'}(\bar{\gamma}) \equiv \sigma_{\gamma'}(\Psi_{\gamma,\gamma'}^{(e)}\bar{\gamma}) .$$

Then the Radon-Nikodym density relative to $d\sigma_\gamma$ is given by

$$(3.9) \quad \frac{d\Psi_{\gamma,\gamma'}^{(e)}\sigma_{\gamma'}}{d\sigma_\gamma} = g_{\gamma,\gamma'}^{(e)} \equiv \prod_{i=-\infty}^0 \frac{\Lambda^*(x_i, T_*^{-i}\gamma)}{\Lambda^*(x'_i, T_*^{-i}\gamma')}$$

with x in γ and $x' = \Psi_{\gamma,\gamma'}^{(e)}x$, where

$$(3.10) \quad \Lambda^*(x, \gamma) = \frac{\{k_1 \cos \varphi + k' \cos \varphi_1 + k_1 k' \tau_1\} b_1(x; du/d\varphi) - k' \tau_1 - \cos \varphi}{\cos \varphi} .$$

Similarly, $\Psi_{\gamma',\gamma}^{(e)}$, $\sigma_{\gamma'}$, σ_γ are defined for increasing curves γ', γ and one has

$$(3.9)' \quad \frac{d\Psi_{\gamma',\gamma}^{(e)}\sigma_\gamma}{d\sigma_{\gamma'}} = g_{\gamma',\gamma}^{(e)} = \prod_{i=0}^{\infty} \frac{\Lambda(x_i, T_*^{-i}\gamma)}{\Lambda(x'_i, T_*^{-i}\gamma')}$$

with x in γ and $x' = \Psi_{\gamma',\gamma}^{(e)}x$, where

$$(3.10)' \quad \Lambda(x, \gamma) = - \frac{\{k_1 \cos \varphi + k' \cos \varphi_1 + k_1 k' \tau_1\} \{du/d\varphi + h\} + k_1 \tau_1 + \cos \varphi_1}{\cos \varphi_1}$$

By Lemmas 6.1, 6.1' and 7.1 in [5], for any $\delta > 0$ there exist an even natural number $\ell_0 = \ell_0(\delta, 1, 1/4)$ and a positive function $\varepsilon_0 = \varepsilon_0(x, \delta, 1)$ which guarantee the following property: For an x not in $\bigcup_{i=-\ell_0}^{\ell_0} T_*^i S$, let G be a K -quadrilateral*) (a domain which is enclosed by four curves such that a pair of opposite curves $\gamma_b(G), \gamma_d(G)$ are K -increasing and the other pair of opposite curves $\gamma_a(G), \gamma_c(G)$ are K -decreasing) in the ε_0 -neighbourhood

*) The notation for G and some properties of G are explained in [5].

$U_{\alpha_0}(x)$ of x . Suppose that $\delta_0 \equiv \theta(\gamma_\alpha(G)) = \theta(\gamma_b(G))^*$ and that $T_*^{-\delta_0}G, T_*^{\delta_0}G$ are K -quadrilaterals. Then there exist subsets $G^{(\epsilon, \delta)}$ and $G^{(\epsilon, \delta)}$ which satisfy the four conditions;

- (C-1) for all x in $G^{(\epsilon, \delta)}$ (resp. $G^{(\epsilon, \delta)}$), $\gamma^{(\epsilon)}(x) \cap G^{(\epsilon, \delta)}$ (resp. $\gamma^{(\epsilon)}(x) \cap G^{(\epsilon, \delta)}$) is a connected segment which joins $\gamma_b(G)$ and $\gamma_a(G)$ (resp. $\gamma_\alpha(G)$ and $\gamma_c(G)$),
- (C-2) $\nu(G^{(\epsilon, \delta)}) \geq (1 - \delta)\nu(G)$ and $\nu(G^{(\epsilon, \delta)}) \geq (1 - \delta)\nu(G)$,
- (C-3) for any K -increasing (resp. K -decreasing) curve γ and γ' in G , the canonical mapping $\Psi_{\gamma', \gamma}^{(\epsilon)}$ (resp. $\Psi_{\gamma', \gamma}^{(\epsilon)}$) is absolutely continuous on $\gamma \cap G^{(\epsilon, \delta)}$ (resp. $\gamma \cap G^{(\epsilon, \delta)}$) with respect to σ_γ and $\sigma_{\gamma'}$,
- (C-4) for any $m \geq 0, T_*^{-m}G^{(\epsilon, \delta)} \cap V_m(\delta_0) = \phi^{**}$ (resp. $T_*^mG^{(\epsilon, \delta)} \cap V_m(\delta_0) = \phi$).




Fig. 1

Define the measure ρ_γ on a K -decreasing (or K -increasing) curve γ by

$$\rho_\gamma(\tilde{\gamma}) \equiv \int_{\tilde{\gamma}} dr \quad \text{for } \tilde{\gamma} \subset \gamma.$$

Let γ be a K -decreasing curve in G which joins $\gamma_a(G)$ and $\gamma_c(G)$, and let $\tilde{\gamma}_0$ be a K -increasing curve which is an extension of $\tilde{\gamma}$ and is given by the equation $r = \tilde{u}(\varphi), \pi/2 \leq \varphi \leq (3/2)\pi$. For given φ and ψ , let $\tilde{\gamma}^{\varphi, \psi}$ be the curve defined by $r = \tilde{u}(\varphi) - \tilde{u}(\psi) - u^\psi(\psi)$, where $r = u^\psi(\psi)$ is the equation of $\gamma^{(\epsilon)}(\iota, \tilde{u}(\varphi), \varphi)$. Then by Lemma 8.3 in [5], the measure ν is expressed in the form

$$(3.11) \quad \nu(B \cap G^{(\epsilon, \delta)}) = \int d\sigma_\gamma(\varphi) \int_{\gamma^{(\epsilon)}(\iota, \tilde{u}(\varphi), \varphi) \cap B \cap G^{(\epsilon, \delta)}} g_0(\varphi, \psi) d\sigma_{\gamma^{(\epsilon)}}(\psi),$$

where $g_0(\varphi, \psi)$ is defined by

*) For monotone connected curve $\gamma, \theta(\gamma)$ denotes the total variation of γ in φ -direction; $\theta(\gamma) \equiv \sigma_\gamma(\gamma) = \int_\gamma d\varphi$.

**) $V_m(\delta_0) \equiv \{x; -\cos \varphi(x) \leq (1 + \eta_1)^{-m/32} \delta_0\}$ with $\eta \equiv k_{\min} |\tau|_{\min}$ and $\eta_1 \equiv \min\{\eta, (1 + \eta)^2, K_{\min}/K_{\max}\}$.

$$(3.12) \quad g_0(\varphi, \psi) = \frac{\nu_0 \cos \varphi}{\chi^{(e)}(\iota, u^\varphi(\psi), \psi)} \prod_{i=0}^{\infty} \frac{\Lambda(T_*^{-i}x; T_*^{-i}\hat{\gamma}_0)}{\Lambda(T_*^{-i}\hat{x}; T_*^{-i}\hat{\gamma}^{\varphi, \psi})}$$

with $x = (\iota, \hat{u}(\varphi), \varphi)$ and $\hat{x} = (\iota, u^\varphi(\psi), \psi)$.

Let $\hat{\gamma}$ be a K -decreasing curve in G which joins $\gamma_b(G)$ and $\gamma_a(G)$, and be defined by the equation $r = \hat{u}(\varphi)$. Then by (3.9), one can see that

$$(3.13) \quad \begin{aligned} &\nu(B \cap G^{(e, \delta)} \cap G^{(e, \delta)}) \\ &= \int_{\hat{\gamma} \cap G^{(e, \delta)}} d\sigma_{\hat{\gamma}}(\varphi) \int_{\hat{\gamma} \cap \Psi_{\hat{\gamma}, \hat{\gamma}'}^{(e)}(B \cap G^{(e, \delta)} \cap \hat{\gamma}')} g_0(\varphi, \Psi_{\hat{\gamma}, \hat{\gamma}'}^{(e)}(\hat{\psi})) g_{\hat{\gamma}, \hat{\gamma}'}^{(e)}(\hat{\psi}) d\sigma_{\hat{\gamma}}(\hat{\psi}) \end{aligned}$$

where $\gamma' = \gamma^{(c)}(\iota, \hat{u}(\varphi), \varphi)$, $\Psi_{\hat{\gamma}, \hat{\gamma}'}^{(e)}(\iota, \hat{u}(\hat{\psi}), \hat{\psi}) = (\iota, u^\varphi(\hat{\varphi}), \hat{\varphi})$ and $\Psi^{(e)}\hat{\psi} \equiv \hat{\varphi}$. One can easily see that for any fixed $m \geq \ell_0$, there exists a positive number $\varepsilon_2 = \varepsilon_2(x, m) < \varepsilon_0$ such that $T_*^j U_{\varepsilon_2}(x) \cap V_0(2(1 + \eta_1)^m \varepsilon_2) = \emptyset$ for $|j| \leq m$.

LEMMA 3.1. *Suppose that $G \subset U_{\varepsilon_2}(x)$ be as above and $T_*^{-m}\hat{\gamma}$ is a K -decreasing curve, then*

$$\begin{aligned} &\exp[-c_{31}(1 + \eta_1)^{-m/2}] \\ &\leq \frac{\nu_0}{\nu(B \cap G^{(e, \delta)} \cap G^{(e, \delta)})} \int_{\hat{\gamma} \cap G^{(e, \delta)}} \cos \varphi d\sigma_{\hat{\gamma}}(\varphi) \int_{\Psi_{\hat{\gamma}, \hat{\gamma}'}^{(e)}(B \cap G^{(e, \delta)} \cap \hat{\gamma}')} d\rho_{\hat{\gamma}}(\hat{\psi}) \\ &\leq \exp[c_{31}(1 + \eta_1)^{-m/2}]. \end{aligned}$$

In particular, if A is a $\zeta^{(c)}|_G$ -measurable subset of G and B is $\zeta^{(e)}|_G$ -measurable subset of G , then

$$\begin{aligned} &\exp[-c_{31}(1 + \eta_1)^{-m/2}] \\ &\leq \frac{\nu_0}{\nu(A \cap B \cap G^{(e, \delta)} \cap G^{(e, \delta)})} \int_{\hat{\gamma} \cap B \cap G^{(e, \delta)}} \cos \varphi d\sigma_{\hat{\gamma}}(\varphi) \int_{\hat{\gamma} \cap A \cap G^{(e, \delta)}} d\rho_{\hat{\gamma}}(\hat{\psi}) \\ &\leq \exp[c_{31}(1 + \eta_1)^{-m/2}]. \end{aligned}$$

In order to prove this lemma, we will prepare two lemmas.

LEMMA 3.2. *Let G be as in Lemma 3.1. Let $\hat{\gamma}$ and $\hat{\hat{\gamma}}$ be K -decreasing curves in G which join $\gamma_b(G)$ and $\gamma_a(G)$. If $T_*^{-m}\hat{\gamma}$ and $T_*^{-m}\hat{\hat{\gamma}}$ are K -decreasing, then for \hat{x} in $\hat{\gamma} \cap G^{(e, \delta)}$*

$$\exp[-c_{32}(1 + \eta_1)^{-m/2}] \leq g_{\hat{\gamma}, \hat{\hat{\gamma}}}^{(e)}(\hat{x}) \leq \exp[c_{32}(1 + \eta_1)^{-m/2}]$$

holds with a positive constant c_{32} .

Proof. Put $\gamma \equiv \gamma^{(e)}(\hat{x}) \cap G$ and $\gamma_j = T_*^{-j}\gamma$. Put $\hat{x}_j = T_*^{-j}\hat{x}$ and $\hat{\hat{x}}_j = T_*^{-j}\hat{\hat{x}}$ with $\hat{\hat{x}} = \Psi_{\hat{\hat{\gamma}}, \hat{\gamma}}^{(e)}(\hat{x})$. Since G is in $U_{\varepsilon_2}(x)$ with an x , $\min\{-\cos \varphi(y); y \in \gamma_j\} \geq 2\varepsilon_2(1 + \eta_1)^m$ holds for $-m \leq j \leq 0$ and $\theta(\gamma) \leq 2\varepsilon_2$ holds. Put $c_2 =$

K_{\max}/K_{\min} . Applying Lemma 5.3 and Lemma 5.4 (i) in [5], one has the estimation

$$\log [A^*(\hat{x}_j, \hat{\gamma}_j)/A^*(\hat{x}_j, \hat{\gamma}_j)] \leq (c_{22} + c_{21} + \log c_2)(1 + \eta_1)^{j-m}$$

for $-m \leq j \leq 0$. Since \hat{x} is in $G^{(e,\delta)}$, it holds that $\min \{-\cos \varphi(y); y \in \gamma_j\} \geq \delta_0(1 + \eta_1)^{j/32}$ for $j \leq 0$ and that $\theta(\gamma) \leq (1 + c_2)\delta_0$. Therefore again one has

$$\log [A^*(\hat{x}_j, \hat{\gamma}_j)/A^*(\hat{x}_j, \hat{\gamma}_j)] \leq 2(c_2 + 1)c_{22}(1 + \eta_1)^{31j/32} + (\log c_2)(1 + \eta_1)^j$$

for $j \leq -m$. These estimates imply Lemma 3.2 by (3.9). Q.E.D.

LEMMA 3.3. *Let G be as in Lemma 3.1 and let $T_*^m \tilde{\gamma}_0$ be K -increasing. Then*

$$\begin{aligned} \exp [-c_{33}(1 + \eta_1)^{-m/2}] &\leq \frac{g_0(\varphi, \psi)}{\nu_0} \frac{\cos \varphi}{\cos \psi} \frac{\chi^{(e)}(\iota, u(\psi), \psi)}{\chi^{(e)}(\iota, \tilde{u}(\varphi), \varphi)} \\ &\leq \exp [c_{33}(1 + \eta_1)^{-m/2}] \end{aligned}$$

with a positive constant c_{33} .

Proof. Put $x = (\iota, \tilde{u}(\varphi), \varphi)$ and $\hat{x} = (\iota, u^e(\psi), \psi)$. Similarly in Lemma 3.2,

$$\sum_{j=0}^{\infty} \left| \log \frac{\Lambda(T_*^{-j} x, T_*^{-j} \tilde{\gamma}_0)}{\Lambda(T_*^{-j} \hat{x}, T_*^{-j} \tilde{\gamma}^e, \psi)} \right| \leq c'_{32}(1 + \eta_1)^{-m}.$$

By Lemma 5.3 in [5], for $(\iota, \hat{u}(\hat{\varphi}), \hat{\varphi}) = \Psi_{\tilde{\gamma}, \gamma^{(e)}(\iota, \tilde{u}(\varphi), \varphi)}(\iota, u^e(\psi), \psi)$

$$(3.14) \quad \left| \log \frac{\chi^{(e)}(\iota, u^e(\psi), \psi)}{d\hat{u}/d\hat{\varphi}} \right| \leq (\pi c_{21} + c_3)(1 + \eta_1)^{-m}.$$

On the other hand, the estimate

$$\left| \log \frac{\cos \varphi}{\cos \psi} \right| \leq 2(1 + \eta_1)^{-m}$$

holds, since G is in $U_{e_2}(x)$. Q.E.D.

Proof of Lemma 3.1. Since $|dr| = |d\hat{u}/d\hat{\varphi}| d\hat{\varphi}$ on $\hat{\gamma}$, Lemma 3.2 and Lemma 3.3 imply the first statement in Lemma 3.1. If A and B are as in Lemma 3.1, then

$$\Psi_{\tilde{\gamma}, \gamma'}^{(e)}(A \cap B \cap G^{(e,\delta)} \cap \gamma') = \begin{cases} B \cap G^{(e,\delta)} \cap \hat{\gamma} & \text{if } \gamma' \subset A \cap G^{(e,\delta)} \\ \phi & \text{if } \gamma' \cap A \cap G^{(e,\delta)} = \phi \end{cases}$$

where $\gamma' = \gamma^{(e)}(y)$ with some y . Further, $\gamma' \subset A \cap G^{(e,\delta)}$ if and only if $\tilde{\gamma} \cap \gamma'$ is in $\tilde{\gamma} \cap A \cap G^{(e,\delta)}$. Therefore one has the second statement. Q.E.D.

§4. The perturbed billiard transformation is isomorphic to a Bernoulli shift

Applying the lemmas in §2 and §3, the following theorems will be shown.

THEOREM 2. *Under the assumptions (H-1)~(H-3), $\alpha^{(c)}$ and $\alpha^{(e)}$ are weak Bernoulli generators for T_* . Thus T_* is isomorphic to a Bernoulli shift.*

Proof. By Theorem 1, it is sufficient for the proof to give a family \mathcal{F} which satisfies (2.3) in Lemma 2.3. For given $\delta > 0$, let m_0 be a natural number such that $\exp c_{31}(1 + \eta_1)^{-m_0/2} < 1 + \delta$ and $m_0 \geq \ell_0 \equiv \ell_0(1, 1/4, \delta)$. For every x not in $\bigcup_{i=-m_0}^{m_0} T^i S$ and for any $\delta_0 > 0$, there exists a K -quadrilateral G in $U_{\varepsilon_0(x, m_0)}(x)$ such that $\theta(\gamma_a(G)) = \theta(\gamma_b(G)) < \delta_0$, G involves the point x and $T_*^{-m_0}G, T_*^{m_0}G$ are K -quadrilaterals. By the covering theorem of Vitali, there exists a finite family \mathcal{G} of such G 's which satisfies $G \cap G' = \emptyset$ for $G \neq G'$ in \mathcal{G} and $\nu(M - \bigcup_{G \in \mathcal{G}} G) < \delta$. Then by Lemmas 6.1, 6.1' and 7.1 in [5], there exist subsets $G^{(c, \delta)}$ and $G^{(e, \delta)}$ which satisfy (C-1), (C-2) and (C-3) in §3. Let A be an element of the partition $\bigvee_{i=0}^n T_*^i \alpha^{(c)}$ and let B be an element of $\bigvee_{i=N'}^{N''} T_*^{-i} \alpha^{(c)}$. Since A is $\zeta^{(c)}$ -measurable and B is $\zeta^{(e)}$ -measurable, Lemma 3.1 is applicable. Put $\hat{\gamma} \equiv \gamma^{(c)}(x) \cap G$ and $\hat{\gamma}' \equiv \gamma^{(e)}(x) \cap G$ for a fixed x in $G^{(c, \delta)} \cap G^{(e, \delta)}$. Then one has

$$(1 + \delta)^{-1} \leq \frac{-\nu_0}{\nu(A \cap B \cap G^{(c, \delta)} \cap G^{(e, \delta)})} \int_{\hat{\gamma} \cap B \cap G^{(e, \delta)}} \cos \varphi d\sigma(\varphi) \times \int_{\hat{\gamma}' \cap A \cap G^{(c, \delta)}} d\rho(\gamma) \leq 1 + \delta.$$

Since A and B are arbitrary, the above inequality holds even if one replaces A to $G^{(c, \delta)}$ (B to $G^{(e, \delta)}$). Hence the estimate

$$(1 + \delta)^{-4} \leq \frac{\nu(A \cap B \cap G^{(c, \delta)} \cap G^{(e, \delta)})\nu(G^{(c, \delta)} \cap G^{(e, \delta)})}{\nu(A \cap G^{(c, \delta)} \cap G^{(e, \delta)})\nu(B \cap G^{(c, \delta)} \cap G^{(e, \delta)})} \leq (1 + \delta)^4$$

is obtained. Therefore the family $\mathcal{F} \equiv \{G^{(c, \delta)} \cap G^{(e, \delta)}; G \in \mathcal{G}\}$ satisfies (2.3). Q.E.D.

COROLLARY 4.1. *A Sinai billiard transformation is isomorphic to a Bernoulli shift. In particular, the natural generators $\alpha^{(c)}$ and $\alpha^{(e)}$ are weakly Bernoullian for T .*

Let $\alpha = \{X_j\}$ be a countable partition. Denote the boundary of X_j by ∂X_j . The union $\partial\alpha \equiv \bigcup_j \partial X_j$ is called the boundary of the partition α .

Let $d(x; \partial\alpha)$ be the distance between a point x in M and the boundary $\partial\alpha$.

LEMMA 4.2. (i) *If $\log d(x; \partial\alpha)$ is integrable, then the entropy of α is finite.*

(ii) *If the boundary $\partial\alpha$ consists of curves whose total arclength is finite, then $\log d(x; \partial\alpha)$ is integrable.*

Proof. Put $R = \sup_{x \in X_j} d(x; \partial X_j)$, then for $x \in X_j$

$$\begin{aligned} \nu(X_j) &\geq -\nu_0 \int_0^R \int_{\pi/2}^{R+\pi/2} \cos \varphi d\varphi dr \geq \nu_0 R^3/4 \\ &\geq \nu_0 \{d(x; \partial X_j)\}^3/4. \end{aligned}$$

This inequality implies $-\sum \nu(X_j) \log \nu(X_j) < \infty$. The second assertion is obvious. Q.E.D.

THEOREM 3. *Under the assumptions (H-1)~(H-3), every countable partition α is weakly Bernoullian for T_* whenever $\log d(x; \partial\alpha)$ is integrable.*

Proof. For a fixed x and $i > 0$, the distance between $T_*^{-i}x$ and $T_*^{-i}\gamma^{(c)}(x) \cap \partial\alpha$ measured along $\gamma^{(c)}(T_*^{-i}x)$ is greater than $d(T_*^{-i}x; \partial\alpha)$, if $T_*^{-i}\gamma^{(c)}(x)$ intersects $\partial\alpha$. Hence

$$d(T_*^{-i}x; \partial\alpha) \leq c_1 \theta(T_*^{-i}\gamma^{(c)}(x)) \leq \pi c_1 (1 + \eta_1)^{-i}$$

holds with $c_1 \equiv (1 + K_{\min}^{-2})^{1/2}$, if $T_*^{-i}\gamma^{(c)}(x)$ intersects $\partial\alpha$. Since $\log d(x; \partial\alpha)$ is integrable, for almost every x , $1/i \log d(T_*^{-i}x; \partial\alpha)$ converges to 0 as $i \rightarrow \infty$, by the Birkhoff ergodic theorem. Thus for almost every x , the boundary $\partial\alpha$ is not intersected by $T_*^{-i}\gamma^{(c)}(x)$ of infinitely many i 's. Hence for almost every x , there exists a natural number $n^{(c)}(x)$ such that for all $i \geq n^{(c)}(x)$ $\gamma^{(c)}(x)$ is included in an element of $T_*^i\alpha$. Further since $\log d(x; \partial\alpha)$ is integrable, the partition of $\gamma^{(c)}(T_*^{-i}x)$ into the connected components of the sets $\{\gamma^{(c)}(T_*^{-i}x) \cap X_j\}_{j=1}^\infty$ is a countable partition. Put

$$\zeta_\alpha^{(c)} \equiv \bigvee_{i=0}^\infty T_*^{-i}\alpha \quad \text{and} \quad \zeta_\alpha^{(e)} \equiv \bigvee_{i=1}^\infty T_*^i\alpha.$$

Then by the above discussions, the restriction of the partition $\zeta_\alpha^{(c)}$ to almost every element $\gamma^{(c)}$ of $\zeta^{(c)}$ is a countable partition, whose elements are countable unions of connected segments of $\gamma^{(c)}$. Let $C^{(c)}(x)$ be the connected component of x in the element of $\zeta_\alpha^{(c)} \vee \zeta^{(c)}$ which contains x . The partition $\zeta_\alpha^{(e)}$ and $C^{(e)}(x)$ are similarly defined. Denote by $\varphi(x)$ the φ -coordinate of $x = (t, r, \varphi)$. Then for almost every x

$$\begin{aligned} \bar{\theta}(C^{(e)}(x); x) &\equiv \sup_{y \in C^{(e)}(x)} \varphi(y) - \varphi(x), & \underline{\theta}(C^{(e)}(x); x) &\equiv \varphi(x) - \inf_{y \in C^{(e)}(x)} \varphi(y) \\ \bar{\theta}(C^{(e)}(x); x) &\equiv \sup_{y \in C^{(e)}(x)} \varphi(y) - \varphi(x), & \underline{\theta}(C^{(e)}(x); x) &\equiv \varphi(x) - \inf_{y \in C^{(e)}(x)} \varphi(y) \end{aligned}$$

are all positive.

For $\delta > 0$, let ℓ_0, m_0 and ε_2 be as in the proof of Theorem 2. Let t be a number which satisfies $\nu(E) > 1 - \delta$ with $E = \{x; \bar{\theta}(C^{(e)}(x); x) > t, \underline{\theta}(C^{(e)}(x); x) > t, \bar{\theta}(C^{(e)}(x); x) > t \text{ and } \underline{\theta}(C^{(e)}(x); x) > t\}$. By the similar way to the proof of Theorem 2, there exists a finite family \mathcal{G}_1 of K -quadrilaterals such that

$$\begin{aligned} \theta(\gamma_a(G)) &= \theta(\gamma_b(G)) < t/(1 + K_{\max}/K_{\min}), \\ \nu(E \cap G) &> (1 - \delta)\nu(G), \\ \nu\left(E - \bigcup_{G \in \mathcal{G}_1} G\right) &< \delta, \\ G \cap G' &= \phi \quad \text{if } G \neq G', \end{aligned}$$

and that $T_*^{-m_0}G, T_*^{m_0}G$ are K -quadrilaterals and there exist subsets $G^{(e, \delta)}$ and $G^{(e, \delta)}$ of G which satisfy the conditions (C-1), (C-2), (C-3), and (C-4) in § 3. For G in \mathcal{G}_1 , put

$$\begin{aligned} \hat{G}^{(e)} &\equiv \{x \in G^{(e, \delta)}; C^{(e)}(x) \text{ intersects both } \gamma_b(G) \text{ and } \gamma_a(G)\}, \\ \hat{G}^{(e)} &\equiv \{x \in G^{(e, \delta)}; C^{(e)}(x) \text{ intersects both } \gamma_a(G) \text{ and } \gamma_c(G)\}. \end{aligned}$$

Then $\hat{G}^{(e)}$ is a $\zeta^{(e)}|_G$ -measurable subset and includes $E \cap G^{(e, \delta)}$. Alternatively, $\hat{G}^{(e)}$ is $\zeta^{(e)}|_G$ -measurable subset and includes $E \cap G^{(e, \delta)}$. Since for any element A of $\bigvee_0^n T_*^i \alpha, A \cap \hat{G}^{(e)}$ is $\zeta^{(e)}|_G$ -measurable and since for any element B of $\bigvee_{i=N'}^{N''} T_*^{-i} \alpha, B \cap \hat{G}^{(e)}$ is $\zeta^{(e)}|_G$ -measurable, Lemma 3.1 is applicable to the subsets A and B . Thus one has the estimates

$$\begin{aligned} \left| \frac{\nu(A \cap B \cap \hat{G}^{(e)} \cap \hat{G}^{(e)})\nu(\hat{G}^{(e)} \cap \hat{G}^{(e)})}{\nu(A \cap \hat{G}^{(e)} \cap \hat{G}^{(e)})\nu(B \cap \hat{G}^{(e)} \cap \hat{G}^{(e)})} - 1 \right| &\leq (1 + \delta)^4 - 1, \\ \nu(\hat{G}^{(e)} \cap \hat{G}^{(e)}) &\geq \nu(E \cap G^{(e, \delta)} \cap G^{(e, \delta)}) \geq (1 - 3\delta)\nu(G), \end{aligned}$$

and

$$\nu\left(\bigcup_{G \in \mathcal{G}_1} (\hat{G}^{(e)} \cap \hat{G}^{(e)})\right) \geq (1 - 2\delta)(1 - 3\delta).$$

Hence the conditions in Lemma 2.3 are fulfilled.

Q.E.D.

§5. K-properties of the flow $\{S_t^f\}$

Let $f(x)$ be a positive function on M and let $\{S_t^f\}$ be a Kakutani-Ambrose flow with the basic space M , the basic transformation T_* and the ceiling function $f(x)$; that is, $\{S_t^f\}$ is defined on the space $W \equiv \{(x, v); 0 \leq v < f(x), x \in M\}$ by

$$(5.1) \quad S_t^f(x, v) \equiv \begin{cases} (T_*^{-k}x, v - t + \sum_{j=1}^k f(T_*^{-j}x)) & \text{if } 0 \leq v - t + \sum_{j=1}^k f(T_*^{-j}x) < f(T_*^{-k}x), k \geq 1, \\ (x, v - t) & \text{if } 0 \leq v - t < f(x), \\ (T_*^{-k}x, v - t - \sum_{j=k+1}^0 f(T_*^{-j}x)) & \text{if } 0 \leq v - t - \sum_{j=k+1}^0 f(T_*^{-j}x) < f(T_*^{-k}x), k \leq -1. \end{cases}$$

Associate the invariant probability measure μ_f with $\{S_t^f\}: d\mu_f = c_f dv dx$. Suppose that the assumptions (H-1)~(H-3) are satisfied. Then $\{S_t^f\}$ is ergodic, since T_* is ergodic. Moreover suppose the following three assumptions (f-1)~(f-3):

(f-1) $f(x)$ is strictly positive and continuously differentiable on each element $X_j^{(e)}$ of $\alpha^{(e)}$,

(f-2) there exists a constant K such that

$$\left\{ \left| \frac{\partial f(t, r, \varphi)}{\partial r} \right| + \left| \frac{\partial f(t, r, \varphi)}{\partial \varphi} \right| \right\} \frac{\cos \varphi - 1}{\tau} \leq K$$

with $(t_{-1}, r_{-1}, \varphi_{-1}) = T_*(t, r, \varphi)$,

(f-3) $f(x) \log |\tau(x)|$ is integrable.

For x not in $\cup_{i=0}^\infty T_*^i S$, put

$$(5.2) \quad \begin{aligned} f^{(+)}(x) &\equiv \sum_{i=1}^\infty \left\{ \frac{1}{\chi^{(e)}(x_i)} \frac{\partial f(x_i)}{\partial r} + \frac{\partial f(x_i)}{\partial \varphi} \right\} \prod_{j=0}^{i-1} [A(x_j, \gamma^{(e)}(x_j))]^{-1}, \\ \hat{f}^{(+)}(x) &\equiv \sum_{i=1}^\infty \left\{ \left| \frac{1}{\chi^{(e)}(x_i)} \frac{\partial f(x_i)}{\partial r} \right| + \left| \frac{\partial f(x_i)}{\partial \varphi} \right| \right\} \prod_{j=0}^{i-1} |A(x_j, \gamma^{(e)}(x_j))|^{-1} \end{aligned}$$

with $x_j \equiv T_*^{-j}x$. For x not in $\cup_{i=0}^\infty T_*^{-i}S$, put

$$(5.3) \quad \begin{aligned} f^{(-)}(x) &\equiv \sum_{i=-\infty}^{-1} \left\{ \frac{1}{\chi^{(e)}(x_i)} \frac{\partial f}{\partial r}(x_i) + \frac{\partial f}{\partial \varphi}(x_i) \right\} \prod_{j=i+1}^0 [A^*(x_j, \gamma^{(e)}(x_j))]^{-1} \\ &\quad + \frac{1}{\chi^{(e)}(x)} \frac{\partial f}{\partial r}(x) + \frac{\partial f}{\partial \varphi}(x), \\ \hat{f}^{(-)}(x) &\equiv \sum_{i=-\infty}^{-1} \left\{ \left| \frac{1}{\chi^{(e)}(x_i)} \frac{\partial f}{\partial r}(x_i) \right| + \left| \frac{\partial f}{\partial \varphi}(x_i) \right| \right\} \prod_{j=i+1}^0 |A^*(x_j, \gamma^{(e)}(x_j))|^{-1} \\ &\quad + \left| \frac{1}{\chi^{(e)}(x)} \frac{\partial f}{\partial r}(x) \right| + \left| \frac{\partial f}{\partial \varphi}(x) \right|. \end{aligned}$$

Then by assumptions (f-1), (f-2) and by Lemma 3.2, Lemma 3.3, the series in (5.2) and (5.3) converge and $f^{(+)}(x)$ (resp. $f^{(-)}(x)$) is continuous at x not in $\bigcup_{i=0}^{\infty} T_*^i S$ (resp. $\bigcup_{i=0}^{\infty} T_*^{-i} S$).

For $w = (\tilde{x}, \tilde{v})$ in W , define curves $\tilde{\gamma}^{(+)}(w)$ and $\tilde{\gamma}^{(-)}(w)$ passing through (\tilde{x}, \tilde{v}) by the following way. Let $r = u^{(\epsilon)}(\varphi; \tilde{x})$ be the equation of the curve $\gamma^{(\epsilon)}(\tilde{x})$ and let $r = u^{(\epsilon)}(\varphi; \tilde{x})$ be the equation of the curve $\gamma^{(\epsilon)}(\tilde{x})$. Let $\tilde{\gamma}^{(+)}(w)$ and $\tilde{\gamma}^{(-)}(w)$ be the curves defined respectively by the equations

$$(5.4) \quad \begin{cases} \iota = \tilde{\iota} \\ r = u^{(\epsilon)}(\varphi; \tilde{x}) \\ v = \tilde{v} - \int_{\tilde{\varphi}}^{\varphi} f^{(-)}(\tilde{\iota}, u^{(\epsilon)}(\varphi; \tilde{x}), \varphi) d\varphi \end{cases} \quad \text{and} \quad \begin{cases} \iota = \tilde{\iota} \\ r = u^{(\epsilon)}(\varphi; \tilde{x}) \\ v = \tilde{v} + \int_{\tilde{\varphi}}^{\varphi} f^{(+)}(\tilde{\iota}, u^{(\epsilon)}(\varphi; \tilde{x}), \varphi) d\varphi \end{cases}$$

for $0 \leq v < f(\iota, r, \varphi)$ with $\tilde{x} = (\tilde{\iota}, \tilde{r}, \tilde{\varphi})$. Then, obviously, $\tilde{\gamma}^{(+)}(w)$ and $\tilde{\gamma}^{(-)}(w)$ are locally transversal fibres; that is,

(i) for w' in $\tilde{\gamma}^{(+)}(w)$ (resp. $\tilde{\gamma}^{(-)}(w)$), $\tilde{\gamma}^{(+)}(w') = \tilde{\gamma}^{(+)}(w)$ (resp. $\tilde{\gamma}^{(-)}(w') = \tilde{\gamma}^{(-)}(w)$),

(ii) $S_i^f \tilde{\gamma}^{(+)}(w)$ coincides with $\tilde{\gamma}^{(+)}(S_i^f w)$ and $S_i^f \tilde{\gamma}^{(-)}(w)$ coincides with $\tilde{\gamma}^{(-)}(S_i^f w)$ in a neighbourhood of $S_i^f w$.

Therefore $\tilde{I}^{(+)}(w) \equiv \bigcup_i S_i^f \tilde{\gamma}^{(+)}(S_i^f w)$ and $\tilde{I}^{(-)}(w) \equiv \bigcup_i S_i^f \tilde{\gamma}^{(-)}(S_i^f w)$ consist of countably many connected curves in W . Further $\tilde{I}^{(+)}(w)$ and $\tilde{I}^{(-)}(w)$ are transversal fibres; that is,

- (i) $\tilde{I}^{(+)}(w') = \tilde{I}^{(+)}(w)$ for $w' \in \tilde{I}^{(+)}(w)$,
 $\tilde{I}^{(-)}(w') = \tilde{I}^{(-)}(w)$ for $w' \in \tilde{I}^{(-)}(w)$,
- (ii) $S_i^f \tilde{I}^{(+)}(w) = \tilde{I}^{(+)}(S_i^f w)$ and $S_i^f \tilde{I}^{(-)}(w) = \tilde{I}^{(-)}(S_i^f w)$.

For each x in M , identify two points $(x, f(x))$ and $(T_* x, 0)$. Under the identification, let $\tilde{\gamma}^{(+)}(w)$ be the connected component of w in $\tilde{I}^{(+)}(w)$. Then $\{\tilde{\gamma}^{(+)}(w); w \in W\}$ gives a partition $\tilde{\zeta}^{(+)}$ of W . Similarly, $\tilde{\gamma}^{(-)}(w)$ and $\tilde{\zeta}^{(-)}$ are given by $\{\tilde{I}^{(-)}(w)\}$.

A curve $\tilde{\gamma}$ in W which is given by the equations $\iota = \tilde{\iota}$, $r = u(\varphi)$ and $v = t(\varphi)$ is said to be K -increasing (resp. K -decreasing), if the curve γ in $M^{(\tilde{\iota})}$ defined by $r = u(\varphi)$ is K -increasing (resp. K -decreasing) and $t(\varphi)$ is locally Lipschitz continuous. For a given K -increasing curve $\tilde{\gamma}$ in W , define a measure $\sigma_{\tilde{\gamma}}$ by

$$(5.5) \quad \sigma_{\tilde{\gamma}}(A) = \int_A |d\varphi|$$

for A in $\tilde{\gamma}$. Put for a subset R of $(-\infty, \infty)$

$$(5.6) \quad \begin{aligned} A^{(+)}[\tilde{\gamma}; t] &\equiv \bigcup_{w \in S_{-t}\tilde{\gamma}} \tilde{\gamma}^{(+)}(w), \\ A^{(+)}[\tilde{\gamma}; R] &\equiv \bigcup_{t \in R} A^{(+)}[\tilde{\gamma}; t]. \end{aligned}$$

Let Π be the natural projection from W to M ; $\Pi(\tilde{x}, \tilde{v}) = \tilde{x}$. Then for sufficiently small R the measure $\mu = \mu_f$ satisfies

$$(5.7) \quad \mu(B \cap A^{(+)}[\tilde{\gamma}; R]) = c_f \int_R \nu(\Pi(A^{(+)}[\tilde{\gamma}; t] \cap B)) dt.$$

Similarly, subsets $A^{(-)}[\tilde{\gamma}; t]$ and $A^{(-)}[\tilde{\gamma}; R]$ are defined. The local fibres $\{\tilde{\gamma}^{(+)}\}$ and $\{\tilde{\gamma}^{(-)}\}$ are called *mutually integrable* (with each other), if for almost every $w = (\tilde{x}, \tilde{v})$ in W and for almost every y in $\Pi(A^{(+)}[\tilde{\gamma}^{(-)}(w); 0]) \cap \Pi(A^{(-)}[\tilde{\gamma}^{(+)}(w); 0])$, the relation

$$A^{(+)}[\tilde{\gamma}^{(-)}(w); 0] \cap \Pi^{-1}(y) = A^{(-)}[\tilde{\gamma}^{(+)}(w); 0] \cap \Pi^{-1}(y)$$

holds.

THEOREM 4. *Under the assumptions (H-1)~(H-3) and (f-1)~(f-3),*

- (i) $S_t^f \tilde{\zeta}^{(+)} > \tilde{\zeta}^{(+)}, S_t^f \tilde{\zeta}^{(-)} < \tilde{\zeta}^{(-)}, t > 0,$
 $\bigvee_t S_t^f \tilde{\zeta}^{(+)} = \bigvee_t S_t^f \tilde{\zeta}^{(-)} = \epsilon,$
 $\bigwedge_t S_t^f \tilde{\zeta}^{(+)} = \bigwedge_t S_t^f \tilde{\zeta}^{(-)} = \pi(\{S_t^f\}),$
- (ii) *the conditional measure $\mu(\cdot | \tilde{\gamma}^{(+)})$ (resp. $\mu(\cdot | \tilde{\gamma}^{(-)})$) is equivalent to $\sigma_{\tilde{\gamma}^{(+)}}$ (resp. $\sigma_{\tilde{\gamma}^{(-)}}$),*
- (iii) $h(S_t^f) = h(S_t^f \tilde{\zeta}^{(+)} | \tilde{\zeta}^{(+)}) = h(S_t^f \tilde{\zeta}^{(-)} | \tilde{\zeta}^{(-)}) = th(T_*) / \int f(x) d\nu,$
- (iv) *if $\{\tilde{\gamma}^{(+)}\}$ and $\{\tilde{\gamma}^{(-)}\}$ are not mutually integrable, then $\pi(\{S_t^f\})$ is the trivial partition, and hence $\{S_t^f\}$ is a K -system,*
- (v) *if $\{S_t^f\}$ has no point spectrum, then $\{S_t^f\}$ is a K -system.*

Proof. By the above discussions and the definitions,

$$S_t^f \tilde{\zeta}^{(+)} > \tilde{\zeta}^{(+)}, S_t^f \tilde{\zeta}^{(-)} < \tilde{\zeta}^{(-)} \quad (t > 0) \quad \text{and} \quad \bigvee_t S_t^f \tilde{\zeta}^{(+)} = \bigvee_t S_t^f \tilde{\zeta}^{(-)} = \epsilon$$

are obvious. Let β be the partition of W given by $\beta \equiv \Pi^{-1}\alpha^{(e)} = \{\Pi^{-1}X_j^{(e)}; X_j^{(e)} \in \alpha^{(e)}\}$. For any countable partition $\alpha = \{Y_j\}$ of W let $\tilde{d}(w; \partial\alpha)$ be the distance between w and the boundaries $\bigcup_j \partial Y_j \cup W_* \cup W^* \cup \Pi^{-1}(S)$ where $W_* \equiv \{(x, 0); x \in M\}$, $W^* \equiv \{(x, f(x)); x \in M\}$ and $\Pi^{-1}(S) \equiv \{(x, v); 0 \leq v < f(x), x \in S\}$. Then $\log \tilde{d}(w; \partial\beta)$ is integrable by virtue of (f-1)~(f-3). Since the flow $\{S_t^f\}$ is ergodic, except for a countable number of t 's the transformation S_t^f is ergodic. Fix such a sufficiently small positive t and suppose that $\log \tilde{d}(w; \partial\alpha)$ is integrable. Then by the same way as in the proof

of Theorem 3, one can see that for almost every element $\tilde{\gamma}^{(+)}$ of $\tilde{\zeta}^{(+)}$, the restriction of $\zeta_\alpha^{(+)} \equiv \bigvee_{k=0}^\infty S_{-kt}^f \alpha$ to $\tilde{\gamma}^{(+)}$ is a countable partition, each element of which is a union of a countable number of segments of $\tilde{\gamma}^{(+)}$. Hence one can see

$$\bigwedge_n \bigvee_{k=0}^\infty S_{(n-k)t}^f \alpha \leq \bigwedge_s S_s^f \tilde{\zeta}^{(+)} \equiv \tilde{\zeta}_\infty^{(+)}$$

Since there exists a sequence of partitions $\{\alpha_n\}$ of W increasing to ϵ such that $\log \tilde{d}(w; \partial\alpha_n)$ are integrable, $\pi(S_t^f) \leq \tilde{\zeta}_\infty^{(+)}$. If $\alpha \geq \beta$, then $\zeta_\alpha^{(+)} \geq \Pi^{-1}\zeta^{(e)} \equiv \{\Pi^{-1}\gamma^{(e)}; \gamma^{(e)} \in \zeta^{(e)}\}$, since $\alpha^{(e)}$ generates $\zeta^{(e)}$. For any $\epsilon > 0$ and for almost every w , $\{S_{-kt}^f w; k \geq 0\}$ visits the set $Y_\epsilon = \{(x, v) \in W; u - \epsilon < v < u, (x, u) \in \bigcup_j \partial Y_j\}$ infinitely many times, since S_t^f is ergodic. Hence $\zeta_\alpha^{(+)} = \bigvee_{k=0}^\infty S_{-kt}^f \alpha \geq \tilde{\zeta}^{(+)}$ if $\alpha \geq \beta$ and if $\log \tilde{d}(w; \partial\alpha)$ is integrable. Hence one obtains

$$\pi(S_t^f) = \pi(\{S_s^f\}) = \bigwedge_s S_s^f \tilde{\zeta}^{(+)}$$

Thus (i) is proved. The second assertion (ii) is obvious by definition and § 3. The third assertion (iii) comes from the theorem of Rohlin and Sinai [10] and a theorem of Abramov [1]. For almost every y in M and for a sufficiently small neighbourhood $U_\epsilon(y)$, there exists a quartet $\{y, y_1, y_2, y_3\}$ in $U_\epsilon(y)$ such that y_1 in $\gamma^{(e)}(y)$, y_2 in $\gamma^{(e)}(y_1)$, y_3 in $\gamma^{(e)}(y_2)$ and y in $\gamma^{(e)}(y_3)$. Then one can define a mapping Ψ of $\Pi^{-1}(y)$ by

$$\Psi w = \tilde{\gamma}^{(-)}(\tilde{\gamma}^{(+)}(\tilde{\gamma}^{(-)}(\tilde{\gamma}^{(+)}(w) \cap \Pi^{-1}(y_1)) \cap \Pi^{-1}(y_2)) \cap \Pi^{-1}(y_3)) \cap \Pi^{-1}(y)$$

for w in $\Pi^{-1}(y)$. Obviously, there exists a real number $a = a(y, y_1, y_2, y_3)$ such that

$$\Psi(y, u) = (y, u + a)$$

for (y, u) in the domain of Ψ . If $\{\tilde{\gamma}^{(+)}\}$ and $\{\tilde{\gamma}^{(-)}\}$ are not mutually integrable, then there exists a subset Y of positive measure such that for all $\delta > 0$ and all y in Y one can choose a quartet $\{y, y_1, y_2, y_3\}$ with $0 < |a(y, y_1, y_2, y_3)| < \delta$. Put $\tilde{\zeta}_{-\infty}^{(-)} \equiv \bigwedge S_t^f \tilde{\zeta}^{(-)}$ and let $h(w)$ be a $\tilde{\zeta}_\infty^{(+)} \wedge \tilde{\zeta}_{-\infty}^{(-)}$ -measurable bounded function. Since $\tilde{\zeta}_\infty^{(+)} \wedge \tilde{\zeta}_{-\infty}^{(-)}$ is $\{S_t^f\}$ -invariant, $h_b(w) = \frac{1}{b} \int_0^b h(S_t^f w) dt$ is again $\tilde{\zeta}_\infty^{(+)} \wedge \tilde{\zeta}_{-\infty}^{(-)}$ -measurable. Then $h_b(y, u)$ is continuous in u and $h_b(w)$ converges to $h(w)$ a.e. w as $b \rightarrow 0$. There exist measurable functions $h_b^{(+)}(w)$ and $h_b^{(-)}(w)$ such that $h_b(w) = h_b^{(+)}(w) = h_b^{(-)}(w)$ for a.e. w and that $h_b^{(+)}(w)$ is constant on $\tilde{\Gamma}^{(+)}$ and $h_b^{(-)}(w)$ is constant on $\tilde{\Gamma}^{(-)}$. Since canonical mappings $\Psi^{(e)}$ and $\Psi^{(e)}$ are absolutely continuous, one can choose

y, y_1, y_2, y_3 such that $h_b(y_i, u) = h_b^{(+)}(y_i, u) = h_b^{(-)}(y_i, u)$ for almost every u in $[0, f(y_i))$, $i = 0, 1, 2, 3$ with $y_0 = y$. Hence one can obtain

$$h_b(y, u) = h_b(y, u + a)$$

for almost every u in $[\delta, f(y) - \delta]$ with small $\delta > 0$. Since δ can be taken arbitrary small and $h_b(y, u)$ is continuous in u , $h_b(y, u)$ is constant in u . Hence $h_b(y, u)$ is constant in a subset with positive measure. Since b and h are arbitrary, one can see that the partition $\zeta_{\infty}^{(+)} \wedge \zeta_{\infty}^{(-)}$ contains an element of positive measure. Since $\{S_i^f\}$ is ergodic and $\zeta_{\infty}^{(+)} \wedge \zeta_{\infty}^{(-)}$ is invariant under $\{S_i^f\}$, the partition $\zeta_{\infty}^{(+)} \wedge \zeta_{\infty}^{(-)}$ is trivial. Thus (iv) was proved. Suppose that $\{\tilde{y}^{(+)}\}$ and $\{\tilde{y}^{(-)}\}$ are mutually integrable and that $\zeta_{\infty}^{(+)} \wedge \zeta_{\infty}^{(-)}$ is not trivial. Since $\bigwedge_k T_*^k \zeta^{(e)} \wedge \bigwedge_k T_*^k \zeta^{(e)}$ is the trivial partition, the factor flow of $\{S_i^f\}$ with respect to $\zeta_{\infty}^{(+)} \wedge \zeta_{\infty}^{(-)}$ is a circle flow. Hence $\{S_i^f\}$ has a point spectrum. Q.E.D.

It is very difficult to check that $\{\tilde{y}^{(+)}\}$ and $\{\tilde{y}^{(-)}\}$ are not mutually integrable for general cases.

LEMMA 5.1. *For a K-quadrilateral G, put*

$$v_f(G) \equiv \int_{r_a} f^{(+)} d\varphi + \int_{r_b} f^{(-)} d\varphi - \int_{r_c} f^{(+)} d\varphi - \int_{r_d} f^{(-)} d\varphi .$$

(i) *If $v_f(G) = 0$ for any G whose lateral sides are segments of $\{\gamma^{(e)}\}$ and $\{\tilde{\gamma}^{(e)}\}$, then $\{\tilde{y}^{(+)}\}$ and $\{\tilde{y}^{(-)}\}$ are mutually integrable.*

(ii) *If $v_f(G) > 0$ for any G in an open set whose lateral sides are segments of $\{\gamma^{(e)}\}$ and $\{\tilde{\gamma}^{(e)}\}$, then $\{\tilde{y}^{(+)}\}$ and $\{\tilde{y}^{(-)}\}$ are not mutually integrable.*

COROLLARY 5.2 ([12]). *A Sinai billiard system is a K-system.*

Proof. Since $f(x) = -\tau(x)$, it holds that

$$\frac{1}{\chi^{(e)}(x)} \frac{\partial f}{\partial r} + \frac{\partial f}{\partial \varphi} = \frac{1}{\chi^{(e)}(x)} \sin \varphi(x) - \frac{1}{\chi^{(e)}(x_1)} \sin \varphi(x_1) \frac{d\varphi_1}{d\varphi} ,$$

and hence $f^{(+)}(x) = \sin \varphi / \chi^{(e)}(x)$ (cf. [5]). Similarly one has $f^{(-)}(x) = -\sin \varphi / \chi^{(e)}(x)$. Hence

$$v_f(G) = - \int_{r_a} \sin \varphi dr - \int_{r_b} \sin \varphi dr + \int_{r_c} \sin \varphi dr + \int_{r_d} \sin \varphi dr = \frac{1}{\nu_0} \nu(G) .$$

Q.E.D.

COROLLARY 5.3. *The flow $\{S_i^f\}$ in § 3 has expanding and contracting*

transversal fibres. Further if $\{S_t^f\}$ has no point spectrum, then $\{S_t^f\}$ is a K -system.

§6. Bernoulli flow

A flow $\{S_t\}$ is called a Bernoulli flow if every S_t ($t \neq 0$) is a Bernoulli shift.

THEOREM 5. *Under the assumptions (H-1)~(H-3) and (f-1)~(f-3), if $\{S_t^f\}$ is a K -system and α is a finite partition such that $\log \tilde{d}(w; \partial\alpha)$ is integrable, then α is very weakly Bernoullian for S_t^f ($t \neq 0$). Furthermore $\{S_t^f\}$ is a Bernoulli flow.*

Proof. For $\varepsilon > 0$, choose sufficiently small $\delta > 0$. Let δ_1 be a positive number with $\mu(E_1) > 1 - \delta$ where $E_1 \equiv \{w \in W; \tilde{d}(w; \partial\alpha) > \delta_1\}$. Fix a positive t . Then S_t^f is an ergodic transformation, since $\{S_t^f\}$ is a K -system by the assumption. Hence by Birkhoff's ergodic theorem, there exist a set E_2 with $\mu(E_2) > 1 - \delta$ and a natural number N_1 such that for all w in E_2 and all $n \geq N_1$

$$\frac{1}{n} \sum_{k=0}^{n-1} I_{E_1^c}(S_{-kt}^f w) \leq 2\delta$$

where $I_{E_1^c}$ is the indicator function of E_1^c . Then there exists a δ_2 with $\mu(E_3) > 1 - \delta$, where $E_3 \equiv \{w \in W; \inf_{0 \leq k \leq N_1-1} \tilde{d}(S_{-kt}^f w; \partial\alpha) > \delta_2\}$. Denote by $C^{(+)}(w)$ and $C^{(-)}(w)$ the connected components of w in the elements of $\bar{\zeta}^{(+)} \vee \zeta_\alpha^{(+)}$ and $\bar{\zeta}^{(-)} \vee \zeta_\alpha^{(-)}$ respectively, where $\zeta_\alpha^{(+)} = \bigvee_{k=0}^\infty S_{-kt}^f \alpha$, $\zeta_\alpha^{(-)} = \bigvee_{k=0}^\infty S_{kt}^f \alpha$ and $\bar{\zeta}^{(+)}$ (resp. $\bar{\zeta}^{(-)}$) is the partition into curves $\{\bar{\gamma}^{(+)}\}$ (resp. $\{\bar{\gamma}^{(-)}\}$). By the same reason in the proof of Theorem 3, there exists a positive number δ_3 such that $\mu(E_4) > 1 - \delta$ where $E_4 \equiv \{w \in W; \bar{\theta}(\Pi(C^{(+)}(w)); \Pi(w)) > \delta_3, \underline{\theta}(\Pi(C^{(+)}(w)); \Pi(w)) > \delta_3, \bar{\theta}(\Pi(C^{(-)}(w)); \Pi(w)) > \delta_3$ and $\underline{\theta}(\Pi(C^{(-)}(w)); \Pi(w)) > \delta_3\}$. There exists a positive δ_4 ($< \delta$) such that $1/\delta_4 > \sup_{x \in E_5} \{|f^{(+)}(x)| + |f^{(-)}(x)|\}$ with $E_5 \equiv \{x \in M; |\cos \varphi(x)| > \delta_4\}$. Note that $\Pi^{-1}(E_5)$ is a subset of E_1 .

For any x not in $\bigcup_{i=-\infty}^\infty T_*^i S$, there exists $\varepsilon_3 = \varepsilon_3(x) > 0$ such that for any y in ε_3 -neighbourhood $U_{\varepsilon_3}(x)$

$$\begin{aligned} |f^{(+)}(x) - f^{(+)}(y)| &< \delta_4 \delta, \quad |f^{(-)}(x) - f^{(-)}(y)| < \delta_4 \delta \\ |f(x) - f(y)| &< \delta_2 \delta \quad \text{and} \quad \left| \frac{\cos \varphi(y)}{\cos \varphi(x)} \right| < 1 + \delta. \end{aligned}$$

Let $\ell_0 = \ell_0(\delta, 1, 1/4)$, $m_0 \geq \ell_0$ and $\varepsilon_2 = \varepsilon_2(x, m_0)$ be as in the proof of Theorem

2. Then for any $w = (\tilde{z}, \tilde{r}, \tilde{\varphi}, \tilde{v})$ in E_3 and for any $\delta_0 > 0$, there exists a subset \tilde{G} of W which is constructed as follows: There exists a K -quadrilateral G in $U_{s_3}(\tilde{x}) \cap U_{s_2}(\tilde{x})$ with $\tilde{x} \equiv (\tilde{z}, \tilde{r}, \tilde{\varphi}) = \Pi(w)$ such that $T_*^{-m_0}$ and $T_*^{m_0}$ are continuous on G , $T_*^{-m_0}G$ and $T_*^{m_0}G$ are K -quadrilaterals, $\gamma_a(G)$ (resp. $\gamma_b(G)$) is a segment of the fibre $\gamma^{(c)}(\tilde{x})$ (resp. $\gamma^{(e)}(\tilde{x})$), and

$$\theta(\gamma_a(G)) = \theta(\gamma_b(G)) < \min \{\delta_0, \delta_1, \delta_2, \delta_3\} / (1 + c_2) .$$

Put $\tilde{\gamma}_a \equiv \tilde{\gamma}^{(+)}(w) \cap \Pi^{-1}(G)$. For \bar{w} in W , put

$$D(\bar{w}) \equiv \left\{ \begin{array}{l} 1 - \delta \leq \frac{\varphi - \bar{\varphi}}{r - \bar{r}} \frac{1}{\chi^{(c)}(x)} \leq 1 + \delta \\ (\tilde{z}, r, \varphi, v) ; \\ f^{(+)}(\tilde{x}) - \delta \leq \frac{v - \bar{v}}{\varphi - \bar{\varphi}} \leq f^{(+)}(\tilde{x}) + \delta \end{array} \right\}$$

with $\bar{w} = (\tilde{z}, \bar{r}, \bar{\varphi}, \bar{v})$. Define \tilde{G} by

$$\tilde{G} = \bigcup_{-a \leq s \leq a} S_s^f \bigcup_{\bar{w} \in \tilde{\gamma}_b} D(\bar{w}) \cap \Pi^{-1}(G) ,$$

with $a = \theta(\gamma_a(G))$. As stated in § 3, there exist subsets $G^{(c,\delta)}$ and $G^{(e,\delta)}$ of G which satisfy (C-1), (C-2) and (C-3). Put

$$\tilde{G}^{(c,\delta)} \equiv \left\{ w \in \Pi^{-1}(G^{(c,\delta)}) \cap \tilde{G}; \tilde{\gamma}^{(+)}(w) \cap \bigcup_{|s| \leq a} S_s^f \tilde{\gamma}_b \neq \phi \right\} .$$

Then one can see the inequality

$$\mu(\tilde{G}^{(c,\delta)} \cap \Pi^{-1}(G^{(e,\delta)})) \geq (1 + 2\delta)^{-4} (1 - 2\delta) \mu(\tilde{G}) ,$$

since $1 - \delta \leq \chi^{(c)}(x) / \chi^{(c)}(\tilde{x}) \leq 1 + \delta$ for x in G . By the covering theorem of Vitali, there exists a finite family \mathcal{G} of \tilde{G} 's which satisfies

$$\begin{aligned} \mu(E_1 \cap E_2 \cap E_3 \cap E_4 \cap \tilde{G}) &\geq (1 - \delta) \mu(\tilde{G}) , \\ \Pi^{-1}(E_5) &\supset \tilde{G} , \\ \mu\left(E_1 \cap E_2 \cap E_3 \cap E_4 \cap \Pi^{-1}(E_5) - \bigcup_{\tilde{G} \in \mathcal{G}} \tilde{G}\right) &< \delta , \\ \tilde{G} \cap \tilde{G}' &= \phi \quad \text{if } \tilde{G} \neq \tilde{G}' . \end{aligned}$$

Put

$$E(\tilde{G}) \equiv \{w \in \tilde{G}^{(c,\delta)}; \tilde{\gamma}^{(+)}(w) \cap E_4 \cap \tilde{G} \neq \phi\} .$$

Then $B \cap E(\tilde{G})$ is $\tilde{\zeta}^{(+)}|_{\mathcal{G}}$ -measurable for every element B of $\bigvee_{k=N'}^{N''} S_{-kt}^f \alpha$ with $N'' > N' \geq 0$. By Lemma 3.1, the estimate

$$\begin{aligned} & \mu(B \cap E(\tilde{G}) \cap \Pi^{-1}(G^{(\epsilon, \delta)})) / (1 + \delta) \\ & \leq \int_{\gamma_a(\tilde{G}) \cap G^{(\epsilon, \delta)}} d\rho(\gamma) \int_{\bar{v}-a-\delta}^{\bar{v}+a+\delta} dt \int_{B \cap E(\tilde{G}) \cap S_{\ell_{\gamma}^{\epsilon}}(G)} - \mu_0 \cos \varphi d\sigma(\varphi) \\ & \leq \mu(B \cap E(\tilde{G}) \cap \Pi^{-1}(G^{(\epsilon, \delta)})) / (1 - \delta) \end{aligned}$$

is obtained. Moreover, for y in $\gamma_a(G) \cap G^{(\epsilon, \delta)}$

$$1 - \delta < \frac{\int_{\bar{v}-a-2\delta}^{\bar{v}+a+2\delta} dt \int_{B \cap E(\tilde{G}) \cap S_{\ell_{\gamma}^{\epsilon}}(y)} \cos \varphi d\varphi}{\int_{\bar{v}-a-\delta}^{\bar{v}+a+\delta} dt \int_{B \cap E(\tilde{G}) \cap S_{\ell_{\gamma}^{\epsilon}}(y)} \cos \varphi d\varphi} < 1 + \delta$$

is obtained. Therefore, there exists a $((1 + \delta)^3 - 1)$ -measure preserving mapping ϕ from $\cup_{\tilde{G} \in \mathcal{G}} (E(\tilde{G}) \cap \Pi^{-1}(G^{(\epsilon, \delta)})) \cap B$ to $\cup_{\tilde{G} \in \mathcal{G}} (E(\tilde{G}) \cap \Pi^{-1}(G^{(\epsilon, \delta)}))$ such that ϕ maps $E(\tilde{G}) \cap \Pi^{-1}(\gamma^{(\epsilon)}(x)) \cap B$ to $E(\tilde{G}) \cap \Pi^{-1}(\gamma^{(\epsilon)}(x))$ for x in $\gamma_a(G) \cap G^{(\epsilon, \delta)}$. Let $\ell_i(w)$ be the name function of $S_{\ell}^f \alpha$. For z in $E_1 \cap E_2 \cap E_3 \cap E(\tilde{G}) \cap B$

$$\ell_i(z) = \ell_i(\phi z) \quad \text{for } 1 \leq i \leq N_1 - 1$$

and for $n \geq N_1$

$$\frac{1}{n} \sum_1^n e(\ell_i(z) - \ell_i(\phi z)) \leq 2\delta$$

hold, since

$$\begin{aligned} & \tilde{d}(S_{\ell}^f z; \partial\alpha) > \delta_2 \quad \text{for } 1 \leq k \leq N_1 - 1, \\ & \frac{1}{n} \sum_{k=0}^{n-1} I_{E_1^c}(S_{\ell}^f z) \leq 2\delta, \\ & (\text{distance of } S_{\ell}^f z \text{ and } S_{\ell}^f \phi z) \leq \min(\delta_1, \delta_2, \delta_3) \end{aligned}$$

for $k \geq 0$. On the other hand, there exist an N_2 and a set E_6 such that $\mu(E_6) > 1 - \delta$ and for all $N', N'' \geq N_2$ and all B in $\bigvee_{k=N'}^{N''} S_{\ell}^f \alpha$, $B \subset E_6$,

$$|\mu(E_1 \cap E_2 \cap E_3 \cap E(\tilde{G})) - \mu(E_1 \cap E_2 \cap E_3 \cap E(\tilde{G})|B)| \leq \delta\mu(\tilde{G})$$

holds, since S_{ℓ}^f is a K -system. Hence

$$\begin{aligned} & \mu\left(\bigcup_{\tilde{G} \in \mathcal{G}} (E_1 \cap E_2 \cap E_3 \cap E(\tilde{G}) \cap B)\right) \\ & \geq \sum_{\tilde{G} \in \mathcal{G}} \mu(E_1 \cap E_2 \cap E_3 \cap E_4 \cap \tilde{G})\mu(B) - \delta\mu(B) \\ & \geq (1 - \delta)\mu\left(\bigcup_{\tilde{G} \in \mathcal{G}} \tilde{G}\right)\mu(B) - \delta\mu(B) \end{aligned}$$

$$\begin{aligned} &\geq (1 - \delta)(\mu(E_1 \cap E_2 \cap E_3 \cap E_4 \cap \Pi^{-1}(E_5)) - \delta)\mu(B) - \delta\mu(B) \\ &\geq [(1 - \delta)(1 - 6\delta) - \delta]\mu(B). \end{aligned}$$

Therefore by Lemma 2.4, the partition α is very weakly Bernoullian.

Since there exists an increasing sequence of finite partitions $\{\alpha_n\}$ such that $\log \tilde{d}(w; \partial\alpha_n)$ is integrable and α_n increases to ϵ as $n \rightarrow \infty$, S_t^f is a Bernoulli shift for fixed $t \neq 0$. Q.E.D.

COROLLARY 6.1. *A Sinai billiard system is a Bernoulli flow.*

COROLLARY 6.2. *If $\{S_t^f\}$ has no point spectrum, then $\{S_t^f\}$ is a Bernoulli flow.*

THEOREM 6. *The flow $\{S_t\}$ given in § 3 is a Bernoulli flow, if the assumptions (H-1)~(H-3) are fulfilled and if $\{S_t\}$ has no point spectrum.*

Appendix

The properties of the partitions $\alpha^{(e)}$ and $\alpha^{(c)}$ have been shown in [5]. Now some of them will be stated. Under the suitable numbering the followings are true, here denote as $a(j) = O(j^b)$ if

$$0 < \liminf_{j \rightarrow \infty} |a(j)|j^{-b} \leq \overline{\lim}_{j \rightarrow \infty} |a(j)|j^{-b} < \infty :$$

- (i) $T_* X_j^{(e)} = X_j^{(c)}$,
- (ii) $\tau(x) = O(j)$ for $x \in X_j^{(e)}$ (resp. $X_j^{(c)}$).

Further the following figure is also true.




Fig. 2

These show that $\log d(x; \partial\alpha^{(e)})$ is integrable. The condition (f-1) and (i) imply that the distance between $w = (x, v)$ in $\Pi^{-1}(X_j^{(e)})$ and the

boundary W^* is greater than $O(j^{-1/2})d(x; \partial\alpha^{(\varepsilon)})^{1/2}(f(x) - v)$ if $f(x) - v \leq O(j^{1/2})d(x; \partial\alpha^{(\varepsilon)})^{1/2}$. Moreover,

$$\sup_{x, y \in X_j^{(\varepsilon)}} (f(x) - f(y)) \leq O(j)$$

is shown. Hence one can easily obtain that $\log \tilde{d}(w; \partial\beta)$ is integrable.

REFERENCES

- [1] L. M. Abramov, On the entropy of a flow, Dokl. Akad. Nauk SSSR **128** (1959), 873–876.=Amer. Math. Soc. Transl. (2) **49** (1966), 167–170.
- [2] N. A. Friedman and D. S. Ornstein, On isomorphism of weak Bernoulli transformations, Advances in Math. **5** (1970), 365–394.
- [3] G. Gallavotti and D. S. Ornstein, Billiards and Bernoulli schemes, Commun. math. Phys. **38** (1974), 83–101.
- [4] Sh. Ito, H. Murata and H. Totoki, Remarks on the isomorphism theorems for weak Bernoulli transformations in general case, Publ. RIMS, Kyoto Univ. **7** (1972), 541–580.
- [5] I. Kubo, Perturbed billiard systems I. The ergodicity of the motion of a particle in a compound central field, Nagoya Math. J. **61** (1976), 1–57.
- [6] —, The ergodicity of the motion of a particle in a potential field, International symposium on mathematical problems in theoretical physics, Jan. 1975 Kyoto Japan, 141–148.
- [7] D. S. Ornstein, Two Bernoulli shifts with infinite entropy are isomorphic, Advances in Math. **5** (1970), 339–348.
- [8] —, Imbedding Bernoulli shifts in flows, Lecture Notes in Math. 160 [L. Sucheston (editor): Contribution to ergodic theory and probability] Springer-Verlag (1970), 178–218.
- [9] D. S. Ornstein and B. Weiss, Geodesic flows are Bernoullian, Israel J. Math. **14** (1973), 184–198.
- [10] V. A. Rohlin and Ya G. Sinai, Construction and properties of invariant measurable partitions, Dokl. Akad. Nauk SSSR **141** (1961), 1038–1041.=Soviet Math. Dokl. **2** (1961), 1611–1614.
- [11] Ya. G. Sinai, On the foundations of the ergodic hypothesis for a dynamical system of statistical mechanics, Dokl. Akad. Nauk SSSR **153** (1963), 1261–1264.=Soviet Math. Dokl. **4** (1963), 1818–1822.
- [12] —, Dynamical systems with elastic reflections, Uspehi Math. Nauk **25** (1970), 141–192.=Russian Math. Surveys **25** (1970), 137–189.

Nagoya University
Hiroshima University