

DIFFERENTIAL OPERATORS ON A HYPERSURFACE

BALWANT SINGH*

Introduction

We study differential operators on an affine algebraic variety, especially a hypersurface, in the context of Nakai's Conjecture. We work over a field k of characteristic zero. Let X be a reduced affine algebraic variety over k and let A be its coordinate ring. Let $\text{Diff}_k^n(A)$ be the A -module of differential operators of A over k of order $\leq n$. Nakai's Conjecture asserts that if $\text{Diff}_k^n(A)$ is generated by $\text{Diff}_k^1(A)$ for every $n \geq 2$ then A is regular. In 1973 Mount and Villamayor [6] proved this in the case when X is an irreducible curve. In the general case no progress seems to have been made on the conjecture, except for a result of Brown [2], where the assertion is proved under an additional hypothesis.⁽¹⁾ An interesting result proved by Becker [1] and Rego [8] says that Nakai's Conjecture implies the Conjecture of Zariski-Lipman, which is still open in the general case and which asserts that if the module of k -derivations of A is A -projective then A is regular.

Write $A = R/J$, where R is a polynomial ring over k and J is an ideal of R . Let $\text{Diff}_k^n(R, A)$ be the A -module of differential operators of R into A over k of order $\leq n$. Since R is a polynomial ring, the structure of $\text{Diff}_k^n(R, A)$ is well-known, and $\text{Diff}_k^n(A)$ can be identified with the A -submodule of those $D \in \text{Diff}_k^n(R, A)$ for which $D(J) = 0$. In this paper we first analyze the condition " $D(J) = 0$ " in some detail, and prove in Proposition (2.10) that for $D(J)$ to be zero it is sufficient (and necessary) that D and certain other differential operators derived from D vanish on a set of generators of J . This is then used to prove that if X is a hypersurface, i.e. if we can write $A = R/J$ with J principal, then $\text{Diff}_k^n(A)$ is *completely*

Received December 27, 1984.

^(*) During the course of this work I was supported generously by the research grant of Leslie Roberts and enjoyed the hospitality of Queen's University. To both of them I express my gratitude.

⁽¹⁾ See additional remarks in the last paragraph of this section.

determined by $\text{Diff}_k^1(A)$ (Theorem (2.13)), although, of course, in general, $\text{Diff}_k^2(A)$ is *not generated* by $\text{Diff}_k^1(A)$. This relation between $\text{Diff}_k^1(A)$ and $\text{Diff}_k^2(A)$ leads us to consider the following question, which is stronger than Nakai's Conjecture: *If $\text{Diff}_k^2(A)$ is generated by $\text{Diff}_k^1(A)$ then is A regular?* We are able to show that the answer is in the affirmative in the following two cases: (1) if X is a plane curve (Theorem (3.3)); (2) if X is a cone in 3-space (Theorem (5.3)). In the process of proving Theorem (3.3) we get the following interesting result proved in Theorem (3.1): If X is a plane curve then the quotient $\text{Diff}_k^2(A)/\text{Diff}_k^1(A)\text{Diff}_k^1(A)$ is isomorphic to $\alpha_1 \cap \alpha_2/\alpha_1\alpha_2$, where α_1, α_2 are certain ideals of A .

The author would like to express his thanks to Sankar Dutta for supplying a proof of Proposition (3.2).

After this work was submitted for publication, the paper [12] of J.-P. Vigué was brought to our attention by the referee.⁽¹⁾ It follows from Proposition 5 of this paper of Vigué that if X is a *normal* cone in the complex affine 3-space and $\text{Diff}_k^2(A)$ is generated by $\text{Diff}_k^1(A)$ then A is regular. It has also come to our notice in the meantime that Y. Ishibashi has recently proved Nakai's Conjecture in case X is a two-dimensional complete intersection cone (with k algebraically closed) [13] and also in case X is the quotient of an affine n -space over k by a finite subgroup of $GL(n, k)$ [14]. There is thus an overlap between our Theorem (5.3) and the results of Vigué and Ishibashi. Our methods, however, are entirely different and more elementary and might therefore be of some interest, since very little seems to be known about the structure of differential operators on X .

§1. Preliminaries

Throughout this paper all rings are assumed to be commutative with 1.

The letter n (resp. m) will denote either an integer or ∞ .

Let k be a ring, let R be a k -algebra and let A be an R -module. The multiplication on A by an element $a \in R$ is denoted a_A . We shall regard $\text{Hom}_k(R, A)$ as an R -module via A , i.e. for $a \in R$ and $D \in \text{Hom}_k(R, A)$ we let $aD = a_A \circ D$. For $a \in R$ and $D \in \text{Hom}_k(R, A)$ the symbol $[D, a]$ denotes, as usual, the element $D \circ a_R - a_A \circ D$ of $\text{Hom}_k(R, A)$. For $n \in \mathbb{Z}$ the R -

⁽¹⁾ The author would like to thank the referee for this and for pointing out some misprints in the manuscript.

submodule $\text{Diff}_k^n(R, A)$ of $\text{Hom}_k(R, A)$ is defined by induction on n : $\text{Diff}_k^n(R, A) = 0$ if $n < 0$, and

$$\text{Diff}_k^n(R, A) = \{D \in \text{Hom}_k(R, A) \mid [D, a] \in \text{Diff}_k^{n-1}(R, A) \text{ for every } a \in R\}.$$

Elements of $\text{Diff}_k^n(R, A)$ are called *differential operators* (of R into A over k) of order $\leq n$.

(1.1) *Remark.* Let $\{a_i\}$ be a set of k -algebra generators of R . It is easily checked that if $D \in \text{Hom}_k(R, A)$ and $[D, a_i] \in \text{Diff}_k^{n-1}(R, A)$ for every i then $D \in \text{Diff}_k^n(R, A)$.

Note that $\text{Diff}_k^0(R, A) = \text{Hom}_R(R, A) \cong A$ and $\text{Diff}_k^n(R, A) \subset \text{Diff}_k^{n+1}(R, A)$ for every n . Put $\text{Diff}_k^\infty(R, A) = \bigcup_{n \in \mathbb{Z}} \text{Diff}_k^n(R, A)$.

Next, recall from Nakai [7] the definition of a high order derivation. For $n \in \mathbb{Z}$, $n \geq 0$, an element D of $\text{Hom}_k(R, A)$ is called an *n th order derivation* (of R into A over k) if for any $n + 1$ elements a_0, \dots, a_n of R we have

$$D(a_0 \cdots a_n) = \sum_{s=1}^n (-1)^{s+1} \sum_{i_1 < \cdots < i_s} a_{i_1} \cdots a_{i_s} D(a_0 \cdots \hat{a}_{i_1} \cdots \hat{a}_{i_s} \cdots a_n),$$

where $\hat{}$ means omitted. Denote by $\text{Der}_k^n(R, A)$ the R -submodule of $\text{Hom}_k(R, A)$ consisting of all n th order derivations. We have $\text{Der}_k^0(R, A) = 0$ and $\text{Der}_k^1(R, A) =$ the module of ordinary k -derivations of R into A . Define $\text{Der}_k^n(R, A) = 0$ if $n < 0$.

We note the relation between $\text{Diff}_k^n(R, A)$ and $\text{Der}_k^n(R, A)$ in the following lemma. For $x \in A$ let x_R denote the R -linear map $a \mapsto ax$ of R into A .

(1.2) LEMMA.

- (1) Let $D \in \text{Hom}_k(R, A)$. Then $D \in \text{Diff}_k^n(R, A)$ if and only if $D - D(1)_R \in \text{Der}_k^n(R, A)$.
- (2) $\text{Der}_k^n(R, A) = \{D \in \text{Diff}_k^n(R, A) \mid D(1) = 0\}$. Consequently, $\text{Der}_k^n(R, A) \subset \text{Der}_k^{n+1}(R, A)$ for every n .
- (3) Put $\text{Der}_k^\infty(R, A) = \bigcup_{n \in \mathbb{Z}} \text{Der}_k^n(R, A)$. For $0 \leq n \leq \infty$ we have an R -linear isomorphism $h: \text{Diff}_k^n(R, A) \rightarrow A \oplus \text{Der}_k^n(R, A)$ given by $h(D) = (D(1), D - D(1)_R)$ and $h^{-1}(x, D) = x_R + D$.

Proof. (1) follows from the formula defining an n th order derivation and a similar characterization of a differential operator given in [4, (IV, 16.8.8)]. (2) and (3) follow from (1) by noting that if $D \in \text{Der}_k^n(A)$ then $D(1) = 0$.

Let S be a multiplicatively closed subset of R . If k is noetherian and R is a finitely generated k -algebra then it follows from [4, (IV, 16.4.15)] that for every n there is a natural isomorphism $S^{-1}(\text{Diff}_k^n(R, A)) \cong \text{Diff}_k^n(S^{-1}R, S^{-1}A)$. We use this isomorphism to identify these two modules. Then $S^{-1}(\text{Der}_k^n(R, A)) = \text{Der}_k^n(S^{-1}R, S^{-1}A)$ by Lemma (1.2).

Assume now that $A = R$. Write $\text{Diff}_k^n(A)$ for $\text{Diff}_k^n(A, A)$ and $\text{Der}_k^n(A)$ for $\text{Der}_k^n(A, A)$. If $D \in \text{Diff}_k^n(A)$ and $D' \in \text{Diff}_k^m(A)$ then $DD' \in \text{Diff}_k^{n+m}(A)$ ([4, (IV, 16.8.9)] or [10, Proposition 1]). Further, if $D' \in \text{Der}_k^m(A)$ then $DD'(1) = 0$, so that $DD' \in \text{Der}_k^{n+m}(A)$.

For an integer $n \geq 2$ let us say that $\text{Diff}_k^n(A)$ is *generated* by $\text{Diff}_k^1(A)$ if $\text{Diff}_k^n(A)$ equals the A -submodule of all finite sums of the products $D_1 \cdots D_n$ with $D_1, \dots, D_n \in \text{Diff}_k^1(A)$. We shall say that $\text{Diff}_k^\infty(A)$ is generated by $\text{Diff}_k^1(A)$ if $\text{Diff}_k^n(A)$ is generated by $\text{Diff}_k^1(A)$ for every $n \geq 2$.

Denote by $\text{Diff}_k^1(A) \text{Diff}_k^{n-1}(A)$ the A -submodule of $\text{Diff}_k^n(A)$ consisting of all finite sums $\sum_i D_i D'_i$ with $D_i \in \text{Diff}_k^1(A)$, $D'_i \in \text{Diff}_k^{n-1}(A)$. Let $\text{Der}_k^1(A) \text{Der}_k^{n-1}(A)$ have a similar meaning.

From Lemma (1.2) we immediately get

(1.3) LEMMA. For $n \geq 2$ we have $\text{Diff}_k^1(A) \text{Diff}_k^{n-1}(A) = \text{Diff}_k^{n-1}(A) + \text{Der}_k^1(A) \text{Der}_k^{n-1}(A)$. Moreover the following three conditions are equivalent:

- (i) $\text{Diff}_k^\infty(A)$ is generated by $\text{Diff}_k^1(A)$.
- (ii) $\text{Diff}_k^n(A) = \text{Diff}_k^1(A) \text{Diff}_k^{n-1}(A)$ for every $n \geq 2$.
- (iii) $\text{Der}_k^n(A) = \text{Der}_k^{n-1}(A) + \text{Der}_k^1(A) \text{Der}_k^{n-1}(A)$ for every $n \geq 2$.

Suppose now that k is a field of characteristic zero and A is a finitely generated k -algebra. Then it follows from [4, (IV, 16.11.2)] that if A is regular then $\text{Diff}_k^\infty(A)$ is generated by $\text{Diff}_k^1(A)$. Nakai's Conjecture asserts the converse:

NAKAI'S CONJECTURE. Let A be a finitely generated algebra over a field k of characteristic zero. If $\text{Diff}_k^\infty(A)$ is generated by $\text{Diff}_k^1(A)$ then A is regular.

§ 2. General results

Let k be a field of characteristic zero. Let $R = k[X_1, \dots, X_r]$ be the polynomial ring in r variables over k . Let J be a proper ideal of R , let $A = R/J$ and let $\eta: R \rightarrow A$ be the natural map. Put $x_i = \eta(X_i)$ for $1 \leq i \leq r$.

Let Z^+ be the set of all non-negative integers and put $V = (Z^+)^r$. For

$\alpha = (\alpha_1, \dots, \alpha_r) \in V$ we use the standard notation: $|\alpha| = \alpha_1 + \dots + \alpha_r$, $\alpha! = \alpha_1! \dots \alpha_r!$, $X^\alpha = X_1^{\alpha_1} \dots X_r^{\alpha_r}$, etc. For $n \in \mathbf{Z}$ let $V_n = \{\alpha \in V \mid |\alpha| \leq n\}$ and $W_n = \{\alpha \in V \mid |\alpha| = n\}$. For $1 \leq i \leq r$ let $e_i = (0, \dots, 1, \dots, 0) \in W_1$ with 1 in the i th place.

For $\alpha \in V$ let $\Delta_\alpha: R \rightarrow A$ denote the composite of $(1/\alpha!) \partial^\alpha / \partial X^\alpha: R \rightarrow R$ and η . Then $\Delta_\alpha \in \text{Diff}_k^{|\alpha|}(R, A)$. It is well-known that every $D \in \text{Diff}_k^\infty(R, A)$ has a unique expression of the form

$$D = \sum_{\alpha \in V} c_\alpha(D) \Delta_\alpha$$

with $c_\alpha(D) \in A$ for all α and $c_\alpha(D) = 0$ for almost all α . Moreover, $D \in \text{Diff}_k^n(R, A)$ if and only if $c_\alpha(D) = 0$ for $|\alpha| > n$. On the other hand, $D \in \text{Diff}_k^n(R, A)$ is also uniquely determined by the values $D(X^\alpha)$ for $\alpha \in V_n$.

For $n \in \mathbf{Z}$ we define a map $\pi_n: \text{Diff}_k^\infty(R, A) \rightarrow \text{Diff}_k^n(R, A)$ as follows: For $D \in \text{Diff}_k^\infty(R, A)$ let $\pi_n(D)$ be the unique element of $\text{Diff}_k^n(R, A)$ determined by $\pi_n(D)(X^\alpha) = D(X^\alpha)$ for all $\alpha \in V_n$.

In (2.1)-(2.3) below, let $D \in \text{Diff}_k^\infty(R, A)$, let $\beta \in V$ and let $j \in \mathbf{Z}$ with $1 \leq j \leq r$.

(2.1) LEMMA.

$$c_\beta([D, X_j]) = c_{\beta + e_j}(D).$$

Proof. Immediate from the observation

$$[\Delta_\beta, X_j] = \begin{cases} \Delta_{\beta - e_j}, & \text{if } \beta_j > 0, \\ 0, & \text{if } \beta_j = 0. \end{cases}$$

(2.2) LEMMA.

$$[\pi_n(D), X_j] = \pi_{n-1}([D, X_j]).$$

Proof. Both sides belong to $\text{Diff}_k^{n-1}(R, A)$. Therefore it is enough to prove that they coincide on X^α for all $\alpha \in V_{n-1}$. Let $\alpha \in V_{n-1}$. Then

$$\begin{aligned} [\pi_n(D), X_j](X^\alpha) &= \pi_n(D)(X_j X^\alpha) - X_j \pi_n(D)(X^\alpha) = D(X_j X^\alpha) - X_j D(X^\alpha) \\ &= [D, X_j](X^\alpha) = \pi_{n-1}([D, X_j])(X^\alpha). \end{aligned}$$

(2.3) PROPOSITION.

$$c_\beta(D) = D(X^\beta) - \pi_{|\beta|-1}(D)(X^\beta).$$

Proof. Induction on $|\beta|$. Evaluating $D = \sum_{\alpha \in V} c_\alpha(D) \Delta_\alpha$ on 1 we get $D(1) = c_0(D)$, which is our assertion for $|\beta| = 0$. Now, let $|\beta| > 0$. Choose

i such that $\beta_i > 0$, and let $\gamma = \beta - e_i$. Then

$$\begin{aligned} c_\beta(D) &= c_\gamma([D, X_i]) && \text{(Lemma (2.1))} \\ &= [D, X_i](X^\gamma) - \pi_{|\gamma|-1}([D, X_i])(X^\gamma) && \text{(Induction)} \\ &= [D, X_i](X^\gamma) - [\pi_{|\beta|-1}(D), X_i](X^\gamma) && \text{(Lemma (2.2))} \\ &= D(X^\beta) - \pi_{|\beta|-1}(D)(X^\beta), \end{aligned}$$

since $\pi_{|\beta|-1}(D)(X^\gamma) = D(X^\gamma)$ by definition.

(2.4) IDENTIFICATION. Let us identify

$$\begin{aligned} \text{Diff}_k^n(A) &= \{D \in \text{Diff}_k^n(R, A) \mid D(J) = 0\}, \\ \text{Der}_k^n(A) &= \{D \in \text{Der}_k^n(R, A) \mid D(J) = 0\} \end{aligned}$$

via the map $D \rightarrow D\eta$.

(2.5) PROPOSITION. Let $D \in \text{Der}_k^1(A)$ and $D' \in \text{Diff}_k^n(A)$. Then for every $\alpha \in W_{n+1}$ we have

$$c_\alpha(DD') = \sum_{i=1}^r \alpha_i D(X_i) c_{\alpha - e_i}(D').$$

(Note that if $\alpha_i = 0$ then $c_{\alpha - e_i}(D')$ is not defined, but then we take the corresponding summand to be zero by standard convention.)

Proof. Induction on n . The assertion being clear for $n < 0$, let $n \geq 0$ and let $\alpha \in W_{n+1}$. We may assume that $\alpha = e_1 + \beta$ with $\beta \in W_n$. By induction we have

$$\begin{aligned} (2.5.1) \quad c_\beta(D[D', X_1]) &= \sum_{i=1}^r \beta_i D(X_i) c_{\beta - e_i}([D', X_1]) \\ &= \sum_{i=1}^r \beta_i D(X_i) c_{\alpha - e_i}(D') \end{aligned}$$

by Lemma (2.1). On the other hand, by Lemma (2.1) again we have

$$(2.5.2) \quad c_\alpha(DD') = c_\beta([DD', X_1]).$$

Thus we need to compare $D[D', X_1]$ and $[DD', X_1]$. Since D is a 1-derivation, we have $[DD', X_1] - D[D', X_1] = D(X_1)D'$. Therefore

$$(2.5.3) \quad c_\beta([DD', X_1] - c_\beta(D[D', X_1])) = c_\beta(D(X_1)D') = D(X_1)c_\beta(D').$$

Now, by (2.5.1), (2.5.2) and (2.5.3) we have

$$\begin{aligned}
 c_\alpha(DD') &= c_\beta([DD', X_1]) = D(X_1)c_\beta(D') + \sum_{i=1}^r \beta_i D(X_i)c_{\alpha-e_i}(D') \\
 &= \sum_{i=1}^r \alpha_i D(X_i)c_{\alpha-e_i}(D').
 \end{aligned}$$

(2.6) DEFINITION. For $D \in \text{Diff}_k^\infty(R, A)$ and $\beta \in V$ define

$$\langle D, X^\beta \rangle = \sum_{\alpha \in V} c_{\alpha+\beta}(D)A_\alpha.$$

Note that $\langle D, 1 \rangle = D$ and if $D \in \text{Diff}_k^n(R, A)$ then $\langle D, X^\beta \rangle \in \text{Diff}_k^{n-|\beta|}(R, A)$.

(2.7) DEFINITION. Let $\Phi: \text{Diff}_k^\infty(R, A) \times V \rightarrow \text{Der}_k^\infty(R, A)$ be the pairing defined by $\Phi(D, \beta) = \langle D, X^\beta \rangle - (\langle D, X^\beta \rangle(1))_R = \langle D, X^\beta \rangle - c_\beta(D)A_0$ (see Lemma (1.2)). Note that Φ is R -linear in the first variable. Further note that Φ is the direct limit of the pairings

$$\Phi_{n,m}: \text{Diff}_k^n(R, A) \times W_m \rightarrow \text{Der}_k^{n-m}(R, A)$$

given by

$$\Phi_{n,m}(D, \beta) = \langle D, X^\beta \rangle - c_\beta(D)A_0.$$

(2.8) PROPOSITION. For $m \leq n$ we have an exact sequence

$$0 \rightarrow \text{Diff}_k^m(R, A) \hookrightarrow \text{Diff}_k^n(R, A) \xrightarrow{\Theta_{n,m}} \bigoplus_{\beta \in W_m} \text{Der}_k^{n-m}(R, A),$$

where $\Theta_{n,m}(D) = (\Phi_{n,m}(D, \beta))_{\beta \in W_m}$.

Proof. We have $\Theta_{n,m}(\text{Diff}_k^m(R, A)) \subset \text{Der}_k^0(R, A) = 0$. Suppose $\Theta_{n,m}(D) = 0$. Then $\langle D, X^\beta \rangle = c_\beta(D)A_0$ for every $\beta \in W_m$. By definition, $\langle D, X^\beta \rangle = \sum_{\alpha \in V} c_{\alpha+\beta}(D)A_\alpha$. Therefore by the uniqueness of this expression we have $c_{\alpha+\beta}(D) = 0$ for $|\alpha| > 0$. This being so for every $\beta \in W_m$, we get $c_\alpha(D) = 0$ for all $\alpha \in V$ with $|\alpha| > m$. This means that $D \in \text{Diff}_k^m(R, A)$.

(2.9) LEMMA. Let $D \in \text{Diff}_k^\infty(R, A)$, let $\beta, \gamma \in V$ and let $j \in \mathbf{Z}$ with $1 \leq j \leq r$. Then:

- (1) $\langle D, X_j \rangle = [D, X_j]$.
- (2) $\langle \langle D, X^\beta \rangle, X^\gamma \rangle = \langle D, X^{\beta+\gamma} \rangle$.
- (3) $\langle [D, X_j], X^\beta \rangle = \langle D, X_j X^\beta \rangle$.
- (4) If $X^\beta = Y_1 \cdots Y_s$ with $Y_1, \dots, Y_s \in \{X_1, \dots, X_r\}$ then $\langle D, X^\beta \rangle = [\cdots [[D, Y_1], Y_2], \dots, Y_s]$.

Proof. (1) follows from Lemma (2.1). For $\alpha \in V$ we have $c_\alpha(\langle \langle D, X^\beta \rangle, X^\gamma \rangle) = c_{\alpha+\gamma}(\langle D, X^\beta \rangle) = c_{\alpha+\gamma+\beta}(D) = c_\alpha(\langle D, X^{\beta+\gamma} \rangle)$, which proves (2). (3) follows from (1) and (2), and (4) follows from (3).

(2.10) PROPOSITION. *Let I be an ideal of R . For $D \in \text{Diff}_k^n(R, A)$ the following five conditions are equivalent:*

- (i) $D(I) \subset IA$.
- (ii) $[D, X_i](I) \subset IA$ for all $i, 1 \leq i \leq r$, and there exists a set of generators $\{f_j\}$ of I such that $D(f_j) \in IA$ for every j .
- (iii) $\langle D, X^\beta \rangle(I) \subset IA$ for every $\beta \in V$.
- (iv) $\langle D, X^\beta \rangle(I) \subset IA$ for every $\beta \in V_{n-1}$.
- (v) *There exists a set of generators $\{f_j\}$ of I such that $\langle D, X^\beta \rangle(f_j) \in IA$ for every $\beta \in V_{n-1}$ and for every j .*

Proof. The implications (i) \Rightarrow (ii) and (iii) \Rightarrow (iv) \Rightarrow (v) are trivial, while the implication (i) \Rightarrow (iii) is immediate from part (4) of the above lemma.

(ii) \Rightarrow (i). It is enough to prove that $D(X^\beta f_j) \in IA$ for every $\beta \in V$ and for every j . For $|\beta| = 0$ this is an assumption in (ii). If $|\beta| > 0$ then we may assume that $\beta = e_i + \gamma$ with $\gamma \in V$. Then $D(X^\beta f_j) = [D, X_i](X^\gamma f_j) + X_i D(X^\gamma f_j)$. Therefore the assertion follows by induction on $|\beta|$.

(v) \Rightarrow (i). Induction on n . The assertion being clear for $n \leq 0$, assume that $n \geq 1$. We have $\langle [D, X_i], X^\beta \rangle = \langle D, X_i X^\beta \rangle$ by Lemma (2.9). Therefore by (v) $\langle [D, X_i], X^\beta \rangle(f_j) \in IA$ for every $\beta \in V_{n-2}$ and for every j . So, by induction, $[D, X_i](I) \subset IA$ for every $i, 1 \leq i \leq r$. Further, by taking $\beta = 0$ in (v) we have $D(f_j) \in IA$ for every j . Therefore, by (ii) \Rightarrow (i), $D(I) \subset IA$.

Applying the proposition with $I = J$ and using the identification (2.4) we get

(2.11) COROLLARY. *For $D \in \text{Diff}_k^n(R, A)$ the following three conditions are equivalent:*

- (i) $D \in \text{Diff}_k^n(A)$.
- (ii) $\langle D, X^\beta \rangle \in \text{Diff}_k^{n-|\beta|}(A)$ for every $\beta \in V$.
- (iii) $\langle D, X^\beta \rangle \in \text{Diff}_k^{n-|\beta|}(A)$ for every $\beta \in V_{n-1}$.

In view of the above corollary the pairings $\Phi_{n,m}$ of Definition (2.7) induce pairings

$$\varphi_{n,m} : \text{Diff}_k^n(A) \times W_m \longrightarrow \text{Der}_k^{n-m}(A).$$

It follows from Proposition (2.8) that for $m \leq n$ we have an exact sequence

$$0 \rightarrow \text{Diff}_k^m(A) \hookrightarrow \text{Diff}_k^n(A) \xrightarrow{\theta_{n,m}} \bigoplus_{\beta \in W_m} \text{Der}_k^{n-m}(A),$$

where $\theta_{n,m}(D) = (\varphi_{n,m}(D, \beta))_{\beta \in W_m}$.

(2.12) DEFINITION. For $n \in \mathbb{Z}$ define

$$\mathcal{D}_k^n(A) = \{(d_\beta)_{\beta \in W_{n-1}} \in \bigoplus_{\beta \in W_{n-1}} \text{Der}_k^1(A) \mid d_\beta(x_i) = d_\gamma(x_j) \text{ whenever } \beta + e_i = \gamma + e_j, \beta, \gamma \in W_{n-1}, 1 \leq i, j \leq r\}.$$

If $D \in \text{Diff}_k^n(A)$ and $\theta_{n,n-1}(D) = (d_\beta)_{\beta \in W_{n-1}}$ then $d_\beta(x_i) = c_{\beta+e_i}(D)$. It follows that $\text{im}(\theta_{n,n-1}) \subset \mathcal{D}_k^n(A)$. Therefore, writing $\theta_n = \theta_{n,n-1}$, we have an exact sequence

$$0 \rightarrow \text{Diff}_k^{n-1}(A) \hookrightarrow \text{Diff}_k^n(A) \xrightarrow{\theta_n} \mathcal{D}_k^n(A).$$

Of particular interest to us is the exact sequence

$$0 \rightarrow \text{Diff}_k^1(A) \hookrightarrow \text{Diff}_k^2(A) \xrightarrow{\theta_2} \mathcal{D}_k^2(A).$$

We note that

$$\mathcal{D}_k^2(A) = \{(d_1, \dots, d_r) \in \bigoplus_{i=1}^r \text{Der}_k^1(A) \mid d_i(x_j) = d_j(x_i) \text{ for all } i, j\}$$

and that if $D \in \text{Diff}_k^2(A)$ then $\theta_2(D) = (d_1, \dots, d_r)$, where $d_i \in \text{Der}_k^1(A)$ is given by $d_i(x_j) = c_{e_i+e_j}(D)$. In the following theorem we prove that if J is principal then θ_2 is surjective.

(2.13) THEOREM. Suppose J is principal. Then the sequence

$$0 \rightarrow \text{Diff}_k^1(A) \hookrightarrow \text{Diff}_k^2(A) \xrightarrow{\theta_2} \mathcal{D}_k^2(A) \rightarrow 0$$

is exact.

Proof. We have only to prove the surjectivity of θ_2 . Let $(d_1, \dots, d_r) \in \mathcal{D}_k^2(A)$. Let $a_{ij} \in R$ be a lift of $d_i(x_j)$ such that $a_{ij} = a_{ji}$ for all i, j . Then, since d_i is a 1-derivation of A we have

$$(E_i) \quad \sum_{j=1}^r a_{ij} \partial f / \partial X_j = g_i f$$

with $g_i \in R$, where f is a generator of J . Differentiating E_i with respect to X_i , adding the results over $1 \leq i \leq r$, dividing the sum by 2 and remembering that $a_{ij} = a_{ji}$, we get

$$\sum_{j=1}^r b_j \partial f / \partial X_j + \frac{1}{2} \sum_{j=1}^r a_j \partial^2 f / \partial X_j^2 + \sum_{i < j} a_{ij} \partial^2 f / \partial X_i \partial X_j = g f$$

with $b_1, \dots, b_r, g \in R$. Define $D = \sum_{\alpha \in V_2} c_\alpha(D) \Delta_\alpha \in \text{Diff}_k^2(R, A)$ by $c_0(D) = 0$,

$c_{e_i}(D) = \eta(b_i)$ and $c_{e_i+e_j}(D) = \eta(a_{ij})$, $1 \leq i, j \leq r$. Then the above equality implies that $D(f) = 0$. Moreover, we have $\langle D, X_i \rangle = \sum_{\alpha \in V_1} c_{\alpha+e_i}(D) \Delta_\alpha = \eta(b_i) \Delta_0 + \sum_{j=1}^r \eta(a_{ij}) \Delta_{e_j}$, so that $\langle D, X_i \rangle(f) = \eta(b_i f + \sum_{j=1}^r a_{ij} \partial f / \partial X_j) = 0$ by (E_i) . This proves that $\langle D, X^\beta \rangle(f) = 0$ for every $\beta \in V_1$. Therefore $D(Rf) = 0$ by Proposition (2.10), and we get $D \in \text{Diff}_k^2(A)$. Since $c_{e_i+e_j}(D) = d_i(x_j)$, we have $\theta_2(D) = (d_1, \dots, d_r)$. This proves that θ_2 is surjective.

(2.14) *Remark.* The above proof actually constructs a k -linear right inverse of θ_2 . For if we define $t(d_1, \dots, d_r) = D$ in the above notation then it is easily checked that $t(d_1, \dots, d_r)$ is independent of the choice of lifts a_{ij} of $d_i(x_j)$ with the condition that $a_{ij} = a_{ji}$.

(2.15) *Remark.* In general, θ_3 is not surjective. *Example:* Take $r = 2$ and $J = Rf$ with $f = X_2^2 - X_1^3$. For $\beta \in W_2$ define $d_\beta \in \text{Der}_k^1(A)$ as follows: $d_{(2,0)} = 8x_1 \Delta_{e_1} + 12x_2 \Delta_{e_2}$, $d_{(1,1)} = 12x_2 \Delta_{e_1} + 18x_1^2 \Delta_{e_2}$, $d_{(0,2)} = 18x_1^2 \Delta_{e_1} + 27x_1 x_2 \Delta_{e_2}$. Then $d = (d_{(2,0)}, d_{(1,1)}, d_{(0,2)}) \in \mathcal{D}_k^3(A)$. We claim that $d \notin \text{im}(\theta_3)$. For suppose $d = \theta_3(D)$ with $D \in \text{Diff}_k^3(A)$. Put $c_{ij} = c_{(i,j)}(D)$. Then $c_{30} = 8x_1$, $c_{21} = 12x_2$, $c_{12} = 18x_1^2$ and $c_{03} = 27x_1 x_2$. By Corollary (2.11) we have $D(f) = 0$, $\langle D, x_1 \rangle(f) = 0$ and $\langle D, x_2 \rangle(f) = 0$. Computing $\langle D, x_2 \rangle(f) - 2x_2 D(f) = 0$ we find that $(7 + 6c_{20})x_1 x_2 \in (x_1^2, x_2^2)$. This implies that $7 + 6c_{20} \in (x_1, x_2)$. From $\langle D, x_1 \rangle(f) = 0$ we get $(6 + 3c_{20})x_1^2 \in (x_2)$, which implies that $6 + 3c_{20} \in (x_1, x_2)$. Thus we get $5 \in (x_1, x_2)$, a contradiction.

§ 3. Differential operators on a plane curve

With the notation of Section 2, assume that $r = 2$ and $J = Rf$ is a non-zero, proper, principal ideal of R . For $i = 1, 2$, put $f_{x_i} = \eta(\partial f / \partial X_i)$ and let $\alpha_i = \{d(x_i) \mid d \in \text{Der}_k^1(A)\}$. Then α_1, α_2 are ideals of A . Recall that in this case we have

$$\mathcal{D}_k^2(A) = \{(d_1, d_2) \in \text{Der}_k^1(A) \oplus \text{Der}_k^1(A) \mid d_1(x_2) = d_2(x_1)\}.$$

Let $\tau: \mathcal{D}_k^2(A) \rightarrow \alpha_1 \cap \alpha_2$ be the map defined by $\tau(d_1, d_2) = d_1(x_2) = d_2(x_1)$. Clearly, τ is A -linear and surjective. Put $\sigma = \tau\theta_2: \text{Diff}_k^2(A) \rightarrow \alpha_1 \cap \alpha_2$.

(3.1) **THEOREM.** *Assume that f_{x_1} and f_{x_2} are non-zero divisors in A . Then τ is an isomorphism, and the sequence*

$$0 \longrightarrow \text{Diff}_k^1(A) \hookrightarrow \text{Diff}_k^2(A) \xrightarrow{\sigma} \alpha_1 \cap \alpha_2 \longrightarrow 0$$

is exact. Moreover, $\sigma(\text{Diff}_k^1(A) \text{Diff}_k^1(A)) = \alpha_1 \alpha_2$. In particular, we have an A -isomorphism

$$\text{Diff}_k^2(A)/\text{Diff}_k^1(A) \text{ Diff}_k^1(A) \cong \alpha_1 \cap \alpha_2/\alpha_1\alpha_2.$$

Proof. If $d \in \text{Der}_k^1(A)$ then $d(x_1)f_{x_1} + d(x_2)f_{x_2} = 0$. Since f_{x_1} and f_{x_2} are non-zero divisors, we have $d(x_1) = 0$ if and only if $d(x_2) = 0$. Consequently, $d = 0$ if and only if $d(x_1) = 0$ if and only if $d(x_2) = 0$. This proves that τ is an isomorphism. The exactness of the sequence follows now from Theorem (2.13). Since $\text{Diff}_k^1(A) \text{ Diff}_k^1(A) = \text{Diff}_k^1(A) + \text{Der}_k^1(A) \text{ Der}_k^1(A)$ by Lemma (1.3) and $\sigma(\text{Diff}_k^1(A)) = 0$, the remaining part of the theorem will follow if we show that $\sigma(\text{Der}_k^1(A) \text{ Der}_k^1(A)) = \alpha_1\alpha_2$. Let $D, D' \in \text{Der}_k^1(A)$ and let $\theta_2(DD') = (d_1, d_2)$. Then $\sigma(DD') = \tau(d_1, d_2) = d_1(x_2) = c_{(1,1)}(DD')$ by the remarks preceding Theorem (2.13). By Proposition (2.5) we have $c_{(1,1)}(DD') = D(x_1)c_{(0,1)}(D') + D(x_2)c_{(1,0)}(D') = D(x_1)D'(x_2) + D(x_2)D'(x_1)$. This proves that $\sigma(\text{Der}_k^1(A) \text{ Der}_k^1(A)) \subset \alpha_1\alpha_2$. Conversely, let $a_i \in \alpha_i, i = 1, 2$. Choose $D, D' \in \text{Der}_k^1(A)$ such that $a_1 = D(x_1), a_2 = D'(x_2)$. Let $b_1 = D(x_2), b_2 = D'(x_1)$. Then $a_1f_{x_1} + b_1f_{x_2} = 0$ and $b_2f_{x_1} + a_2f_{x_2} = 0$. Therefore, since f_{x_1} is a non-zero divisor, we get $a_1a_2 = b_1b_2$. Now, by the above computation we have $\sigma(DD') = a_1a_2 + b_1b_2 = 2a_1a_2$. This proves that $\alpha_1\alpha_2 \subset \sigma(\text{Der}_k^1(A) \text{ Der}_k^1(A))$, and the theorem is proved.

(3.2) PROPOSITION. Let I_1, I_2 be ideals of R containing f . Assume that R/I_1 and R/I_2 are of finite length and that $I_1 + I_2 \neq R$. Then $\text{Tor}_1^A(R/I_1, R/I_2) \neq 0$.

Proof (S. Dutta). Localizing at a maximal ideal of R containing $I_1 + I_2$, we may assume that R is a regular local ring of dimension two containing k . By [3, Chapter XV, Section 5, Case C' and Chapter XVI, Section 5, Case 1] we have an exact sequence

$$\text{Tor}_1^R(R/I_1, A) \otimes_R R/I_2 \longrightarrow \text{Tor}_1^R(R/I_1, R/I_2) \longrightarrow \text{Tor}_1^A(R/I_1, R/I_2) \longrightarrow 0.$$

Since f is a non-zero divisor in R , we have $\text{Tor}_1^R(R/I_1, A) \cong R/I_1$. Therefore the exact sequence becomes

$$R/I_1 \otimes_R R/I_2 \longrightarrow \text{Tor}_1^R(R/I_1, R/I_2) \longrightarrow \text{Tor}_1^A(R/I_1, R/I_2) \longrightarrow 0.$$

Put $t = \text{length}(\text{Tor}_1^A(R/I_1, R/I_2))$ and $t_i = \text{length}(\text{Tor}_i^R(R/I_1, R/I_2))$. Then t, t_i are non-negative integers and $t_i = 0$ for $i \neq 0, 1, 2$. From the exact sequence we have $t - t_1 + t_0 \geq 0$. Since $\dim(R/I_1) + \dim(R/I_2) = 0 < 2$, we have $t_0 - t_1 + t_2 = 0$ by [9, Chapter V, Section 3]. Thus we get $t \geq t_2$, and it is now enough to prove that $t_2 > 0$. Let

$$0 \longrightarrow R^n \xrightarrow{\lambda} R^{n+1} \longrightarrow R \longrightarrow R/I_2 \longrightarrow 0$$

be a minimal free resolution of R/I_2 . Let \mathfrak{m} be the maximal ideal of R . Choose $a \in R$ such that $a \notin I_1$ and $\mathfrak{m}a \subset I_1$. Let $b = \bar{a} \otimes (1, 0, \dots, 0) \in R/I_1 \otimes_R R^n$, where \bar{a} is the natural image of a in R/I_1 . Then $b \neq 0$ and, since $\text{im}(\lambda) \subset \mathfrak{m}R^{n+1}$, $b \in \ker(1_{R/I_1} \otimes \lambda) = \text{Tor}_2^R(R/I_1, R/I_2)$. This proves that $t_2 > 0$.

(3.3) THEOREM. *Let J be a proper, principal ideal of $R = k[X_1, X_2]$ and let $A = R/J$. Assume that A is reduced.⁽¹⁾ Then A is regular if and only if $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \text{Diff}_k^1(A)$.*

Proof. If A is regular then the equality $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \text{Diff}_k^1(A)$ follows from [4, (IV, 16.11.2)]. Conversely, suppose $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \text{Diff}_k^1(A)$. We may assume that $J \neq 0$. Let $J = Rf$ and let $f = f_1 \cdots f_s$ be the prime factorization of f in R . Since k is infinite, we may make a linear change of variables to assume that $\partial f_i / \partial X_j \neq 0$ for all $i = 1, \dots, s, j = 1, 2$. Then, since f is without multiple factors, f_{x_1} and f_{x_2} are non-zero divisors in A . Since $f_{x_2} \mathcal{A}_{(1,0)} - f_{x_1} \mathcal{A}_{(0,1)} \in \text{Der}_k^1(A)$, we have $f_{x_2} \in \alpha_1$ and $f_{x_1} \in \alpha_2$. Therefore A/α_1 and A/α_2 are of finite length. Let I_j be the ideal of R containing f such that $\alpha_j = I_j/Rf, j = 1, 2$. Since $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \text{Diff}_k^1(A)$, we have, by Theorem (3.1), $0 = \alpha_1 \cap \alpha_2/\alpha_1 \alpha_2 \cong \text{Tor}_1^A(A/\alpha_1, A/\alpha_2) = \text{Tor}_1^A(R/I_1, R/I_2)$. Therefore $I_1 + I_2 = R$ by Proposition (3.2). Now, let \mathfrak{m} be a maximal ideal of A . Then α_1 or α_2 , say α_1 , is not contained in \mathfrak{m} . This means that there exists $D \in \text{Der}_k^1(A)$ such that $D(x_1) \notin \mathfrak{m}$. Since $f_{x_2} \neq 0, x_1$ is transcendental over k . Since A/\mathfrak{m} is algebraic over k, \mathfrak{m} contains a non-zero polynomial $g(x_1) \in k[x_1]$. Choose such g of least degree. Then $D(g) = \eta(\partial g / \partial X_1) D(x_1) \notin \mathfrak{m}$. Thus $D(\mathfrak{m}) \not\subset \mathfrak{m}$. Let B be the $\mathfrak{m}A_{\mathfrak{m}}$ -adic completion of $A_{\mathfrak{m}}$ and let \mathfrak{n} be the maximal ideal of B . Then D extends to a 1-derivation \hat{D} of B such that $\hat{D}(\mathfrak{n}) \not\subset \mathfrak{n}$. Therefore by Zariski's Lemma [11, Lemma 4] B is of the form $B = B_0[[Y]]$ with Y analytically independent over B_0 . Since B , hence B_0 , is reduced and $\dim B_0 = 0, B_0$ is regular. Therefore B is regular. This proves that $A_{\mathfrak{m}}$ is regular for every maximal ideal \mathfrak{m} of A .

§ 4. $A[T]$ and $A[T, T^{-1}]$

In this section let k be a noetherian ring, let A be a finitely generated k -algebra and let T be an indeterminate over A . Let $u: A \hookrightarrow A[T]$ be the natural inclusion. For $i \in \mathbb{Z}, i \geq 0$, let $p_i: A[T] \rightarrow A$ be the A -linear map defined by $f = \sum_{i \geq 0} (T - 1)^i p_i(f)$ for $f \in A[T]$. Define $q_i: \text{Diff}_k^\infty(A[T])$

⁽¹⁾ See note added in proof.

→ Diff_k[∞](A) by q_i(D) = p_iDu for D ∈ Diff_k[∞](A[T]). Note that p_iDu ∈ Diff_k[∞](A), since p_i is A-linear. Note also that q_i maps Diff_kⁿ(A[T]) into Diff_kⁿ(A) for every n. Further, we have Du = ∑_{i≥0} (T - 1)ⁱuq_i(D) for D ∈ Diff_k[∞](A[T]).

Regard Diff_k[∞](A, A[T]) as an A[T]-module in a natural way. For D ∈ Diff_k[∞](A, A[T]) let μ(D) denote the k[T]-linear endomorphism of A[T] obtained from D by extension of scalars, i.e. for ∑_{i≥0} a_iTⁱ ∈ A[T] with a_i ∈ A, μ(D)(∑_{i≥0} a_iTⁱ) = ∑_{i≥0} D(a_i)Tⁱ. Using Remark (1.1) it is easily checked that this gives us an A[T]-linear map

$$\mu: \text{Diff}_k^\infty(A, A[T]) \longrightarrow \text{Diff}_{k[T]}^\infty(A[T]) \subset \text{Diff}_k^\infty(A[T])$$

with μ(Diff_kⁿ(A, A[T])) ⊂ Diff_{k[T]}ⁿ(A[T]) ⊂ Diff_kⁿ(A[T]) for every n. Identify Diff_k[∞](A) as an A-submodule of Diff_k[∞](A, A[T]) via u, and denote by λ the restriction of μ to Diff_k[∞](A).

- (4.1) LEMMA. (1) ∑_{i≥0} (T - 1)ⁱ uq_iμ = identity on Diff_k[∞](A, A[T]).
 (2) For every non-negative integer s, q₀T^sλ = identity on Diff_k[∞](A).
 (3) If D, D' ∈ Diff_kⁿ(A) then λ(DD') = λ(D)λ(D'). Consequently,
 λ(Diff_k^m(A) Diff_kⁿ(A)) ⊂ Diff_{k[T]}^m(A[T]) Diff_{k[T]}ⁿ(A[T])

for all m, n.

- (4) q₀(Diff_k^m(A[T]) Diff_kⁿ(A[T])) ⊂ Diff_k^m(A) Diff_kⁿ(A) for all m, n.

Proof. (1) and (2) follow from the definition. (3) follows from the fact that each of λ(DD') and λ(D)λ(D') is k[T]-linear and coincides with DD' on A. To prove (4), let D ∈ Diff_k^m(A[T]), D' ∈ Diff_kⁿ(A[T]). By (1) we have D'u = ∑_{i≥0} (T - 1)ⁱuq_i(μ(D'u)) = ∑_{i≥0} (T - 1)ⁱuq_i(D'). Therefore

$$DD'u = \sum_{i \geq 0} ([D, (T - 1)^i] + (T - 1)^i D) u q_i(D')$$

and so

$$\begin{aligned} q_0(DD') &= \sum_{i \geq 0} p_0([D, (T - 1)^i] + (T - 1)^i D) u q_i(D') \\ &= \sum_{i \geq 0} q_0([D, (T - 1)^i] + (T - 1)^i D) q_i(D'), \end{aligned}$$

which belongs to Diff_k^m(A) Diff_kⁿ(A). This proves (4).

In the following proposition we have identified Diff_k[∞](A[T]) as an A[T]-submodule of Diff_k[∞](A[T, T⁻¹]) via the natural map Diff_k[∞](A[T]) → (Diff_k[∞](A[T]))_T = Diff_k[∞](A[T, T⁻¹]), which is injective because T is a non-zero divisor in A[T].

- (4.2) PROPOSITION. For D ∈ Diff_kⁿ(A) the following three conditions are equivalent:

- (i) $D \in \text{Diff}_k^1(A) \text{Diff}_k^{n-1}(A)$.
- (ii) $\lambda(D) \in \text{Diff}_k^1(A[T]) \text{Diff}_k^{n-1}(A[T])$.
- (iii) $\lambda(D) \in \text{Diff}_k^1(A[T, T^{-1}]) \text{Diff}_k^{n-1}(A[T, T^{-1}])$.

Proof. The implication (i) \Rightarrow (ii) follows from Lemma (4.1) (3), while the implication (ii) \Rightarrow (iii) is clear. Now, assume (iii). Then $\lambda(D) = \sum_j D_j D'_j$ (finite sum) with $D_j \in \text{Diff}_k^1(A[T, T^{-1}])$, $D'_j \in \text{Diff}_k^{n-1}(A[T, T^{-1}])$. Since A is a finitely generated k -algebra, there exists a positive integer s such that $T^s D_j \in \text{Diff}_k^1(A[T])$ and $T^s D'_j \in \text{Diff}_k^{n-1}(A[T])$. Put $\Delta = \sum_j T^s [D_j, T^s] D'_j \in \text{Diff}_k^{n-1}(A[T, T^{-1}])$. We have $\sum_j (T^s D_j)(T^s D'_j) = \Delta + T^{2s} \lambda(D)$. Therefore $\Delta \in \text{Diff}_k^{n-1}(A[T])$. Now, $q_0(\sum_j (T^s D_j)(T^s D'_j)) \in \text{Diff}_k^1(A) \text{Diff}_k^{n-1}(A)$ by Lemma (4.1) (4), and $q_0(\Delta) \in \text{Diff}_k^{n-1}(A) \subset \text{Diff}_k^1(A) \text{Diff}_k^{n-1}(A)$. Therefore $q_0(T^{2s} \lambda(D)) \in \text{Diff}_k^1(A) \text{Diff}_k^{n-1}(A)$, and now (i) follows by Lemma (4.1) (2).

(4.3) COROLLARY. *If $\text{Diff}_k^n(A[T]) = \text{Diff}_k^1(A[T]) \text{Diff}_k^{n-1}(A[T])$ or $\text{Diff}_k^n(A[T, T^{-1}]) = \text{Diff}_k^1(A[T, T^{-1}]) \text{Diff}_k^{n-1}(A[T, T^{-1}])$ then $\text{Diff}_k^n(A) = \text{Diff}_k^1(A) \text{Diff}_k^{n-1}(A)$.*

§ 5. Differential operators on a cone in 3-space

Let us return now to the notation of Section 2. For $\alpha \in V$ and $g \in R$, put $g_{x^\alpha} = \eta(\partial^\alpha g / \partial X^\alpha)$.

(5.1) LEMMA. *Assume that $J = Rf$ and that x_1, f_{x_r} is an A -regular sequence. Let $D \in \text{Diff}_k^n(A)$. Suppose $c_\alpha(D) \in x_1 A$ for all those $\alpha \in V_n$ for which $\alpha_r = 0$. Then there exists $D' \in \text{Diff}_k^n(A)$ such that $D = x_1 D'$.*

Proof. Since x_1 is a non-zero divisor in A , it is enough to prove that $c_\alpha(D) \in x_1 A$ for every $\alpha \in V_n$. We do this by lexicographic induction on $(n - |\alpha|, \alpha_r)$. Let $\alpha \in V_n$. The result being given for $\alpha_r = 0$, we may assume that $\alpha_r \geq 1$. Put $\beta = \alpha - e_r$. By Corollary (2.11) $\langle D, X^\beta \rangle \in \text{Diff}_k^{n-|\beta|}(A)$. Therefore, since $c_\beta(D) \Delta_0(f) = 0$, we have

$$0 = \langle D, X^\beta \rangle(f) = \sum_{\substack{\gamma \in V_{n-|\beta|} \\ |\gamma| > 0}} c_{\gamma+\beta}(D) \Delta_\gamma(f).$$

Let $\gamma \in V_{n-|\beta|}$ with $|\gamma| > 0$ and $|\gamma + \beta| \leq |\alpha|$. Then $|\gamma| = 1$, $|\gamma + \beta| = |\alpha|$ and either $\gamma = e_r$ or $\gamma_r = 0$. It follows that for every index $\gamma \neq e_r$ appearing in the above summation we have $(n - |\gamma + \beta|, \gamma_r + \beta_r) < (n - |\alpha|, \alpha_r)$ lexicographically. Therefore by induction we get $c_\alpha(D) \Delta_{e_r}(f) \in x_1 A$. Since $\Delta_{e_r}(f) = f_{x_r}$ and x_1, f_{x_r} is A -regular, we get $c_\alpha(D) \in x_1 A$.

(5.2) LEMMA. Let $H \in k[Y, Z]$ be a homogeneous polynomial in two variables Y, Z over k , of degree ≥ 2 , monic in Z and without multiple factors. Then $(\partial H/\partial Z)^2$ does not belong to the ideal $(Y^2\partial H/\partial Y, Y^2\partial H/\partial Z, H)$ of $k[Y, Z]$.

Proof. Replacing k by its algebraic closure, we may assume that $H = \prod_{i=1}^n (Z - t_i Y)$ with $n = \deg(H)$ and t_1, \dots, t_n distinct elements of k . Suppose $(\partial H/\partial Z)^2 = FY^2\partial H/\partial Y + GY^2\partial H/\partial Z + KH$ with $F, G, K \in k[Y, Z]$, F, G homogeneous of degree $n - 3$, and K homogeneous of degree $n - 2$. Putting $Y = 1$ and $Z = t_i$ in this identity for a fixed i and dividing the result by $\prod_{j \neq i} (t_i - t_j)$ we get $\prod_{j \neq i} (t_i - t_j) = g_0 + (g_1 - f_0)t_i + \dots + (g_{n-3} - f_{n-3})t_i^{n-3} - f_{n-3}t_i^{n-2}$ for $1 \leq i \leq n$, where $f_j, g_j \in k$ are defined by $F = \sum_{j=0}^{n-3} f_j Y^{n-3-j} Z^j$, $G = \sum_{j=0}^{n-3} g_j Y^{n-3-j} Z^j$. It follows that if we define $C = (c_{ij})_{1 \leq i, j \leq n}$ with $c_{i1} = \prod_{j \neq i} (t_i - t_j)$ and $c_{ij} = t_i^{j-2}$ for $j \geq 2$ then $\det(C) = 0$. On the other hand, it is easily checked that, since t_1, \dots, t_n are distinct, $\det(C) \neq 0$. This contradiction proves the lemma.

(5.3) THEOREM. Let J be a homogeneous, proper, principal ideal of $R = k[X_1, X_2, X_3]$ and let $A = R/J$. Assume that A is reduced.⁽¹⁾ Then A is regular if and only if $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \text{Diff}_k^1(A)$.

Proof. If A is regular then the equality $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \text{Diff}_k^1(A)$ follows from [4, (IV, 16.11.2)]. Conversely, suppose $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \cdot \text{Diff}_k^1(A)$. Let T be a homogeneous element of A of degree one. Then $\text{Diff}_k^2(A_T) = \text{Diff}_k^1(A_T) \text{Diff}_k^1(A_T)$. Since $A_T = A_{(T)}[T, T^{-1}]$ with T algebraically independent over $A_{(T)}$, it follows from Corollary (4.3) that

$$(5.3.1) \quad \text{Diff}_k^2(A_{(T)}) = \text{Diff}_k^1(A_{(T)}) \text{Diff}_k^1(A_{(T)}).$$

Now, since $\deg(T) = 1$, $A_{(T)}$ is a reduced ring of the form $k[Y_1, Y_2]/(f(Y_1, Y_2))$. Therefore by (5.3.1) and Theorem (3.3) $A_{(T)}$ is regular. This proves that the curve $\text{Proj}(A)$ is non-singular. Therefore the only possible singularity of the cone $\text{Spec}(A)$ is its vertex. Also, A is Cohen-Macaulay. Therefore A is normal.

Suppose A is not regular. Let $J = RF$ with $F = F(X_1, X_2, X_3)$ homogeneous, and let $n = \deg(F)$. Then $n \geq 2$. Since $\text{char}(k) = 0$, we can arrange the following by a homogeneous change of variables: First, $F = X_3^n + \sum_{j=1}^n F_j X_3^{n-j}$ with $F_j \in k[X_1, X_2]$ homogeneous of degree j for every

⁽¹⁾ See note added in proof.

j , and $F_1 = 0$. Then, since A is reduced, $\text{Disc}_{X_3}(F)$, the discriminant of F with respect to X_3 , is non-zero. We choose X_1 such that $\text{Disc}_{X_3}(F) \notin X_1 k[X_1, X_2]$. Then $\text{Disc}_{X_3}(F(0, X_2, X_3)) \neq 0$ and it follows that $F(0, X_2, X_3)$ and $\partial F(0, X_2, X_3)/\partial X_3$ have no common factors, and $F(0, X_2, X_3)$ has no multiple factors.

Having arranged the above, we *claim* the following:

- (1) $\partial F/\partial X_2 \in (X_1, X_2)$.
- (2) $(\partial F/\partial X_3)^2 \in (X_1, X_2^2 \partial F/\partial X_2, X_2^2 \partial F/\partial X_3, F)$.
- (3) $F, X_1, \partial F/\partial X_3$ is an R -regular sequence.
- (4) $(\partial F/\partial X_3)^2 \in (F) + (X_1, X_2)^2$.

Note that (1) and (4) follow from the fact that $F_1 = 0$. (2) follows from Lemma (5.2) applied to $H = F(0, X_2, X_3)$. To prove (3), first note that $X_1, F, \partial F/\partial X_3$ is an R -regular sequence by our choice of variables. Then, since these three elements are homogeneous, any permutation of these is again R -regular. This proves our claim.

Now, given $D \in \text{Diff}_k^\infty(R, A)$, put $\epsilon(D) = \sum_{\alpha \in V, \alpha_3=0} c_\alpha(D) \Delta_\alpha$. In particular, if $D \in \text{Diff}_k^\infty(A)$ then we have $\epsilon(D) \in \text{Diff}_k^\infty(R, A)$ (see (2.4)).

Since A is normal, it follows from [5, p. 893] that $\text{Der}_k^1(A)$ is generated as an A -module by d, d_1, d_2, d_3 where $d = x_1 \Delta_{e_1} + x_2 \Delta_{e_2} + x_3 \Delta_{e_3}$, $d_1 = F_{x_3} \Delta_{e_2} - F_{x_2} \Delta_{e_3}$, $d_2 = F_{x_3} \Delta_{e_1} - F_{x_1} \Delta_{e_3}$ and $d_3 = F_{x_2} \Delta_{e_1} - F_{x_1} \Delta_{e_2}$. Consider the elements $d^2, d_1 d, d_1^2$ of $\text{Diff}_k^2(A)$. Using Proposition (2.5) it is easily checked that we have

$$\begin{aligned} \epsilon(d^2) &= x_1 \Delta_{e_1} + x_2 \Delta_{e_2} + 2x_1^2 \Delta_{2e_1} + 2x_1 x_2 \Delta_{e_1+e_2} + 2x_2^2 \Delta_{2e_2}, \\ \epsilon(d_1 d) &= F_{x_3} \Delta_{e_2} + x_1 F_{x_3} \Delta_{e_1+e_2} + 2x_2 F_{x_3} \Delta_{2e_2}, \\ \epsilon(d_1^2) &= a \Delta_{e_2} + 2(F_{x_3})^2 \Delta_{2e_2}, \end{aligned}$$

where $a = F_{x_2 x_3} F_{x_3} - F_{x_3^2} F_{x_2}$. By (4) we can write $(F_{x_3})^2 = b_1 x_1 + b_2 x_2^2$ with $b_1, b_2 \in A$ homogeneous and $\text{deg}(b_2) = 2n - 4$. By (1) we can write $a = u F_{x_3} + v_1 x_1 + v_2 x_2$ with $u, v_1, v_2 \in A$. Put $D = (b_2 - v_2) d - u d_1 - b_2 d^2 + d_1^2 \in \text{Diff}_k^2(A)$. Then an easy computation shows that

$$\epsilon(D) = -v_2 x_1 \Delta_{e_1} + v_1 x_1 \Delta_{e_2} - 2x_1^2 b_2 \Delta_{2e_1} - 2x_1 x_2 b_2 \Delta_{e_1+e_2} + 2x_1 b_1 \Delta_{2e_2}.$$

Thus $c_\alpha(D) \in x_1 A$ for all those $\alpha \in V_2$ for which $\alpha_3 = 0$. Therefore it follows from (3) and Lemma (5.1) that there exists $D' \in \text{Diff}_k^2(A)$ such that $D = x_1 D'$. We have $c_{2e_1}(D') = x_1^{-1} c_{2e_1}(D) = -2x_1 b_2$. Let

$$\alpha = \{y \in A \mid \exists \delta \in \text{Diff}_k^2(A) \text{ with } c_{2e_1}(\delta) = y\}.$$

Then α is an ideal of A and $x_1 b_2 \in \alpha$. Now, we have $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \cdot \text{Diff}_k^1(A)$ by assumption. So $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) + \text{Der}_k^1(A) \text{Der}_k^1(A)$ by Lemma (1.3). Therefore, since $\text{Der}_k^1(A)$ is generated by d, d_1, d_2, d_3 , it follows from Proposition (2.5) that α is generated by $x_1^2, x_1 F_{x_2}, x_1 F_{x_3}, (F_{x_2})^2, F_{x_2} F_{x_3}, (F_{x_3})^2$. All these elements are homogeneous and $\deg((F_{x_2})^2) = \deg(F_{x_2} F_{x_3}) = \deg((F_{x_3})^2) = 2n - 2 > 2n - 3 = \deg(x_1 b_2)$. It follows that $x_1 b_2 \in (x_1^2, x_1 F_{x_2}, x_1 F_{x_3})$. Therefore, since x_1 is a non-zero divisor, $b_2 \in (x_1, F_{x_2}, F_{x_3})$. Since $(F_{x_3})^2 = b_1 x_1 + b_2 x_2^2$, we get $(F_{x_3})^2 \in (x_1, x_2^2 F_{x_2}, x_2^2 F_{x_3})$. This contradicts (2), and the theorem is proved.

§ 6. A remark on the non-reduced case

In proving our results in Sections 3 and 5 we have assumed that A is reduced. We believe that this condition can be dropped.⁽¹⁾ Let us show this in the case $\dim(A) = 0$. With the notation of Section 2 let $r = 1$, and omit the subscript 1. Assume that $J = Rf$ is non-zero, proper and principal. Then $A = k[x] = k[X]/(f)$ and $\Delta = \eta\partial/\partial X$. Put $f_x = \eta(\partial f/\partial X)$.

(6.1) PROPOSITION. *If $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \text{Diff}_k^1(A)$ then A is reduced, hence regular.*

Proof. Let

$$0 \rightarrow \text{Diff}_k^1(A) \hookrightarrow \text{Diff}_k^2(A) \xrightarrow{\theta_2} \text{Der}_k^1(A) \longrightarrow 0$$

be the exact sequence of Theorem (2.13). Recall that in this case $\theta_2(D) = c_2(D)\Delta$. Identify $\text{Der}_k^1(A)$ with the ideal $\alpha = \{d(x)|d \in \text{Der}_k^1(A)\}$ of A by sending $d \in \text{Der}_k^1(A)$ to $d(x)$. Then it follows from Proposition (2.5) that $\theta_2(\text{Der}_k^1(A) \text{Der}_k^1(A)) = \alpha^2$. By our assumption and Lemma (1.3) we have $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) + \text{Der}_k^1(A) \text{Der}_k^1(A)$. Therefore $\alpha = \alpha^2$. Now, $\alpha = \{a \in A | af_x = 0\}$. It follows that if $f = f_1^{t_1} \cdots f_m^{t_m}$ is the prime factorization of f with f_1, \dots, f_m mutually coprime then $\alpha = (\eta(f_1 \cdots f_m))$. Thus the equality $\alpha = \alpha^2$ implies that $f_1 \cdots f_m$ belongs to the ideal $(f, f_1^2 \cdots f_m^2)$ of R . This is possible only if $t_1 = \cdots = t_m = 1$, i.e., if A is reduced.

NOTE ADDED IN PROOF. It was recently proved by the author jointly with D. P. Patil that if X is a hypersurface and $\text{Diff}_k^2(A) = \text{Diff}_k^1(A) \text{Diff}_k^1(A)$ then A is reduced. Thus, in Theorems (3.3) and (5.3) we can drop the assumption that A be reduced.

⁽¹⁾ See note added in proof.

REFERENCES

- [1] J. Becker, Higher derivations and integral closure, *Amer. J. Math.*, **100** (1978), 495–521.
- [2] W. C. Brown, Higher derivations on finitely generated integral domains, *Proc. Amer. Math. Soc.*, **42** (1974), 23–27.
- [3] H. Cartan and S. Eilenberg, *Homological Algebra*, Princeton Univ. Press, Princeton 1956.
- [4] A. Grothendieck, *Éléments de Géométrie Algébrique*, *Publ. Math. IHES*, **32** (1967).
- [5] J. Lipman, Free derivation modules on algebraic varieties, *Amer. J. Math.*, **87** (1965), 874–898.
- [6] K. R. Mount and O. E. Villamayor, On a conjecture of Y. Nakai, *Osaka J. Math.*, **10** (1973), 325–327.
- [7] Y. Nakai, High order derivations I, *Osaka J. Math.*, **7** (1970), 1–27.
- [8] C. J. Rego, Remarks on differential operators on algebraic varieties, *Osaka J. Math.*, **14** (1977), 481–486.
- [9] J.-P. Serre, *Algèbre locale—Multiplicités*, Springer-Verlag Lecture Notes in Math., No. 11, Berlin-Heidelberg-New York, 1975.
- [10] J.-P. Vigué, Opérateurs différentiels sur les espaces analytiques, *Invent. math.*, **20** (1973), 313–336.
- [11] O. Zariski, Studies in equisingularity I, *Amer. J. Math.*, **87** (1965), 507–536.
- [12] J.-P. Vigué, Opérateurs différentiels sur les cônes normaux de dimension 2, *C. R. Acad. Sc. Paris*, **278**, Série A, (1974), 1047–1050.
- [13] Y. Ishibashi, Remarks on a Conjecture of Nakai, *J. Algebra*, **95** (1985), 31–45.
- [14] ———, Nakai's conjecture for invariant subrings, *Hiroshima Math. J.*, **15** (1985), 429–436.

Tata Institute of Fundamental Research
Homi Bhabha Road,
Bombay 400 005,
India