



# Lethal and sublethal concentrations of spiroadiclofen stress may increase the adaptation of *Panonychus citri* (Acari: Tetranychidae)

## Research Paper

**Cite this article:** Wang H, Wang H, Wen K, Xie T, Luo S, Wu J, Xia B (2024). Lethal and sublethal concentrations spiroadiclofen stress may increase the adaptation of *Panonychus citri* (Acari: Tetranychidae). *Bulletin of Entomological Research* 1–7. <https://doi.org/10.1017/S0007485324000087>

Received: 10 July 2023

Revised: 6 October 2023

Accepted: 27 January 2024

### Keywords:

biological traits; enzymatic parameters; lethal and sublethal effects; tolerance; Vitellogenin

### Corresponding author:

Bin Xia;

Email: [xiabin9@163.com](mailto:xiabin9@163.com)

Hongyan Wang, Haifeng Wang, Kexin Wen, Tao Xie, Shigan Luo, Jiawei Wu and Bin Xia

College of Life Science, Nanchang University, Nanchang 330031, China

### Abstract

*Panonychus citri* is one of the most destructive pests in citrus orchards, exhibiting varying degrees of tolerance to numerous insecticides, such as spiroadiclofen. To effectively manage pests, this study explores the response of *P. citri* to spiroadiclofen stress from the perspectives of life history, enzymatic parameters, and reproduction. The effects of two concentrations ( $LC_{30}$  and  $LC_{50}$ ) of spiroadiclofen on the biological parameters of *P. citri* were evaluated by the life table method. The results showed that the development duration, fecundity, oviposition days, and lifespan were shortened, though the pre-oviposition period of two treatments was prolonged in comparison with the control. A significant decrease was recorded in the net reproductive rate ( $R_0$ ) and the mean generation time ( $T$ ) for the two treatments. Nevertheless, the intrinsic rate of increase ( $r$ ) and the rate of increase ( $\lambda$ ) were not significantly affected in the  $LC_{30}$  treatment, whereas they declined in the  $LC_{50}$  treatment. The enzyme activity assay resulted in higher activities of catalase (CAT), superoxide dismutase (SOD), peroxidase (POD), and carboxylesterase (CarE), among the treatments than the control. In contrast, the treatments recorded lower cytochrome P450 (CYP450) and Glutathione S-transferase (GST) activities than the control. Furthermore, the study detected that relative mRNA expression of Vitellogenin (Vg) and Vitellogenin receptor (VgR) for two treatments were lower than the control. In summary, two concentrations of spiroadiclofen inhibited progeny growth and fecundity of *P. citri*. Additionally, the results of this study may support further research on tolerance of *P. citri* in response to spiroadiclofen stress.

## Introduction

*Panonychus citri* (Acarina: Tetranychidae) is distributed worldwide and mainly feeds on the leaves of citrus (Pan *et al.*, 2020). Presently, the key method for the control of *P. citri* is chemical control (Cheng *et al.*, 2022). The enhanced application of pesticides prevents further damage, but it has several undesirable consequences, such as impacting natural enemies and non-target organisms. Meanwhile, the increased use of insecticides has led to widespread resistance of *P. citri* to insecticides (Quesada and Sadof, 2019). This emphasises the need to adopt effective acaricides for the control of *P. citri* (Andrade *et al.*, 2018).

Spiroadiclofen is an acaricide that belongs to the spirocyclic tetraelectron acid family and works by interfering with lipid biosynthesis (Amaral *et al.*, 2020). Lipids constitute approximately 30–40% of the dry weight in insects (Ziegler and Antwerpen, 2006), with a higher proportion in mites. Consequently, this acaricide is widely used worldwide for controlling *P. citri* and mites in other crops, making it one of the most commonly used acaricides in China (Claudiane *et al.*, 2021). However, its extensive use has also increased the frequency of misuse and overuse of insecticides. This improper usage has led to insects being exposed to lethal and sub-lethal concentrations of the insecticides, thereby becoming a key factor in accelerating pest resistance development. Previous research has shown that field populations of *P. citri* have relatively low resistance levels to spiroadiclofen, but there is a possibility of developing higher levels of resistance in the future (Alavijeh *et al.*, 2020). Field populations of phytophagous mites have also been confirmed to possess tolerance to spiroadiclofen in regions such as the United States, Germany, Brazil, China, Turkey, and Iran (Cheng *et al.*, 2022). Therefore, exposure of *P. citri* to lethal and sub-lethal concentrations of spiroadiclofen in the field cannot be avoided. Assessing the potential lethal and sub-lethal effects of spiroadiclofen on *P. citri* is crucial for a comprehensive analysis of the effectiveness of this insecticide.

Sublethal effects are defined as impacts (either physiological or behavioural) on survival individuals when exposed to a toxicant at low or sublethal concentration/dose (Desneux *et al.*, 2007). The sublethal effects of pesticides on pests are not only reflected in the direct

killing, but also affect their reproduction, lifespan, and physiological characteristics (Wang *et al.*, 2016; Dong *et al.*, 2017), as well as changes in their detoxification and antioxidant enzymes (Ma *et al.*, 2019). Hormesis, the stimulation of performance when an organism is exposed to low levels of agents, is regarded as the main mechanism for pest population resurgence (Cordeiro *et al.*, 2013). For example, some researchers have suggested that the sublethal concentrations of insecticides might incite insect outbreaks (Bartle, 1968). Currently, the lethal and sublethal effects of spiroadicofen on *P. citri* have not been determined.

Here, we used an age-stage, life table to evaluate the lethal and sublethal effects of the spiroadicofen on the life table parameters of *P. citri*, including development time, survival rate, longevity, fecundity, and hatchability. The results will help to provide a comprehensive assessment of this new insecticide for integrated pest management (IPM) and chemical applications in the field. Furthermore, understanding the activities of several key antioxidant enzymes and detoxification enzymes in *P. citri* can provide the basis for mitigating or reducing resistance to potent acaricides (Rasheed *et al.*, 2020).

## Materials and methods

### Mite and pesticide

In 2019, a stable population of *P. citri* was established by collecting a colony from Nanchang University of Nanchang, Jiangxi Province, China and maintaining them through continuous breeding. The population of *P. citri* were fed citrus tender leaves and were kept in an artificial climate box with the following environmental conditions: a temperature of  $26 \pm 1^\circ\text{C}$ , a relative humidity (RH) of  $70 \pm 5\%$ , and a photoperiod of 16:8 hours. The population was not exposed to any pesticides until the present study (in 2021), which was considered the acaricide-susceptible strain (SS).


The acaricide used in the bioassay was spiroadicofen (24%SC), which was purchased from Bayer Crop Science (China) Co. Ltd.

### Bioassay

The modified leaf dipping method was used for the bioassay on the *P. citri* population (Yamamoto *et al.*, 1995). The adult females were transferred to citrus leaves ( $n = 45$ , three repetitions per treatment). Based on the pre-experiment, the acaricide was diluted to six concentrations (fig. 1). The control was performed on leaves with adult females soaked in triton X-100 solution. Immerse the leaves with *P. citri* in the prepared insecticide for 5 s (Wang *et al.*, 2021). The treated citrus leaves were placed on the prepared Petri dishes ( $d = 15$  cm) in an air-conditioned room. After 24 h, the mortality of the *P. citri* population was recorded, and if their legs did not move, they were considered dead. The mortality data were used for  $\text{LC}_{30}$  and  $\text{LC}_{50}$  determination. All treated-individuals are raised through fresh citrus leaves (not exposed to any insecticides).

### Changes in life table parameters

The adult females ( $n = 100$ , three repetitions per treatment) were transferred to the leaf dish and soaked with two spiroadicofen concentrations ( $\text{LC}_{30}$  and  $\text{LC}_{50}$ ) by the leaf dipping method refer to the method described 2.2 (fig. 1A), the control was performed on leaves with adult females soaked in triton X-100 solution. Following 24 h, survivors (newly emerged 3-day-old adult females) were selected to continue feeding on the fresh-rearing platform. Meanwhile, adult males (without insecticide treated) were selected for mating. Eggs produced were cultured independently 12 h after mating ( $n = 30$ ). The adult females (with insecticide treatment) were mated with adult males as soon as they



**Figure 1.** (A) Bioassay method schematic diagram (B) An observation of the life history of adult females of *Panonychus citri*.

reached maturity, and they were observed every 24 h until died (fig. 1B).

**Enzyme activity assay**

3-day-old adult females were exposed to two concentrations of spirodiclofen (LC<sub>30</sub> and LC<sub>50</sub>) and control for 24 h, the prepared samples (n = 300, three biological repetitions per treatment) were frozen in liquid nitrogen and stored at -80°C in one 1.5 mL centrifuge tube to create the crude enzyme solution.

After calculation of protein concentration (BCA Protein Assay Kit- A045, Nanjing Jiancheng Bioengineering Institute, China), the supernatant of required volume was used to measure the enzyme activities of SOD, CAT, POD, CarE, GST, and CYP450 by commercial assay kits A001, A007, A084, A133, A004, and H303(Nanjing Jiancheng Bioengineering Institute, China). According to the instruction of CAT, SOD, POD test kit (Nanjing Jiancheng Bioengineering Institute, Nanjing, China), anti-oxidant enzymes activities: CAT, SOD, POD and detoxifying enzyme activities: CarE, GST, CYP450 were measured (Qi *et al.*, 2020). Using a photometer set at 450 nm, 550 nm, 420 nm, 450 nm, 412 nm, 450 nm, the amount of CAT, SOD, POD, CarE, GST, CYP450 in *P. citri* was determined. The change of absorbance value is the measured value, repeat for three times, and take the average value. Calculate the activity of six enzyme activities.

**Gene expression analysis**

The StepOnePlus real-time quantitative PCR machine was used to determine the mRNA level of some genes (such as Vg and VgR genes) after two concentrations of spirodiclofen exposure.

The 200 adult females were used to extract total RNA by Trizol reagent.(Shenggong Biological Technology Co. Ltd., Shanghai, China) following the manufacturer’s specifications, and quantification was then performed using a NanoDrop 2000 spectrophotometer. In the following step, reverse transcription of RNA is performed using PrimeScript RT Reagent Kit with gDNA Eraser in a 20 µl reaction volume (TaKaRa, Shiga, Japan). Every RT-qPCR was carried out in a 20-µL mixture. The following were the qPCR cycling parameters: 95°C for 10 min, then 40 cycles of 95°C for 30 s and 60°C for 30 s. Primers were designed as shown in table 1, with the GAPDH and EF-1FA genes as internal reference genes. Relative quantification was calculated using the comparative 2<sup>-ΔΔCt</sup> method (Zhao *et al.*, 2018b).

**Statistical analysis**

The life history parameters were counted, including age-stage specific rate (s<sub>xj</sub>), Age-specific survival (l<sub>x</sub>), age-specific reproduction

(m<sub>x</sub>), APOP (adult pre-ovipositional period), TPOP total pre-ovipositional period (from newborn egg to first oviposition). Population parameters including R<sub>0</sub>, rm, λ, and T were also calculated as follows:

$$R_0 = \sum l_x m_x; \tag{1}$$

$$rm: \sum e - rm(x + 1) l_x m_x = 1 \tag{2}$$

$$T:T = \ln R_0 / rm; \tag{3}$$

$$\lambda:\lambda = erm. \tag{4}$$

The standard errors of raw data were calculated by using the bootstrap method with 100,000× resamplings, the paired bootstrap test was used to compare differences (Akkopru *et al.*, 2015). The TWSEXMSChart program was used to analyse the raw data and population parameters (Chi and Liu, 1985).

**Results**

**Toxicity of spirodiclofen in the adult females of Panonychus citri**

Acute toxicity of spirodiclofen (24%SC) was observed in adult females *P. citri* (table 2). Concentrations resulting in 30 and 50% mortality were 3.898 g l<sup>-1</sup> and 5.215 g l<sup>-1</sup>.

**Responses of spirodiclofen on population parameters of Panonychus citri**

Through bioassay of *P. citri*, the low lethal concentration (LC<sub>30</sub>) and median concentration (LC<sub>50</sub>) were obtained. tables 3 and 4 present the lethal and sublethal effects on the F<sub>1</sub> generation of *P. citri*. The average immaturity time (The development stage of adult females before spawning) of *P. citri* treated with LC<sub>30</sub> was not significantly different from the control (table 3). However, LC<sub>50</sub> treatment was prolonged in egg duration. Compared to the control, LC<sub>30</sub> and LC<sub>50</sub> treatments significantly prolonged both oviposition period and total pre-ovipositional period. Compared with control in maturity (13.930 d), longevity (22.310 d), and fecundity (5.350 eggs/female), there were significant reductions in maturity (10.510 d and 5.620 d), longevity (21.430 d and 17.650 d), and fecundity (4.310 eggs/female and 2.680 eggs/female) after exposure two concentrations (LC<sub>30</sub> and LC<sub>50</sub>) of spirodiclofen.

**Table 1.** Primers for real-time quantitative PCR.

| Genes                          | GenBank No.  | Primer sequences(5' to 3')                                     |
|--------------------------------|--------------|--|
| <i>PcVg</i>                    | KC978893     | Vg-F: GCCTCAAACGAAGCTCAATC<br>Vg-R: AGCCAAAGCGTCGAGTAAAA       |
| <i>PcVgR</i>                   | KC978894     | VgR-F: TTGTTTCGATGGCTCAGATG<br>VgR-R: TCACCGTGTGGACAATCAGT     |
| <i>GAPDH</i> (Reference gene1) | HM582445     | GAP-F: CTTFGGCCAAGGTCATCAAT<br>GAP-R: CGGTAGCGGCAGGTATAATG     |
| <i>ELFIA</i> (Reference gene2) | LOC100037748 | ELFIA-F: GGCACITCGTCITCCACITC<br>ELFIA-R: ATGATTCGTGGTGCATCTCA |

**Table 2.** Acute toxicity of spiroticlofen to adult females of *Panonychus citri*.

| Acaricide     | Concentration g/L (95% CL) |                     | LC-P equation           | $\chi^2$ | R    |
|---------------|----------------------------|---------------------|-------------------------|----------|------|
|               | LC <sub>30</sub>           | LC <sub>50</sub>    |                         |          |      |
| Spiroticlofen | 3.898 (3.461–4.273)        | 5.215 (4.823–5.566) | $y = -0.5219 + 1.3878x$ | 17.43    | 0.94 |

The  $R_0$  and  $T$  of adult females were significantly reduced on LC<sub>30</sub> and LC<sub>50</sub> treatments compared to the control. Adult females *P. citri* treated with LC<sub>30</sub> showed no significant difference in  $rm$  and  $\lambda$  from the control, whereas they decreased in the LC<sub>50</sub> treatment (table 4). It is clear that the survival rates at different ages are overlapping, as eggs are more likely to survive up to age  $x$  and develop to stage  $j$  when they occur at different age stages (fig. 2). Compared with the control, adult females survival rates in the LC<sub>30</sub> and LC<sub>50</sub> groups are relatively low. *Panonychus citri* survival rate ( $l_x$ ), fecundity ( $m_x$ ), and maternity rates ( $l_x m_x$ ) after exposure to two concentrations of spiroticlofen are shown in fig. 3. The  $l_x$  and  $m_x$  of the treatment groups are lower than the control. Furthermore, both  $l_x$  and  $m_x$  decrease with the increase of concentration.

### Responses of spiroticlofen on enzymatic activity of *Panonychus citri*

CAT, SOD, and POD activities were measured 24 h after spiroticlofen treatment. Difference in CAT between control and LC<sub>30</sub> group was not significant (fig. 4). However, LC<sub>30</sub> treatment significantly increased SOD activity compared with control, but both CAT and SOD activities of the LC<sub>50</sub> treatment group showed a significant decline. The POD activity of the LC<sub>30</sub> group has no significant difference which was compared with the control, whereas POD activity was significantly higher in the LC<sub>50</sub> group. The activities of CarE, GST, and CYP450 of *P. citri* treated with spiroticlofen were determined. The activity of CarE in LC<sub>30</sub> treatment increased compared with control, while it decreased significantly with increasing concentrations (fig. 5). Comparatively to the control group, all treatment groups showed a decrease in GST and CYP450 activities.

### Vg and VgR relative expression

To determine how spiroticlofen affects the contents of Vg and VgR in *P. citri*, mRNA-relative expression of Vg and VgR were measured. Compared with CK, Vg and VgR expression were significantly decreased and showed a more significant decrease with increasing concentrations (fig. 6). In addition, significant concentration effects were also observed between the LC<sub>30</sub> and LC<sub>50</sub> treatments.

### Discussion

The study of life table is a key aspect of insect population dynamics (Desneux *et al.*, 2007). The field of insect population dynamics is characterised by changes that are affected by many factors, including diet, temperature, light, and especially chemical pesticides (Mousavi *et al.*, 2020). Fecundity,  $rm$ ,  $\lambda$ , and  $R_0$  are several important parameters for assessing population dynamics (Papachristos and Milonas, 2008; Rahmani and Bandani, 2013).

The toxicity test indicated that fecundity and population parameters including  $rm$ ,  $\lambda$  and  $R_0$  of *P. citri* were decreased in the treated groups, and similar findings were observed in other insecticide-treated pests (Zhao *et al.*, 2018c). Chlorfenapyr, for example, inhibits *Tetranychus urticae* Koch growth and reproduction at low lethal concentrations (LC<sub>20</sub> and LC<sub>30</sub>) (Sani *et al.*, 2018). At low lethal concentrations (LC<sub>30</sub>), cyantraniliprole significantly inhibits the fertility of *Helicoverpa assulta* (Dong *et al.*, 2017). In contrast, there have been several studies showing that low levels of pesticides can stimulate fertility. For instance, treatment with spinetoram LC<sub>10</sub> and LC<sub>20</sub> shortens the time for *T. urticae* to develop from egg to adult and increases their fecundity (Wang *et al.*, 2016). Chlorfenapyr stimulated *Bradysia odoriphaga* reproduction (Sani *et al.*, 2018). However, in the present study, the growth of *P. citri* can be effectively

**Table 3.** Spiroticlofen effects on *Panonychus citri* developmental time.


| Parameter                | Treatment      |                  |                  |
|--------------------------|----------------|------------------|------------------|
|                          | Control        | LC <sub>30</sub> | LC <sub>50</sub> |
| Egg duration(d)          | 5.30 ± 0.20 b  | 5.96 ± 0.19 ab   | 6.58 ± 0.06 a    |
| Larvae duration(d)       | 1.48 ± 0.15 a  | 1.41 ± 0.12 a    | 1.74 ± 0.04 a    |
| Nymph duration(d)        | 2.49 ± 0.15 a  | 2.44 ± 0.11 a    | 2.78 ± 0.06 a    |
| Immaturity period (d)    | 13.93 ± 0.23 a | 10.51 ± 0.24 b   | 5.62 ± 0.06 c    |
| Longevity(d)             | 22.31 ± 0.13 a | 21.43 ± 0.41 b   | 17.65 ± 0.09 c   |
| APOP(d)                  | 1.61 ± 0.12 b  | 1.81 ± 0.08 ab   | 2.22 ± 0.04 a    |
| TPOP(d)                  | 10.18 ± 0.16 c | 11.92 ± 0.22 b   | 13.25 ± 0.09 a   |
| Oviposition period(d)    | 12.50 ± 0.37 a | 9.47 ± 0.29 b    | 6.62 ± 0.06 c    |
| Fecundity(eggs/female/d) | 5.35 ± 0.25 a  | 4.31 ± 0.16 b    | 2.68 ± 0.03 c    |

The bootstrap paired test does not detect significant differences between means in a row following the same letter ( $P > 0.05$ ). APOP, adult pre-ovipositional period; TPOP, total pre-ovipositional period (from newborn egg to first oviposition).


**Table 4.** Effects of spiroadiclofen on population parameters of *Panonychus citri* offspring.

| Population parameters                              | Treatment      |                  |                  |
|--|----------------|------------------|------------------|
|  | Control        | LC <sub>30</sub> | LC <sub>50</sub> |
| Net reproductive rate( $R_0$ ) ( $d^{-1}$ )        | 54.80 ± 3.12 a | 35.28 ± 1.54 b   | 11.21 ± 0.04 c   |
| Mean generation time(T) (d)                        | 1623 ± 0.28 a  | 14.23 ± 0.18 b   | 13.74 ± 0.05 b   |
| Intrinsic rate of increase( $r_m$ ) ( $d^{-1}$ )   | 0.35 ± 0.03 a  | 0.27 ± 0.02 a    | 0.18 ± 0.03 b    |
| Finite rate of increase ( $\lambda$ ) ( $d^{-1}$ ) | 1.42 ± 0.05 a  | 1.19 ± 0.03 a    | 1.07 ± 0.01 b    |

By the bootstrap paired test ( $P > 0.05$ ), means in a row followed by the same letter are not significantly different.



**Figure 2.** The age-stage specific rate ( $s_{ij}$ ) of *Panonychus citri* at two concentrations of spiroadiclofen.




**Figure 3.** Age-specific survival ( $l_x$ ), age-specific reproduction ( $m_x$ ) and age-specific motherhood ( $l_x m_x$ ) of *Panonychus citri* after exposure to two concentrations of spiroadiclofen.

inhibited by two concentrations of spiroadiclofen. As a consequence, different insects react differently to insecticide stress.


The fertility of the *P. citri* F1 generation decreased significantly. Adverse effects of drugs such as shorter lifespans and reduced fecundity are often associated with their development (Zhang *et al.*, 2018). Adaptive costs of insecticide resistance, or the dominant disadvantage that accompanies the development of insecticide resistance, are associated with susceptible insects in a population. To gain more insight into this mechanism, our study measured the relative expression levels of two genes associated with growth and reproduction. Vitellogenin, a protein associated with reproduction, has traditionally been considered a suitable parameter for assessing the fertility of female insects (Zhao *et al.*, 2018a). For instance, the down-regulation of Vg levels adversely affected fertility in *Chilo suppressalis* and *Apolysus lucorum* (Huang *et al.*, 2016). The vitellogenin receptor is the main receptor of Vg function; notably, the down-regulation of VgR can inhibit Vg function. There was a significant reduction

in mRNA expression of Vg and VgR in the experimental group compared to the control group. Combined with the significantly reduced fecundity of the treatments, these data suggest that the reduced expression of Vg mRNA in spiroadiclofen-treatment may bear a significant effect on *P. citri* fecundity.


Various parameters related to insect population and development are affected by insecticide stress (Zhao *et al.*, 2018c; Zhang *et al.*, 2019, 2020; Ullah *et al.*, 2020). When insects are repeatedly exposed to the same pesticide, pesticide resistance will develop. Furthermore, the activities of the protective enzymes are influenced (Van *et al.*, 2006). Antioxidant enzymes are essential components of the insect immune system, preventing oxidative damage caused by foreign organisms. Toxin-induced reactive oxygen species (ROS) are removed by POD, SOD, and CAT antioxidant enzymes. As a result, the SOD-CAT-POD system acts as a first line of defence against ROS.  $H_2O_2$  is produced when the body responds to chemical stress by generating ROS, upregulating the SOD activity, and activating defence systems of



**Figure 4.** Response of protective enzyme activities of *Panonychus citri* to spiroticlofen stress (data are Means  $\pm$  SE of three biological replications; different letters above each bar indicate statistically significant difference by ANOVA followed by the Duncan's multiple range test) \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .



**Figure 5.** Response of detoxification enzyme activities of *Panonychus citri* to spiroticlofen stress. (data are Means  $\pm$  SE of three biological replications; different letters above each bar indicate statistically significant difference by ANOVA followed by Duncan's multiple range test). \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .



**Figure 6.** Relative expression level of Vg and VgR genes in adult females of *Panonychus citri* exposed to two concentrations of spiroticlofen. The bars represent the Means  $\pm$  SE of three replications.

the body. A relative balance of the body needs to be regulated by the decomposition of CAT and POD. An increase in pesticide concentration weakened the self-defence mechanism and inhibited the function of the protective enzymes (Bolter and Chefurka, 1990). At low concentration, avermectin stimulates SOD activity. Conversely, at high concentrations SOD activity was inhibited. The same trend was observed for CAT activity. The study results showed that SOD and CAT have an inhibitory effect on LC<sub>50</sub> treatment compared to lower concentrations (Ma *et al.*, 2014; Liu *et al.*, 2021).

Insecticide resistance is frequently caused by physiological changes that increase detoxifying enzymes like CarE, GST, and CYP450 (Zhao *et al.*, 2018c; Zhang *et al.*, 2020). In present study, when exposed to low lethal concentrations (LC<sub>30</sub>) of

spiroticlofen, CarE activity increased significantly, but decreased when near death. As concentrations increased, CYP450 and GST activities decreased (fig. 5). Toxins introduced into the body are dealt with by detoxifying enzymes in arthropods. CarE is phase II detoxification enzyme important in chemical metabolism and detoxification (Papachristos and Milonas, 2008). The activities of CarE was upregulated after 48 h of treatment with low concentrations of buprofezin, and the activity decreased as insecticide concentration increased (Zhao *et al.*, 2018c). However, several hydrophobic toxins fail to be modified by CarE, causing lipid peroxidation in cell membranes. Therefore, the CYP450 and GST are important in the emergence of resistance genes (Ullah *et al.*, 2020). The production of ROS by insecticides induces oxidative stress in a wide variety of animal cells (Döker *et al.*, 2021). ROS attack causes oxidative damage to proteins and lipids, destroys structural integrity, and reduces enzyme activity (Goel *et al.*, 2005). For instance, in locusts exposed to different sublethal doses of chlorpyrifos, CYP450 enzyme activity decreased with increasing concentrations of the insecticide (Van *et al.*, 2006). Our study showed a significant decrease in the CYP450 enzyme activity with increasing spiroticlofen concentrations. The study assumed that GST and CYP450 activities in *P. citri* were a cause of tolerance against spiroticlofen.

In summary, spiroticlofen not only shows acute toxicity to *P. citri* but also exhibits sublethal effects. The present study showed that the population parameters of *P. citri* had adverse effects under two concentrations of spiroticlofen stress. Furthermore, reduced expression of Vg mRNA under spiroticlofen markedly affects *P. citri* reproduction. Thus, as a part of pest management programme, the responses of spiroticlofen stress show an effective way to control *P. citri*. However, the mechanism should be studied

further in the future to clarify the effects against gene expression levels of detoxification enzymes and antioxidant enzymes.

**Acknowledgements.** This research was supported by the key project of the Ministry of Agriculture and Rural Affairs of China 'Investigation on the Occurrence, Harm and Diffusion Risk of Agricultural Alien Invasive Species' (13220141), The Graduate Innovation Foundation of Jiangxi Province, China (YC2022—B024).

**Competing interests.** None.

## References

- Akkopru EP, Atlihan R, Okut H and Chi H (2015) Demographic assessment of plant cultivar resistance to insect pests: a case study of the dusky-veined walnut aphid (Hemiptera: Callaphididae) on five walnut cultivars. *Journal of Economic Entomology* **108**, 378–387.
- Alavijeh E, Khajehali J, Snoeck S, Panteleri R, Ghadamyari M, Jonckheere W, Bajda S, Saalwaechter C, Geibel S, Douris V, Vontas J, Van LT and Dermauw W (2020) Molecular and genetic analysis of resistance to METI-I acaricides in Iranian populations of the citrus red mite *Panonychus citri*. *Pesticide Biochemistry and Physiology* **164**, 73–84.
- Amaral I, Melville CC, Rocha CM, Della VJF, Prado TJ and Andrade DJ (2020) Sublethal effects of spirodiclofen on biological and demographic parameters of the citrus leprosis mite *Brevipalpus yothersi* (Acari: Tenuipalpidae). *Pest Management Science* **76**, 1874–1880.
- Andrade DJ, Lorencon JR, Siqueira DS, Novelli VM and Bassanezi RB (2018) Space-time variability of citrus leprosis as strategic planning for crop management. *Pest Management Science* **74**, 1798–1803.
- Bartle B (1968) Outbreaks of two-spotted spider-mites and cotton aphids following Pesticide Treatment. I. Pest stimulation vs. natural enemy destruction as the cause of outbreaks. *Journal of economic entomology* **61**, 297–303.
- Bolter CJ and Chefurka W (1990) Extramitochondrial release of hydrogen peroxide from insect and mouse liver mitochondria using the respiratory inhibitors cyphosphine, myxothiazol, and antimycin and spectral analysis of inhibited cytochromes. *Archives of Biochemistry Biophysics* **278**, 65–72.
- Cheng L, Hou D, Sun Q, Yu S, Li S, Liu H, Cong L and Ran C (2022) Biochemical and molecular analysis of field resistance to Spirodiclofen in *Panonychus citri* (McGregor). *Insects* **13**, 1011.
- Chi H and Liu H (1985) Two new methods for study of insect population ecology. *Bulletin of the Institute of Zoology* **24**, 225–240.
- Claudiane MR, Jaqueline FDV, Patrice JS, Celso O and Daniel JA (2021) Resistance to spirodiclofen in *Brevipalpus yothersi* (Acari: Tenuipalpidae) from Brazilian citrus groves: detection, monitoring, and population performance. *Pest Management Science* **77**, 3099–3106.
- Cordeiro EM, de Moura IL, Fadini MA and Guedes RN (2013) Beyond selectivity: are behavioral avoidance and hormesis likely causes of pyrethroid-induced outbreaks of the southern red mite *Oligonychus ilicis*? *Chemosphere* **93**, 1111–1116.
- Desneux N, Decourtye A and Delpuech JM (2007) The sublethal effects of pesticides on beneficial arthropods. *Annual Review Of Entomology* **52**, 81–106.
- Döker İ, Kazak C and Ay R (2021) Resistance status and detoxification enzyme activity in ten populations of *Panonychus citri* (Acari: Tetranychidae) from Turkey. *Crop Protection* **141**, 105488.
- Dong J, Wang K, Li Y and Wang S (2017) Lethal and sublethal effects of cyantraniliprole on *Helicoverpa assulta* (Lepidoptera: Noctuidae). *Pesticide Biochemistry and Physiology* **136**, 58–63.
- Goel A, Dani V and Dhawan DK (2005) Protective effects of zinc on lipid peroxidation, antioxidant enzymes and hepatic histoarchitecture in chlorpyrifos-induced toxicity. *Chemico-Biological Interactions* **156**, 131–140.
- Huang L, Lu M, Han G, Du Y and Wang J (2016) Sublethal effects of chlorantraniliprole on development, reproduction and vitellogenin gene (CsVg) expression in the rice stem borer, *Chilo suppressalis*. *Pest Management Science* **72**, 2280–2286.
- Liu Y, Wang C, Qi S, He J and Bai Y (2021) The sublethal effects of ethiprole on the development, detoxification mechanisms, and immune pathways of honeybees (*Apis mellifera* L.). *Environmental Geochemistry and Health* **43**, 461–473.
- Ma J, Zhou C and Li Y (2014) Biochemical responses to the toxicity of the biocide abamectin on the freshwater snail *Physa acuta*. *Ecotoxicology and Environmental Safety* **101**, 31–35.
- Ma K, Tang Q, Xia J, Lv N and Gao X (2019) Fitness costs of sulfoxaflor resistance in the cotton aphid, *Aphis gossypii* Glover. *Pesticide Biochemistry and Physiology* **158**, 40–46.
- Mousavi MG, Maroofpour and Y and Nariman (2020) Insecticidal activity and sublethal effects of *Beauveria bassiana* (Bals.-Criv.) Vuill. isolates and essential oils against *Aphis gossypii* Glover, 1877 (Hemiptera: Aphididae). *Acta agriculturae Slovenica* **115**, 463–472.
- Pan D, Dou W, Yuan GR, Zhou QH and Wang JJ (2020) Monitoring the resistance of the citrus red mite (Acari: Tetranychidae) to four acaricides in different citrus orchards in China. *Journal of Economic Entomology* **113**, 918–923.
- Papachristos DP and Milonas PG (2008) Adverse effects of soil applied insecticides on the predatory coccinellid *Hippodamia undecimnotata* (Coleoptera: Coccinellidae). *Biological Control* **47**, 77–81.
- Qi S, Niu X, Wang DH, Wang C, Zhu L, Xue X, Zhang Z and Wu L (2020) Flumethrin at sublethal concentrations induces stresses in adult honey bees (*Apis mellifera* L.). *Science of the Total Environment* **700**, 134500.
- Quesada CR and Sadof CS (2019) Field evaluation of insecticides and application timing on natural enemies of selected armored and soft scales. *Biological Control* **133**, 81–90.
- Rahmani S and Bandani AR (2013) Sublethal concentrations of thiamethoxam adversely affect life table parameters of the aphid predator, *Hippodamia variegata* (Goeze) (Coleoptera: Coccinellidae). *Crop Protection* **54**, 168–175.
- Rasheed MAK, Hafeez MM, Zhao M, Islam J, Ali Y, Ur-Rehman S, Hani USE and Zhou X (2020) Lethal and sublethal effects of Chlorpyrifos on biological traits and feeding of the aphidophagous predator *Harmonia axyridis*. *Insects* **11**, 491–505.
- Sani B, Hamid G and Elham R (2018) Sublethal effects of chlorfenapyr on the life table parameters of two-spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae). *Systematic & Applied Acarology* **23**, 1342–.
- Ullah F, Gul H, Desneux N, Said F, Gao X and Song D (2020) Fitness costs in chlorfenapyr-resistant populations of the chive maggot, *Bradysia odoriphaga*. *Ecotoxicology* **29**, 407–416.
- Van LT, Van PS and Tirry L (2006) Biochemical analysis of a chlorfenapyr-selected resistant strain of *Tetranychus urticae* Koch. *Pest Management Science* **62**, 425–433.
- Wang L, Zhang Y, Xie W, Wu Q and Wang S (2016) Sublethal effects of spinetoram on the two-spotted spider mite, *Tetranychus urticae* (Acari: Tetranychidae). *Pesticide Biochemistry and Physiology* **132**, 102–107.
- Wang H, Xin T, Wang J, Zou Z, Zhong L and Xia B (2021) Sublethal effects of bifentazate on biological traits and enzymatic properties in the *Panonychus citri* (Acari: Tetranychidae). *Scientific Reports* **11**, 20934.
- Yamamoto A, Yoneda H, Hatano R and Asada M (1995) Genetic analysis of Hexythiazox resistance in the citrus red mite, *Panonychus citri* (MCGREGOR). *Journal of Pesticide Science* **20**, 513–519.
- Zhang XL, Mao KK, Liao X, He B, Jin RH, Tang T, Wan H and Li JH (2018) Fitness cost of nitenpyram resistance in the brown planthopper *Nilaparvata lugens*. *Journal of Pest Science* **91**, 1145–1151.
- Zhang Y, Guo L, Atlihan R, Chi H and Chu D (2019) Demographic analysis of progeny fitness and timing of resurgence of *Laodelphax striatellus* after insecticides exposure. *Entomologia Generalis* **39**, 221–230.
- Zhang S, Wang X, Gu F, Gong C, Chen L, Zhang Y, Hasnain A, Shen L and Jiang C (2020) Sublethal effects of triflumezopyrim on biological traits and detoxification enzyme activities in the small brown planthopper *Laodelphax striatellus* (Hemiptera: Delphacidae). *Frontiers in Physiology* **11**, 261.
- Zhao J, Sun Y, Xiao L, Tan Y, Jiang Y and Bai L (2018a) Vitellogenin and vitellogenin receptor gene expression profiles in *Spodoptera exigua* are related to host plant suitability. *Pest Management Science* **74**, 950–958.
- Zhao X, Wang SQ, Wang XY and Cui K (2018b) A multi-state shock model with mutative failure patterns. *Reliability Engineering and System Safety* **178**, 1–11.
- Zhao Y, Wang Q, Ding J, Wang Y, Zhang Z, Liu F and Mu W (2018c) Sublethal effects of chlorfenapyr on the life table parameters, nutritional physiology and enzymatic properties of *Bradysia odoriphaga* (Diptera: Scliaridae). *Pesticide Biochemistry and Physiology* **148**, 93–102.
- Ziegler R and Antwerpen RV (2006) Lipid uptake by insect oocytes. *Insect Biochemistry and Molecular Biology* **36**, 264–272.