

- 36** Proudfoot J, Goldberg A, Mann A, Everitt B, Marks I, Gray JA. Computerized, interactive, multimedia cognitive behavioral program for anxiety and depression in general practice. *Psychol Med* 2003; **33**: 217–27.
- 37** Cavanagh K, Shapiro DA. Computer treatment for common mental health problems. *J Clin Psychol* 2004; **60**: 239–51.
- 38** Clarke G, Eubanks D, O'Connor E, DeBar LL, Kelleher C, Lynch F, et al. Overcoming Depression on the Internet (ODIN): a randomised controlled trial of an Internet depression skills intervention program. *J Med Internet Res* 2002; **4**: e16.
- 39** Patten SB. Prevention of depressive symptoms through the use of distance technologies. *Psychiatr Serv* 2003; **54**: 396–8.
- 40** Clarke G, Eubanks D, Reid E, Kelleher C, O'Connor E, DeBar LL, et al. Overcoming depression on the internet (ODIN) (2): a randomized trial of a self-help depression skills program with reminders. *J Med Internet Res* 2005; **7**: e16.
- 41** Carlbring P, Gunnarsdóttir M, Hedensjö L, Andersson G, Ekselius L, Furmark T. Treatment of social phobia: randomised trial of internet-delivered cognitive-behavioural therapy with telephone support. *Br J Psychiatry* 2007; **190**: 123–8.
- 42** Kenwright M, Marks I, Graham C, Franses A, Mataix-Cols D. Brief scheduled phone support from a clinician to enhance computer-aided self-help for obsessive-compulsive disorder: randomized controlled trial. *J Clin Psychol* 2005; **61**: 1499–508.
- 43** Vuorilehto M, Melartin TK, Ryttsä HJ, Isometsä E. Do characteristics of patients with major depressive disorder differ between primary and psychiatric care? *Psychol Med* 2007; **37**: 893–904.
- 44** Hamilton KE, Dobson KS. Cognitive therapy of depression: pretreatment patient predictors of outcome. *Clin Psychol Rev* 2002; **22**: 875–93.
- 45** Walsh BT, Seidman SN, Sysko R, Gould M. Placebo response in studies of major depression: variable, substantial, and growing. *JAMA* 2002; **287**: 1840–7.
- 46** Bialik RJ, Ravindran AV, Bakish D, Lapierre YD. A comparison of placebo responders and nonresponders in subgroups of depressive disorder. *J Psychiatry Neurosci* 1995; **20**: 265–70.
- 47** Posternak MA, Miller I. Untreated short-term course of major depression: a meta-analysis of outcomes from studies using wait-list control groups. *J Affect Disord* 2001; **66**: 139–46.
- 48** DeRubeis RJ, Hollon SD, Amsterdam JD, Shelton RC, Young PR, Salomon RM, et al. Cognitive therapy vs medications in the treatment of moderate to severe depression. *Arch Gen Psychiatry* 2005; **62**: 409–16.
- 49** Dimidjian S, Hollon SD, Dobson KS, Schmaling KB, Kohlenberg RJ, Addis ME, et al. Randomized trial of behavioral activation, cognitive therapy, and antidepressant medication in the acute treatment of adults with major depression. *J Consult Clin Psychol* 2006; **74**: 658–70.
- 50** Katon W, Von Korff M, Lin E, Walker E, Simon GE, Bush T, et al. Collaborative management to achieve treatment guidelines. Impact on depression in primary care. *JAMA* 1995; **273**: 1026–31.
- 51** Scogin FR, Hanson A, Welsh D. Self-administered treatment in stepped-care models of depression treatment. *J Clin Psychol* 2003; **59**: 341–9.
- 52** McCrone P, Knapp M, Proudfoot J, Ryden C, Cavanagh K, Shapiro DA, et al. Cost-effectiveness of computerised cognitive-behavioural therapy for anxiety and depression in primary care: randomised controlled trial. *Br J Psychiatry* 2004; **185**: 55–62.
- 53** Kalenthaler E, Sutcliffe P, Parry G, Beverly C, Rees A, Ferriter M. The acceptability to patients of computerized cognitive behaviour therapy for depression: a systematic review. *Psychol Med* 2008; **38**: 1521–30.
- 54** Paykel ES, Tylee A, Wright A, Priest RG. The Defeat Depression Campaign: psychiatry in the public arena. *Am J Psychiatry* 1997; **154**: 59–65.
- 55** Butcher JN, Perry J, Hahn J. Computers in clinical assessment: historical developments, present status, and future challenges. *J Clin Psychol* 2004; **60**: 331–45.
- 56** Butcher JN, Perry JN, Atlis MM. Validity and utility of computer-based test interpretation. *Psychol Assess* 2000; **12**: 6–18.
- 57** Schulenberg SE, Yutrzenka BA. Equivalence of computerized and conventional versions of the Beck Depression Inventory-II (BDI-II). *Current Psychology* 2001; **20**: 216–30.
- 58** Bijl RV, Ravelli A, van Zessen G. Prevalence of psychiatric disorder in the general population: results of the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Soc Psychiatry Psychiatr Epidemiol* 1998; **33**: 587–95.
- 59** Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA* 2003; **289**: 3095–105.
- 60** Andersson G, Cuijpers P. Pros and cons of online cognitive-behavioural therapy. *Br J Psychiatry* 2008; **193**: 270–1.

Stigma

**100
words**

Peter Byrne

Stigma is a prejudice (negative attitude) based on stereotypes usually leading to discrimination. Familiar mental illness stereotypes (weak, violent, comic) drive prejudice in society. Discrimination ranges from simple avoidance through exit life events (relationship, employment and housing losses) and institutional discrimination (denial of health interventions, insurance, jury service, visa *inter alia*). Stigma-discrimination cannot occur without a power differential. When a person with mental health problems shares societal prejudices, their self-stigma contributes to further morbidity and status loss. Reducing stigma requires multiple interventions: a language of inclusion (no more 'schizophrenics'), legal and organisational reforms, and cultural changes based on empowerment.

The British Journal of Psychiatry (2009)
195, 80. doi: 10.1192/bjp.195.1.80