

This is an Accepted Manuscript for Parasitology. This version may be subject to change during the production process. DOI: 10.1017/S0031182024001331

## Assessment of farmers' knowledge, attitudes and control practices (KAP) to mitigate acaricide resistance and tick borne diseases

Sunita Jamra<sup>1</sup>, Mukesh Shakya<sup>1</sup>, Anant K. Jayraw<sup>1</sup>, Vivek Agrawal<sup>1</sup>, Mamta Singh<sup>2</sup>, Anil Kumar Sharma<sup>3</sup>, Gajendra N. Bhangale<sup>4</sup>, Gaya P. Jatav<sup>5</sup> and Nirmala Jamra<sup>1</sup>

<sup>1</sup>Department of Veterinary Parasitology, College of Veterinary Science and Animal Husbandry (NDVSU), Mhow-453446, Madhya Pradesh, India

<sup>2</sup>Department of Veterinary Medicine, College of Veterinary Science and Animal Husbandry (NDVSU), Mhow-453446, Madhya Pradesh, India

<sup>3</sup>School of Agriculture, Uttarakhand University, Dehradun-248007, Uttarakhand, India

<sup>4</sup>Department of Veterinary Parasitology, College of Veterinary Science and Animal Sciences, Parbhani (MAFSU), Maharashtra, India

<sup>5</sup>Department of Veterinary Pathology, College of Veterinary Science and Animal Husbandry (NDVSU), Mhow-453446, Madhya Pradesh, India

**Corresponding author:** Mukesh Shakya, Email: [drmukeshsadvet@gmail.com](mailto:drmukeshsadvet@gmail.com)

This is an Open Access article, distributed under the terms of the Creative Commons Attribution - NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

## Abstract

A KAP-based study on ticks and tick-borne diseases (TTBD) and resistance development in ticks was conducted in Dhar district of Madhya Pradesh covering 200 livestock owners using a questionnaire. Based on our scoring criteria, results indicated only 25% (19.16-31.60) respondents possessing basic knowledge of TTBDs while 75% (68.40-80.84) respondents were not aware of TTBDs. Due to lack of proper awareness of TTBDs, about 1.28 times more respondents (OR 95% CI = 0.42–3.86) were having heavy tick infestations in their animals. However, about 36.5% (29.82-43.58) respondents showed a favorable attitude towards the adoption of different tick control practices consequently, their animals showed low level infestation. Amongst various feeding system for animals, a mixed type of feeding system was mostly adopted by 57.5% respondents followed by manger system (37.5%) while grazing was least adopted method (5%). Results indicated that the grazing animals were six times (OR 95% CI = 2.93–12.28) more susceptible to ticks and possessed heavy tick infestation. Resistance status of collected tick isolates of *Rhipicephalus microplus* and *Hyalomma anatolicum* was assessed and revealed that both tick species were found resistant to deltamethrin (DLM). The goals of this study were to assess some of the underlying causes of ticks and TBD in livestock in Dhar district of Madhya Pradesh state using the KAP survey and resistance characterization of ticks.

**Keywords:** Acaricide, Deltamethrin, Epidemiology, Fipronil, *Hyalomma anatolicum*, *Rhipicephalus microplus*, Resistance

## Introduction

Tick infestation and its impact are significant challenges that limit livestock output in cattle rearing communities mainly in tropical and subtropical countries. It is reported that about 80% of cattle population globally are adversely affected by ticks and the tick-borne diseases (de Castro *et al.*, 1997). In India, the key species of cattle ticks are *Rhipicephalus microplus* and *Hyalomma anatolicum* (Ghosh *et al.*, 2006) which serve as a vector of fatal diseases like anaplasmosis, babesiosis and theileriosis. Globally, ticks cause economic annual losses of US\$22–30 billion in cattle by transmitting tick-borne diseases (TBDs) (Hurtado and Giraldo-Rios *et al.*, 2018). In Brazil, *R. microplus* alone causes a loss of \$ 32.4 million per year (Grisi *et al.*, 2014). However, losses estimated due to TTBDs varies by country such as \$ 3.0 million (Graham and Hourrigan, 1977) in USA, \$ 573.16 million in Mexico, \$ 168.0 million in Colombia (Rodríguez-Vivas *et al.*, 2017), \$ 250.0 million in Australia (Meat and Livestock Australia report 2020), \$ 364.0 million in Tanzania (Kivaria, 2006), \$ 6.7 million in Puerto Rico, and \$ 5.0 million in Zambia (Senbill *et al.*, 2018). In India, economic impact of TTBDs was estimated over \$ 787.63 million per annum (Singh *et al.*, 2022).

No specific study was focused to create a more efficient, long-lasting, and comprehensive tick control approach, or to assess the performance of existing control strategies beyond the traditional application of acaricides (Jongejan and Uilenberg, 2004). In India, generally, four chemical classes of acaricides i.e. organophosphates, synthetic pyrethroids, amidines, and avermectins (Fular *et al.*, 2021) are commonly used for tick management. Some of these chemicals are not effectively working against ticks in many parts of the country due to the development of acaricidal resistance (Bisht *et al.*, 2022; Shakya *et al.*, 2023). The use of acaricides on animals is prevented due to increase in resistant tick populations, their high cost, negative effects on unintended species, and acaricidal residues in animal products (Singh *et al.*, 2022). The success of tick control program is based on

comprehensive farmers' knowledge on TBDs, their perspective on efficient control methods, and the socio-cultural environment in which the program is carried out. The information is usually collected using the commonly used knowledge, attitude and practices (KAP) survey (Launiala, 2009). The method sets the initial standard for future evaluation and analysis of the impact of knowledge, attitude, and practice on modifying TBD-related issues. It proposes an intervention approach that takes into account the unique local conditions and the cultural variables that shape them, and designs activities that are appropriate for the particular community concerned (Gumicio *et al.*, 2011). Despite criticism for generalized data of a large population for planning purposes, KAP surveys on TBDs have played a significant role in developing effective intervention strategies (Butler *et al.*, 2016; Zoldi *et al.*, 2017; Niesobecki *et al.*, 2019; Gupta *et al.*, 2021).

The animal owners of Dhar district of Madhya Pradesh face problem of resistance development in ticks and their management due to lack of KAP-based data. This data is essential to formulate suitable strategy to manage resistant ticks and to improve livestock health and the income of the marginal animal owners. Thus, to tackle the problem in the targeted region, a KAP-based study was conducted to assess the influence of TBDs on livestock productivity and determination of resistance status of tick populations, and the control strategies adopted by livestock owners. The collected data will aid in creating efficient animal health initiatives to boost livestock output and to enhance the socioeconomic status of livestock owners of targeted region.

## **Materials and methods**

### *Study area*

Dhar district is located in Malwa region of Madhya Pradesh of India and was selected for conducting the KAP survey. It possesses a diverse terrain with altitudes ranging from 150 to

600 m above sea level, influenced by the Vindhya Range. The vegetation consists of dry deciduous forests, including teak, sal, and bamboo, with more dense forest cover in the hilly regions. The semi-arid climate and topographical variations contribute to the presence of grasslands and scrub forests in lower areas. The geographic locations of different subdivisions of Dhar district are Dhar (DHA, 75.32°N, 22.61°E), Manawar (MAN, 75.08°N, 22.23°E), Sardarpur (SAR, 74.97°N, 22.65°E), Kukshi (KUK, 74.75°N, 22.20°E) and Gandhwani (GAN, 75.08°N, 22.23°E). Cattle and buffaloes are primarily reared for milk production, contributing significantly to the livelihoods of the local population. However, challenges such as limited access to quality feed, veterinary services, and water resources can affect the productivity of milch animals in the region. The organized farms included more than minimum ten milch animals, well maintained shelter with proper cemented flooring infrastructure for animals. On the other hand, unorganized farm included households animals which had mud flooring and no proper amenities and only two or three animals were maintained for personal purpose.

### *Questionnaire survey*

A systematic questionnaire was designed to gather data on several aspects associated with cattle productivity and TTBDs. A questionnaire proforma was designed in a multiple-choice form as per the guidelines (Thrusfield, 2018), with modifications made via both informal and formal testing processes. The questionnaire proforma contained several subjects like socio-demographic information, animal sheds, animal feeding methods, shed conditions, farming practices, methods of acaricidal application, risk factors, etc. The survey was carried out bi-monthly from February 2022 to January 2023 to monitor seasonal variations in cattle productivity and the prevalence of tick-borne diseases (TTBDs) and the questionnaire was provided to livestock owners at the surveyed places. The study authors conducted the survey

through face-to-face interviews with the owner. There were no specific inclusion and exclusion criteria for the participants in the study. The survey was intended for household heads, however, if there were other persons involved in livestock rearing in the household, they were asked to be in the survey as well. The questionnaire underwent pilot testing to ensure its effectiveness in gathering correct information (Williams, 2003). Prior to providing the questionnaire to the targeted participants, it was reviewed by a number of experienced investigators of epidemiological study. Then, the data was carefully collected, analyzed and screened for accuracy. The farmers selected for the investigation were chosen for their willingness to participate and operational convenience. They were owners of ruminant herds consisting of 5 to 15 animals (Hussain *et al.*, 2021). About 200 individual interviews were conducted with livestock owners using a developed questionnaire.

#### *Tick collection and processing*

The biological samples of *Rhipicephalus microplus* ticks were collected from different regions of Dhar district following a randomized sampling procedure. Tick samples were collected from cattle and buffaloes of the households and well managed dairy farms. Engorged female ticks were collected in labeled sample bottles covered with cotton cloth, and brought to the research centre. In the laboratory, at least 100–150 engorged female ticks collected from each sub-division were pooled and placed in Petri plates (five ticks per plate) and maintained in the laboratory (Ghosh and Azhahianambi, 2007). They were then kept at 28 °C and 85±5% relative humidity for normal oviposition. The ticks procured from Dhar, Manawar, Sardarpur, Kukshi and Gandhwani sub-divisions were encoded as DHA, MAN, SAR, KUK, and GAN isolates, respectively. The eggs laid by the female ticks from each sub-division were combined and identified as a representative sample of that sub-division. The eggs of each sub-division were pooled collected and stored in tick rearing tubes. Once

hatched, the larvae were placed in an incubator set at 28°C and 85±5% relative humidity for 8–10 days for larval-based experiments.

#### *Identification of ticks*

The collected tick samples of both the sexes were observed morphologically under stereomicroscope. The specific characters of *R. microplus* and *H. anatolicum* were identified with the help of book ‘Helminths, Arthropods and Protozoa of Domesticated Animals’ (Soulsby, 1982) and then characterized them.

#### *Reference tick*

The reference susceptible IVRI-I strain of *R. microplus* was used as the reference tick for resistance characterization. The IVRI-I strain is already characterized as susceptible to most of the chemical acaricides in the Entomology laboratory of Indian Veterinary Research Institute, Izatnagar.

#### *Chemical acaricides*

Technical grade deltamethrin and fipronil were procured from Sigma Aldrich (St Louis, MO, USA) and their stock solutions of 5000 and 1000 ppm, respectively, were prepared in methanol. Working concentrations of deltamethrin (60, 90, 120, 150 and 180 ppm) and fipronil (10, 15, 20, 25 and 30 ppm) were prepared in distilled water from their stock solutions and were tested for resistance characterization in collected tick samples.

## *Resistance characterization*

### *Larval packet test (LPT)*

A modified version of the larval packet test (LPT) as recommended by the FAO (2004) was used. The packets were prepared in triangular shape from Whatmann filter paper No. 1 measuring 5.5 cm × 5 cm. These packets were soaked with 0.7 ml solution of acaricide and then dried at 37 °C in hot air oven. After drying, one side of the packets was sealed with adhesive tape. Then, about 150-200 larvae of 7-10 days old were introduced in the packets and sealed using a “bulldog” clips. The packets were then kept in for 24h in a biological oxygen demand (BOD) chamber at 28°C and 85 ± 5% relative humidity. After 24 h, these packets were removed from the BOD, and opened on white paper sheet under electric lamp to observe dead and alive larvae. The larvae only moving their legs were considered as dead while running larvae were counted as alive. Accordingly, the mortality percentage of larvae was determined by counting the number of alive and dead larvae. Three replications were maintained for each concentration of acaricides along with control with distilled water.

### *Larval immersion test (LIT)*

Shaw (1966) was the initial developer of the larval immersion test. The Shaw's immersion sandwich method involves larval immersion in an acaricide solution or suspension. For the assay, more than 300 larvae were transferred into 1.5 ml microcentrifuge tubes (three repetitions per dilution) with the help of drawing brush and then an amount of 0.75 ml of working solution of the acaricide was poured in these tubes. The larvae were submerged for 10 minutes and agitated intermittently. After opening the tubes, approximately 100 larvae were transferred to filter paper packets and sealed with "bulldog" clips. The packets were kept at 28 °C and a relative humidity of 85 ± 5% for 24 h. Control groups of each acaricide



were also immersed in distilled water in the same way. After 24 h, larval mortality was assessed as mentioned in LPT.

The resistance status of field isolates was determined on the basis of resistance ratio (RR). The resistance ratio (RR50) is the ratio of LC50 value of an acaricide for field ticks and LC50 value of the acaricide for reference susceptible IVRI-I strain (Castro-Janer *et al.*, 2009). Ticks were then classified according to various resistance levels as per the method of Sharma *et al.* (2012).

### *Statistical analysis*

The questionnaire data from 200 respondents were transferred to the Microsoft Excel 2010 sheet for proper management and analysis. The proportions of variables recorded in questionnaires were analyzed following descriptive statistics (frequencies and percentages). The data were analyzed by Epi Info™ software (Centers for Disease Control and Prevention, Atlanta, Georgia, US). Association of socio-demographic characteristics to level of tick infestation was analysed by Chi-square test. Simple logistic regression analysis through R-software package (dplyr) was also performed to observe the association of respondent's knowledge and level of tick infestations (Wickham *et al.*, 2021). The dose–response data of LPT and LIT were subjected to probit analysis (Finney, 1971) using GraphPad Prism v.5 statistical software (GraphPad Software, San Diego, California USA) to determine LC50 values of each acaricide.

## **Results**

### *Collection of ticks and farm management practices*

The tick isolates were collected from the households and well managed dairy farms located in five sub-divisions of Dhar district of Madhya Pradesh. There were both cross-bred and native

breeds of cattle and buffaloes in the district. The surveyed animals were found to have a moderate (>50–100 ticks/animal) to high (>150–200 ticks/animal) level of tick infestations. Despite repeated applications of different synthetic acaricides such as cypermethrin, deltamethrin, ivermectin and amitraz, a significant number of farmers reported the failure of tick control. During the survey, it was noticed that the application of fipronil in the dairy farms is not frequent and almost lacking. However, the application schedule for other synthetic acaricides was not properly maintained and the animals were treated whenever tick infestation was visible on animals. The frequency of acaricidal treatment of household animals was comparatively lower than those maintained in well managed dairy farms. The targeted area was highly dominated by the tribal population where animals were kept in small to big huts made of mud, concrete, and thatched roofs with no proper acaricidal dose and application. It was noticed that the farmers rarely applied insecticides in the animal sheds to eradicate off the host tick stages (Table 1).

#### *Analysis of socio-demographic characteristics of the respondents*

The study comprised 200 farmers including literate and illiterate from five sub-divisions of Dhar district and more than 90% famers belonged to rural areas. It was interesting to mention that farmers showed their interest to adopt new techniques and technologies for the management of TBDs. Face-to-face interviews revealed that about 40% (33.15-47.15) respondents were literate and 60% (52.85-66.85) were illiterate, out of which 55% (47.82-62.02) were using uncemented floors and 45% (37.98-52.18) were using cemented floors for their animals. Respondents adopted different feeding methods for their animals as observed during the survey. It was found that about 37.5 % (31.25-45.11) respondents fed their animals in manger, only 5% (2.42-9.00) adopted grazing system and the rest (42.5%) adopted a mixed type of feeding system. Only 25% (19.16-31.60) respondents had knowledge about TTBDs

while 75% (68.40-80.84) were not aware of it. During the study, we observed that the cypermethrin was preferred for animal application by the livestock owners i.e. [35% (28.41-42.05)] followed by deltamethrin [29% (22.82-35.82)] and ivermectin [15% (10.35-20.72)] while only 9% (5.42-13.85) farmers were applying amitraz for tick control. The respondents [12% (7.84-17.33)] were using more than one acaricide without maintaining any fixed application pattern and hence they were considered in the mixed category. . A favorable attitude towards different tick control methods was showed by 36.5% (29.82-43.58) respondents in which manual hand picking as well as chemical control methods were most preferred methods (Table 2).

*Association of socio-demographic characteristics to level of tick infestation by Chi-square test*

Analysis by Chi-square test revealed that the animals of literate respondents were significantly ( $p < 0.05$ ) less susceptible to tick infestation as compared to those of illiterate respondents ( $p = 0.0180$ ). The animals kept in the uncemented floor of shed exhibited a high intensity of tick infestation (0.0029). The animals having manger feeding were observed to be less susceptible to ticks as compared to grazing and mixed feeding animals ( $p < 0.0335$ ) indicating the significant association ( $p < 0.05$ ). Amongst 200 respondents, only 50 exhibited knowledge regarding TBDs. The data obtained was found to be statistically significant at a 5% level ( $p = 0.0063$ ) (Table 3). Insignificant differences were observed between tick infestation level and acaricides used for tick control by respondents.

*Association of socio-demographic characteristics to level of tick infestation by logistic regression analysis*

The data analyzed by logistic regression revealed that the animals of female livestock owners were 2.20 times (OR 95% CI = 1.13-4.27) more likely than male livestock owners to experience tick infestation. Although the level of tick infestation was not considerably impacted by respondents' educational levels, their attitudes towards various tick-control strategies were greatly influenced. Similarly, the livestock owners without having knowledge of TBDs were 1.28 times (OR 95% CI = 0.42–3.86) more likely to have a high level of tick infestation (Table 4). Moreover, the low level infestation was recorded in the animals of respondents having a favorable attitude towards different tick control methods (OR=1.04, 95% CI = 0.4–2.66). The respondents who practiced grazing as a sole method of feeding for their animals were likely to be more susceptible by six times (OR 95% CI = 2.93–12.28) to ticks and had a heavy tick infestation as compared to mangers and mixed feeding practices. No significant difference between level of tick infestation and acaricides used for tick control was observed (Table 5).

*Association of socio-demographic characteristics to level of tick infestation by R software analysis (Table 6)*

The multiple logistic regression analysis showed that shed floor type, feeding system, and acaricides used for tick control were the significant variables in this model. Respondents having uncemented animal sheds were 5.16 times [OR = 5.16] more likely to have a high level of tick infestation in their animals as compared to those having cemented floor of sheds. The respondents adopted three types of feeding systems: grazing, manger, and mixed feeding. The respondents adopting a grazing system showed that their animals were 4.10 times [OR = 4.10] more likely to have a high level infestation as compared to those kept in a manger or

mixed feeding system while the other variable in the model is held constant. The acaricides commonly used by respondents also significantly affected the tick infestation level [OR = 1.77]. The interaction term (i.e., Sex, Literacy, Knowledge about TTBDs and Attitude towards tick control) was not significant in this analysis (Fig. 1, Table 6).

### *Resistance status of deltamethrin and fipronil*

To identify the generation of acaricide resistance in tick population collected from different sites of a district, the larval based assays i.e. LPT and LIT were conducted in the laboratory against deltamethrin (DLM) and fipronil (FIP) and the larval mortality was recorded after 24 h. In case of *R. microplus*, the tested isolates were highly resistant to deltamethrin (RF = 33.9–39.9) as observed by LPT (Table 7). Surprisingly, in case of LIT, a low level of resistance against deltamethrin was detected (RF = 1.2–4.3) (Table 7). All the isolates were susceptible to fipronil by LPT (RR= 0.17–0.24) and LIT (RR= 0.48–0.51). The LC50 values were ranging from 400.69- 471.6 ppm and 15.02-51.42 ppm against DLM in LPT and LIT format. The lower mortality slopes were observed in all isolates as compared to reference susceptible IVRI-I strain ( $3.42 \pm 0.49$ ) indicating presence of more heterogeneous deltamethrin resistant population of *R. microplus*. The results indicated that the ticks of this area developed resistance against deltamethrin (Fig. 2, Table 7).

The LPT and LIT based resistance data of *H. anatolicum* against deltamethrin and fipronil is documented in table 8. Results revealed that all the field isolates were resistant to DLM at level II (RF = 11.1–16.6) by LPT. Similarly, in case of LIT, an initiation of resistance to DLM was detected (RF = 1.5–2.3). Like *R. microplus* all the isolates of *H. anatolicum* were also found susceptible to FIP. The LC50 values were in the range from 132.17 - 194.90 ppm and 18.53 - 28.04 ppm against DLM in LPT and LIT, respectively. The lower mortality slopes were seen in all the samples in comparison to IVRI-I strain ( $3.42 \pm 0.49$ ) except Dhar

isolate ( $5.002 \pm 1.23$ ) indicating the presence of more heterogeneous deltamethrin resistant populations of *H. anatolicum* (Fig. 3).

## Discussion

This research is the first to investigate the knowledge, attitudes, and behaviors of livestock owners in Madhya Pradesh, India, about ticks and tick control measures. The report also covers the perceptions of stakeholders and livestock farmers in the study region about these limitations. Historically, most of the animal health researches worldwide were focused on pastoral regions (Catley *et al.*, 2014; Queenan *et al.*, 2017). Pastoral communities predominate in the majority of African nations; in contrast, mixed crop-livestock farming practices and production systems are widespread in India (Hemme *et al.*, 2013). More than half of the surveyed respondents did not know how their livestock become infested with ticks or where ticks are typically located in the environment, despite the fact that every respondent had encountered a tick problem. According to some respondents, ticks are less prevalent in the winter. As temperature is an important factor in several tick developmental processes, including moulting, oviposition, and questing, low temperature in the winter is typically expected to slow down these processes (Estrada-Peña, 2020). Ticks are only known to search for hosts at temperatures over 7°C (Süss *et al.*, 2008; Namgyal *et al.*, 2021a). However, winter in the Dhar district of Madhya Pradesh is typically dry and cold, with a mean temperature of 15–20°C. In this part of Madhya Pradesh, ticks are more prevalent in rainy seasons due to high temperatures and humidity conditions.

The present study revealed a lack of knowledge among livestock owners on TBDs. The similarities in education and livestock-rearing techniques suggest that these findings may be applicable to other areas and countries (Chakraborty *et al.*, 2023). Ticks and TBDs lead to extensive veterinary and public health issues, particularly in India. TBDs and severe tick

infestations have been linked to reduced milk, meat, and other animal product output in several developing nations, along with increased animal sickness and death. Ticks transmit a greater number of illnesses than any other blood-feeding arthropod globally, posing a threat to people, their pets, and cattle (Rahman *et al.*, 2019; Ngnindji-Youdje *et al.*, 2022). Indigenous cow breeds are often believed to have high resistance to ticks and may be reared without proper attention on tick management (Phanchung *et al.*, 2007; Minjauw and McLeod, 2003; Jonsson *et al.*, 2014).

Face-to-face interviews in the present study helped to understand farmers' knowledge, their attitudes towards TBDs, and acaricide application patterns in fields. The current data indicate majority of farmers to be illiterate and lacking awareness of the TBDs. Most of the farmers used traditional uncemented sheds to maintain their livestock. Besides, the majority of animal owners were using chemical acaricides on their animals without adhering to suitable tick management methods and dosage regime. Animals were classified as having low, moderate, or high levels of tick infestation based on the presence of 25, 100, and 150 ticks, as documented by Chigure *et al.* (2018). We observed that several farms were severely infected with ticks, leading to a decrease in total productivity. A high negative association was seen between the frequency of acaricide usage and the proportion of tick-infested animals. This suggests that the frequent and effective use of acaricides is a significant factor contributing to the variation in tick prevalence across various farms. Indian researchers determined deltamethrin as the most commonly used acaricide in the field, followed by cypermethrin, amitraz, and ivermectin and observed that farms experienced high tick infestation, possibly due to owners' lack of awareness about the correct use of acaricides and the resistance of ticks to the products being used (Ghosh *et al.*, 2015; Chigure *et al.*, 2018; Shakya *et al.*, 2020; Upadhaya *et al.* (2020). According to, Husain *et al.* (2021), out of the livestock owners in the survey, 51 (45.5%) used acaricides frequently, but 49 (43.8%) did not

have appropriate disposal methods for spent acaricidal bottles and unused goods, opting to dispose of them in general waste streams, including farm drainage systems. Thirty-four livestock owners, accounting for 30.4% of the total, did not use any acaricides in the year before to our visit, although they had used them previously. Regarding application techniques, 26 farmers (23.2%) used systemic acaricide, while 34 farmers (30.4%) employed topical treatments for tick control. Tesfaye and Abate (2023), noted that the respondents estimated the amount of the acaricide instead of monitoring doses (whether sprayed or injected) before treatment. In our findings, researchers conducted an investigation that revealed that native breeds were allowed to graze outside, but cross-bred animals were kept confined in a shed. Native breeds have a lower tick infection rate compared to cross-breeds. In many countries that are developing, herd owners acquire acaricide use information from persons without expertise, leading to improper acaricide practices. In the present surveyed places, rural veterinary stores and shop workers with little technical knowledge serve as the primary source of information for farmers, leading to inadequate and improper acaricidal practices. In a prior research conducted in Kenya (Mugambi *et al.*, 2012), it was shown that many herd owners get information on acaricide administration from untrained vet shop attendants. This lack of sufficient training in animal health care might result in herd owners engaging in harmful practices. Recommendations to farmers were given to rotate the use of acaricides in cattle to reduce acaricide resistance and for cost-effective treatment due to the high frequency of TTBDs (Ghosh and Azhahianambi, 2007).

Most respondents and farmers lack awareness of TBDs and expressed unfavorable attitudes about tick management during face-to-face interviews. Similarly, researchers worldwide shared their views on the knowledge and attitudes of respondents. For instance, Lontsi-Demano *et al.* (2021) conducted a cross-sectional survey to evaluate farmers' knowledge and practices regarding ticks and the management of tick borne diseases. They



found that herd managers possessed a fundamental understanding of ticks and their impact on animals. Namgyal *et al.* (2021b) observed that 128 out of 246 respondents (52%) had sufficient information regarding ticks as carriers of illnesses in people and animals. Hussain *et al.* (2021) studied how cattle producers perceive and handle tick infestation. In another study, Hussain *et al.* (2021) determined that 47.3% of cattle owners were knowledgeable about tick-borne diseases and used sandy flooring, indicating awareness of the related risk factors. In the present study, the most popular animal feeding system is mixed type (57.5%) followed by stall feeding (37.5%). Similarly, Hussain *et al.* (2021) noted that 25% of farmers used stall feeding, and 53.6% embraced both methods. Tesfaye and Abate (2023) reported that the prevalence of tick infestation sometimes increased. Insufficient grazing habitat has caused animal herds to cluster in some areas, resulting in a higher spread of tick infestation.

The present study used the LPT, initially developed by Stone and Haydock (1962), and the LIT, developed by Shaw (1966), to identify and monitor resistance to acaricides. In this study, we found that, all five isolates of *R. microplus* and *H. anatolicum* collected from five different sub-divisions of Dhar district were found to be resistant to deltamethrin which may be due to extensive use of synthetic pyrethroid compounds and easy availability of this compound. Accordingly, the DLM resistance in both the tick species has been reported across the country. For example, Jyothimol *et al.* (2014) reported comparatively low level of resistance in field tick larvae collected from two districts of Kerala. Shyma *et al.* (2015) and Gaur *et al.* (2016) also reported DLM resistance in field ticks collected from Haryana, Rajasthan and Gujarat states of India. Similarly, Kumar *et al.* (2017) reported tick larvae from six districts of Andhra Pradesh state and reported RF of 1.05 to 8.78. The ineffectiveness of DLM was also reported from the states like Uttar Pradesh, Assam, and Maharashtra at resistance levels I-IV (Chigure *et al.*, 2018; Updhayay *et al.*, 2020; Khating *et al.*, 2024). DLM resistance has also been reported in *R. microplus* from West Africa (Adehan

*et al.*, 2016; Yessinou *et al.*, 2018), Mexico (Rosario-Cruz *et al.*, 2009), and Australia (Gurrero *et al.*, 2012). Besides *R. microplus*, Becker *et al.* (2019) reported resistance in *R. sanguineus* isolate collected from eight Porto Alegre metropolitan areas, Brazil with RF 1.18 to 5.67.

Earlier, country specific discriminating concentration ( $DC=2 \times LC_{99}$ ) of FIP was determined as 9.6 ppm using LPT against reference susceptible IVRI-I strain of *R. microplus* (Kumar *et al.*, 2016) for differentiating between susceptible and resistant ticks. In the present study, all the collected isolates of *R. microplus* and *H. anaticum* were found susceptible to fipronil. This may be due to the high cost and comparatively less use of fipronil for tick control. Recently, Shakya *et al.* (2020) characterized twenty-five isolates collected from six states (Madhya Pradesh, Uttarakhand, Meghalaya, Assam, Gujarat and Haryana) and reported RF in the range of 0.39 to 10.9. Analyzing the data, it is observed that fipronil is not widely adopted in most of the countries for the management of ticks, and therefore, reports on development of fipronil resistance in tick population are not frequently available in the literature.

## Conclusion

The results provide useful insights to aid in the development of educational and outreach programs that may go beyond the research region. The proper knowledge of TBDs among the animal owners is essential for effective management of tick infestation and improvement of animal health and productivity. The present study to mitigate acaricide resistance and TBDs revealed significant gaps in awareness and proper management strategies. Some farmers showed a basic understanding of tick control, the majority lacked comprehensive knowledge of acaricide resistance and effective disease prevention. Future research should focus on developing targeted educational programs to enhance farmers' knowledge and attitudes

towards sustainable tick control practices. Further, studies should explore other alternatives to chemical acaricides, to minimize acaricide resistance and TBDs in livestock.

**Acknowledgements.** The authors are grateful to the Hon'ble Vice-Chancellor, Director Research of Nanaji Deshmukh Veterinary Science University, Jabalpur, Dean and Department of Parasitology, College of Veterinary Science & Animal Husbandry (NDVSU), Mhow, Indore (Madhya Pradesh), for providing essential facilities required for the successful completion of the research work.

**Author's contributions.** SJ: Conceptualization, methodology, MS: formal analysis; investigation, resources, acquisition, original draft preparation. AKJ: Conceptualization, supervision, VA, MS, AKS: writing—review and editing. GNB: data analysis, GPJ and NJ: writing, review and editing. All authors have read and agreed to the published version of the manuscript.

**Financial support.** This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

**Competing interests.** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Ethical standards.** The present study was approved by the Institutional Ethical Committee of the College of Veterinary Science & Animal Husbandry (NDVSU), Mhow, Madhya Pradesh, India. All the activities involving individuals were performed following ethical regulations to

protect the rights and well-being of the participants (Lodhi *et al.*, 2016) and the data were gathered with consent of the participants.

Accepted Manuscript

## References

- Adehan, SB, Biguezoton, A, Adakal, H, Assogba, MN, Gbaguidi, AM, Tonouhewa, A, Kand, Ā, S, Achi, L, Kagone, H, Adehan, R and Mensah, GA** (2016) Acaricide resistance of *Rhipicephalus microplus* ticks in Benin. *African Journal of Agricultural Research* **11**(14), 1199-1208.
- Becker, S, Webster, A, Doyle, RL, Martins, JR, Reck, J and Klafke, GM** (2019) Resistance to deltamethrin, fipronil and ivermectin in the brown dog tick, *Rhipicephalus sanguineus sensu stricto*, Latreille (Acari: Ixodidae). *Ticks and Tick Borne Diseases* **10**(5), 1046-1050.
- Bisht, N, Kumar, S, Sharma, AK, Nandi, A, Singh, K, Fular, A, Nagar, G and Ghosh, S** (2021) Comparative susceptibility of *Rhipicephalus microplus* collected from the northern state of India to coumaphos, malathion, deltamethrin, ivermectin, and fipronil. *Tropical Animal Health and Production* **53**, 1-10.
- Butler, AD, Sedghi, T, Petrini, JR and Ahmadi, R** (2016) Tick-borne disease preventive practices and perceptions in an endemic area. *Ticks and Tick Borne Diseases* **7**(2), 331-337.
- Castro-Janer, E, Rifran, L, Piaggio, J, Gil, A, Miller, RJ, Schumaker, TTS** (2009) In vitro tests to establish LC50 and discriminating concentrations for fipronil against *Rhipicephalus (Boophilus) microplus* (Acari: ixodidae) and their standardization. *Veterinary Parasitology* **162**(1–2), 120–128.
- Catley, A, Admassu, B, Bekele, G and Abebe, D** (2014) Livestock mortality in pastoralist herds in Ethiopia and implications for drought response. *Disasters* **38**, 500–516.
- Chakraborty, S and Steckler, TL** (2023) Gronemeyer P, Mateus-Pinilla N, Smith RL. Farmers' knowledge and practices about ticks and tickborne diseases in Illinois. *Journal of Agromedicine* **28**(4), 756-68.

- Chigure, GM, Sharma, AK, Kumar, S, Fular, A, Sagar, SV, Nagar, G, Upadhaya, D, Saravanan, BC, Kumar, R and Ghosh, S** (2018) Role of metabolic enzymes in conferring resistance to synthetic pyrethroids, organophosphates, and phenylpyrazole compounds in *Rhipicephalus microplus*. *International Journal of Acarology* **44**(1), 28-34.
- de Castro, JJ, James, AD, Minjauw, B, Di Giulio, GU, Permin, A, Pegram, RG, Chizyuka, GB and Sinyangwe, P** (1997) Long-term studies on the economic impact of ticks on Sanga cattle in Zambia. *Experimental & Applied Acarology* **21**, 3-19.
- Estrada-Peña, A and Fernández-Ruiz, N** (2020) A retrospective assessment of temperature trends in Northern Europe reveals a deep impact on the life cycle of *Ixodes ricinus* (Acari: Ixodidae). *Pathogens* **9**(5), 345.
- FAO** (2004) Resistance management and integrated parasite control in ruminants - guidelines. Animal Production and Health Division, 25–77.
- Finney, DJ** (1971) Probit Analysis. Cambridge University Press, Cambridge, UK.
- Fular, A, Sharma, AK, Upadhaya, D, Nandi, A, Nagar, G, Bisht, N, Shakya, M, Kumar, S, Kumar, S, Kumar, R and Ghosh, S** (2021) Evaluation of acaricidal resistance status of *Rhipicephalus microplus* ticks from the hilly state (Uttarakhand) of India and evaluation of efficacy of a natural formulation for the management of resistant ticks. *Experimental and Applied Acarology* **85**, 355-377.
- Gaur, RS, Sangwan, AK, Sangwan, N and Kumar, S** (2016) Acaricide resistance in *Rhipicephalus (Boophilus) microplus* and *Hyalomma anatolicum* collected from Haryana and Rajasthan states of India. *Experimental and Applied Acarology* **69**, 487-500.
- Ghosh, S and Azhahianambi, P** (2007) Laboratory rearing of *Theileria annulata*-free *Hyalomma anatolicum anatolicum* ticks. *Experimental and Applied Acarology* **43**, 137-146.

- Ghosh, S, Azhahianambi, P and de la Fuente, J** (2006) Control of ticks of ruminants, with special emphasis on livestock farming system in India—present and future possibilities for integrated control: A review. *Experimental and Applied Acarology* **40**, 49–66.
- Ghosh, S, Tiwari, SS, Kumar, B, Srivastava, S, Sharma, AK, Kumar, S, Bandyopadhyay, A, Julliet, S, Kumar, R and Rawat, AKS** (2015) Identification of potential plant extracts for anti-tick activity against acaricide resistant cattle ticks, *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae). *Experimental and Applied Acarology* **66**, 159-171.
- Graham, OH and Hourrigan, JL** (1977) Eradication programs for the arthropod parasites of livestock. *Journal of Medical Entomology* **13**(6), 629-658.
- Grisi, L, Leite, RC, Martins, JRDS, Barros, ATMD, Andreotti, R, Cançado, PHD, León, AAPD, Pereira, JB and Villela, HS** (2014) Reassessment of the potential economic impact of cattle parasites in Brazil. *Revista Brasileira de Parasitologia Veterinária* **23**, 150-156.
- Guerrero, FD, Miller, RJ and de León, AAP** (2012) Cattle tick vaccines: many candidate antigens, but will a commercially viable product emerge? *International Journal of Parasitology* **42**(5), 421-427.
- Gumicio, S, Merica, M, Luhman, N, Fauvel, G, Zompi, S and Ronsse, A** (2011) The KAP survey model (Knowledge, Attitudes & Practices). *Medicins du Monde*, 1-73.
- Gupta, S, Gupta, S and Kumar, S** (2021) Cypermethrin resistance in *Hyalomma anatolicum* and *Rhipicephalus microplus* ticks of arid and semi-arid zone of Haryana, a northern state of India. *International Journal of Tropical Insect Science* **41**, 703-709.
- Hemme, T, Ndambi, A and Schröer-Merker, E** (2013) Overview on Milk Prices and Production Costs Worldwide; International Farm Comparison Network (IFCN): Milwaukee, MI, USA.

- Hurtado, OJB and Giraldo-Ríos, C** (2018) Economic and health impact of the ticks in production animals. *Ticks and Tick Borne Diseases* **9**, 1-9.
- Hussain, S, Hussain, A, Ho, J, Li, J, George, D, Rehman, A, Zeb, J and Sparagano, O** (2021) An epidemiological survey regarding ticks and tick-borne diseases among livestock owners in Punjab, Pakistan: A one health context. *Pathogens* **10**, 361.
- Jongejan, F and Uilenberg, G** (2004) The global importance of ticks. *Parasitology* **129**, S1–S12
- Jonsson, NN, Piper, EK and Constantinoiu, CC** (2014) Host resistance in cattle to infestation with the cattle tick *Rhipicephalus microplus*. *Parasite Immunology* **36**(11), 553-559.
- Jyothimol, G, Ravindran, R, Juliet, S, Ajithkumar, KG, Suresh, NN, Vimalkumar, MB, Lenka, DR, Varghese, S and Ghosh, S** (2014) Low level deltamethrin resistance in ticks from cattle of Kerala, a south Indian state. *Veterinary Parasitology* **204**(3-4), 433-438.
- Khating, S, Jadhav, N, Vijay, M, Sharma, AK, Srivastava, A, Jadhao, S, Kumar, S, Kalwaghe, S, Siddiqui, MF, Narawade, M, Dhabale, A and Chigure, G** (2024) Current profile of phenotypic pyrethroid resistance in *Rhipicephalus microplus* (Acari: Ixodidae) populations sampled from Marathwada region of Maharashtra state, India. *Parasitology Research* **123**. <https://doi.org/10.1007/s00436-024-08306-x>
- Kivaria, FM** (2006) Estimated direct economic costs associated with tick-borne diseases on cattle in Tanzania. *Tropical Animal Health and Production* **38**, 291-299.
- Kumar, S, Sharma, AK, Nagar, G, Rawat, SS, Tiwari, SS, Kumar, R, Dhakad, ML, Sharma, RK, Saxana, RK, Mehraniya, RS and Singh, RS** (2016) Characterization of acaricide resistance in tick isolates collected from Rajasthan, India. *Indian Journal of Animal Research* **86**(1), 14-23.



- Kumar, SS, Rayulu, VC, Rao, KS and Kumar, NV** (2017) Acaricidal resistance in *Rhipicephalus (Boophilus) microplus* ticks infesting cattle of Andhra Pradesh. *Journal of Entomology and Zoology Studies* **5**(6), 580-584.
- Launiala, A** (2009) How much can a KAP survey tell us about people's knowledge, attitudes and practices? Some observations from medical anthropology research on malaria in pregnancy in Malawi. *Anthropology Matters* **11**(1). DOI:[10.22582/AM.V11I1.31](https://doi.org/10.22582/AM.V11I1.31)
- Lodhi, LA, Muhammad, G, Iqbal, Z, Nazir, A, Ali, M, Adil, SA, Shamim, R and Sajid, MS** (2016) *Animal Care Handbook*; University Press, University of Agriculture: Faisalabad, Pakistan.
- Lontsi-Demano, M, Laroche, M, Ngnindji, YC, Djikolmbairangar, JE, Mamoudou, A and Tchuinkam, T** (2021) Breeder's Knowledge on Ticks and Tick-Borne Diseases and Management Strategies in Menoua Division (Western Region of Cameroon). *International Journal of Veterinary Sciences and Animal Husbandry* **6**, 12-21.
- Minjauw, B and McLeod, A** (2003) Tick-borne diseases and poverty: the impact of ticks and tick-borne diseases on the livelihoods of small-scale and marginal livestock owners in India and eastern and southern Africa. Research Report, DFID-AHP. UK: Centre for Tropical Veterinary Medicine, University of Edinburgh 2003; p. 116.
- Mugambi, JM, Wesonga, FD and Ndungu, SG** (2012) Ticks and tick-borne disease control in a pastoral and an agro-pastoral farming systems in Kenya. *Livestock Research for Rural Development* **24**(5), 1-8.
- Namgyal, J, Lysyk, TJ, Couloigner, I, Checkley, S, Gurung, RB, Tenzin, T, Dorjee, S and Cork, SC** (2021a) Identification, distribution, and habitat suitability models of Ixodid tick species in cattle in Eastern Bhutan. *Tropical Medicine and Infectious Disease* **6**(1), 27.


- Namgyal, J, Tenzin, T, Checkley, S, Lysyk, TJ, Rinchen, S, Gurung, RB, Dorjee, S, Couloigner, I and Cork, SC (2021b)** A knowledge, attitudes, and practices study on ticks and tick-borne diseases in cattle among farmers in a selected area of eastern Bhutan. *PloS one*. **16**(2), 0247302.
- Ngnindji-Youdje Y, Diarra AZ, Lontsi-Demano M, Tchuinkam T and Parola P (2022)** Detection of Tick-Borne Pathogens in Ticks from Cattle in Western Highlands of Cameroon. *Microorganisms* **10**, 1957.
- Niesobecki, S, Hansen, A, Rutz, H, Mehta, S, Feldman, K, Meek, J, Niccolai, L, Hook, S and Hinckley, A (2019)** Knowledge, attitudes, and behaviors regarding tick-borne disease prevention in endemic areas. *Ticks and Tick Borne Diseases* **10**(6), 101264.
- Phanchung, Dorji, P, Sonam, T and Pelden, K (2007)** Smallholder Dairy Farming in Bhutan: Characteristics, Constraints, and Development Opportunities. Pp 19-35
- Queenan, K, Mangesho, P, Ole-Neselle, M, Karimuribo, E, Rweyemamu, M, Kock, R and Häslér, B (2017)** Using local language syndromic terminology in participatory epidemiology: Lessons for One Health practitioners among the Maasai of Ngorongoro, Tanzania. *Preventive Veterinary Medicine* **139**, 42–49.
- Rehman, A, Conraths, FJ, Sauter-Louis, C, Krücken, J and Nijhof, AM (2019)** Epidemiology of tick-borne pathogens in the semi-arid and the arid agro-ecological zones of Punjab province, Pakistan. *Transboundary and Emerging Diseases* **66**, 526–536.
- Rodríguez-Vivas, RI, Grisi, L, Pérez de Leon, AA, Villela, HS, Torres-Acosta, JF, Fragoso Sanchez, H, Romero Salas, D, Rosario Cruz, R, Saldierna, F and García Carrasco, D (2017)** Potential economic impact assessment for cattle parasites in Mexico. Review. *Revista Mexicana de Ciencias Pecuarias* **8**, 61–74.

- Rosario-Cruz, R, Almazan, C, Miller, R, Dominguez-Garcia, DI, Hernandez-Ortiz, R and de la Fuente, J** (2009) Genetic basis and impact of tick acaricide resistance. *Frontiers in Bioscience* **14**, 2657-2665
- Senbill, H, Hazarika, LK and Baruah, A** (2018) Life cycle of the southern cattle tick, *Rhipicephalus (Boophilus) microplus* Canestrini 1888 (Acari: ixodidae) under laboratory conditions. *Systematic and Applied Acarology* **23**(6), 1169-1179.
- Shakya, M, Kumar, S, Fular, A, Upadhaya, D, Sharma, AK, Bisht, N, Nandi, A and Ghosh, S** (2020) Emergence of fipronil resistant *Rhipicephalus microplus* populations in Indian states. *Experimental and Applied Acarology* **80**, 591-602.
- Shakya, M, Sharma, AK, Kumar, S, Upadhaya, D, Nagar, G, Singh, K, Sankar, M and Ghosh, S** (2023) Acaricides resistance in *Rhipicephalus microplus* and expression profile of ABC-transporter genes in the sampled populations. *Experimental Parasitology* **252**, 108584.
- Shaw, RD** (1966) Culture of an organophosphorus-resistant strain of *Boophilus microplus* (Can.) and an assessment of its resistance spectrum. *Bulletin of Entomological Research* **56**(3), 389-405.
- Shyma, KP, Gupta, JP and Singh, V** (2015) Breeding strategies for tick resistance in tropical cattle: a sustainable approach for tick control. *Journal of Parasitic Diseases* **39**, 1-6.
- Singh, K, Kumar, S, Sharma, AK, Jacob, SS, Verma, MR, Singh, NK, Shakya, M, Sankar, M and Ghosh, S** (2022) Economic impact of predominant ticks and tick-borne diseases on Indian dairy production systems. *Experimental Parasitology* **243**. 108408. <https://doi.org/10.1016/j.exppara.2022.108408>
- Soulsby, E JL** (1982) Helminths, Arthropods and Protozoa of Domesticated animals, 7th Ed. Bailliere Tindall, London, pp. 486–487.

- Stone, BF and Haydock, KP** (1962) A method for measuring the acaricide-susceptibility of the cattle tick *Boophilus microplus* (Can.). *Bulletin of Entomological Research* **53**(3), 563-578.
- Süss, J** (2008) Tick-borne encephalitis in Europe and beyond—the epidemiological situation as of 2007. *Euro surveillance* **13** (26). 18916. PMID: 18761916.
- Tesfaye, T and Abate, A** (2023) Knowledge, attitude and practices study of acaricide usage and tick control in South Omo Zone pastoral areas, South-Western Ethiopia. *Heliyon* **9**(6). E17212. <https://doi.org/10.1016/j.heliyon.2023.e17212>
- Thrusfield, M** (2018) *Veterinary Epidemiology*; John Wiley & Sons: Hoboken, NJ, USA.
- Upadhaya, D, Kumar, B, Kumar, S, Sharma, AK, Fular, A, Bisht, N, Srivastava, S, Boruah, RR, Nagar, G, Shakya, M and Nath, T** (2020) Characterization of acaricide resistance in *Rhipicephalus microplus* populations infesting cattle in northeastern India and assessment of local plant extracts for tick management. *Veterinary Parasitology* **277**, 109011. doi: 10.1016/j.vetpar.2019.109011.
- Wickham, H, François, R, Henry, L, Müller, K** (2021) dplyr: A Grammar of Data Manipulation. R package version 1.0. 4 h ps. CRAN. [R-project.org/package= dplyr](https://cran.r-project.org/package=dplyr) [Acesso 09/01/2021].
- Williams, A** (2003) How to write and analyse a questionnaire. *Journal of Orthodontics* **30**, 245–252.
- Yessinou, RE, Akpo, Y, Sidick, A, Adoligbe, C, Karim, IYA, Akogbeto, M and Farougou, S** (2018) Evidence of multiple mechanisms of alphacypermethrin and deltamethrin resistance in ticks *Rhipicephalus microplus* in Benin, West Africa. *Ticks and Tick. Borne Diseases* **9**(3), 665-671.


**Zöldi, V, Turunen, T, Lyytikäinen, O and Sane, J** (2017) Knowledge, attitudes, and practices regarding ticks and tick-borne diseases, Finland. *Ticks and Tick Borne Diseases* **8**(6), 872-877.

Accepted Manuscript




**Figure 1.** Variable Importance model for each contributing factor in tick infestation.

Accepted



**Figure 2.** Regression curves showing probit mortality in larval packet test (LPT, A & B) and larval immersion test (LIT, C & D) against log concentration of chemical acaricides (deltamethrin and fipronil) in five field isolates of *Rhipicephalus microplus* from Dhar district, Madhya Pradesh, India: *DHR* Dhar, *GAN* Gandhwani, *KUK* Kukshi, *MAN* Manawar, *SAR* Sardarpur



**Figure 3.** Regression curves showing probit mortality in larval packet test (LPT, A & B) and larval immersion test (LIT, C & D) against log concentration of chemical acaricides (deltamethrin and fipronil) in five field isolates of *Hyalomma anatolicum* from Dhar district, Madhya Pradesh, India: *DHR* Dhar, *GAN* Gandhwani, *KUK* Kukshi, *MAN* Manawar, *SAR* Sardarpur



**Table 1.** Questionnaire data collected from surveyed places for determination of pattern of acaricidal application in fields

Blocks	Farm type	No. of farms visited	Commonly used acaricides	Mode of application	Frequency of application <sup>a</sup>	Shed treatment
DHR	Unorganised, Organised	40	Deltamethrin, Cypermethrin, Amitraz, Ivermectin	Pour on/ Injection/ Swabs	Occasional, Frequent	Rarely
GAN	Unorganised, Organised	40	Deltamethrin, Cypermethrin, Ivermectin	Pour on/ Injection	Occasional	Rarely
KUK	Unorganised	40	Deltamethrin, Cypermethrin, Ivermectin	Pour on/ Injection/ swabs	Occasional, Frequent	Rarely
MAN	Unorganised, Organised	40	Deltamethrin, Cypermethrin, Ivermectin	Pour on/ Injection/ Swabs	Frequent	Rarely
SAR	Unorganised, Organised	40	Deltamethrin, Cypermethrin, Amitraz, Ivermectin	Pour on/ Injection/ Swabs	Frequent	Rarely

<sup>a</sup>Application frequency: frequent = 10–14 applications/tick active season; occasional = 4-8 application/tick active season

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

**Table 2.** Socio-demographic characteristics of the respondents

<b>Variables</b>	<b>Categories</b>	<b>Frequencies</b>	<b>Percentage (95% CL)</b>
Gender	Male	126	63.00(55.91-69.70)
	Female	74	37.00 (33.30-44.09)
Education	Literate	80	40.00 (33.15-47.15)
	Illiterate	120	60.00 (52.85-66.85)
Type of housing	Kachha floor	110	55.00 (47.82-62.02)
	Pakka floor	90	45.00 (37.98-52.18)
Feeding system	Grazing	10	05.00 (2.42-9.00)
	Manger	75	37.50 (31.25-45.11)
	Mixed	115	57.50 (49.83-63.96)
Knowledge about TTBDs	Yes	50	25.00 (19.16-31.60)
	No	150	75.00 (68.40-80.84)
Attitude towards tick control	Favorable	73	36.50 (29.82-43.58)
	Unfavorable	127	63.50 (56.42-70.18)
Commonly used acaricides	Cypermethrin	70	35.00 (28.41-42.05)
	Deltamethrin	58	29.00 (22.82-35.82)
	Ivermectin	30	15.00 (10.35-20.72)
	Amitraz	18	09.00 (5.42-13.85)
	Mixed	24	12.00 (7.84-17.33)

**Table 3.** Association between socio-demographic characteristics of the respondents (n=200) and tick infestation

Variables	Categories	Respondents	Tick infestation level		*P-value
			High	Low	
Education	Literate	80	15	65	0.0180
	Illiterate	120	45	75	
Shed floor type	Kachha	110	47	63	0.0029
	Pakka	90	18	72	
Feeding system	Grazing	10	6	4	0.0335
	Manger	75	13	62	
	Mixed	115	43	72	
Knowledge about TTBDs	Yes	50	15	35	0.0063
	No	150	84	66	
Commonly used acaricides	Cypermethrin	70	29	41	-
	Deltamethrin	58	23	35	
	Ivermectin	30	11	19	
	Amitraz	18	7	11	
	Mixed	24	05	19	

\*Significant at  $p < 0.05$  \*\*Significant at  $p < 0.01$

**Table 4.** Simple logistic regression analysis for the estimation of the association between socio-demographic variables of respondents and binary outcome.

Variable	Category		Estimated±S.E	Adjusted OR (95% CI)	p-Values
Sex	Female	Male	0.7909±0.33	2.20 (1.13-4.27)	0.0191
	74	126			
Education	Literate	Illiterate	-1.074±0.56	0.34 (0.11-1.04)	0.0592
	80	120			
Type of animal shed floor	Kachha	Pakka	-1.267±0.36	0.28 (0.13-0.57)	0.0005
	110	90			
Knowledge about TTBDs	Yes	No	0.2527±0.56	1.28 (0.42-3.86)	0.6525
	50	150			
Attitude towards tick control	Favorable	Unfavorable	0.0459±0.47	1.04 (0.4-2.66)	0.9231
	73	127			
Constant	-	-	0.0289±0.26	-	0.9134

Significant at  $p < 0.05$ , OR: odd ratio

**Table 5.** Multiple logistic regression analysis for the estimation of the association between practices of respondents with level of tick infestation.

Variable	Category	Respondents	Estimated±S.E	Adjusted OR (95% CI)	p-Values
Feeding	Grazing	10	1.7931±0.36	6.00(2.93-12.28)	0.0000
	Manger	75			
	Mixed	115			
Acaricides Used	Cypermethrin	70	0.0194±0.13	1.01(0.78-1.32)	0.8845
	Deltamethrin	58			
	Ivermectin	30			
	Amitraz	18			
	Mixed	24			
Constant	-	-	-3.3805±0.72	-	0.0000

Significant at  $p < 0.05$

**Table 6.** Association between socio-demographic variables of respondents and tick infestation level by R software analysis with different R packages.

Coefficient	Estimate	Std. Error	Z value	p-Value	Odds Ratio
Intercept	2.33544	0.83395	2.800	0.005103 **	10.33
Sex (Female vs. Male)	0.39980	0.44346	0.902	0.367293	1.491
Literacy (Illiterate vs. Literate)	1.25456	0.74840	-1.676	0.093676	3.51
Floor (Kachha vs. Pakka)	1.64023	0.49015	3.346	0.000819 ***	5.16
Feeding system (Grazing vs. Manger/Mixed)	1.41199	0.44494	3.173	0.001506 **	4.10
Knowledge about TTBDs (Yes vs. No)	-0.04726	0.81512	0.058	0.953761	0.95
Acaricides Used (Cypermethrin, Deltamethrin, Ivermectin, Amitraz & Mixed)	0.57355	0.25822	2.221	0.026340 *	1.77
Attitude towards tick control (Unfavorable vs. Favorable)	0.17804	0.58454	0.305	0.760681	1.20

The odds ratio is the “Exponential” of the estimate obtained in glm model (log regression), Significant at \*\*\* $p < 0.001$ , \*\* $p < 0.01$ , \* $p < 0.05$

**Table 7.** Mortality slope, R<sup>2</sup>, LC<sub>50</sub> with 95%CI and RR50 values of deltamethrin and fipronil against larvae of *R. microplus* by using LPT and LIT

Acaricide	LPT						LIT					
	Tick isolates	Mortality (slope ± SE)	R <sup>2</sup>	LC50 (95% CI)	RR50	RL	Mortality (slope ± SE)	R <sup>2</sup>	LC50 (95% CI)	RR50	RL	
Deltamethrin	DHR	2.769 ± 0.39	0.94	441.43 (405.06 - 481.06)	37.4	III	2.455 ± 0.56	0.86	21.20(19.23 - 23.36)	1.7	I	
	GAN	3.117 ± 0.35	0.96	471.6 (436.93 - 509.01)	39.9	III	1.499 ± 0.24	0.92	51.42 (43.87 - 60.26)	4.3	I	
	KUK	3.561 ± 0.22	0.98	400.69 (374.77 - 428.39)	33.9	III	1.740 ± 0.25	0.94	20.70 (18.05 - 23.73)	1.7	I	
	MAN	2.904 ± 0.33	0.96	435.85 (401.5 - 473.08)	36.9	III	1.487 ± 0.23	0.92	15.02 (14.57 - 15.48)	1.2	S	
	SAR	2.685 ± 0.40	0.93	428.57 (392.21 - 468.29)	36.3	III	2.528 ± 0.61	0.84	22.80 (20.74 - 25.05)	1.9	I	
	IVRI-I	3.42 ± 0.49	0.87	11.8 (11.6 -12.0)	1.0	S	3.42 ± 0.49	0.87	11.80 (11.6 -12.0)	1.0	S	
Fipronil	DHR	1.847 ± 0.19	0.96	0.56 (0.49 - 0.63)	0.23	S	2.804 ± 0.22	0.98	1.21 (1.11 - 1.31)	0.50	S	
	GAN	1.222 ± 0.10	0.97	0.47 (0.38 - 0.57)	0.19	S	2.873 ± 0.20	0.98	1.24(1.14 - 1.34)	0.51	S	
	KUK	1.261 ± 0.03	0.99	0.49 (0.40 - 0.59)	0.20	S	2.852 ± 0.24	0.98	1.22 (1.12 - 1.32)	0.50	S	
	MAN	1.176 ± 0.05	0.99	0.42 (0.34 - 0.51)	0.17	S	2.865 ± 0.35	0.96	1.18 (1.08 - 1.28)	0.49	S	
	SAR	1.391 ± 0.11	0.97	0.59 (0.49 - 0.69)	0.24	S	2.760 ± 0.27	0.97	1.16 (1.06 - 1.26)	0.48	S	
	IVRI-I	7.67 ± 2.4	0.84	2.4 ( 2.38 - 2.42)	1.00	S	7.67 ± 2.4	0.84	2.4 ( 2.38 - 2.42)	1.00	S	

DHA Dhar, GAN Gandhwani, KUK Kukshi, MAN Manawar, SAR Sardarpur, IVRI-I reference susceptible tick strain, RR50 (median) resistance ratio, RL resistance level (susceptible [S]=RR < 1.4; level I: 1.5 < RR < 5; level II: 5.1 < RR < 25; level III: 26 < RR < 40; level IV: RR > 41.

**Table 8.** Mortality slope, R<sup>2</sup>, LC<sub>50</sub> with 95%CI and RR50 values of deltamethrin and fipronil against larvae of *H. anatolicum* by using LPT and LIT

Acaricide	LPT						LIT					
	Tick isolates	Mortality (slope ± SE)	R <sup>2</sup>	LC50 (95% CI)	RR50	RL	Mortality (Slope ± SE)	R <sup>2</sup>	LC50 (95% CI)	RR50	RL	
Deltamethrin	DHR	5.002 ± 1.23	0.84	132.17 (126.02 - 138.61)	11.1	II	3.048 ± 0.44	0.95	20.63 (19.07 - 22.31)	1.7	I	
	GAN	2.915 ± 0.33	0.96	182.88 (168.53 - 198.44)	15.4	II	2.928 ± 1.09	0.78	26.46 (24.39 - 28.70)	2.2	I	
	KUK	3.448 ± 0.30	0.97	194.90 (181.89 - 208.83)	16.5	II	2.616 ± 0.63	0.89	18.53 (16.91 - 20.29)	1.5	I	
	MAN	3.379 ± 0.21	0.98	180.91 (168.60 - 194.11)	15.3	II	3.436 ± 0.74	0.91	28.04 (26.16 - 30.05)	2.3	I	
	SAR	5.374 ± 1.19	0.87	145.61 (139.30 - 152.20)	12.3	II	3.257 ± 0.86	0.87	27.54 (25.59 - 29.62)	2.3	I	
	IVRI-I	3.42 ± 0.49	0.87	11.8 (11.6 - 12.0)	1.0	S	3.42 ± 0.49	0.87	11.8 (11.6 - 12.0)	1.0	S	
Fipronil	DHR	3.125 ± 0.35	0.96	1.41 (1.30 - 1.52)	0.58	S	3.144 ± 0.24	0.98	1.07 (0.99 - 1.15)	0.44	S	
	GAN	3.320 ± 0.49	0.93	1.58 (1.47 - 1.69)	0.65	S	3.240 ± 0.26	0.98	1.15 (1.06 - 1.23)	0.47	S	
	KUK	2.948 ± 0.38	0.95	1.97 (1.81 - 2.13)	0.82	S	3.239 ± 0.19	0.99	1.11 (1.03 - 1.19)	0.46	S	
	MAN	3.410 ± 0.31	0.97	1.88 (1.75 - 2.01)	0.78	S	3.201 ± 0.26	0.98	1.09 (1.01 - 1.17)	0.45	S	
	SAR	3.144 ± 0.20	0.98	1.85 (1.71 - 1.99)	0.77	S	3.142 ± 0.29	0.98	1.11 (1.02 - 1.19)	0.46	S	
	IVRI-I	7.67 ± 2.4	0.84	2.4 (2.38 - 2.42)	1.00	S	7.67 ± 2.4	0.84	2.4 (2.38 - 2.42)	1.00	S	

DHA Dhar, GAN Gandhwani, KUK Kukshi, MAN Manawar, SAR Sardarpur, IVRI-I reference susceptible tick strain, RR50 (median) resistance ratio, RL resistance level (susceptible [S]= RR < 1.4; level I: 1.5 < RR < 5; level II: 5.1 < RR < 25; level III: 26 < RR < 40; level IV: RR > 41.