

THE JOURNAL OF AGRICULTURAL SCIENCE

EDITED BY

G. D. H. BELL, C.B.E., PH.D., F.R.S., Plant Breeding Institute, Cambridge

K. L. BLAXTER, PH.D., N.D.A. (HONS.), D.SC., F.R.S.E.,
Rowett Research Institute, Bucksburn, Aberdeen

G. W. COOKE, PH.D., F.R.I.C., Rothamsted Experimental Station, Harpenden

JOHN HAMMOND, Jr., M.A., School of Agriculture, Cambridge

PROF. SIR J. B. HUTCHINSON, C.M.G., SC.D., F.R.S., School of Agriculture,
Cambridge

H. L. PENMAN, O.B.E., PH.D., F.R.S., Rothamsted Experimental Station, Harpenden

H. H. ROGERS, B.SC., DIP.AG.SCI., Plant Breeding Institute, Cambridge

PROF. E. W. RUSSELL, C.M.G., PH.D., F.INST.P., Department of Soil Science,
University of Reading

F. YATES, C.B.E., SC.D., F.R.S., Rothamsted Experimental Station, Harpenden



CAMBRIDGE UNIVERSITY PRESS

BENTLEY HOUSE, 200 EUSTON ROAD, LONDON, N.W. 1

AMERICAN BRANCH: 32 EAST 57TH STREET, NEW YORK, N.Y. 10022

Price 32s. 6d. net

(U.S.A. \$5.50)

INSTRUCTIONS TO AUTHORS

Papers intended for publication should be submitted to Dr G. D. H. Bell, C.B.E., F.R.S., Plant Breeding Institute, Cambridge, or to one of the Associate Editors. It must be understood that, if accepted by the Editorial Board, the paper will not be published elsewhere in the same form.

MANUSCRIPTS. Papers, written in English, are accepted from any country and should be typed in double-spacing on one side of the paper with a margin at least 4 cm wide on the left-hand side. Authors should instruct typists on the style of manuscript required. A top copy and one carbon copy should be submitted.

TABLES must be self-explanatory. They should be typed on separate sheets, numbered consecutively and carry an appropriate title. Wherever possible, tables should be arranged so that they can be printed in the normal orientation of the text and without the need for rules.

LINE DRAWINGS (with photocopies) should be 25 cm. wide and drawn in black waterproof ink on Bristol board, graph paper with blue lines or tracing paper. Legends should be typed on a separate copy and numbering inserted lightly and clearly in soft pencil on the drawing. Tables and figures should not reproduce the same data.

The approximate position of tables and figures should be noted in the text.

PLATES should make a definite contribution to the value of the paper and the number submitted should be kept to a minimum. They should be good quality, unmounted, glossy prints and be lightly numbered in pencil on the reverse side. If several, or coloured, plates are submitted the author may be asked to contribute to the cost of reproducing them.

TITLE. The title must be specific and suitable for indexing by the mechanical methods now being employed. The full name and address of the institution in which the research has been carried out should be stated. Change of address may be given as a footnote. A short title, not exceeding 50 characters, should be provided for the running headlines.

STYLE. Experimental details and results should be recorded in the past tense and there should be no unnecessary repetition or loose phrases. Manuscripts are likely to be returned for modification if the presentation is not clear and precise.

LAYOUT. Authors are recommended to study '*General Notes on the Preparation of Scientific Papers*'

(Royal Society, London, 2nd edn., 1965). The Editorial Board do not insist upon a rigid format but it is usually convenient to divide the paper into sections, e.g. Introduction, Materials and Methods, Results and Discussion. An excess of headings and sub-headings should be avoided. A Summary should always be included.

Authors are advised to note the following points: a detailed review of literature is not necessary; relevant details should be given of the plant or animal material, the experimental design and chemical or other techniques employed; mean results with their relevant standard errors should be presented rather than detailed data; the statistical methods used should be clearly stated; the discussion should relate the author's experiments to other work on the subject and give the author's conclusions; the summary should be factual and suitable for use in abstracting journals. Footnotes should be avoided.

REFERENCES. The bibliography must be given in the form—Surname of authors, initials, year of publication (in parentheses), title of paper, name of journal (abbreviated according to the *World List of Scientific Periodicals*, 4th edn, Butterworths, London), volume and pages of reference. References should be in alphabetical order. In the text a reference should be quoted by the author's name and date (in parentheses). Where there are more than two authors, the initial reference in the text should include the names of all authors but subsequent citations, should be in the form—first author followed by *et al.* Authors should check that all references in the text appear at the end of the paper and vice versa, and that names and dates correspond in the two places.

PROOF CORRECTION. Standard proof correction marks (British Standard 1219) should be made as legibly as possible in ink, not pencil. Directions to the printer which are not to be set up in type should be encircled. Captions to illustrations and all references should be checked. Queries marked by the printer should be answered. Proofs are provided in order that authors can check the correctness of the type-setting—excessive alterations may be charged to the author.

OFFPRINTS. Contributors will receive 25 copies of their papers free and can order others when they receive the proofs.

Failure to comply with the 'Instructions to Authors' may delay publication.

JOURNAL OF REPRODUCTION AND FERTILITY

The Journal publishes original papers, abstracts of Proceedings, and occasionally reviews and bibliographies, dealing with the morphology, physiology, biochemistry and pathology of reproduction in man and animals, and with the biological, medical and veterinary problems of fertility and lactation. Numbers are issued bimonthly to give two volumes a year.

Papers to appear in 1966 include:

- R. G. WALES and T. O'SHEA. Effect of low levels of carbon dioxide on the metabolism of ram and bull spermatozoa.
- R. G. ZIMBELMAN and L. W. SMITH. Control of ovulation in cattle with melen-gestrol acetate. 1. Effects of dosage and route of administration. 2. Effects of follicular size and activity.
- L. E. A. ROWSON and R. M. MOOR. Embryo transfer in the sheep: the signifi-cance of synchronising oestrus in the donor and recipient animal.
- H. PIGOŃ, M. MARCHUT and N. NESTOROV. Urinary oestrogens in the mature male mule.
- H. PETERS and E. LEVY. Cell dynamics of the ovarian cycle.
- C. DESJARDINS, K. T. KIRTON and H. D. HAFS. Some chemical, immunochemical and electrophoretic properties of bovine follicular fluid.
- H. P. VAN KREY, F. C. OGASAWARA and F. W. LORENZ. Distribution of sperma-tozoa in the oviduct and fertility in domestic birds. IV. Fertility of sperma-tozoa from infundibular and uterovaginal glands.
- T. O'SHEA and R. G. WALES. The aerobic metabolism of ram spermatozoa.
- G. H. KIRACOFÉ and H. G. SPIES. Length of maintenance of naturally formed and experimentally induced corpora lutea in hysterectomized ewes.
- C. V. HULET. Effects of various treatments with 6-methyl-17-acetoxypogesterone (MAP) on subsequent fertility in ewes.
- P. HEALEY and R. M. F. S. SADLEIR. The construction of rectal electrodes for electro-ejaculation.
- R. M. MOOR and L. E. A. ROWSON. Local uterine mechanisms affecting luteal function in the sheep.
- L. E. A. ROWSON and R. M. MOOR. Non-surgical transfer of cow eggs.

Editorial correspondence should be sent to Professor A. S. Parkes, Journal of Reproduction and Fertility, 8 Jesus Lane, Cambridge, England, and business correspondence to the publishers.

*The subscription rate is £6 per volume or £12 per year (two volumes).
U.S.A. and Canada \$20.00 per volume or \$40.00 per year*

**BLACKWELL SCIENTIFIC PUBLICATIONS
OXFORD**

ANIMAL PRODUCTION

JOURNAL OF THE BRITISH SOCIETY OF ANIMAL PRODUCTION

Contents of Volume 8, Part 3, October 1966 include

- JONES, MADDEVER, COURT and PHILLIPS. The time taken by cows to eat concentrates.
- BICHARD and COOPER. Analysis of production records from a lowland sheep flock: I.
- MOAV. Specialized sire and dam lines. III: Choice of the most profitable parental combination.
- MOAV and HILL. Specialized sire and dam lines. IV. Selection within lines.
- BROADBENT and WATSON. Factors governing the shape of the growth curve of Suffolk \times Welsh lambs.
- WIESER, PRESTON and MACDEARMID. Intensive beef production. 8. The effect of chlortetracycline on barley beef cattle.
- HINKS. Selection practices in dairy herds. I and II.
- DOORNENBAL and MARTIN. The association between commercial cut-out yield and gross chemical composition in market pigs.
- LODGE, ELSLEY and MACPHERSON. The effects of level of feeding of sows during pregnancy. II.
- ELSLEY, ANDERSON, McDONALD, MACPHERSON and SMART. A comparison of the live-weight changes, nitrogen retention and carcass composition of pregnant and non-pregnant gilts.
- SLEE. Variation in the responses of shorn sheep to cold exposure.
- HUNTER. The effect of superovulation on fertilization and embryonic survival in the pig.

The Journal is published three times a year, but **COMMENCING 1967 WILL BE PUBLISHED QUARTERLY IN FEBRUARY, MAY, AUGUST and NOVEMBER.**

The 1966 annual subscription rate is 65s. (U.S.A. \$11.00), single copy 25s. (U.S.A. \$4.00). **COMMENCING 1967 THE SUBSCRIPTION RATE WILL BE 90s. (U.S.A. \$14.00).** Orders and subscriptions may be sent to any bookseller or direct to:

OLIVER & BOYD LTD.

Tweeddale Court, 14 High Street, Edinburgh, 1

FULL PARTICULARS OF THE

JOURNALS

PUBLISHED BY THE

CAMBRIDGE
UNIVERSITY PRESS

MAY BE HAD FROM


THE MANAGER
CAMBRIDGE UNIVERSITY PRESS
BENTLEY HOUSE
200 EUSTON ROAD
LONDON, N.W.1

AGRONOMY JOURNAL

This official organ of the American Society of Agronomy is a bimonthly publication of up-to-date reports of general agronomic research. Workers in the fields of forages and pastures, crop improvement, cultural practices, soil fertility, and allied areas of investigation will find articles of lasting interest in *Agronomy Journal*. Publication is open to members of the American Society of Agronomy.

\$14.00 per year in U.S. and Canada.
\$15.00 per year elsewhere

American Society of Agronomy
677 S. Segoe Rd.
Madison, Wisconsin, U.S.A., 53711



**NEW
FROM
SHELL**

**END APHIDS, MITES, THRIPS
ON VEGETABLES, COTTON, FRUIT WITH
SYSTEMIC, PERSISTENT, ACTIVE BIDRIN**

Bidrin is Shell's new organo-phosphorus insecticide. It is systemic and has particular advantages for many crops. For example, Bidrin's extremely wide spectrum of activity includes the easy control of aphids, mites and thrips on vegetables, citrus fruits, cotton and seed-crops. Bidrin is equally useful in dealing with stem borers and leaf hoppers on rice – scale insects on citrus crops – and even the elm bark beetle. Bidrin kills on contact then continues to control biting and sucking insects for up to two to three weeks. Bidrin may be used to within four weeks before harvest, by then it has completely broken down and crops are ready for consumption. Current trials are proving that Bidrin may have many other valuable uses.

For further details about Bidrin contact your Shell company.



EXPERIMENTAL AGRICULTURE

Contents of Volume 2, No. 3, July 1966

- A. R. Sen, A. R. Sarkar and R. P. Chakrabarty.** Sample Surveys of Pests and Diseases of Tea in North-East India.
- D. P. Gibbon.** Experiments with Maize in Trinidad.
- N. Okafor.** Microbial Rotting of Stered Yams (*Dioscorea* spp) in Nigeria.
- G. D. Anderson, B. G. Houston and P. J. Northwood.** Effects of Soil Cultivation History and Weather on Responses of Wheat to Fertilizers in Northern Tanzania.
- D. Jowett.** Use of Rank Correlation Methods to Determine Food Preferences.
- E. S. Clayton.** Book Review. Agrarian Development in Peasant Economics.
- Anuwar Bin Mahmud and C. Devendra.** Repeatability of Milk Yield and Birth Weights of Goats in Malaya. 1. Milk Yield and Length of Lactation.
- P. Gillard.** Responses to Grazing Intensity on the Transvaal Highveld.
- H. K. Pande and V. M. Bhan.** Effect of Depth of Tillage on Yield of Upland Paddy and on Associated Weeds.
- J. E. Miltimore.** Factors Influencing the Survival of Propagules of Tall Fescue (*Festuca arundinacea* Schreb.) I. Effects of Daylength, Length and Weight of Propagules and Number of Culms.
- P. P. Courtenay.** Book Review. Plantation Agriculture.
- J. E. Miltimore.** Factors Influencing the Survival of Propagules of Tall Fescue (*Festuca arundinacea* Schreb.) II. Relations of Leaf, Root and Planting Weight to Survival and Hereditability of Survival.

Experimental Agriculture is a quarterly journal published by Cambridge University Press. It is the successor to *The Empire Journal of Experimental Agriculture*.

The subscription price is £3. 15s. (U.S.A. \$13.50) including postage; single parts are available at £1. 5s. each (U.S.A. \$4.50). Orders may be sent to any bookseller or subscription agent, or direct to Cambridge University Press.

CAMBRIDGE UNIVERSITY PRESS

Bentley House, 200 Euston Road, London, N.W.1

American Branch: 32 East 57th Street, New York, N.Y.10022, U.S.A.

A New Book

Just Ready

Recent Advances

36s.
(Paperback)

in

ANIMAL NUTRITION

Edited by **J. T. ABRAMS, Ph.D., M.A., M.Sc.,**
School of Veterinary Medicine, University of Cambridge

This inexpensive volume gives a concise account of a wide survey of animal feeding problems, covering nutritional techniques and foodstuffs. The contributors have been very carefully selected so that the work will be of value to students, lecturers and those engaged in research.

Emphasis is given to the influence of nutrition on the health of livestock on which man depends so much. The book provides a good cross-section of what is important in present-day nutrition.

Of interest to veterinary surgeons, research workers in all fields of biology, biochemists in the animal food industry and pedigree breeders.

Sm. Demy 8vo.

272 pp.

28 Illustrations.

J. & A. CHURCHILL LTD., 104 Gloucester Place, London, W.1

MOISTURE & DENSITY MEASURING EQUIPMENT NE 8401

SURFACE AND DEPTH MEASUREMENTS

For soil moisture and companion studies, evaporation measurements and irrigation problems.

With the Nuclear Enterprises NE 8401 Moisture and Density Measuring Equipment **rapid 'on site'** determinations of the moisture content and density in many bulk materials and on many surfaces can be made accurately with **reproducibility better than 1%**. Relatively inexperienced operators can use this easily transportable equipment. It is **fully transistorised** and consists of two depth probes, a surface unit, portable scaler, battery and shields. **Depth** probes and the **surface** unit may be purchased separately.

In addition to being in use in road, coal and agricultural research establishments and in industry in Britain, NE8401



Taking moisture measurements with NE 8401

Equipments are operating in Australia, Ghana, Borneo, Holland, Rumania, Sweden, Austria, Finland, Germany, India, France, Canada, Spain, Italy, Pakistan, Philippines and Turkey.

For full details of operation and prices, write to:

NUCLEAR ENTERPRISES (G.B.) LIMITED
SIGHTHILL, EDINBURGH 11, SCOTLAND, Tel: CRAiglockhart 4060,
Cable: NUCLEAR, EDINBURGH

Canadian Associate: Nuclear Enterprises Ltd., 550 Berry Street, Winnipeg, 21

U.S. Representative: Nuclear Enterprises Division, h-nu systems inc., 470 San Antonio Road, Palo Alto, California





ALDRIN PROTECTS CROPS FROM THE GROUND UP!

The root system is the vital link between the plant and its food in the soil—a link that is in constant danger of damage from soil insects. Aldrin gives economic control of the complete range of soil insect pests. It can be used for row-treatment, as an overall application—for which aldrin-fertiliser mixtures are particularly suitable, or seed treatment. It is of particular use with cereals, especially maize and for sugar cane, sugar beet, potatoes and tobacco.

Aldrin is complementary to endrin and dieldrin. Together they control a large variety of major destructive pests, ranging from soil insects to foliage pests. They have excellent residual action, are stable in the presence of other chemicals, do not affect germination and do not cause taint. Furthermore their use, according to recommendation is completely safe to the applicator and to the consumer of treated crops. Aldrin, dieldrin and endrin are available as technical materials and high concentrate dusts for further formulation, and also in forms suitable for the grower: emulsifiable concentrates, wettable powders, field-strength dusts and granules.

For details of approved uses and advice on application, consult the Shell Company in your country.

