

Conclusions: Teaching is challenging and yet one of the most rewarding professions, but several factors correlate with stress, burnout, anxiety, and depression among teachers. Highlighting these factors is the first step in recognizing the magnitude of the issues encountered by those in the teaching profession. Implementation of a school-based awareness and intervention program is crucial to resolve the early signs of teacher stress and burnout to avoid future deterioration.

Disclosure of Interest: None Declared

EPV0699

Effects of cumulative trauma from multiple natural disasters

B. Agyapong*, R. Shalaby, E. Eboreime, G. Obuobi-Donkor, E. Owusu, M. Adu, W. Mao, F. Oluwasina and V. Agyapong

Department of Psychiatry, University of Alberta, Edmonton, Canada
*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2011

Introduction: Fort McMurray, a city in northern Alberta, Canada, has experienced multiple traumatic events in the last five years, including the 2016 wildfire, the 2020 floods, and the COVID-19 pandemic. Traumatic events often lead to increased mental health burdens in affected communities.

Objectives: To assess if the number of traumatic events experienced by residents of Fort McMurray correlates with the prevalence and severity of mental health issues experienced.

Methods: A cross-sectional study using an online survey questionnaire was used to gather demographic, trauma (wildfire, flooding, and COVID-19), and clinical information from the resident of Fort McMurray between April 24 to June 2 2021. Likely Generalized Anxiety Disorder (GAD), Major Depressive Disorder (MDD), Post-Traumatic Stress Disorder (PTSD) and low resilience were measured using standardized rating scales. Data were analyzed with SPSS version 26 using Chi-Square tests and multivariate regression analysis.

Results: Respondents who experienced COVID-19 and either flood or wildfire traumas ($N = 101$) were eleven times more likely to have GAD symptoms (OR: 11.39; 95% CI: 1.43-91.04), four times more likely to have likely MDD, (OR: 3.85; 95% CI: .995-14.90), ten times more likely to have likely PTSD (OR: 10.47; 95% CI: 1.28-85.67), and low resilience (OR: 10.56; 95% CI: 1.21-92.17). Respondents who experienced COVID-19, flooding, and wildfire traumas ($N = 47$) were eighteen times more likely to express GAD symptoms (OR: 18.30; 95% CI: 2.20-152.45) and more than eleven times likely to have likely PTSD (OR: 11.41; 95% CI: 1.34-97.37) in comparison to the respondents who experienced COVID-19 only trauma ($N = 19$).

Conclusions: Measures to reduce climate change and associated natural disasters could reduce the impact of cumulative trauma and associated mental health burden in vulnerable populations. It is essential that more mental health resources are mobilized to support communities impacted by multiple natural disasters.

Disclosure of Interest: None Declared

EPV0700

Predictors of suicidal ideation among female residents in a community impacted by multiple natural disasters.

B. Agyapong*, R. Shalaby, E. Eboreime, Y. W. and V. Agyapong

Department of Psychiatry, University of Alberta, Edmonton, Canada
*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2012

Introduction: Suicidal ideation and thoughts of self-harm continue to be challenging public health problems. It's presently unknown what the prevalence and correlates of suicidal thoughts and self-harm are in female residents of Fort McMurray, a city that has endured wildfires, flooding, and the COVID-19 pandemic in the last five years.

Objectives: This study aimed to determine the prevalence and correlates of suicidal ideation and thoughts of self-harm among female residents of Fort McMurray.

Methods: A cross-sectional study using an online survey questionnaire was used to collect sociodemographic and clinical information from the resident of Fort McMurray between April 24 to June 2 2021. Suicidal ideation and thoughts of self-harm among females were assessed using the ninth question of the Patient Health Questionnaire-9, a validated screening tool used to assess depression symptoms. Likely Generalized Anxiety Disorder (GAD), Major Depressive Disorder (MDD), Post-Traumatic Stress Disorder (PTSD) and low resilience were measured using standardized rating scales. Data were analyzed with SPSS version 25 using Chi-Square tests and multivariate logistic regression analysis.

Results: Among Fort McMurray residents, 249 accessed the online survey, while 186 ultimately completed it, yielding a survey completion rate of 74.7%. Of these, 159 (85%) were females. After controlling for other variables in the regression model, respondents who expressed a desire to receive mental health counselling were more than seven times more likely to report suicidal ideation and thoughts of self-harm compared to the respondents who didn't desire to receive mental health counselling (OR: 7.29; 95% CI: 1.19 – 44.58). Similarly, respondents who reported having abused alcohol in the past year were nearly four times more likely to report suicidal ideation or thoughts of self-harm compared to the respondents who said they had not abused alcohol in the past year (OR: 3.91; 95% CI: 1.05 – 14.57).

Conclusions: High prevalence of suicidal thoughts and thoughts of self-harm were reported among female residents of Fort McMurray. Timely access to adequate mental health support should be offered to female residents of communities impacted by multiple natural disasters, particularly residents who self-report alcohol abuse or desire to receive mental health counselling

Disclosure of Interest: None Declared

EPV0701

Mental Health Impacts of Wildfire, Flooding and COVID-19: on educators: A Comparative Study

B. Agyapong*, E. Eboreime, R. Shalaby, H. Pazderka, G. Obuobi-Donkor, M. Adu, W. Mao, F. Oluwasina, E. Owusu, A. Greenshaw and V. Agyapong

¹Department of Psychiatry, University of Alberta, Edmonton, Canada
*Corresponding author.

doi: 10.1192/j.eurpsy.2023.2013