







Palmer amaranth (*Amaranthus palmeri*) control in furrow-irrigated rice with fluridoneTanner A. King<sup>1</sup> , Jason K. Norsworthy<sup>2</sup> , Thomas R. Butts<sup>3</sup> , L. Tom Barber<sup>4</sup> , Gerson L. Drescher<sup>5</sup>  and Amar S. Godar<sup>6</sup> 

## Research Article

**Cite this article:** King TA, Norsworthy JK, Butts TR, Barber LT, Drescher GL, Godar AS (2025) Palmer amaranth (*Amaranthus palmeri*) control in furrow-irrigated rice with fluridone. *Weed Technol.* **39**(e8), 1–11. doi: [10.1017/wet.2024.91](https://doi.org/10.1017/wet.2024.91)

Received: 15 July 2024  
Revised: 7 October 2024  
Accepted: 5 November 2024

**Associate Editor:**

Connor Webster, Louisiana State University Agricultural Center

**Nomenclature:**

Florpyrauxifen-benzyl; fluridone; Palmer amaranth, *Amaranthus palmeri* (S.) Watson; cotton, *Gossypium hirsutum* L.; rice, *Oryza sativa* L.

**Keywords:**

chemical control; crop injury; herbicides; seed production; weed control

**Corresponding author:**

Tanner A. King; Email: [tak196@msstate.edu](mailto:tak196@msstate.edu)

<sup>1</sup>Graduate Research Student, Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, Fayetteville, AR, USA; <sup>2</sup>Distinguished Professor and Elms Farming Chair of Weed Science, Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, Fayetteville, AR, USA; <sup>3</sup>Clinical Assistant Professor and Extension Specialist, Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN, USA; <sup>4</sup>Professor and Extension Specialist; Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, Lonoke, AR, USA; <sup>5</sup>Assistant Professor of Soil Fertility, Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, Fayetteville, AR, USA and <sup>6</sup>Post Doctoral Fellow, Department of Crop, Soil, and Environmental Sciences, University of Arkansas System Division of Agriculture, Fayetteville, AR, USA

**Abstract**

Herbicide-resistant Palmer amaranth is creating additional challenges for producers who choose to adopt a furrow-irrigated rice production system due to the absence of a sustained flood, enabling extended weed emergence. Fluridone has been shown to effectively control Palmer amaranth in cotton production systems and was recently registered for use in rice. Experiments were initiated in 2022 and 2023 1) to evaluate Palmer amaranth control and rice tolerance to preemergence- and postemergence-applied fluridone at 0.5× (84 g ai ha<sup>-1</sup>) and 1× (168 g ai ha<sup>-1</sup>) rates on a silt loam soil; and 2) assess the effect of various herbicide programs that contain fluridone on Palmer amaranth biomass, seed production, and rough rice grain yield. Preemergence applications of fluridone at a 1× rate in combination with clomazone resulted in 84% control of Palmer amaranth 21 d after treatment (DAT). Fluridone, in combination with clomazone preemergence, caused up to 36% rice injury 21 DAT; however, early season injury did not negatively affect rice yields. Palmer amaranth biomass and fecundity were reduced with herbicide programs that included fluridone plus florpyrauxifen-benzyl, and, in some instances, there was no Palmer amaranth biomass or seed production following multiple applications of both herbicides. Fluridone- and florpyrauxifen-benzyl-based herbicide programs achieved effective control of Palmer amaranth when applied timely, but injury to hybrid rice is enhanced with preemergence applications of fluridone that are not permitted with the current label.

**Introduction**

In the mid-southern United States, rice is typically produced in flooded fields, which requires straight or contour levees to help maintain a permanent flood throughout the growing season (Hardke et al. 2022). However, furrow-irrigated rice has gained popularity in recent years, and in 2022, 18% of the total rice hectares in Arkansas were furrow-irrigated. While flood-irrigated rice involves establishing a continuous flood at the V5 stage of rice until maturity, furrow-irrigated rice is made possible by creating raised beds with furrows between the hipped rows, allowing the movement of water via gravity from the top end of the field (Counce et al. 2000; Lunga et al. 2021). Not only have producers lauded the idea of furrow-irrigated rice potentially decreasing water and equipment use (Chlapecka et al. 2021), but it also makes for an efficient transition into a rice-soybean [*Glycine max* (L.) Merr.] crop rotation, which is a frequent practice in Arkansas (Nalley et al. 2022). A major benefit of crop rotation is being able to integrate herbicide programs targeting troublesome grass species in a broadleaf crop (Burgos et al. 2021). However, the different water management practices associated with these two systems influence the emergence pattern and spectrum of weeds within a field (Kraehmer et al. 2016).

As of 2022, the most troublesome weed species in flood-irrigated rice included barnyardgrass [*Echinochloa crus-galli* (L.) P. Beauv.], sedge species (*Cyperus* spp.) and weedy rice (*Oryza sativa* L.), whereas barnyardgrass, Palmer amaranth, and sedges were among the most problematic weeds in a furrow-irrigated rice system (Butts et al. 2022). The permanent flood established by conventional paddy rice can help alleviate weed emergence, especially from terrestrial weeds that cannot survive in anaerobic conditions (Bagavathiannan et al. 2011). Furrow-irrigated rice allows typical upland crop weeds, such as Palmer amaranth, to thrive throughout the growing season due to the aerobic conditions that create a favorable environment for weed emergence and survival (Beesinger et al. 2022; Norsworthy et al. 2011).

© University of Arkansas, 2024. This is a work of the US Government and is not subject to copyright protection within the United States. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



Palmer amaranth has historically been among the five most troublesome weeds in major row crop production systems in mid-southern states (Norsworthy et al. 2014; Van Wychen 2022; Webster and Nichols 2012). Although the influence of Palmer amaranth emergence on rice yields is unknown, some studies have demonstrated its negative impact on soybean, cotton, and corn (*Zea mays* L.) yields (Klingaman and Oliver 1994; Massinga et al. 2001). Previous research has shown that Palmer amaranth densities  $\leq 8$  plants  $m^{-2}$  of row can reduce soybean and cotton yields by 78% and 70%, respectively (Bensch et al. 2003; Rowland et al. 1999). Considering Palmer amaranth has the potential to severely affect crop yields and furrow-irrigated rice hectares are increasing, more research is needed to create management strategies aimed at reducing the growth and development of the weed.

With herbicide-resistant populations of Palmer amaranth being widespread, effective chemical control options for Palmer amaranth are scarce. Typically, herbicides are used to control the most problematic weeds because they are easily accessible and convenient to apply (Priess et al. 2022). However, in Arkansas, Palmer amaranth has evolved resistance to eight herbicide sites of action, which is why it is recommended that producers overlap the use of multiple herbicide chemistries to help slow the evolution of herbicide-resistant weed species (Bagavathiannan et al. 2013; Barber et al. 2015; Heap 2024). Among those sites of action previously mentioned, microtubule assembly inhibitors (categorized as a Group 3 herbicide by the Herbicide Resistance Action Committee [HRAC] and Weed Science Society of America [WSSA]), acetolactate synthase (ALS) inhibitors (HRAC/WSSA Group 2), and protoporphyrinogen oxidase (PPO) inhibitors (HRAC/WSSA Group 14) are used in rice production but are no longer effective due to confirmation of herbicide-resistant Palmer amaranth populations (Bond et al. 2006; Gossett et al. 1992; Varanasi et al. 2019). Florpyrauxifen-benzyl (Loyant™; Corteva Agriscience, Wilmington, DE) and 2,4-D, both synthetic auxins (HRAC/WSSA Group 4), are generally effective at controlling Palmer amaranth in a furrow-irrigated rice system, but there is also confirmed resistance to 2,4-D (Hwang et al. 2023). Additionally, strict application regulations have been placed on both herbicides due to injury to adjacent susceptible crops from off-target movement (ASPB 2020; Barber et al. 2023; Wright et al. 2020). The adoption of novel herbicide sites of action for Palmer amaranth control would be beneficial given the status of current common chemical weed control options.

Fluridone (Brake; SePRO Corporation, Carmel, IN), a phytoene desaturase inhibitor (WSSA Group 12), has been commonly used as a soil-applied, preemergence herbicide for control of broadleaf and grass species in cotton crops (Hill et al. 2016). Fluridone is highly effective at controlling Palmer amaranth, especially on silt loam soils (Banks and Merkle 1979). The previous study also showed fluridone having prolonged activity in clay soils, with the herbicide being detected up to 250 d after application. Since most Arkansas rice hectares consist of silt loam soils (Hardke 2022), fluridone has potential value in furrow-irrigated rice production systems. However, before recommending fluridone for use on furrow-irrigated rice, rice tolerance and herbicidal efficacy must be assessed.

Since 2016, studies have been conducted to evaluate rice injury from fluridone carryover, with one study showing that fluridone applied at 224 g ai  $ha^{-1}$  prior to cotton emergence caused no more than 5% injury to rice the following year (Hill et al. 2016). Conversely, a 16% and 22% reduction in rice stand was observed

**Table 1.** Herbicide treatment, timing, and rate for the different programs evaluated in the single fluridone application experiment in 2022 and 2023<sup>a,b</sup>.

Herbicide treatment	Timing	Rate
		g ai/ae $ha^{-1}$
Clomazone +	PRE	336
Fluridone	PRE	168
Clomazone +	PRE	336
Fluridone	PRE	168
Florpyrauxifen-benzyl	MPOST	15
Clomazone +	PRE	336
Fluridone	PRE	168
Florpyrauxifen-benzyl	MPOST	30
Clomazone	PRE	336
Florpyrauxifen-benzyl +	EPOST	15
Florpyrauxifen-benzyl	MPOST	15
Clomazone	PRE	336
Fluridone +	EPOST	168
Florpyrauxifen-benzyl	EPOST	15
Florpyrauxifen-benzyl	MPOST	15
Clomazone	PRE	336
Florpyrauxifen-benzyl	EPOST	15
Fluridone +	MPOST	168
Florpyrauxifen-benzyl	MPOST	15

<sup>a</sup>Abbreviations: EPOST, early postemergence; MPOST, mid-postemergence; PRE, preemergence.

<sup>b</sup>All florpyrauxifen-benzyl applications included 0.58 L  $ha^{-1}$  methylated seed oil.

with fluridone applied at 448 and 900 g ai  $ha^{-1}$ , respectively. As of 2023, fluridone was labeled for postemergence use in dry-seeded rice production beginning at the 3-leaf rice growth stage at a maximum annual use rate of 168 g ai  $ha^{-1}$  (Anonymous 2023). Therefore, the objectives of this research were to 1) assess Palmer amaranth control and rice tolerance to preemergence- and postemergence-applied fluridone at 0.5 $\times$  and 1 $\times$  label rates on a silt loam soil, and 2) evaluate the effect of various herbicides combinations, including fluridone, on Palmer amaranth biomass, seed production, and rice grain yield.

## Materials and Methods

### Palmer Amaranth Control and Rice Injury with Single Applications of Fluridone

Field experiments were initiated in 2022 and 2023 at the Pine Tree Research Station near Colt, Arkansas (35.10887°N, 90.94066°W), on a Calhoun silt loam (fine-silty, mixed, active, thermic Typic Glossaqualfs) consisting of 11% sand, 68% silt, 21% clay, and 1.6% organic matter, pH 7.2. In 2022 and 2023, an additional site was located at the Milo J. Shult Research and Extension Center in Fayetteville, Arkansas (36.09366°N, 94.17344°W), on a Leaf silt loam (fine, mixed, active, thermic Typic Albaquults) consisting of 18% sand, 69% silt, 13% clay, and 1.6% organic matter, with a pH of 6.7. The field experiment was designed to evaluate rice tolerance and Palmer amaranth control with a single fluridone application at a 1 $\times$  label rate (168 g ai  $ha^{-1}$ ) in combination with clomazone (Command 3ME; FMC Corporation, Philadelphia, PA) applied preemergence or florpyrauxifen-benzyl applied postemergence in a furrow-irrigated rice system (Table 1). The experiment was conducted as a randomized complete block design with four replications. Before planting, the test site was field-cultivated and hipped into 91-cm and 76-cm wide beds in Fayetteville and Colt, respectively. The trials were kept free of grass weeds and sedge species using fenoxaprop (Ricestar HT; Gowan Company, Yuma, AZ) and halosulfuron (Permit®; Gowan Company), and hand-

**Table 2.** Planting and herbicide application dates<sup>a</sup>.

Location	Planting	PRE	EPOST	MPOST
Fayetteville	June 1, 2022	June 1, 2022	June 23, 2022	July 1, 2022
	May 2, 2023	May 3, 2023	June 1, 2023	June 13, 2023
Colt	May 17, 2022	May 18, 2022	June 9, 2022	June 15, 2022
	May 3, 2023	May 3, 2023	May 26, 2023	June 6, 2023

<sup>a</sup>Abbreviations: EPOST, early postemergence; MPOST, mid-postemergence; PRE, preemergence.

weeding when necessary. Three split nitrogen applications as urea (460 g N kg<sup>-1</sup>), were applied at 135 kg N ha<sup>-1</sup> in 2-wk intervals following the V5 growth stage. Other nutrients were supplied preplant based on recommendations from the University of Arkansas System Division of Agriculture Marianna Soil Test Laboratory. Unless rainfall occurred, the experimental area was irrigated twice weekly following the V5 rice stage.

At all sites, a hybrid, long-grain rice cultivar 'Full Page RT7321FP' (RiceTec, Alvin, TX) was planted at 35 kg ha<sup>-1</sup> at a 1-cm depth and 19 cm between rows. Plot dimensions were 3.7 m (four beds) wide by 5.2 m long and 3.1 m wide (four beds) by 5.2 m long in Fayetteville and Colt, respectively. The experiment consisted of seven treatments, including a nontreated control for comparison, with application timings occurring preemergence, early postemergence at 3-leaf rice, and mid-postemergence at rice tillering. Herbicides were applied using a CO<sub>2</sub>-pressurized backpack sprayer calibrated to deliver 140 L ha<sup>-1</sup> at 276 kPa using four TeeJet 110015 AIXR nozzles (Spraying Systems Co., Glendale Heights, IL) spaced 48 cm apart (Table 2).

Visible estimations of rice injury and Palmer amaranth control were rated on a 0 to 100 scale, with zero being no plant symptomatology and 100 representing complete plant mortality. Ratings were recorded 21 d after preemergence, 7 d after early postemergence, and 14 and 28 d after mid-postemergence. Before rice harvest, Palmer amaranth aboveground biomass was collected from a 1-m<sup>2</sup> quadrat within the center of each plot. All harvested biomass was placed in an oven at 66 C for 2 wk, dried to constant mass, and dry biomass weight was recorded. Afterward, each female Palmer amaranth plant was threshed, and the ground plant material was separated from the seeds using a 20-mesh sieve and a vertical air column seed cleaner (Miranda et al. 2021). After cleaning, a 200-seed subsample from three random plots was weighed, and the average weight was used to calculate the number of seeds produced in the square-meter quadrat. Rough rice grain yield was determined by harvesting the center of each plot using an 8-XP small-plot combine (Kincaid, Haven, KS) equipped with a 1.8-m-wide header. The grain yield was adjusted to 12% moisture.

### Palmer Amaranth Control and Rice Injury with Single and Multiple Applications of Fluridone

A field experiment was conducted at the Lon Mann Cotton Research Station in Marianna, AR (34.72549°N, 90.73423°W), in 2022 and the Milo J. Shult Research and Extension Center in Fayetteville, AR, in 2023, to assess Palmer amaranth control and rice injury with single and multiple applications of fluridone in combination with clomazone applied preemergence or florpyrauxifen-benzyl applied postemergence. The experiment was conducted as a randomized complete block design with four replications. The soil at the Marianna site was a Convent silt loam consisting of 9% sand, 80% silt, 11% clay, and 1.8% organic matter, pH 6.5. In Fayetteville, the soil was a Leaf silt loam consisting of

18% sand, 69% silt, 13% clay, and 1.6% organic matter, with a pH of 6.6. Before planting, the soil was tilled and hipped into 96-cm-wide and 91-cm wide beds in Marianna and Fayetteville, respectively.

In both years, a hybrid, long-grain rice cultivar 'Full Page RT7321FP' (RiceTec, Alvin, TX) was sown at 35 kg ha<sup>-1</sup> at a 1-cm depth with 19 cm between rows. Plot dimensions were 1.9 m (two beds) wide by 5.2 m long and 1.8 m (two beds) wide by 5.2 m long in Marianna and Fayetteville, respectively. A natural population of Palmer amaranth was allowed to germinate after rice planting was completed. The trials were kept free of other undesirable weed species using applications of fenoxaprop or mechanical methods if necessary. The soil for each trial was amended for fertility preplant based on soil test values from the University of Arkansas System Division of Agriculture Marianna Soil Test Laboratory fertility recommendations. Once the rice reached the V5 growth stage, irrigation water was delivered every 2 d unless rainfall occurred. Nitrogen, as urea (460 g N kg<sup>-1</sup>), was applied at 135 kg N ha<sup>-1</sup> in three separate applications in 2-wk intervals beginning at the 5-leaf stage of rice. This trial consisted of nine treatments, including a nontreated control, with application timings occurring preemergence, mid-postemergence, and late postemergence at 42 d after planting (DAP) (Table 3). Herbicide treatments were made using a CO<sub>2</sub>-pressurized backpack sprayer calibrated to deliver 140 L ha<sup>-1</sup> at 276 kPa using four 110015 AIXR nozzles spaced 48 cm apart (TeeJet Technologies, Glendale Heights, IL) (Table 4).

Visible Palmer amaranth control and rice injury ratings were recorded 21 d after preemergence, 14 d after mid-postemergence, and 14 and 28 d after late postemergence. Visual ratings were based on the same 0 to 100 scale mentioned in the previous experiment. Likewise, Palmer amaranth biomass, weed seed production, and rice grain yield data collection used the same parameters as the single fluridone application experiment.

### Data Analysis

All data were analyzed using R studio software (v. 4.3.2) (R Core Team) using the *glmmTMB* function (GLMMTMB package; Brooks et al. 2017). Data were checked to determine whether the assumptions of normality and homogenous variance were met using the Shapiro-Wilks test and Levene's test. Rice injury, Palmer amaranth control, Palmer amaranth biomass, Palmer amaranth seed production, and rice grain yield were fitted to a generalized linear mixed-effect model (Stroup 2015). All injury and control data were analyzed using a beta distribution by evaluation timing, whereas Palmer amaranth biomass and seed production at harvest were analyzed using a Poisson distribution (Gbur et al. 2012). After the residuals failed to violate the Shapiro-Wilks and Levene's tests, rice grain yield was analyzed using a Gaussian or normal distribution. ANOVA was performed on each model using the CAR package with a Type III Wald chi-square test (Fox and Weisberg 2019). For each herbicide program, estimated marginal means (Searle et al. 1980) were obtained using the EMMEANS

**Table 3.** Herbicide treatments, timings, and rates evaluated for the multiple fluridone application experiment in 2022 and 2023<sup>a,b</sup>.

Herbicide treatment	Timing	Rate
		g ai/ae ha <sup>-1</sup>
Clomazone	PRE	336
Florpyrauxifen-benzyl	MPOST	15
Florpyrauxifen-benzyl	LPOST	15
Clomazone +	PRE	336
Fluridone	PRE	84
Florpyrauxifen-benzyl	MPOST	15
Florpyrauxifen-benzyl	LPOST	15
Clomazone +	PRE	336
Fluridone	PRE	168
Florpyrauxifen-benzyl	MPOST	15
Florpyrauxifen-benzyl	LPOST	15
Clomazone +	PRE	336
Fluridone	PRE	84
Florpyrauxifen-benzyl +	MPOST	15
Fluridone	MPOST	84
Florpyrauxifen-benzyl	LPOST	15
Clomazone +	PRE	336
Fluridone	PRE	84
Florpyrauxifen-benzyl	MPOST	15
Florpyrauxifen-benzyl +	LPOST	15
Fluridone	LPOST	84
Clomazone +	PRE	336
Fluridone	PRE	84
Florpyrauxifen-benzyl	LPOST	15
Clomazone +	PRE	336
Fluridone	PRE	84
Florpyrauxifen-benzyl +	LPOST	15
Fluridone	LPOST	84

<sup>a</sup>Abbreviations: LPOST, late postemergence; MPOST, mid-postemergence; PRE, preemergence.

<sup>b</sup>All florpyrauxifen-benzyl applications included 0.58 L ha<sup>-1</sup> methylated seed oil.

package. The Sidak method was used to adjust for multiple pairwise comparisons (Midway et al. 2020), and the MULTCOMP package was used to generate a compact letter display to visually distinguish differences among treatments (Hothorn et al. 2008). In both experiments, site year and block nested within site year were treated as random effects to draw general conclusions over diverse environments (McLean et al. 1991; Midway 2022), and herbicide treatment was considered a fixed effect (Blouin et al. 2011).

## Results and Discussion

### Palmer Amaranth Control and Rice Injury with Single Applications of Fluridone

There were differences among herbicide treatments for visible estimations of Palmer amaranth control 21 d after preemergence when averaged over four site-years ( $P = 0.0002$ , Table 5). At 21 d after preemergence, Palmer amaranth control ranged from 40% to 86% among herbicide treatments. Across all site years, Palmer amaranth control was  $\geq 83\%$  when clomazone + fluridone was applied at 21 d after preemergence, which is similar to other research reporting 70% to 94% Palmer amaranth control 3 wk after cotton planting with preemergence applications of fluridone at rates ranging from 224 to 448 g ai ha<sup>-1</sup> (Hill et al. 2017). However, the fluridone rates in the study by Hill et al. (2017) are higher than the maximum annual use rate of 168 g ai ha<sup>-1</sup> in rice. In most instances, preemergence treatments that included clomazone +

fluridone provided greater Palmer amaranth control than clomazone applied alone 21 d after preemergence. Norsworthy et al. (2008) also reported reduced control of Palmer amaranth when clomazone was applied alone preemergence.

Similarly, rice injury differed among herbicide treatments 21 d after preemergence ( $P < 0.0001$ ; Table 5). Both clomazone and fluridone disrupt pigment formation in the plant cells of susceptible species, leading to a bleached appearance on plant leaves (Anderson and Roberston 1960; Duke et al. 1991). Although rice exhibits acceptable tolerance to clomazone, severe rice injury may still occur under various climatic conditions (Zhang et al. 2005); hence, the bleaching symptomology was greater when fluridone was mixed with clomazone. As a result, preemergence treatments containing clomazone alone caused 10% to 16% injury to rice, whereas fluridone with clomazone resulted in 29% to 33% injury 21 d after preemergence. These findings are similar to those observed by Martin et al. (2018), who reported that clomazone and fluridone, applied individually at higher rates, caused 12% and 18% injury to rice, respectively, 4 wk after treatment. Based on results, preemergence treatments containing fluridone + clomazone caused a 2-fold increase in rice injury.

Herbicide treatment was significant for Palmer amaranth control ratings taken 7 d after early postemergence ( $P < 0.0001$ ; Table 6). Palmer amaranth control ranged from 76% to 92%, and preemergence treatments containing clomazone and fluridone provided greater control than clomazone applied preemergence followed by florpyrauxifen-benzyl applied early postemergence. Waldrep and Taylor (1976) reported that fluridone provides prolonged residual control of *Amaranthus* species; hence, greater control of Palmer amaranth was observed at this timing even in the absence of a postemergence herbicide application. Additionally, applying fluridone with florpyrauxifen-benzyl early postemergence did not enhance Palmer amaranth control, which is likely attributed to fluridone having minimal postemergence activity on established weeds (Waldrep and Taylor 1976).

At 7 d after early postemergence, rice injury evaluations differed among herbicide treatments ( $P < 0.0001$ ; Table 6). Preemergence treatments that included clomazone + fluridone were more injurious to rice than clomazone applied preemergence followed by florpyrauxifen-benzyl early postemergence. Based on the injury data from similar treatments, clomazone + fluridone applied preemergence caused 27% to 30% injury to rice, whereas injury from clomazone applied preemergence followed by an application of florpyrauxifen-benzyl early postemergence ranged from 8% to 13%. Moreover, fluridone with florpyrauxifen-benzyl applied early postemergence did not cause additional injury to rice compared with herbicide treatments that contained clomazone applied preemergence followed by florpyrauxifen-benzyl applied early postemergence. Hence, rice injury can be minimized with an early postemergence application of fluridone while achieving greater Palmer amaranth control.

The final treatment was applied mid-postemergence, and visible estimations of Palmer amaranth control were significant at 14 d ( $P = 0.0169$ ) and 28 d ( $P < 0.0001$ ) after treatment (Table 7). Palmer amaranth control was similar or comparable (90% to 92%) for all herbicide programs containing a postemergence application of florpyrauxifen benzyl 14 DAT. A previous study also reported similar Palmer amaranth control levels 21 DAT with florpyrauxifen-benzyl applied at 15 and 30 g ae ha<sup>-1</sup> (Beesinger et al. 2022). At 28 DAT, herbicide programs that included either a single postemergence application of florpyrauxifen-benzyl at 30 g ae ha<sup>-1</sup> or sequential early postemergence and



**Table 4.** Planting and herbicide application dates<sup>a</sup>.

Location	Planting	PRE	MPOST	LPOST
Marianna	May 10, 2022	May 10, 2022	June 6, 2022	June 21, 2022
Fayetteville	May 2, 2023	May 3, 2023	June 1, 2023	June 13, 2023

<sup>a</sup>Abbreviations: LPOST, late postemergence; MPOST, mid-postemergence; PRE, preemergence.

**Table 5.** Visible Palmer amaranth control and rice injury for the single fluridone application experiment at 21 d after preemergence, averaged over four site-years<sup>a,b</sup>.

Herbicides	Rate g ai ha <sup>-1</sup>	AMAPA control		Rice injury
		— % of nontreated —		
Clomazone + fluridone	336 + 168	86 a		30 a
Clomazone + fluridone	336 + 168	83 a		33 a
Clomazone + fluridone	336 + 168	84 a		29 a
Clomazone	336	45 b		10 b
Clomazone	336	40 b		16 b
Clomazone	336	69 ab		10 b
P-value		0.0002		<0.0001

<sup>a</sup>Abbreviation: AMAPA, Palmer amaranth.

<sup>b</sup>Means within a column followed by the same letter are not different according to the Sidak method ( $\alpha = 0.05$ ).

mid-postemergence applications of the herbicide at 15 g ae ha<sup>-1</sup> provided 97% to 98% Palmer amaranth control. Clomazone applied preemergence followed by an early postemergence application of fluridone and sequential applications of florpyrauxifen-benzyl at early postemergence and mid-postemergence provided 98% control of Palmer amaranth, indicating that control can be optimized with labeled fluridone applications with florpyrauxifen-benzyl. Additionally, these findings support the need for effective postemergence herbicides in combination with soil-applied residuals to achieve extended Palmer amaranth control, as noted by Hill et al. (2017).

Similarly, rice injury differed among herbicide programs at 14 d ( $P = 0.0034$ ) and 28 d ( $P = 0.0232$ ) after the final application (Table 7). At 14 DAT, clomazone applied preemergence followed by florpyrauxifen-benzyl applied early postemergence, followed by fluridone + florpyrauxifen-benzyl applied mid-postemergence resulted in 7% injury; clomazone + fluridone applied preemergence resulted in 15% injury; and clomazone + fluridone applied preemergence followed by florpyrauxifen-benzyl at 15 g ae ha<sup>-1</sup> applied mid-postemergence resulted in 17% injury. These results indicate that minimal injury to rice occurs with mid-season fluridone applications mixed with florpyrauxifen-benzyl. As a result, producers can achieve residual Palmer amaranth control later in the growing season with postemergence fluridone applications while mitigating severe injury to the crop caused by a preemergence application. At 28 DAT, clomazone + fluridone applied preemergence followed by florpyrauxifen-benzyl at 30 g ae ha<sup>-1</sup> applied mid-postemergence caused 8% injury to rice. Rice injury was reduced to 3% when fluridone was applied mid-postemergence or excluded from the herbicide program, suggesting that rice is more tolerant to fluridone at later growth stages and when applied to foliage (Waldrep and Taylor 1976).

Over four site-years, all herbicide programs affected rough rice grain yield ( $P = 0.0044$ ; Table 8). Rice yield in the nontreated control was lowest than all other evaluated treatments. For all other herbicide programs, rice yields were similar and ranged from 6,380 to 6,600 kg ha<sup>-1</sup>. High levels of Palmer amaranth control were achieved early in the season; thus, there was a lack of interference

from Palmer amaranth and subsequent impact on grain yield throughout the remainder of the growing season. The yield data observed here indicates that postemergence fluridone applications to rice do not affect rough rice grain yield; hence, Palmer amaranth control can be achieved with labeled rates of fluridone while not affecting grain development.

Conversely, Palmer amaranth biomass production differed as a function of the herbicide program averaged over the four site-years, and weed biomass ranged from 0.2 to 350 g m<sup>-2</sup> ( $P < 0.0001$ ; Table 8). Compared with the nontreated control, Palmer amaranth biomass production was reduced by all the herbicide programs. However, postemergence treatments that contained florpyrauxifen-benzyl or florpyrauxifen-benzyl + fluridone caused the greatest reduction in Palmer amaranth biomass production, indicating the importance of applying effective postemergence herbicides to rice crops. The preemergence application of clomazone + fluridone followed by florpyrauxifen-benzyl at 30 g ae ha<sup>-1</sup> mid-postemergence allowed Palmer amaranth to produce similar quantities of biomass compared with a preemergence application of clomazone with sequential postemergence applications of florpyrauxifen-benzyl at 15 g ae ha<sup>-1</sup>. These findings indicate that preemergence applications of fluridone may not be necessary to effectively suppress Palmer amaranth biomass production in a furrow-irrigated rice system.

The quantity of weed seed produced by Palmer amaranth was affected by the herbicide program ( $P < 0.0001$ ; Table 8). Compared with the nontreated control, clomazone + fluridone applied preemergence allowed Palmer amaranth to produce the highest quantity of seed at approximately 23,500 seed m<sup>-2</sup>. Sequential postemergence applications had the greatest impact on Palmer amaranth seed production relative to treatments that included a single postemergence application. The seed production data reported here is supported by Beesinger et al. (2022), who found that sequential florpyrauxifen-benzyl applications reduced Palmer amaranth seed production to a greater extent than a single application. Additionally, weed seed production was similar among programs in which fluridone was applied either early postemergence or mid-postemergence with florpyrauxifen-benzyl, which may be attributed to the herbicide preventing additional weed emergence and subsequent offspring additions to the soil seedbank.

### Palmer Amaranth Control and Rice Injury with Single and Multiple Applications of Fluridone

Averaged over site-year, herbicide treatment affected Palmer amaranth control when it was recorded 21 d after preemergence ( $P < 0.0001$ ; Table 9). At 21 d after preemergence, Palmer amaranth control improved when fluridone was applied at either 84 or 168 g ai ha<sup>-1</sup> in a mixture with clomazone, providing 84% to 91% control across the herbicide treatments we evaluated. Weed control was greatest when clomazone was applied preemergence with fluridone at 84 and 168 g ai ha<sup>-1</sup>. Herbicide combinations that contained clomazone + fluridone at 84 or 168 g ai ha<sup>-1</sup> resulted in

**Table 6.** Visible Palmer amaranth control and rice injury for the single fluridone application experiment 7 d after early postemergence, averaged over four site-years<sup>a,b,c</sup>.

Herbicides	Timing	Rate g ai/ae ha <sup>-1</sup>	AMAPA control		Rice injury
			% of nontreated		
Clomazone + Fluridone	PRE	336 168	92 a		30 a
Clomazone + Fluridone	PRE	336 168	91 a		32 a
Clomazone + Fluridone	PRE	336 168	88 ab		27 a
Clomazone	PRE	336	76 b		12 b
Florpyrauxifen-benzyl Clomazone	EPOST PRE	15 336	82 ab		13 b
Fluridone + Florpyrauxifen-benzyl	EPOST EPOST	168 15	77 b		8 b
Clomazone Florpyrauxifen-benzyl	PRE EPOST	336 15	<0.0001		<0.0001
P-value					

<sup>a</sup>Abbreviations: AMAPA, Palmer amaranth; EPOST, early postemergence; PRE, preemergence.

<sup>b</sup>Means within a column followed by the same letter are not different according to the Sidak method ( $\alpha = 0.05$ ).

<sup>c</sup>All flurpyrauxifen-benzyl applications included 0.58 L ha<sup>-1</sup> methylated seed oil.

**Table 7.** Visible Palmer amaranth control and rice injury for the single fluridone application experiment 14 and 28 d after mid-postemergence, averaged over four site-years<sup>a,b</sup>.

Herbicides	Timing	Rate g ai/ae ha <sup>-1</sup>	AMAPA control		Rice injury	
			14 DAT	28 DAT	14 DAT	28 DAT
Clomazone + Fluridone	PRE	336 168	69 b		73 c	
Clomazone + Fluridone	PRE	336 168	90 a		94 b	
Clomazone + Fluridone	PRE	336 168	91 a		97 ab	
Florpyrauxifen-benzyl Clomazone	MPOST PRE	15 336	91 a		97 ab	
Fluridone + Florpyrauxifen-benzyl	EPOST EPOST	168 15	92 a		98 a	
Clomazone Fluridone + Florpyrauxifen-benzyl	PRE EPOST MPOST	336 168 15	90 a		97 ab	
Fluridone + Florpyrauxifen-benzyl	EPOST MPOST	168 15	0.0169		<0.0001	
P-value					0.0034	

<sup>a</sup>Abbreviations: AMAPA, Palmer amaranth; DAT, days after treatment preemergence; EPOST, early postemergence; MPOST, mid-postemergence; PRE, preemergence.

<sup>b</sup>Means within a column followed by the same letter are not different according to the Sidak method ( $\alpha = 0.05$ ).

84% to 91% Palmer amaranth control, while clomazone applied alone resulted in only 43% control. These results indicate that applying fluridone along with clomazone preemergence increases Palmer amaranth control relative to clomazone applied alone, as an approximate 2-fold increase in control was observed 21 DAT.

Visible injury to rice was also different among herbicide treatments 21 d after preemergence ( $P = 0.0472$ ; Table 9). Relative to the nontreated control, rice injury was greatest with treatments that contained clomazone + fluridone at 168 g ai ha<sup>-1</sup>. When applied preemergence, clomazone + fluridone at 168 g ai ha<sup>-1</sup> caused 32% to 36% injury to rice, which was greater than the 18% injury caused by clomazone applied alone. These results indicate that preemergence applications of fluridone, sprayed at a 1× rate, have a greater effect on rice than treatments that did not include

the herbicide. Although rice injury from treatments that contained clomazone + fluridone at 84 g ai ha<sup>-1</sup> was not statistically different from the injury that occurred when clomazone was used alone, producers should not apply fluridone until rice reaches the V3 growth stage (Anonymous 2023).

At 14 d after mid-postemergence, visible evaluations of Palmer amaranth control differed among herbicide treatments ( $P < 0.0001$ ; Table 10). Clomazone + fluridone at 84 g ai ha<sup>-1</sup> applied preemergence followed by flurpyrauxifen-benzyl was superior to clomazone alone followed by flurpyrauxifen-benzyl in controlling Palmer amaranth, with each treatment providing 93% and 87% control on average, respectively. In the absence of a postemergence application of flurpyrauxifen-benzyl, Palmer amaranth control ranged from 69% to 94%. Based on the data,

**Table 8.** The influence of herbicide combinations using single fluridone applications on Palmer amaranth seed production, Palmer amaranth biomass, and rice grain yield, averaged over four site-years<sup>a,b,c</sup>.

Herbicides	Timing	Rate	SP	Biomass	Yield
		g ai/ae ha <sup>-1</sup>	seed m <sup>-2</sup>	g m <sup>-2</sup>	kg ha <sup>-1</sup>
Nontreated	–	–	72,380 a	350 a	4,020 b
Clomazone + Fluridone	PRE	336	23,496 b	218 b	6,600 a
Clomazone + Fluridone	PRE	336	1,285 c	9 c	6,560 a
Florpyrauxifen-benzyl	MPOST	15			
Clomazone + Fluridone	PRE	336	921 d	0.2 e	6,520 a
Florpyrauxifen-benzyl	MPOST	30			
Clomazone	PRE	336	520 e	2.0 de	6,380 a
Florpyrauxifen-benzyl	EPOST	15			
Florpyrauxifen-benzyl	MPOST	15			
Clomazone	PRE	336	151 f	3.6 d	6,540 a
Fluridone + Florpyrauxifen-benzyl	EPOST	168			
Florpyrauxifen-benzyl	EPOST	15			
Florpyrauxifen-benzyl	MPOST	15			
Clomazone	PRE	336	86 f	5.8 cd	6,590 a
Florpyrauxifen-benzyl	EPOST	15			
Fluridone + Florpyrauxifen-benzyl	MPOST	168			
P-value			<0.0001	<0.0001	0.0044

<sup>a</sup>Abbreviations: EPOST, early postemergence; MPOST, mid-postemergence; PRE, preemergence; SP, seed production.

<sup>b</sup>Means within the same column followed by the same letter are not different according to the Sidak method ( $\alpha = 0.05$ ); the absence of letters indicates no treatment difference was present.

<sup>c</sup>All florpyrauxifen-benzyl applications included 0.58 L ha<sup>-1</sup> methylated seed oil.

**Table 9.** Visible Palmer amaranth control and rice injury for the multiple fluridone application experiment 21 d after preemergence, averaged over 2022 and 2023<sup>a,b</sup>.

Herbicides	Rate	AMAPA control	Rice injury
	g ai ha <sup>-1</sup>	—% of nontreated—	
Clomazone	336	43 b	18 b
Clomazone + fluridone	336 + 84	85 a	29 ab
Clomazone + fluridone	336 + 168	88 a	32 a
Clomazone + fluridone	336 + 84	84 a	26 ab
Clomazone + fluridone	336 + 84	85 a	28 ab
Clomazone + fluridone	336 + 84	85 a	27 ab
Clomazone + fluridone	336 + 168	90 a	36 a
Clomazone + fluridone	336 + 84	91 a	27 ab
P-value		<0.0001	0.0472

<sup>a</sup>Abbreviations: AMAPA, Palmer amaranth; PRE, preemergence.

<sup>b</sup>Means within a column followed by the same letter are not different according to the Sidak method ( $\alpha = 0.05$ ); the absence of letters indicates no treatment difference was present.

integrating fluridone into herbicide programs will add value when targeting Palmer amaranth. However, fluridone is not currently labeled for preemergence use in rice production systems. Additionally, these results indicate the need for effective postemergence herbicides, such as florpyrauxifen-benzyl, for enhanced Palmer amaranth control in a furrow-irrigated rice system.

Herbicide treatment also affected rice injury 14 d after mid-postemergence, with injury ranging from 13% to 39% ( $P < 0.0001$ ; Table 10). Rice injury was greatest when fluridone was applied preemergence at 168 g ai ha<sup>-1</sup> in combination with clomazone. Fluridone was less injurious to rice when applied at a rate of 84 g ai ha<sup>-1</sup>, even if the treatment included a sequential application of the herbicide mid-postemergence at the same rate. A previous study conducted on a precision-leveled field also reported greater rice injury when fluridone was applied to 3-leaf rice at a 1× rate vs. a 0.5× rate 4 wk after application (Butts et al. 2024). Additionally,

preemergence applications of fluridone at the 0.5× rate in combination with clomazone followed by florpyrauxifen-benzyl applied mid-postemergence does not increase rice injury compared with preemergence treatments that did not contain fluridone. There were also no differences in injury with clomazone + fluridone preemergence treatments compared with an identical preemergence treatment followed by an mid-postemergence application of florpyrauxifen-benzyl. Overall, greater rice injury was observed with fluridone applied at a 1× rate, which may be attributed to the extensive persistence of the herbicide in the soil, as reported by others (Banks et al. 1979).

The final herbicide application occurred late postemergence, and visible Palmer amaranth control ratings differed as a function of herbicide program 14 and 28 d after late postemergence (Table 11). At 14 d after late postemergence, herbicide combinations that contained sequential florpyrauxifen-benzyl applications achieved 90% to 96% Palmer amaranth control. Relative to those treatments, preemergence applications of clomazone with fluridone followed by a single late postemergence application of florpyrauxifen-benzyl were less effective. These results suggest that multiple applications of fluridone or florpyrauxifen-benzyl should be used to optimize Palmer amaranth control. At 28 d after late postemergence, herbicide programs that entailed multiple applications of fluridone and florpyrauxifen-benzyl achieved 97% to 98% control of Palmer amaranth. Hence, applying multiple herbicides with different modes of action is among the best management practices when targeting problematic weed species, such as Palmer amaranth (Norsworthy et al. 2012).

Herbicide programs that contained multiple applications of fluridone and florpyrauxifen-benzyl caused up to 29% rice injury 14 DAT, a percentage that was comparable to all other programs evaluated in the experiment (Table 11). Similarly, sequential fluridone applications did not exacerbate rice injury compared with single applications of fluridone and florpyrauxifen-benzyl 28 DAT. These injury data suggest that compared to commercial

**Table 10.** Visible Palmer amaranth control and rice injury for the multiple fluridone application experiment 14 d after mid-postemergence, averaged over 2022 and 2023<sup>a,b,c</sup>.

Herbicides	Timing	Rate g ai/ae ha <sup>-1</sup>	AMAPA control		Rice injury
			% of nontreated		
Clomazone	PRE	336	87 b		13 c
Florpyrauxifen-benzyl	MPOST	15			
Clomazone + fluridone	PRE	336 + 84	94 ab		16 bc
Florpyrauxifen-benzyl	MPOST	15			
Clomazone + fluridone	PRE	336 + 168	96 a		39 a
Florpyrauxifen-benzyl	MPOST	15			
Clomazone + fluridone	PRE	336 + 84	92 ab		25 b
Florpyrauxifen-benzyl + fluridone	MPOST	15 + 84			
Clomazone + fluridone	PRE	336 + 84	92 ab		17 bc
Florpyrauxifen-benzyl	MPOST	15			
Clomazone + fluridone	PRE	336 + 84	68 c		14 bc
Clomazone + fluridone	PRE	336 + 168	94 ab		37 a
Clomazone + fluridone	PRE	336 + 84	75 c		20 bc
P-value			<0.0001		<0.0001

<sup>a</sup>Abbreviations: AMAPA, Palmer amaranth; MPOST, mid-postemergence PRE, preemergence.

<sup>b</sup>Means within a column followed by the same letter are not different according to the Sidak method ( $\alpha = 0.05$ ).

<sup>c</sup>All florpyrauxifen-benzyl applications included 0.58 L ha<sup>-1</sup> methylated seed oil.

**Table 11.** Main effect of herbicide program on visible Palmer amaranth control and rice injury 14 and 28 DALPOST, averaged over 2022 and 2023<sup>a,b,c</sup>.

Herbicides	Timing	Rate g ai/ae ha <sup>-1</sup>	AMAPA control		Rice injury	
			14 DAT	28 DAT	14 DAT	28 DAT
Clomazone	PRE	336	90 abc	92 bc	16 b	12 b
Florpyrauxifen-benzyl	MPOST	15				
Florpyrauxifen-benzyl	LPOST	15				
Clomazone +	PRE	336	92 abc	96 ab	13 b	20 ab
Fluridone	PRE	84				
Florpyrauxifen-benzyl	MPOST	15				
Florpyrauxifen-benzyl	LPOST	15				
Clomazone +	PRE	336	96 a	96 ab	35 a	26 ab
Fluridone	PRE	168				
Florpyrauxifen-benzyl	MPOST	15				
Florpyrauxifen-benzyl	LPOST	15				
Clomazone +	PRE	336	94 ab	97 a	29 ab	25 ab
Fluridone	PRE	84				
Florpyrauxifen-benzyl +	MPOST	15				
Fluridone	MPOST	84				
Florpyrauxifen-benzyl	LPOST	15				
Clomazone +	PRE	336	94 ab	98 a	26 ab	17 ab
Fluridone	PRE	84				
Florpyrauxifen-benzyl	MPOST	15				
Florpyrauxifen-benzyl +	LPOST	15				
Fluridone	LPOST	84				
Clomazone +	PRE	336	80 d	87 c	20 ab	13 b
Fluridone	PRE	84				
Florpyrauxifen-benzyl	LPOST	15				
Clomazone +	PRE	336	85 cd	93 abc	36 a	33 a
Fluridone	PRE	168				
Florpyrauxifen-benzyl	LPOST	15				
Clomazone +	PRE	336	88 bcd	93 abc	17 b	18 ab
Fluridone	PRE	84				
Florpyrauxifen-benzyl +	LPOST	15				
Fluridone	LPOST	84				
P-value			<0.0001	0.0022	0.0267	0.0409

<sup>a</sup>Abbreviations: AMAPA, Palmer amaranth; DALPOST, days after late postemergence; DAT, days after treatment; LPOST, late postemergence; MPOST, mid-postemergence; PRE, preemergence.

<sup>b</sup>Means within a column followed by the same letter are not different according to the Sidak method ( $\alpha = 0.05$ ).

<sup>c</sup>All florpyrauxifen-benzyl applications included 0.58 L ha<sup>-1</sup> methylated seed oil.

standards, extended weed control can be achieved with more herbicide applications without causing additional rice injury.

Averaged across site-years, rough rice yields differed among the herbicide programs (Table 12). Rice yield was lowest from the

nontreated control at 3,580 kg ha<sup>-1</sup>. Rice yields were similar for all other herbicide programs and ranged from 7,540 to 8,260 kg ha<sup>-1</sup>.

The consistent grain yield across treatments is likely attributed to each program having an late postemergence application of



**Table 12.** Influence of different herbicide combinations using single and multiple applications of fluridone on Palmer amaranth seed production, Palmer amaranth biomass, and rice grain yield averaged over 2022 and 2023<sup>a,b,c</sup>.

Herbicides	Timing	Rate	SP	Biomass	Yield
		g ai/ae ha <sup>-1</sup>	seed m <sup>-2</sup>	g m <sup>-2</sup>	kg ha <sup>-1</sup>
Nontreated	–	–	30,700 a	166 a	3,580 b
Clomazone	PRE	336	8,273 b	39 b	7,890 a
Florpyrauxifen-benzyl	MOST	15			
Florpyrauxifen-benzyl	LPOST	15			
Clomazone +	PRE	336	8,216 b	25 bc	7,860 a
Fluridone	PRE	84			
Florpyrauxifen-benzyl	MPOST	15			
Florpyrauxifen-benzyl	LPOST	15			
Clomazone +	PRE	336	8,019 b	17 cd	7,740 a
Fluridone	PRE	168			
Florpyrauxifen-benzyl	MPOST	15			
Florpyrauxifen-benzyl	LPOST	15			
Clomazone +	PRE	336	0 f	0 d	7,540 a
Fluridone	PRE	84			
Florpyrauxifen-benzyl +	MPOST	15			
Fluridone	MPOST	84			
Florpyrauxifen-benzyl	LPOST	15			
Clomazone +	PRE	336	0 f	0 d	8,110 a
Fluridone	PRE	84			
Florpyrauxifen-benzyl	MPOST	15			
Florpyrauxifen-benzyl +	LPOST	15			
Fluridone	LPOST	84			
Clomazone +	PRE	336	1,355 e	12 cd	8,120 a
Fluridone	PRE	84			
Florpyrauxifen-benzyl	LPOST	15			
Clomazone +	PRE	336	1,722 d	13 cd	7,580 a
Fluridone	PRE	168			
Florpyrauxifen-benzyl	LPOST	15			
Clomazone +	PRE	336	5,642 c	41 b	8,260 a
Fluridone	PRE	84			
Florpyrauxifen-benzyl +	LPOST	15			
Fluridone	LPOST	84			
P-value			<0.0001	<0.0001	<0.0001

<sup>a</sup>Abbreviations: LPOST, late postemergence; MPOST, mid-postemergence; PRE, preemergence; SP, seed production.

<sup>b</sup>Means within the same column followed by the same letter are not different according to the Sidak method ( $\alpha = 0.05$ ).

<sup>c</sup>All florpyrauxifen-benzyl applications included 0.58 L ha<sup>-1</sup> methylated seed oil.

florpyrauxifen-benzyl. Another study also observed maximum rice grain yield with late postemergence applications of florpyrauxifen-benzyl in a furrow-irrigated rice system (Wright et al. 2020). Since fluridone provides effective early-season control of Palmer amaranth (Grichar et al. 2020; Hill et al. 2017), rice will likely have a competitive advantage in suppressing additional Palmer amaranth seedlings due to canopy formation. Hence, fewer postemergence herbicides may be required to control the weed sufficiently. Additionally, the visual injury observed with both fluridone and florpyrauxifen-benzyl did not have a long-term effect on rice yield by the end of the growing season.

Palmer amaranth biomass accumulation differed among the herbicide programs (Table 12). Compared with the nontreated control, all herbicide programs successfully reduced Palmer amaranth biomass production. Additionally, those programs that included a preemergence and postemergence application of fluridone and sequential postemergence applications of florpyrauxifen-benzyl had the greatest impact on Palmer amaranth biomass production, allowing no weeds to escape in those plots at the time of rice harvest in both site-years. These results are unsurprising, considering fluridone and florpyrauxifen-benzyl both exhibited noteworthy control of Palmer amaranth when applied preemergence and postemergence, respectively.

Likewise, Palmer amaranth seed production was affected by herbicide program (Table 12), and the nontreated control

produced the greatest number of Palmer amaranth seed at 30,700 seed m<sup>-2</sup>. In recent years, a zero-tolerance threshold approach has been widely recommended for long-term management of Palmer amaranth and preserving herbicide efficacy (Bagavathiannan and Norsworthy 2012; Norsworthy et al. 2012), in which no weeds are allowed to escape control and produce seed (Norris 2007; Norsworthy et al. 2014). Considering there was no Palmer amaranth present in plots where fluridone and florpyrauxifen-benzyl were applied multiple times, no offspring were produced at rice harvest after those treatments. Conversely, Beesinger et al. (2022) found that sequential florpyrauxifen-benzyl applications still allowed Palmer amaranth to produce viable seed by rice harvest. Therefore, findings from this research indicate that multiple applications of fluridone in combination with sequential postemergence applications of florpyrauxifen-benzyl will be advantageous in reducing the number of weed seeds returned to the soil seedbank, further supporting the zero-tolerance threshold approach.

### Practical Implications

This research highlights the ability of fluridone to provide excellent Palmer amaranth control when integrated into herbicide programs targeting the weed in a furrow-irrigated rice system. However, preemergence fluridone applications caused greater rice injury

than herbicide treatments that excluded the herbicide; therefore, producers should not apply fluridone until the V3 rice growth stage, as stated on the herbicide label. Both experiments were placed at the higher end of the field where the soil is drier relative to other portions of the field; therefore, rice injury may be less where there is less moisture compared to the bottom end of a field. Due to its high persistence on a silt loam soil, severe injury from fluridone may occur in areas of a field where water from irrigation or rainfall collects (Banks et al. 1979) and in fields that have been previously precision-leveled (Butts et al. 2024). Overall, the results reported here display the flexibility of applying fluridone as a postemergence, residual herbicide along with floryprauxifen-benzyl for Palmer amaranth control on a silt loam soil. Furthermore, using fluridone at labeled rates does not appear to translate into persistent, season-long rice injury, yet it effectively controls Palmer amaranth while preserving rice yield. Although Palmer amaranth biomass and seed production was reduced with herbicide programs that included fluridone, the weed was still present at harvest in most instances; hence, producers must remain aware of the offspring and sufficient seedbank replenishment potential of the weed, which facilitates the spread of herbicide-resistant genes in future growing seasons.

**Acknowledgments.** This research was conducted in cooperation with SePRO Corporation. The University of Arkansas System Division of Agriculture supplied facilities and equipment.

**Funding.** SePRO Corporation and the Arkansas Rice Research and Promotion Board funded this research.

**Competing Interests.** The authors declare they have no competing interests.

## References

- [ASPB] Arkansas State Plant Board (2020) Arkansas pesticide use and application act and rules. Act 389. Little Rock: Arkansas Department of Agriculture, Arkansas State Plant Board. 21 p
- Anderson IC, Roberston DS (1960) Role of carotenoids in protecting chlorophyll from photodestruction. *Plant Physiol* 35:531–534
- Anonymous (2023) Brake® herbicide product label. Carmel, IN, US: SePRO Corporation. [https://www3.epa.gov/pesticides/chem\\_search/ppls/067690-00078-20230124.pdf](https://www3.epa.gov/pesticides/chem_search/ppls/067690-00078-20230124.pdf). Accessed: April 18, 2024
- Bagavathiannan MV, Norsworthy JK (2012) Late-season seed production in arable weed communities: management implications. *Weed Sci* 60:325–334
- Bagavathiannan MV, Norsworthy JK, Scott RC (2011) Comparison of weed management program for furrow-irrigated and flooded hybrid rice production in Arkansas. *Weed Technol* 25:556–562
- Bagavathiannan MV, Norsworthy JK, Scott RC, Barber LT (2013) Answers to frequently asked questions on herbicide resistance management. Little Rock: University of Arkansas Division of Agriculture Fact Sheet FSA2172. <https://www.uaex.uada.edu/publications/PDF/FSA-2172.pdf>. Accessed: September 12, 2023
- Banks PA, Merkle MG (1979) Field evaluations of the herbicidal effects of fluridone on two soils. *Agron J* 71:759–762
- Barber LT, Butts TR, Boyd JW, Cunningham K, Selden G, Norsworthy JK, Burgos NR, and Bertucci M (2023) Pages 80–115 in MP44: Recommended chemicals for weed and brush control. Little Rock: University of Arkansas System Division of Agriculture Cooperative Extension Service
- Barber LT, Smith KL, Scott RC, Norsworthy JK, Vangilder AM (2015) Zero tolerance: a community-based program for glyphosate-resistant Palmer amaranth management. University of Arkansas Cooperative Extension. <https://www.uaex.uada.edu/publications/pdf/FSA2177.pdf>. Accessed: September 12, 2023
- Beesinger JW, Norsworthy JK, Butts TR, Roberts TL (2022) Palmer amaranth control in furrow-irrigated rice with floryprauxifen-benzyl. *Weed Technol* 36:490–496
- Bensch CN, Horak MJ, Peterson D (2003) Interference of redroot pigweed (*Amaranthus retroflexus*), Palmer amaranth (*A. palmeri*), and common waterhemp (*A. rudis*) in soybean. *Weed Sci* 51:37–43
- Blouin DC, Webster EP, Bond JA (2011) On the analysis of combined experiments. *Weed Technol* 25:165–169
- Bond JA, Oliver LR, Stephenson DO IV (2006) Response of Palmer amaranth (*Amaranthus palmeri*) accessions to glyphosate, fomesafen, and pyriproxyfen. *Weed Technol* 20:885–892
- Brooks ME, Kristensen K, Van Benthem KJ, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Machler M, Bolker BM (2017) GlmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *R J* 9:378–400
- Burgos NR, Butts TR, Werle IS, Bottoms S, Mauromoustakos A (2021) Weed rice update in Arkansas, USA, and adjacent locales. *Weed Sci* 69:514–525
- Butts TR, Kouame KB-J, Norsworthy JK, Barber LT (2022) Arkansas rice: herbicide resistance concerns, production practices, and weed management costs. *Front Agron* 4:881667
- Butts TR, Souza MCCR, Norsworthy JK, Barber JK, Hardke JT (2024) Rice response to fluridone following topsoil removal on a precision-leveled field. *Agrosyst Geosci Environ* 7:e20541
- Chlapeccka JL, Hardke JT, Roberts TL, Mann MG, Ablao A (2021) Scheduling rice irrigation using soil moisture thresholds for furrow irrigation and intermittent flooding. *Agron J* 113:1258–1270
- Counce PA, Keisling TC, Mitchell AJ (2000) A uniform, objective, and adaptive system for expressing rice development. *Crop Sci* 40:436–443
- Duke SO, Paul RN, Becerril JM, Schmidt JH (1991) Clomazone causes accumulation of sesquiterpenoids in cotton (*Gossypium hirsutum* L.). *Weed Sci* 39:339–346
- Fox J, Weisberg S (2019) Nonlinear Regression, Nonlinear Least Squares, and Nonlinear Mixed Models in R. An R Companion to Applied Regression. 3rd ed. Thousand Oaks, CA: Sage Publications 608 p
- Gbur EE, Stroup WW, McCarter KS, Durham SL, Young LJ, Christman MC, West M, Kramer M (2012) Analysis of generalized linear mixed models in the agricultural and natural resources sciences. Madison, WI: American Society of Agronomy, Soil Science Society of America, Crop Science Society of America
- Gossett BJ, Murdock EC, Toler JE (1992) Resistance of Palmer amaranth (*Amaranthus palmeri*) to the dinitroaniline herbicides. *Weed Technol* 6:587–591
- Grichar WJ, Dotray P, McGinty J (2020) Using fluridone herbicide systems for weed control in Texas cotton (*Gossypium hirsutum* L.). *J Adv Agric* 11:1–14
- Hardke JT (2022) Trends in Arkansas rice production, 2022. B.R. Wells Arkansas Rice Research Studies 2022. Little Rock: University of Arkansas System Division of Agriculture, Cooperative Extension Service
- Heap I (2024) International survey of herbicide-resistant weed database. <http://www.weedscience.org/Pages/Case.aspx?ResistID=18156>. Accessed: April 29, 2024
- Hill ZT, Norsworthy JK, Barber LT, Gbur EE (2017) Assessing the potential for fluridone to reduce the number of postemergence herbicide applications in glyphosate-resistant cotton. *J Cotton Sci* 21:175–182
- Hill ZT, Norsworthy JK, Barber LT, Roberts TL, Gbur EE (2016) Assessing the potential for fluridone carryover to six crop rotated with cotton. *Weed Technol* 30:346–354
- Hothorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models. *Biom J* 50:346–363
- Hwang JI, Norsworthy JK, Piveta LB, Souza MCCR, Barber LT, Butts TR (2023) Metabolism of 2,4-D in resistant *Amaranthus palmeri* S. Wats. (Palmer amaranth). *Crop Prot* 165:106169
- Klingaman TE, Oliver LR (1994) Palmer amaranth (*Amaranthus palmeri*) interference in soybean (*Glycine max*). *Weed Sci* 42:525–527
- Kraehmer H, Jabran K, Mennan H, Chauhan BS (2016) Global distribution of rice weeds – a review. *Crop Prot* 80:73–86
- Lunga DD, Brye KR, Henry CG, Slayden JM (2021) Plant productivity and nutrient uptake as affected by tillage and site-position in furrow-irrigated rice. *Agron J* 113:2374–2386

- Martin SM, Norsworthy JK, Scott RC, Hardke J, Gbur E (2018) Effect of thiamethoxam on injurious herbicide in rice. *Adv Crop Sci Technol*. doi: [10.4172/2329-8863.1000351](https://doi.org/10.4172/2329-8863.1000351)
- Massinga RA, Currie RS, Horak MJ, Boyer J (2001) Interference of Palmer amaranth in corn. *Weed Sci* 49:202–208
- McLean RA, Sanders WL, Stroup WW (1991) A unified approach to mixed linear models. *Am Stat* 45:54–64
- Midway S (2022) Random Effects. Page 174 in *Data Analysis in R*. [https://bookdown.org/steve\\_midway/DAR/random-effects.html#fixed-and-random-effects](https://bookdown.org/steve_midway/DAR/random-effects.html#fixed-and-random-effects)
- Midway S, Robertson M, Flinn S, Kaller M (2020) Comparing multiple comparisons: practical guidance for choosing the best multiple comparisons test. *Peer J* 8:e10387 <https://doi.org/10.7717/peerj.10387>
- Miranda JWA, Jhala AJ, Bradshaw J, Lawrence NC (2021) Palmer amaranth (*Amaranthus palmeri*) interference and seed production in dry edible bean. *Weed Technol* 35:996–1006
- Nalley LL, Massey J, Durand-Morat A, Shew A, Parajuli R, Tsiboe F (2022) Comparative economic and environmental assessments of furrow- and flood-irrigated rice production systems. *Agr Water Manage* 274:107964
- Norris RF (2007) Weed fecundity: current status and future needs. *Crop Prot* 26:182–188
- Norsworthy JK, Griffith GM, Griffin T, Bagavathiannan M, Gbur EE (2014) In-field movement of glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) and its impact on cotton lint yield: evidence supporting a zero-threshold strategy. *Weed Sci* 62:237–249
- Norsworthy JK, Griffith GM, Scott RC (2008) Imazethapyr use with and without clomazone for weed control in furrow-irrigated, imidazolinone-tolerant rice. *Weed Technol* 22:217–221
- Norsworthy JK, Scott RC, Bangarwa SK, Griffith GM, Wilson MJ, McClelland M (2011) Weed management in a furrow-irrigated imidazolinone-resistant hybrid rice production system. *Weed Technol* 25:25–29
- Norsworthy JK, Ward SM, Shaw DR, Llewellyn RS, Nichols RL, Webster TM, Bradley KW, Frisvold G, Powles SB, Burgos NR, Witt WW, Barrett M (2012) Reducing the risks of herbicide resistance: best management practices and recommendations. *Weed Sci* 60(SP1):31–62
- Priess GL, Norsworthy JK, Godara N, Mauromoustakos A, Butts TR, Roberts TL, Barber T (2022) Confirmation of glufosinate-resistant Palmer amaranth and response to other herbicides. *Weed Technol* 36:368–372
- R Core Team (2021) R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing
- Rowland MW, Murray DS, Verhalen LM (1999) Full-season Palmer amaranth (*Amaranthus palmeri*) interference with cotton (*Gossypium hirsutum*). *Weed Sci* 47:305–309
- Searle SR, Speed FM, Milliken GA (1980) Population marginal means in the linear model: an alternative to least squares means. *Am Stat* 34:216–221
- Stroup WW (2015) Rethinking the analysis of non-normal data in plant and soil science. *Agron J* 107:811–827
- Van Wychen L (2022) Survey of the most common and troublesome weeds in broadleaf crops, fruits, and vegetables in the United States and Canada. Westminster, CO: Weed Science Society of America. [http://wssa.net/wp-content/uploads/2022-Weed-Survey\\_Broadleaf-crops.xlsx](http://wssa.net/wp-content/uploads/2022-Weed-Survey_Broadleaf-crops.xlsx). Accessed: September 11, 2023
- Varanasi VK, Brabham C, Korres NE, Norsworthy JK (2019) Nontarget site resistance in Palmer amaranth [*Amaranthus palmeri* (S.) Wats.] confers cross-resistance to protoporphyrinogen oxide-inhibiting herbicides. *Weed Technol* 33:349–354
- Waldrep TW, Taylor HM (1976) 1-Methyl-3-phenyl-5-[3-(trifluoromethyl)phenyl]-4(1H)-pyridinone, a new herbicide. *J Agric Food Chem* 24:1250–1251
- Webster TM, Nichols RL (2012) Changes in the prevalence of weed species in the major agronomic crops of the southern United States: 1994/1995 to 2008/2009. *Weed Sci* 60:145–157
- Wright HE, Norsworthy JK, Roberts TL, Scott RC, Hardke JT, Gbur EE (2020) Use of florypyrauxifen-benzyl in non-flooded rice production systems. *CFTM* 7:e20081. doi: [10.1002/cft2.20081](https://doi.org/10.1002/cft2.20081)
- Zhang W, Webster EP, Blouin DC (2005) Response of rice and barnyardgrass (*Echinochloa crus-galli*) to rates and timings of clomazone. *Weed Technol* 19:528–531