

QUANTUM MARKOV CHAINS AND CLASSICAL RANDOM SEQUENCES

YUN-GANG LU

§1. Introduction

Quantum Markov chain introduced by Accardi (cf. [1,2,3]) is one of natural generalization of classical Markov chain. It has many interesting applications in physics and the most important one is given by the paper of Fannes-Nachtergaele-Werner ([4]), where an application of quantum Markov chain's technique enables us to understand the Valence bond states well.

In this note, by the terminology of *quantum Markov chain* we follow Accardi's definition (cf. [1] or [3]):

Let be given a C^* -algebra $\mathcal{B} = \mathbf{B}(H)$ for some fixed separable Hilbert space H and $\mathcal{A} := \bigotimes_{\mathbf{N}} \mathcal{B}$ (where, \otimes means the C^* -tensor induced by the usual tensor product of Hilbert space). For each $n \in \mathbf{N}$, let be given a homomorphism

$$j_n: \mathcal{B} \hookrightarrow 1_0 \otimes 1_1 \otimes \cdots \otimes 1_{n-1} \otimes \mathcal{B} \otimes 1_{n+1} \otimes 1_{n+2} \otimes \cdots =: 1_n \otimes \mathcal{B} \otimes 1_n$$

such that

$$j_n(\alpha) := 1_n \otimes \alpha \otimes 1_n, \quad \forall \alpha \in \mathcal{B}.$$

A bilinear map \mathcal{E} from $\mathcal{B} \otimes \mathcal{B}$ to \mathcal{B} is called a transition expectation if it is completely positive and identity preserving.

With above notations, quantum Markov chain (or quantum Markovian state) is defined on $\mathcal{A} := \bigotimes_{\mathbf{N}} \mathcal{B}$ as the following: a state ϕ on \mathcal{A} is called Markovian state if there exists a state ϕ_0 (initial state) on \mathcal{B} and a transition expectation \mathcal{E} such that

$$(1.1) \quad \phi(j_0(a_0)j_1(a_1)\cdots j_n(a_n)) = \phi_0(\mathcal{E}(a_0) \otimes \mathcal{E}(a_1) \otimes \cdots \otimes \mathcal{E}(a_{n-1}) \otimes \mathcal{E}(a_n \otimes 1) \cdots).$$

It is proved (cf. [3]) that any transition expectation \mathcal{E} has the form

Received August 5, 1992.
 Revised October 4, 1994.

$$(1.2) \quad \mathcal{E}(x) = \text{Tr}_2(\sum_j K_j^* x K_j)$$

where, Tr_2 means the partial trace with respect to the second factor; $K_j \in \mathcal{B} \otimes \mathcal{B}$, $j = 1, 2, \dots$. Moreover, the index j runs over a finite subset of \mathbf{N} if H is a finite dimensional Hilbert space.

Throughout this note we restrict ourselves to the case of $\dim(H) = d < \infty$. Since for each orthonormal basis (o.n.b.) $\{e_n\}_{n=1}^d$ of H , $\mathbf{B}(H)$ can be considered as the algebra of all $d \times d$ matrices, i.e. $\mathbf{B}(H) \cong M_d$, the further assumption is made: $\mathcal{B} = M_d$.

For each $K \in \mathcal{B} \otimes \mathcal{B}$, we have the expression

$$(1.3) \quad K = \sum_{h,h'} |e_h\rangle\langle e_{h'}| \otimes K_{h,h'}$$

where $|e_h\rangle\langle e_{h'}|$ means the matrix whose (h, h') -element is equal to 1 and elsewhere zero. In the following by the notation \mathcal{D} we shall denote the diagonal subalgebra of M_d (corresponding to $\{e_n\}_{n=1}^d$).

With above definitions and notations, each $\{1, \dots, d\}$ -valued classical Markov chain $\{X_n\}_{n=1}^\infty$ on a propability space $(\mathbf{P}, \mathcal{Q})$ with initial distribution \mathbf{p} and transition matrix $\mathbf{P} = (p_{i,j})$ can be understood as a quantum Markov chain in the following canonical way:

i) each bounded function on $\{1, \dots, d\}$, say f , will be regarded as an element of \mathcal{D} :

$$(1.4) \quad \begin{pmatrix} f(1) & 0 & \cdots & 0 \\ 0 & f(2) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & f(d) \end{pmatrix}$$

ii) for each $n \in \mathbf{N}$, let us define a homomorphism

$$(1.5) \quad j_n(f) := f(X_n), \quad \forall f \in \mathcal{D}$$

iii) define

$$(1.6) \quad K_{1,h,h'} := \begin{pmatrix} \sqrt{p_{h,1}} & 0 & \cdots & 0 \\ 0 & \sqrt{p_{h,2}} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sqrt{p_{h,d}} \end{pmatrix}$$

$$(1.7a) \quad K_{1,h,h'} := 0, \quad \text{if } h \neq h'$$

$$(1.7b) \quad K_{n,h,h'} := 0, \quad \forall n \geq 2$$

iv) define initial state ϕ_0 by

$$(1.8) \quad \phi_0(\cdot) := \text{Tr} \left(\begin{pmatrix} p_1 & 0 & \cdots & 0 \\ 0 & p_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & p_d \end{pmatrix} \cdot \right).$$

It is easy to check that

$$(1.9) \quad \mathcal{E}_1(x) := \text{Tr}_2(K_1^* \cdot K_1)$$

where $K_1 := \sum_h |e_h\rangle\langle e_h| \otimes K_{1,h,h}$, is a transition expectation with the property:

$$(1.10) \quad \mathcal{E}_1(f \otimes g) = f \cdot \mathcal{E}_1(1 \otimes g) \quad \forall f, g \in \mathcal{D}$$

and ϕ_0 is a state on M_d . Therefore (ϕ_0, \mathcal{E}_1) is a quantum Markov chain and moreover the restriction of this quantum Markov chain to the diagonal subalgebra \mathcal{D} is nothing but the given classical Markov chain.

From above discussion one can see that the choice of (ϕ_0, \mathcal{E}) is not unique. In fact we can replace the initial density matrix

$$\begin{pmatrix} p_1 & 0 & \cdots & 0 \\ 0 & p_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & p_d \end{pmatrix}$$

by any density matrix ω_0 which has diagonal elements $\{p_1, p_2, \dots, p_d\}$. Of course there are many such density matrices. Moreover we can replace $K_{1,h,h}$ by any $K_{h,h}$ which has only to possess the property:

$$(1.11) \quad K_{h,h}K_{h,h}^*(j, j) = p_{h,j}, \quad \forall j = 1, 2, \dots, d$$

Let denote

$$K_0 := \sum_h |e_h\rangle\langle e_h| \otimes K_{h,h}$$

and define

$$\mathcal{E}_0(\cdot) := \text{Tr}_2(K_0^* \cdot K_0).$$

Then $(\text{Tr}(\omega_0 \cdot), \mathcal{E}_0(\cdot))$ is also a quantum Markov chain obtained from the same classical Markov chain $(p, P = (p_{i,j}))$ and moreover, by restricting the two quantum Markov chains $(\text{Tr}(\omega_0 \cdot), \mathcal{E}_0(\cdot))$ and (ϕ_0, \mathcal{E}_1) on \mathcal{D} , one obtains the same classical Markov chain $(p, P = (p_{i,j}))$.

Now a natural problem can be asked: starting from a classical Markov chain $(p, P = (p_{i,j}))$, what is the most general form of the quantum form of the quantum Markov chains whose restriction to the diagonal subalgebra gives the classical Markov chain? One can also ask the problem reversely: which kind of quantum Markov chain gives, by restricting it to diagonal subalgebra, a classical Markov chain?

Moreover, it is usually believed that some non-Markovian stochastic processes (classical) are quantum Markov chain. Then a natural question is this: which kind of classical stochastic processes can be a quantum Markov chain? In order to answer this question, first of all we have to understand which kind of quantum Markov chain, by restricting it to diagonal subalgebra \mathcal{D} , gives a classical stochastic process in the sense of \mathcal{E} maps $\mathcal{D} \otimes \mathcal{D}$ to \mathcal{D} ?

Remark. Notice that if

$$(1.12) \quad \mathcal{E}(\mathcal{D} \otimes \mathcal{D}) \subset \mathcal{D}$$

then we can consider the initial density matrix ω_0 as a diagonal one since its non-diagonal elements don't play any rule in this case.

In this note we give the answer of above questions. Our main results are stated in the next section.

§2. The main results

In the section 1, we opened the problems and gave their motivations. Now we shall state the answer of the problems. The proof of our main results is the contents of the next section.

PROPOSITION (2.1). *A transition expectation \mathcal{E} maps $\mathcal{D} \otimes \mathcal{D}$ to \mathcal{D} if and only if for each $r, r', h \in \{1, \dots, d\}$ with $r \neq r'$, the matrix $\sum_j K_{j,h,r} K_{j,h,r'}^*$ has only zero diagonal elements.*

DEFINITION (2.2). We say that an $\{1, \dots, d\}$ -valued classical stochastic process $\{X_n\}_{n=0}^\infty$ on probability space (\mathbf{P}, Ω) is the quantum Markov chain (ϕ_0, \mathcal{E}) if \mathcal{E} satisfies (1.12) and the joint distributions are the same, i.e. for each $n \in \mathbf{N}$, $\{i_h\}_{h=0}^n \subset \{1, \dots, d\}$

$$(2.1) \quad \mathbf{P}(X_0 = i_0, X_1 = i_1, \dots, X_n = i_n)$$

$$\begin{aligned}
 &= \phi(|e_{i_0}\rangle\langle e_{i_0}| \otimes |e_{i_1}\rangle\langle e_{i_1}| \otimes \cdots \otimes |e_{i_n}\rangle\langle e_{i_n}|) \\
 &= \phi_0(\mathcal{E}(|e_{i_0}\rangle\langle e_{i_0}| \otimes \mathcal{E}(|e_{i_1}\rangle\langle e_{i_1}| \otimes \cdots \otimes \mathcal{E}(|e_{i_{n-1}}\rangle\langle e_{i_{n-1}}| \otimes \mathcal{E}(|e_{i_n}\rangle\langle e_{i_n}| \otimes 1)) \cdots))).
 \end{aligned}$$

Now we can state the results which replies the question: which kind of classical stochastic process can be a quantum Markov chain?

THEOREM (2.3). *Let $\{X_n\}_{n=1}^\infty$ be an $\{1, \dots, d\}$ -valued classical stochastic process on a probability space (\mathbf{P}, Ω) . Then it is a quantum Markov chain if and only if there exist a probability measure p_0 on $\{1, \dots, d\}$ and a 3-indices cubic matrix $(T_{j,i,k})$ with the following properties:*

$$(2.2a) \quad T_{j,i,k} \geq 0, \quad \forall j, i, k \in \{1, \dots, d\}$$

$$(2.2b) \quad \sum_{i,k} T_{j,i,k} = 1, \quad \forall j \in \{1, \dots, d\}$$

$$(2.2c) \quad \mathbf{P}(X_0 = j) = \sum_{i,k} p_0(i) T_{j,i,k}, \quad \forall j \in \{1, \dots, d\}$$

such that the joint distribution is given by:

$$\begin{aligned}
 (2.2d) \quad &\mathbf{P}(X_0 = i_0, X_1 = i_1, \dots, X_n = i_n) \\
 &= \sum_{j_0, j_1, \dots, j_n} p_0(j_0) T_{j_0, i_0, j_0} T_{j_0, i_1, j_1} \cdots T_{j_{n-2}, i_{n-1}, j_{n-1}} T_{j_{n-1}, i_n, j_n}
 \end{aligned}$$

THEOREM (2.4). *Let $\{X_n\}_{n=0}^\infty$ be an $\{1, \dots, d\}$ -valued classical stochastic process on a probability space (\mathbf{P}, Ω) . Then it is a quantum Markov chain if and only if there exists an $\{1, \dots, d\}^2$ -valued classical Markov chain $\{(Z_n, Y_n)\}_{n=1}^\infty$ on a probability space (\mathbf{P}', Ω') with the properties:*

i) the transition probability

$$(2.3a) \quad p_{(j,i),(j',i')} := \mathbf{P}'((Z_n, Y_n) = (j', i') \mid (Z_{n-1}, Y_{n-1}) = (j, i))$$

is independent of i ;

ii) there exists a probability distribution p_0 on $\{1, \dots, d\}$ and the initial distribution of $\{(Z_n, Y_n)\}_{n=0}^\infty$ is given by

$$(2.3b) \quad \mathbf{P}'((Z_0, Y_0) = (j, i)) = \sum_k p_0(k) T_{k,i,j}$$

(where $(T_{i,j,k})$ is determined by the process $\{X_n\}_{n=0}^\infty$ as stated in Theorem (2.3)) such that

$$(2.4) \quad \mathbf{P}(X_0 = i_0, X_1 = i_1, \dots, X_n = i_n) = \mathbf{P}'(Y_0 = i_0, Y_1 = i_1, \dots, Y_n = i_n).$$

COROLLARY. *If the quantum Markov chain (ϕ_0, \mathcal{E}) on $\otimes_{\mathbb{N}} M_d$ has the property that \mathcal{E} maps $\mathcal{D} \otimes \mathcal{D}$ to \mathcal{D} . Then the classical stochastic process, obtained by restricting (ϕ_0, \mathcal{E}) to the diagonal subalgebra \mathcal{D} , is stochastically equivalent to the second component of an $\{1, \dots, d\}^2$ -valued classical Markov chain $\{(Z_n, Y_n)\}_{n=0}^\infty$ which has the properties i), ii) in Theorem (2.4).*

This Corollary is a direct conclusion of Theorems (2.3) and (2.4).

DEFINITION (2.5). We say that the quantum Markov chain (ϕ_0, \mathcal{E}) is a classical Markov chain on the diagonal subalgebra \mathcal{D} if (1.12) is valid and

$$(2.5) \quad \mathcal{E}(\alpha \otimes b) = a \cdot \mathcal{E}(1 \otimes b), \quad \forall a, b \in \mathcal{D}.$$

THEOREM (2.6). *The quantum Markov chain (ϕ_0, \mathcal{E}) is a classical Markov chain on the diagonal subalgebra if and only if the operators $\{K_j\}$ (see (1.2)) have the form:*

$$(2.6) \quad K_j = \sum_h |e_h\rangle\langle e_h| \otimes K_{j,h,h}, \quad \forall j = 1, \dots, d$$

Remark. The formula (2.6) means that each K_j is a *block-diagonal* (not necessarily diagonal) $d^2 \times d^2$ matrix, i.e.

$$(2.7) \quad \begin{pmatrix} K_{j,1,1} & 0 & \cdots & 0 \\ 0 & K_{j,2,2} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & K_{j,d,d} \end{pmatrix}$$

where, each $K_{j,l,l}$ ($l = 1, \dots, d$) is a $d \times d$ matrix.

§3. The proof of the main results

This section is devoted to prove our results stated in the preceding section. The basic step is to make transition expectation in more clear form. From (1.3), one knows that for each $b \in M_d$,

$$\begin{aligned} (3.1) \quad \mathcal{E}(|e_i\rangle\langle e_m| \otimes b) &= \sum_j \text{Tr}_2(K_j^*(|e_i\rangle\langle e_m| \otimes b)K_j) \\ &= \sum_j \sum_{h,h',r,r'} \text{Tr}_2((|e_{h'}\rangle\langle e_r|^* \otimes K_{j,h',r}^*) (|e_i\rangle\langle e_m| \otimes b) (|e_h\rangle\langle e_r| \otimes K_{j,h,r})) \\ &= \sum_{h,h',r,r'} |e_r\rangle\langle e_{h'}| \cdot |e_i\rangle\langle e_m| \cdot |e_h\rangle\langle e_r| \text{Tr}(\sum_j K_{j,h',r}^* b K_{j,h,r}) \\ &= \sum_{r,r'} |e_r\rangle\langle e_r| \text{Tr}(\sum_j K_{j,i,r}^* b K_{j,m,r}). \end{aligned}$$

Since $\{|e_i\rangle\langle e_m| \in M_d : i, m \in \{1, \dots, d\}\}$ is a basis of M_d , we find that for any $a = (a(i, m)) \in M_d$,

$$(3.2) \quad \mathcal{E}(a \otimes b) = \sum_{i,m} a(i, m) \left[\sum_{r,r'} |e_{r'}\rangle\langle e_r| \operatorname{Tr}(\sum_j K_{j,i,r'}^* b K_{j,m,r}) \right].$$

By (3.2) and the fact that \mathcal{E} is identity preserving, one obtains that

$$(3.3) \quad \operatorname{Tr}(\sum_j K_{j,i,r'}^* b K_{j,i,r}) = \delta_{r,r'}, \quad \forall i \in \mathbf{N}.$$

Proof of Proposition (2.1). For any $|e_i\rangle\langle e_m|, |e_r\rangle\langle e_{m'}|$, (3.1) shows that

$$(3.4) \quad \mathcal{E}(|e_i\rangle\langle e_m| \otimes |e_r\rangle\langle e_{m'}|) = \sum_{r,r'} |e_r\rangle\langle e_r| \operatorname{Tr}(\sum_j K_{j,i,r'}^* \cdot |e_r\rangle\langle e_{m'}| K_{j,m,r}).$$

If \mathcal{E} maps $\mathcal{D} \otimes \mathcal{D}$ to \mathcal{D} , then for any m, i

$$(3.5) \quad \begin{aligned} \mathcal{E}(|e_m\rangle\langle e_m| \otimes |e_i\rangle\langle e_i|) &= \sum_{r,r'} |e_r\rangle\langle e_r| \operatorname{Tr}(\sum_j K_{j,m,r}^* \cdot |e_i\rangle\langle e_i| \cdot K_{j,m,r}) = \\ &= \sum_r |e_r\rangle\langle e_r| \operatorname{Tr}(\sum_j K_{j,m,r} K_{j,m,r}^* \cdot |e_i\rangle\langle e_i|). \end{aligned}$$

Therefore for any $r \neq r'$

$$(3.6) \quad \operatorname{Tr}(\sum_j K_{j,m,r} K_{j,m,r'}^* \cdot |e_i\rangle\langle e_i|) = \sum_j K_{j,m,r} K_{j,m,r'}^*(i, i) = 0.$$

This shows that \mathcal{E} maps $\mathcal{D} \otimes \mathcal{D}$ to \mathcal{D} only if for any $m, r \neq r'$, the matrix $\sum_j K_{j,m,r} K_{j,m,r'}^*$ has only zero diagonal elements.

On the other hand, if for any $m, r \neq r'$, the matrix $\sum_j K_{j,m,r} K_{j,m,r'}^*$ has only zero diagonal elements, (3.4), (3.5) and (3.6) guarantee that $\mathcal{E}(|e_m\rangle\langle e_m| \otimes b) \in \mathcal{D}$ for each $b \in \mathcal{D}$. Since $\{|e_m\rangle\langle e_m|\}_{m \in \mathbf{N}}$ is a basis of \mathcal{D} , it is obvious that $\mathcal{E}(a \otimes b) \in \mathcal{D}$ for each $a, b \in \mathcal{D}$. These arguments complete the proof.

Proof of Theorem (2.3). If classical stochastic process $\{X_n\}_{n=0}^\infty$ on a probability space $(\mathbf{P}, \mathcal{Q})$ is a quantum Markov chain, then by definition, there exists a pair (ϕ_0, \mathcal{E}) such that (1.12) and (2.1) are valid.

First of all, let see the initial distribution. By (1.3) and (2.1), for any $i_0 \in \{1, \dots, d\}$,

$$(3.7) \quad \mathbf{P}(X_0 = i_0) = \phi_0(\mathcal{E}(|e_{i_0}\rangle\langle e_{i_0}| \otimes 1)) = \phi_0(\sum_{r,r'} |e_r\rangle\langle e_r|) \cdot \sum_j \operatorname{Tr}(K_{j,i_0,r} K_{j,i_0,r}^*).$$

By (1.12), we know that (3.7) is equal to

$$(3.8) \quad \phi_0(\sum_r |e_r\rangle\langle e_r|) \cdot \sum_j \text{Tr}(K_{j,i_0,r} K_{j,i_0,r}^*).$$

Denote

$$(3.9) \quad \phi_0(\cdot) := \text{Tr}(\omega_0 \cdot)$$

with the density matrix $\omega_0 = (\omega_{k,j})$, (3.8) becomes

$$(3.10) \quad \sum_r \omega_{r,r} \cdot \sum_j \text{Tr}(K_{j,i_0,r} K_{j,i_0,r}^*) = \sum_r \omega_{r,r} \cdot \sum_j \sum_k (K_{j,i_0,r} K_{j,i_0,r}^*)(k, k).$$

Define

$$(3.11a) \quad p_0(r) := \omega_{r,r}, \quad \forall r \in \{1, \dots, d\}$$

and

$$(3.11b) \quad T_{r,i,k} := \sum_j (K_{j,i,r} K_{j,i,r}^*)(k, k), \quad \forall r, i, k \in \{1, \dots, d\}.$$

Then p_0 is a probability measure on $\{1, \dots, d\}$ and (2.2a,b,c) are satisfied.

Our second step is to compute the joint distribution. In order to do this we must know the explicit form of the matrix

$$(3.12) \quad \mathcal{E}(|e_{i_0}\rangle\langle e_{i_0}| \otimes \mathcal{E}(|e_{i_1}\rangle\langle e_{i_1}| \otimes \dots \otimes \mathcal{E}(|e_{i_{n-1}}\rangle\langle e_{i_{n-1}}| \otimes \mathcal{E}(|e_{i_n}\rangle\langle e_{i_n}| \otimes 1)) \dots)).$$

By the formulae (3.1) and (1.12), we find that (3.12) is equal to

$$(3.13) \quad \sum_{r_0} |e_{r_0}\rangle\langle e_{r_0}| \sum_{j_0} \text{Tr}(K_{j_0,i_0,r_0} K_{j_0,i_0,r_0}^* \mathcal{E}(|e_{i_1}\rangle\langle e_{i_1}| \otimes \dots \otimes \mathcal{E}(|e_{i_{n-1}}\rangle\langle e_{i_{n-1}}| \otimes \mathcal{E}(|e_{i_n}\rangle\langle e_{i_n}| \otimes 1)) \dots)).$$

Applying again the formulae (3.1) and (1.12), we are able to rewrite (3.13) as

$$(3.13a) \quad \sum_{r_0} |e_{r_0}\rangle\langle e_{r_0}| \sum_{j_0} \text{Tr}(K_{j_0,i_0,r_0}^{*2} \sum_{r_1} |e_{r_1}\rangle\langle e_{r_1}| \sum_{j_0} \text{Tr}(K_{j_1,i_1,r_1}^{*2} \mathcal{E}(|e_{i_1}\rangle\langle e_{i_1}| \otimes \dots \otimes \mathcal{E}(|e_{i_{n-1}}\rangle\langle e_{i_{n-1}}| \otimes \mathcal{E}(|e_{i_n}\rangle\langle e_{i_n}| \otimes 1)) \dots))) \\ = \sum_{r_0, r_1} |e_{r_0}\rangle\langle e_{r_0}| \sum_{j_0} \text{Tr}(K_{j_0,i_0,r_0}^{*2}(r_1, r_1) \sum_{j_1} \text{Tr}(K_{j_1,i_1,r_1}^{*2} \mathcal{E}(|e_{i_1}\rangle\langle e_{i_1}| \otimes \dots \otimes \mathcal{E}(|e_{i_{n-1}}\rangle\langle e_{i_{n-1}}| \otimes \mathcal{E}(|e_{i_n}\rangle\langle e_{i_n}| \otimes 1)) \dots))).$$

By using the symbol introduced in (3.11b), the right hand side of (3.13a) becomes

$$(3.13b) \quad \sum_{r_0, r_1} |e_{r_0}\rangle\langle e_{r_0}| T_{r_0,i_0,r_1} \sum_{j_1} \text{Tr}(K_{j_1,i_1,r_1}^{*2} \mathcal{E}(|e_{i_1}\rangle\langle e_{i_1}| \otimes \dots \otimes \mathcal{E}(|e_{i_{n-1}}\rangle\langle e_{i_{n-1}}| \otimes \mathcal{E}(|e_{i_n}\rangle\langle e_{i_n}| \otimes 1)) \dots))).$$

Repeating above discussion, we find that (3.13) is equal to

$$(3.14) \quad \sum_{r_0, r_1, \dots, r_n, r} |e_{r_0}\rangle \langle e_{r_0}| T_{r_0, i_0, r_1} T_{r_1, i_1, r_2} \cdots T_{r_{n-1}, i_{n-1}, r_n} T_{r_n, i_n, r}$$

Introducing new indices:

$$(3.15) \quad j := r_0, j_0 := r_1, \dots, j_{n-1} := r_n, j_n := r$$

(3.14) can be rewritten as

$$(3.14a) \quad \sum_{j, j_0, j_1, \dots, j_n} |e_j\rangle \langle e_j| T_{j, i_0, j_0} T_{j_0, i_1, j_1} \cdots T_{j_{n-2}, i_{n-1}, j_{n-1}} T_{j_{n-1}, i_n, j_n}$$

By acting the initial state $\phi_0(\cdot) = \text{Tr}(\omega_0 \cdot)$ to (3.14a), one obtains (2.2d) with $p_0(j) := \omega_0(j, j)$.

Now let be given a classical stochastic process $\{X_n\}_{n=1}^\infty$ with the properties (2.2a,b,c,d), we have to construct a quantum Markov chain $\{\phi_0, \mathcal{E}\}$ such that (1.12) and (2.1) are satisfied.

Taking the initial density matrix ω_0 as

$$(3.15) \quad \begin{pmatrix} p_0(1) & 0 & \cdots & 0 \\ 0 & p_0(2) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & p_0(d) \end{pmatrix}$$

$\phi_0(\cdot) := \text{Tr}(\omega_0 \cdot)$ is, clearly, a state on M_d .

Define

$$(3.16) \quad K := \sum_{h,r} |e_h\rangle \langle e_r| \otimes K_{h,r}$$

$$(3.17) \quad K_{i,j}^* := (\sqrt{T_{j,i,1}}e_j, \sqrt{T_{j,i,2}}e_j, \dots, \sqrt{T_{j,i,d}}e_j).$$

We have

$$(3.18a) \quad K_{i,j}K_{i,j'}^* = 0, \quad \forall i, j \neq j' \in \{1, \dots, d\}$$

and

$$(3.18b) \quad K_{i,j}K_{i,j}^*(k, k) = T_{j,i,k}, \quad \forall i, j, k \in \{1, \dots, d\}.$$

Thus, a quantum Markov chain (ϕ_0, \mathcal{E}) is obtained and these end the proof of Theorem (2.3).

Proof of Theorem (2.4). Let classical stochastic process $\{X_n\}_{n=0}^\infty$ on a probability space (\mathbf{P}, Ω) be a quantum Markov chain. Then by Theorem (2.3), we know

that $p_0, (T_{j,i,k})$ satisfy (2.2a,b,c,d).

Define a probability measure $p(\cdot, \cdot)$ on $\{1, \dots, d\}^2$ by

$$(3.19) \quad p(j, i) := \sum_r p_0(r) T_{r,i,j}$$

and a transition matrix $(p_{(j,i),(j',i')})$ by

$$(3.20) \quad p_{(j,i),(j',i')} := T_{j,i',j'}, \quad \forall j, i, j', i' \in \{1, \dots, d\}.$$

Then with a canonical way we can construct a probability space (P', Ω') and an $\{1, \dots, d\}^2$ -valued Markov chain $\{(Z_n, Y_n)\}_{n=1}^\infty$ with the initial distribution:

$$(3.21) \quad P'((Z_0, Y_0) = (j, i)) = p(j, i) = \sum_k p_0(k) T_{k,i,j}$$

and the transition probability:

$$(3.22) \quad P'((Z_n, Y_n) = (j', i') \mid (Z_{n-1}, Y_{n-1}) = (j, i)) = p_{(j,i),(j',i')} = T_{j,i',j'}$$

Thus by a simple computation we know that the joint distribution $P'(Y_0 = i_0, \dots, Y_1 = i_1)$ is given by the right hand side of (2.2d) and this implies (2.4).

On the other hand, let be given an $\{1, \dots, d\}^2$ -valued Markov chain $\{(Z_n, Y_n)\}_{n=1}^\infty$ on a probability space (P', Ω') . If it has the properties i) and ii) in Theorem (2.4), then following conclusions are obvious:

- 1) p_0 is a probability measure on $\{1, \dots, d\}$ and $p_0, (T_{j,i,k})$ satisfy (2.2a,b,c);
- 2) the joint distribution of $\{Y_n\}_{n=1}^\infty$ is given by the right hand side of (2.2d).

These end the proof.

Proof of Theorem (2.6). For any $m \in \{1, \dots, d\}$, (3.5) shows that

$$(3.23) \quad \mathcal{E}(|e_m\rangle\langle e_m| \otimes 1) = \sum_r |e_r\rangle\langle e_r| \text{Tr}(\sum_j K_{j,m,r} K_{j,m,r}^*).$$

On the other hand, (2.5) makes sure that

$$(3.24) \quad \mathcal{E}(|e_m\rangle\langle e_m| \otimes 1) = |e_m\rangle\langle e_m|.$$

Comparing (3.23) and (3.24), we have

$$(3.25) \quad \text{Tr}(\sum_j K_{j,m,r} K_{j,m,r}^*) = \delta_{m,r}$$

and this gives that

$$(3.26) \quad K_{j,m,r} = 0, \quad \forall m \neq r \in \{1, 2, \dots, d\}$$

i.e. K_j has the special form as (2.6).

If any K_j ($j \in \{1, \dots, d\}$) has the form (2.6), we shall prove that

- 1) for each $a, b \in \mathcal{D}$, $\mathcal{E}(a \otimes b) = a \cdot \mathcal{E}(1 \otimes b)$;
- 2) for each $b \in \mathcal{D}$, $\mathcal{E}(1 \otimes b) \in \mathcal{D}$.

It is clear that with 1), the condition 2) is equivalent to

- 2)' for each $a, b \in \mathcal{D}$, $\mathcal{E}(a \otimes b) \in \mathcal{D}$.

The formula (3.5) shows, since any K_j has the form (2.6), that

$$(3.27) \quad \mathcal{E}(1 \otimes b) = \sum_r |e_r\rangle \langle e_r| \text{Tr}(\sum_j K_{j,r,r}^* b K_{j,r,r}), \quad \forall b \in \mathcal{D}$$

and this belongs to \mathcal{D} . Moreover

$$(3.28) \quad \mathcal{E}(a \otimes b) = \sum_m a(m, m) [|e_m\rangle \langle e_m| \text{Tr}(\sum_j K_{j,m,m}^* b K_{j,m,m})]$$

$$= \begin{pmatrix} a(1,1) & 0 & \cdots & 0 \\ 0 & a(2,2) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a(d, d) \end{pmatrix} \\ \times \begin{pmatrix} \text{Tr}(\sum_j K_{j,1,1}^* b K_{j,1,1}) & 0 & \cdots & 0 \\ 0 & \text{Tr}(\sum_j K_{j,2,2}^* b K_{j,2,2}) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \text{Tr}(\sum_j K_{j,d,d}^* b K_{j,d,d}) \end{pmatrix} \\ = a \cdot \mathcal{E}(1 \otimes b).$$

This finishes the proof.

REFERENCES

- [1] L. Accardi, On the noncommutative Markovian property, *Funct. Anal. Appl.*, **9** (1975), 1–8.
- [2] —, Noncommutative Markov Chains, In: *International School of Math. Phys. Camerino*, (1974), 268–295.
- [3] —, Topics in quantum probability, *Phy. Reports*, **77**, No. 3 (1981), 169–192.
- [4] M. Fannes, B. Nachtergaele, R. F. Werner, Valence bond states on quantum spin chains as ground states with spectral gap, *J. Phys. A: Math. Gen.*, **24** (1991), 185–190.

*Dipartimento di Matematica
Università di Bari
Centro V. Volterra
Università di Roma II*

Current address:
*Università degli Studi di Bari
Dipartimento di Matematica
Campus Universitario
Via E. Orabona, 4*