

THE LINE BUNDLES ON THE MODULI STACK OF PRINCIPAL BUNDLES ON FAMILIES OF CURVES

ROBERTO FRINGUELLI¹ AND FILIPPO VIVIANI²

¹*Dipartimento di Matematica, Università di Roma ‘La Sapienza’, Roma, Italy*

²*Dipartimento di Matematica, Università di Roma ‘Tor Vergata’, Roma, Italy*

Corresponding author: Roberto Fringuelli, email: rob.fringuelli@gmail.com

(Received 5 January 2024)

Abstract Given a connected reductive algebraic group G over an algebraically closed field, we investigate the Picard group of the moduli stack of principal G -bundles over an arbitrary family of smooth curves.

Keywords: moduli of principal bundles; algebraic curves; Picard group

2020 Mathematics Subject Classification: Primary: 14H60

Secondary: 14D20; 14C22; 20G07; 14H10

1. Introduction

The aim of this paper, which is a sequel of the papers [21] and [22], is to study the Picard group of the *moduli stack of (principal) G -bundles* $\mathrm{Bun}_G(C/S)$, which parametrizes G -bundles, where G is a connected and smooth linear algebraic group over a field $k = \bar{k}$, over an arbitrary family $\pi : C \rightarrow S$ of (connected, smooth and projective) k -curves of genus $g = g(C/S) \geq 0$.

The moduli stack $\mathrm{Bun}_G(C)$ of (principal) G -bundles, where G is a complex reductive group and C is a complex curve, has been deeply studied because of its relation to the Wess–Zumino–Witten (=WZW) model associated to G , which form a special class of rational conformal field theories, see [3], [41] and [4] for nice surveys. In the WZW-model associated to a simply connected group G , the spaces of conformal blocks can be interpreted as spaces of generalized theta functions, that is spaces of global sections of suitable line bundles (e.g. powers of determinant line bundles) on $\mathrm{Bun}_G(C)$, see [5, 16, 28, 31, 37]. The above application to conformal field theory leads naturally to the study of the Picard group of $\mathrm{Bun}_G(C)$ (for G reductive over $k = \bar{k}$), which was computed in increasing degree of generality, thanks to the effort of many mathematicians: the case of SL_r dates back to pioneering work of Drezet–Narasimhan in the late eighties [15]; the case of a simply connected, almost simple G is dealt with in [11, 17, 29, 31, 42]; the case of a semisimple almost simple G is dealt with in [6, 30, 43]; the case of an arbitrary reductive group was finally established by Biswas–Hoffmann [7].



Motivated by the long-term project of generalizing the study of the vector bundle of conformal blocks (also known as Verlinde bundle) behind the simply connected case, the authors in [21] and [22] have computed the Picard group of $\text{Bun}_G(\mathcal{C}_{g,n}/\mathcal{M}_{g,n})$, for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ over the moduli stack of n -pointed curves of genus g , as well as the restriction homomorphism to the Picard group of the fibres of $\text{Bun}_G(\mathcal{C}_{g,n}/\mathcal{M}_{g,n}) \rightarrow \mathcal{M}_{g,n}$, generalizing the previous works of Melo–Viviani [33] for $G = \mathbb{G}_m$ and Fringuelli [18] for $G = \text{GL}_r$.

In this follow-up paper, we study the Picard group of $\text{Bun}_G(C/S)$ for an arbitrary family C/S of curves over an algebraic stack S (which, for technical reasons, it is often assumed to be an integral and regular quotient stack), in which case very little was known before. A remarkable exception is the work of Faltings [17], which provides a functorial identification $\text{Pic Bun}_G(C/S) \cong \mathbb{Z}$ for a simply connected and almost simple group G and an arbitrary family C/S admitting a section. Our Theorem B can be seen as a generalization of Faltings’ result (see Corollary C and the discussion following it).

We can now explain our main results focusing, in this introduction, on families C/S of positive genus $g(C/S) > 0$ and referring to the body of the paper for the slightly different, and easier, case $g(C/S) = 0$.

Recall that the stack $\text{Bun}_G(C/S)$ is an algebraic stack, locally of finite type and smooth over S and its relative connected components are in functorial bijection with the fundamental group $\pi_1(G)$ (see Fact 2.6). We will denote the connected components and the restriction of the forgetful morphism by

$$\Phi_G^\delta : \text{Bun}_G^\delta(C/S) \rightarrow S \quad \text{for any } \delta \in \pi_1(G).$$

The pull-back morphism at the level of the Picard groups is injective, so will often focus onto the relative Picard group

$$\text{RPic Bun}_G^\delta(C/S) := \frac{\text{Pic Bun}_G^\delta(C/S)}{(\Phi^\delta)^*(\text{Pic } S)}.$$

We proved in [21, Thm. A] that if $\text{red} : G \rightarrow G^{\text{red}}$ is the reductive quotient of G , i.e. the quotient of G by its unipotent radical, then for any $\delta \in \pi_1(G) \xrightarrow[\cong]{\pi_1(\text{red})} \pi_1(G^{\text{red}})$, the pull-back homomorphism

$$\text{red}_\#^* : \text{Pic}(\text{Bun}_{G^{\text{red}}}^\delta(C/S)) \xrightarrow{\cong} \text{Pic}(\text{Bun}_G^\delta(C/S))$$

is an isomorphism. Hence, throughout this paper, we will restrict to the case of a *reductive group* G .

A first source of line bundles on $\text{Bun}_G^\delta(C/S)$ comes from the determinant of cohomology $d_{\hat{\pi}}(-)$ and the Deligne pairing $\langle -, - \rangle_{\hat{\pi}}$ of line bundles on the universal curve $\hat{\pi} : C \times_S \text{Bun}_G^\delta(C/S) \rightarrow \text{Bun}_G^\delta(C/S)$. To be more precise, any character $\chi \in \Lambda^*(G) := \text{Hom}(G, \mathbb{G}_m)$ gives rise to a morphism of S -stacks

$$\chi_\# : \text{Bun}_G^\delta(C/S) \rightarrow \text{Bun}_{\mathbb{G}_m}^{\chi(d)}(C/S),$$

and by pulling back via $\chi_{\#}$ the universal \mathbb{G}_m -bundle (i.e. line bundle) on the universal curve $C \times_S \text{Bun}_{\mathbb{G}_m}^{\chi(\delta)}(C/S) \rightarrow \text{Bun}_{\mathbb{G}_m}^{\chi(\delta)}(C/S)$, we get a line bundle \mathcal{L}_χ on $C \times_S \text{Bun}_G^\delta(C/S)$. Then, using these line bundles \mathcal{L}_χ and the line bundles $\{p_1^*(M) : M \in \text{Pic}(C)\}$ pulled-back from the family C/S , we define the following line bundles, that we call *tautological line bundles*, on $\text{Bun}_G^\delta(C/S)$

$$\begin{aligned} & \{d_{\widehat{\pi}}(\mathcal{L}_\chi(M)) : \chi \in \Lambda^*(G), M \in \text{Pic}(C)\}, \\ & \{(\mathcal{L}_\chi(M), \mathcal{L}_\mu(N))_{\widehat{\pi}} : \chi, \mu \in \Lambda^*(G), M, N \in \text{Pic}(C)\}, \end{aligned}$$

where we set $\mathcal{L}_\chi(M) := \mathcal{L}_\chi(p_1^*(M))$ for simplicity.

Note that since any character of G factors through the abelianization of $\text{ab} : G \rightarrow G^{\text{ab}}$ of G , the tautological line bundles lie in the subgroup

$$\text{Pic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S) \xrightarrow{\text{ab}_{\#}^*} \text{Pic Bun}_G^\delta(C/S),$$

where $\delta^{\text{ab}} := \pi_1(\text{ab})(\delta)$.

The next theorem describes, for an arbitrary torus T and an arbitrary $d \in \pi_1(T)$ (which includes the case $T = G^{\text{ab}}$ and $\delta^{\text{ab}} = d$), the structure of the *tautological subgroup*

$$\text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S) \subseteq \text{Pic Bun}_T^d(C/S)$$

generated by the tautological line bundles (or, in other words, it encodes all the relations among tautological line bundles), and it describes when it coincides with the full Picard group.

Theorem A. (see Theorems 3.8 and 3.9) *Let $C \rightarrow S$ be a family of curves of genus $g > 0$ on a regular and integral quotient stack. Let T be a torus and fix $d \in \Lambda(T) = \pi_1(T)$.*

- (1) *The relative tautological Picard group of $\text{Bun}_T^d(C/S)$ sits in the following exact sequence, functorial in S and T :*

$$0 \rightarrow \Lambda^*(T) \otimes \text{RPic}(C/S) \xrightarrow{i_T^d} \text{RPic}^{\text{taut}} \text{Bun}_T^d(C/S) \xrightarrow{\gamma_T^d} \text{Bil}^s(\Lambda(T)) \rightarrow 0,$$

where $\text{RPic}(C/S)$ is the relative Picard group of C/S , and the homomorphisms i_T^d and γ_T^d are defined by

$$i_T^d(\chi \otimes M) = \langle \mathcal{L}_\chi, M \rangle_{\widehat{\pi}} \quad \text{and} \quad \gamma_T^d(d_{\widehat{\pi}}(\mathcal{L}_\chi(M))) = \chi \otimes \chi.$$

- (2) *If $\text{End}(J_{C_{\overline{\eta}}}) = \mathbb{Z}$ for a geometric generic point $\overline{\eta} \rightarrow S$ and the natural map $\text{Pic}(C) \rightarrow \text{Pic}_{C/S}(S)$ is surjective, then*

$$\text{Pic Bun}_T^d(C/S) = \text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S).$$

Moreover, in Theorem 3.8, we give two alternative presentations of the relative tautological Picard group of $\text{Bun}_T^d(C/S)$.

The case of families of genus zero curves is easier, and it is dealt with in Theorem 3.6. In particular, we will show that the Picard group of $\text{Bun}_T^d(C/S)$ is always generated by tautological line bundles if $g(C/S) = 0$.

The above Theorem A was known:

- for a curve C over $k = \bar{k}$ by Biswas–Hoffmann [7] (see also [21, Prop. 4.1.1]), in which case the hypothesis that $\text{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$ is also necessary for part (2) to hold (while the second assumption is always satisfied);
- for the universal family $\mathcal{C}_{g,n}/\mathcal{M}_{g,n}$ over the stack of n -pointed curves of genus $g > 0$ (which satisfies the assumption of part (2), see Remarks 2.3 and 2.5(1)), by work of the authors [21, Thm. B] and [22, Thm. A], generalizing previous results for $T = \mathbb{G}_m$, $g \geq 2$ and $n = 0$ by Melo–Viviani [33].

The other source of line bundles on $\text{Bun}_G^\delta(C/S)$, for G non-abelian, is the *transgression map* introduced in [21, Thm. 5.0.1, Rmk. 5.2.2]:

$$\tau_G^\delta = \tau_G^\delta(C/S) : (\text{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}_G} \rightarrow \text{Pic Bun}_G^\delta(C/S),$$

which is the homomorphism, functorial in S and G , uniquely characterized the following commutative diagram:

$$\begin{array}{ccc} (\text{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}_G} & \xrightarrow{\tau_G^\delta} & \text{Pic Bun}_G^\delta(C/S), \\ \downarrow & & \downarrow \iota_\pi^* \\ \text{Sym}^2 \Lambda^*(T_G) & \xrightarrow{\tau_{T_G}^d} & \text{Pic Bun}_{T_G}^d(C/S), \\ & & \\ \chi \cdot \chi' & \longrightarrow & \langle \mathcal{L}_\chi, \mathcal{L}_{\chi'} \rangle_{\widehat{\pi}} \end{array}$$

where $\iota : T_G \hookrightarrow G$ is a fixed maximal torus of G , $d \in \pi_1(T_G)$ is any lift of $\delta \in \pi_1(G)$, $\mathscr{W}_G := \mathcal{N}(T_G)/T_G$ is the Weyl group of G and $(\text{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}_G}$ is identified with the lattice $\text{Bil}^{\text{s, ev}}(\Lambda(T_G))^{\mathscr{W}_G}$ of \mathscr{W}_G -invariant even symmetric bilinear forms on the lattice $\Lambda(T_G)$ of cocharacters of T_G (see Lemma 3.13 for an explicit description).

The next theorem describes when the Picard group of $\text{Bun}_G^\delta(C/S)$ is generated by the image of the transgression map and the pull-back of the line bundles on $\text{Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S)$.

Theorem B. (see Theorems 3.14 and 3.16) *Let $C \rightarrow S$ be a family of curves of genus $g > 0$ on a regular and integral quotient stack. Let G be a reductive group and denote by $\text{ab} : G \rightarrow G^{\text{ab}}$ its abelianization. Fix $\delta \in \pi_1(G)$ and set $\delta^{\text{ab}} := \pi_1(\text{ab})(\delta) \in \pi_1(G^{\text{ab}})$.*

If one of the following two hypothesis is satisfied:

- the derived subgroup $\mathcal{D}(G)$ of G is simply connected;*
- $\text{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$ for a geometric generic point $\bar{\eta} \rightarrow S$, the natural map $\text{Pic}(C) \rightarrow \text{Pic}_{C/S}(S)$ is surjective and $\text{RPic}^0(C/S)$ is torsion-free;*

then the following diagram, which is functorial in G and S ,

$$\begin{CD}
 \mathrm{Sym}^2 \Lambda^*(G^{\mathrm{ab}}) @<\mathrm{Sym}^2 \Lambda_{\mathrm{ab}}^*><< \mathrm{Sym}^2 \Lambda^*(T_G) @>\mathscr{W}_G>> \\
 @V\tau_{G^{\mathrm{ab}}(C/S)}^{\delta^{\mathrm{ab}}}& & V\tau_G^{\delta}(C/S) \\
 \mathrm{Pic} \mathrm{Bun}_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}}(C/S) @<\mathrm{ab}_{\sharp}^*><< \mathrm{Pic} \mathrm{Bun}_G^{\delta}(C/S) @.
 \end{CD}$$

is a push-out diagram.

In particular, there exists an exact sequence, functorial in S and G ,

$$0 \rightarrow \mathrm{Pic} \mathrm{Bun}_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}}(C/S) \xrightarrow{\mathrm{ab}_{\sharp}^*} \mathrm{Pic} \mathrm{Bun}_G^{\delta}(C/S) \rightarrow \mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_{\mathscr{D}(G)})|\Lambda(T_{G^{\mathrm{ss}}}))^{\mathscr{W}_G} \rightarrow 0,$$

where $\mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_{\mathscr{D}(G)})|\Lambda(T_{G^{\mathrm{ss}}}))^{\mathscr{W}_G}$ is the lattice of \mathscr{W}_G -invariant symmetric even bilinear form on $\Lambda(T_{\mathscr{D}(G)})$ that are integral on $\Lambda(T_{\mathscr{D}(G)}) \otimes \Lambda(T_{G^{\mathrm{ss}}})$, where G^{ss} is the semisimplification of G .

For a family of curves C/S of positive genus such that $\mathrm{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$, the map $\mathrm{Pic}(C) \rightarrow \mathrm{Pic}_{C/S}(S)$ is surjective and satisfying one of the two assumptions of **Theorem B**, we give two alternative presentations of the relative Picard group of $\mathrm{Bun}_G^{\delta}(C/S)$ in **Theorem 3.18**.

The case of families of genus zero curves is easier, and it is treated in detail in **Theorem 3.20** and **Corollary 3.21**. In particular, we will show that **Theorem B** holds true for families of genus zero curves under the assumption that $\mathscr{D}(G)$ is simply connected (see **Theorem 3.14**).

For a semisimple group G , the above theorem gives the following:

Corollary C. *Let $C \rightarrow S$ be a family of curves of genus $g > 0$ on a regular and integral quotient stack. Let G be a semisimple group and fix $\delta \in \pi_1(G)$.*

If one of the following two hypothesis is satisfied:

- (a) G is simply connected,
- (b) $\mathrm{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$ for a geometric generic point $\bar{\eta} \rightarrow S$, the natural map $\mathrm{Pic}(C) \rightarrow \mathrm{Pic}_{C/S}(S)$ is surjective and $\mathrm{RPic}^0(C/S)$ is torsion-free,

then the transgression map

$$\tau_G^{\delta} = \tau_G^{\delta}(C/S) : (\mathrm{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}_G} \xrightarrow{\cong} \mathrm{Pic} \mathrm{Bun}_G^{\delta}(C/S)$$

is an isomorphism, functorial in S and G .

Note that, for G semisimple, we have that $(\mathrm{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}_G} \cong \mathbb{Z}^s$, where s is the number of almost simple factors of the universal cover G^{sc} of G (see **Lemma 3.13**). In particular, if G is simply connected and almost simple (i.e. $s = 1$), then the above Corollary recovers (in the case where C/S has a section) the isomorphism $\mathrm{Pic} \mathrm{Bun}_G^{\delta}(C/S) \cong \mathbb{Z}$ established by Faltings [17].

The above **Theorem B** (and hence **Corollary C**) was known:

- for G semisimple and simply connected and a family C/S admitting a section, by Faltings [17];
- for a curve C over $k = \bar{k}$ if $\mathcal{D}(G)$ is simply connected, by Biswas–Hoffmann [7];
- for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ (which satisfies the assumption (b) by Remarks 2.3 and 2.5(1), and the Franchetta’s conjecture) by the work of the authors [21, Thm. C(2)] and [22, Cor. 3.4], generalizing the work of the first author for if $G = \mathrm{GL}_r$, $g \geq 2$ and $n = 0$ in [18, Thm. A].

Note that Theorem B does not (necessarily) hold for a curve C over $k = \bar{k}$ (which clearly does not satisfies assumption (b)) if the group $\mathcal{D}(G)$ is not simply connected, as it follows from the work of Biswas–Hoffmann [7].

In § 4 of the paper, we consider the rigidification

$$\nu_G^\delta : \mathrm{Bun}_G^\delta(C/S) \rightarrow \mathrm{Bun}_G^\delta(C/S) // \mathcal{Z}(G) := \mathfrak{Bun}_G^\delta(C/S)$$

of the stack $\mathrm{Bun}_G^\delta(C/S)$ by the centre $\mathcal{Z}(G)$ of G , which acts functorially on any G -bundle.

The Leray spectral sequence for the cohomology of the group \mathbb{G}_m with respect to the $\mathcal{Z}(G)$ -gerbe ν_G^δ gives the exact sequence

$$0 \rightarrow \mathrm{Pic}(\mathfrak{Bun}_G^\delta(C/S)) \xrightarrow{(\nu_G^\delta)^*} \mathrm{Pic}(\mathrm{Bun}_G^\delta(C/S)) \xrightarrow{\mathrm{wt}_G^\delta} \Lambda^*(\mathcal{Z}(G)) := \mathrm{Hom}(\mathcal{Z}(G), \mathbb{G}_m),$$

where the weight homomorphism wt_G^δ is given by restricting the line bundles on $\mathrm{Bun}_G^\delta(C/S)$ to the fibres of ν_G^δ , and its cokernel measures the triviality of the $\mathcal{Z}(G)$ -gerbe ν_G^δ (see § 4 for more details).

In the next theorem, we describe the Picard group of the rigidification $\mathfrak{Bun}_G^\delta(C/S)$ in terms of the Néron–Severi group $\mathrm{NS}(\mathfrak{Bun}_G^\delta)$ (see Definition 4.1) and the cokernel of the weight homomorphism wt_G^δ in terms of the cokernel of the evaluation homomorphism $\mathrm{ev}_{\mathcal{D}(G)}^\delta$ (see Definition 4.1).

Theorem D. (see Theorems 4.2 and 4.3) *Let $C \rightarrow S$ be a family of curves of genus $g > 0$ on a regular and integral quotient stack. Assume that both the following conditions hold:*

- (i) $\mathrm{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$ for a geometric generic point $\bar{\eta} \rightarrow S$ and the natural map $\mathrm{Pic}(C) \rightarrow \mathrm{Pic}_{C/S}(S)$ is surjective;
- (ii) either $\mathrm{RPic}^0(C/S)$ is torsion-free or $\mathcal{D}(G)$ is simply connected.

(1) *We have an exact sequence, functorial in S and G :*

$$0 \rightarrow \Lambda^*(G^{\mathrm{ab}}) \otimes \mathrm{RPic}^0(C/S) \xrightarrow{j_G^\delta} \mathrm{RPic}(\mathfrak{Bun}_G^\delta(C/S)) \xrightarrow{\gamma_G^\delta} \mathrm{NS}(\mathfrak{Bun}_G^\delta),$$

such that

$$Im(\overline{\gamma_G^\delta}) = \left\{ b \in NS(\mathfrak{Bun}_G^\delta) : \begin{array}{l} b(\delta \otimes x) + (g-1)b(x \otimes x) \text{ is a multiple of } \delta(C/S) \\ \text{for any } x \in \Lambda(T_G) \end{array} \right\}.$$

(2) We have an exact sequence, functorial in S and G :

$$0 \rightarrow \text{coker}(\overline{\gamma_G^\delta}) \xrightarrow{\partial_G^\delta} \text{Hom} \left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{\delta(C/S)\mathbb{Z}} \right) \xrightarrow{\widetilde{\Lambda}_{\text{ab}}^*} \text{coker}(\text{wt}_G^\delta) \\ \xrightarrow{\widetilde{\Lambda}_{\mathcal{D}}^*} \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta) \rightarrow 0,$$

where $\delta(C/S)$ is the minimum relative degree of a relative ample line bundle on C/S (and zero if such a line bundle does not exist).

The Néron–Severi group $NS(\mathfrak{Bun}_G^\delta)$ and the evaluation homomorphism $\text{ev}_{\mathcal{D}(G)}^\delta$ are defined in Definition 4.1. We make explicit Theorem D(2) in the special cases $\delta(C/S) = 1$ or $G = T$ a torus in Corollary 4.4.

The case of families of genus zero curves is easier, and it is treated in Theorem 4.5. Theorem D(1) was known:

- for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ (which satisfies the assumptions by Remarks 2.3 and 2.5(1), and the Franchetta’s conjecture) by the work of the authors [22, Thm. 1.3], generalizing [33, Thm. B(i) and 1.5] (see also [26]) for $G = \mathbb{G}_m$, $n = 0$ and [18, Thm. B(i) and Thm. A.2] (see also [27]) if $G = \text{GL}_r$ and $n = 0$.

Theorem D(2) was known:

- for a curve C over $k = \bar{k}$ if $\mathcal{D}(G)$ is simply connected, by Biswas–Hoffmann [8] (where also the case of an arbitrary reductive group is treated);
- for $G = \mathbb{G}_m$ and an arbitrary family C/S in [35];
- for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ by the work of the authors [22, Thm. 1.4], generalizing [33, Thm. 6.4] for $G = \mathbb{G}_m$ and $n = 0$, [18, Cor. 3.3.2(i)] for $G = \text{GL}_r$ and $n = 0$ and [20, Thm. B(i)] for $G = \text{GL}_r$.

The computation of the cokernel of the weight homomorphism wt_G^δ (which coincides with the kernel of the obstruction homomorphism, see Equation (4.1)) carried over in Theorem D(2) will be used in our upcoming work [23], where we will compute the (cohomological) Brauer groups of $\text{Bun}_G^\delta(C/S)$ and $\mathfrak{Bun}_G^\delta(C/S)$ for certain reductive groups, extending the work of Biswas–Holla [9] for a fixed curve and a complex semisimple group, as well as the work of Fringuelli–Pirisi [20] for the universal family $\mathcal{C}_{g,n}/\mathcal{M}_{g,n}$ and $G = \text{GL}_r$.

Notations.

1.1. We denote by $k = \bar{k}$ an algebraically closed field of arbitrary characteristic $p \geq 0$. An algebraic stack will always be assumed locally of finite type over k and quasi-separated,

i.e. with quasi-compact and separated diagonal. In particular, all the schemes and algebraic spaces must be quasi-separated. Furthermore, any algebraic space is generically a scheme (see [44, Proposition 06NH])

2. Preliminaries

In this section, we collect some definitions and preliminaries results that we will need throughout the paper.

2.1. Family of curves

A *curve* is a connected, smooth and projective scheme of dimension one over an algebraically closed field. The genus of a curve C is $g(C) := \dim H^0(C, \omega_C)$.

A *family of curves* $\pi : C \rightarrow S$ is a proper, flat and representable morphism of algebraic stacks whose geometric fibres are curves. If all the geometric fibres of π have the same genus g , then we say that $\pi : C \rightarrow S$ is a family of curves of genus g (or a family of curves with relative genus g) and we set $g(C/S) := g$. Note that any family of curves $\pi : C \rightarrow S$ with S connected is a family of genus g curves for some $g \geq 0$. For technical reasons, we will often restrict to families of curves $\pi : C \rightarrow S$ over a regular and integral quotient stack S over k .

Given $g, n \geq 0$, we denote by $\mathcal{M}_{g,n}$ the algebraic stack parametrizing families of curves of genus g with n ordered and pairwise disjoint sections and by $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ the *universal family of n -marked curves of genus g* . We omit the n from the notation when it is zero. Note that $\mathcal{M}_{g,n}$ is a DM (=Deligne–Mumford) stack if and only if $2g - 2 + n > 0$.

A family of curves $\pi : C \rightarrow S$ is *locally projective* if, Zariski-locally on the base, it is a closed subscheme of \mathbb{P}_S^n for some $n \geq 1$. Equivalently, $\pi : C \rightarrow S$ is locally projective if there exists a π -relatively ample line bundle on C , or equivalently a line bundle on C having π -relative positive degree. Note that a family of curves $\pi : C \rightarrow S$ is locally projective if either $g(C/S) \neq 1$ (because the relative dualizing line bundle ω_π has relative degree $2g(C/S) - 2$) or π has a section $\sigma : S \rightarrow C$. However, there are examples of families of genus one curves without sections that are not locally projective, e.g. the universal family $\mathcal{C}_1 \rightarrow \mathcal{M}_1$ of curves of genus one (see Remark 2.2(2), and also [39, XIII, 3.2] and [46] for other examples).

In the results of this paper, an important role is played by the following invariant of the family C/S :

Definition 2.1. *For a family of curves C/S , we define*

$$\delta(C/S) := \begin{cases} \min\{\deg_\pi(M) > 0 : M \in \text{Pic}(C)\}, & \text{if } \pi \text{ is locally projective;} \\ 0, & \text{otherwise.} \end{cases}$$

Equivalently, if π is locally projective, then $\delta(C/S)$ is the non-negative generator of the non-zero image of the relative degree map

$$\begin{aligned} \deg_\pi : \text{Pic}(C) &\longrightarrow \mathbb{Z}, \\ M &\mapsto \deg_\pi(M). \end{aligned} \tag{2.1}$$

Remark 2.2.

- (1) Since the relative dualizing line bundle ω_π of $\pi : C \rightarrow S$ has relative degree equal to $2g(C/S) - 2$, it follows that

$$\delta(C/S) \text{ divides } 2g(C/S) - 2.$$

- (2) It follows from the weak Franchetta conjecture (see [2], [40], [24]) that, for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ of n -marked curves of genus g , we have that

$$\delta(\mathcal{C}_{g,n}/\mathcal{M}_{g,n}) = \begin{cases} |2g - 2| & \text{if } n = 0, \\ 1 & \text{if } n > 0. \end{cases}$$

Note that $\delta(\mathcal{C}_1/\mathcal{M}_1) = 0$, which implies that $\mathcal{C}_1 \rightarrow \mathcal{M}_1$ is not locally projective.

- (3) The invariant $\delta(C/S)$ has been computed for many interesting families of curves:
- If C/S is the universal family of (smooth) complete intersections curves in \mathbb{P}^r of type (d_1, \dots, d_{r-1}) , then $\delta(C/S) = d_1 \dots d_{r-1}$ (see [13, Thm. 4.4]). In particular, if C/S is the universal family of (smooth) plane curves of degree d , then $\delta(C/S) = d$ (see [13, Thm. 4.2]).
 - If C/S is the universal family of hyperplanes sections of the universal family of polarized genus $g \geq 3$ K3 surfaces, then $\delta(C/S) = 2g - 2$ (see [13, Prop. 4.5]).
 - If C/S is the universal family of one of the following:
 - the unique component with general moduli of the Hilbert scheme of curves in \mathbb{P}^r with $r \geq 3$ of genus g and degree d ;
 - the Severi variety of curves of genus g together with a birational embedding as a nodal plane curve of degree d ;
 - the Hurwitz scheme of degree d covers of \mathbb{P}^1 of genus g ;
 and we assume that $\rho := g - (r + 1)(g + r - d) \geq 2$, then $\delta(C/S) = \gcd\{2g - 2, d\}$ (see [14, Thm. 1.1] and [13, Thm. 5.3]).
 - If C/S is the universal family of hyperelliptic genus g curves, then (see [35, Thm. 3.5])

$$\delta(C/S) = \begin{cases} 4 & \text{if } g \text{ is odd,} \\ 2 & \text{if } g \text{ is even.} \end{cases}$$

- (4) If C/S is a family of curves of genus $g = 0$, then

$$\delta(C/S) = \begin{cases} 1 & \text{if } C/S \text{ is Zariski-locally trivial,} \\ 2 & \text{if } C/S \text{ is not Zariski-locally trivial.} \end{cases}$$

Note also that C/S is Zariski locally trivial if and only if C/S admits a section.

We will denote by $\text{Pic}_{C/S}$ the relative Picard functor of the family $C \rightarrow S$, which is an algebraic stack representable and locally of finite presentation over S (by Artin’s representability theorem, see e.g. [10, § 8.3]). We have an exact sequence (see [10, § 8.1, Prop. 4])

$$0 \rightarrow \text{Pic}(S) \xrightarrow{\pi^*} \text{Pic}(C) \rightarrow \text{Pic}_{C/S}(S) \rightarrow \text{Br}(S) \xrightarrow{\pi^*} \text{Br}(C),$$

where $\text{Br}(-) = H^2(-, \mathbb{G}_m)$ denote the cohomological Brauer group. In particular, we get an injective homomorphism

$$\text{RPic}(C/S) := \frac{\text{Pic}(C)}{\text{Pic}(S)} \hookrightarrow \text{Pic}_{C/S}(S). \tag{2.2}$$

Several of our results are restricted to families such that the inclusion (2.2) is an equality, or equivalently such that the natural map $\text{Pic}(C) \rightarrow \text{Pic}_{C/S}(S)$ is surjective. In the next Remark, we describe several families where this assumption holds true.

Remark 2.3. The inclusion (2.2) is an equality in the following cases:

- (1) if $\pi : C \rightarrow S$ admits a section (because $\pi^* : \text{Br}(S) \rightarrow \text{Br}(C)$ is injective); in particular, if $C/S = \mathcal{C}_{g,n}/\mathcal{M}_{g,n}$ with $n > 0$. Therefore, equality in Equation (2.2) can always be achieved after an étale cover of the base S .
- (2) if $\text{Br}(S) = 0$, e.g. if S is the spectrum of a field $k = \bar{k}$, or the spectrum of strictly henselian valuation ring, or a smooth curve over $k = \bar{k}$.
- (3) if $C/S = \mathcal{C}_g/\mathcal{M}_g$ and $g \geq 2$ by the strong Franchetta’s conjecture (see [34]).
- (4) if the relative Jacobian $J_{C/S}^d := \text{Pic}_{C/S}^d \rightarrow S$ in degree d admits a Poincaré line bundle for every $d \in \mathbb{Z}$.
- (5) if $\delta(C/S) = 1$ (see [35, Lemma 2.2]).

Several of our results are restricted to families of curves $\pi : C \rightarrow S$ over an integral base S , which satisfy the following condition:

$$\text{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}, \tag{2.3}$$

where $\bar{\eta}$ is a geometric generic point of S .

In the next result, we collect some sufficient and necessary conditions for Equation (2.3) to hold.

Proposition 2.4. *Let $\pi : C \rightarrow S$ a family of curves over an integral base S (locally of finite type over $k = \bar{k}$) and let $\bar{\eta} \rightarrow S$ be a geometric generic point.*

- (1) *If $\text{End}(J_{C_s}) = \mathbb{Z}$ for some $s \in S(k)$, then $\text{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$.*
- (2) *If k is uncountable and $\text{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$, then $\text{End}(J_{C_s}) = \mathbb{Z}$ for s very general in $S(k)$.*

- (3) If $k \not\subseteq \overline{\mathbb{F}_p}$, then $\text{End}(J_{C_{\overline{\eta}}}) = \mathbb{Z}$ if and only if there exists $s \in S(\overline{k})$ such that $\text{End}(J_{C_s}) = \mathbb{Z}$.

Note that part (3) is false if $k = \overline{k} = \overline{\mathbb{F}_p}$, see [32, Rmk. 1.12].

Proof. Parts (1) and (2) follow from the well-known fact (see the proof of [32, Prop. 1.13]) that the jumping locus

$$S(k)_{\text{jump}} = \{s \in S(k) : \text{End}(J_{C_{\overline{\eta}}}) \hookrightarrow \text{End}(J_{C_s}) \text{ is an isomorphism}\}$$

is the set of k -points of a union of countably many proper substacks of S .

Part (3) follows from the non-trivial result (proved in [32], [12], [1]) that $S(k)_{\text{jump}} \neq S(k)$ if $k \neq \overline{\mathbb{F}_p}$. □

Remark 2.5.

- (1) The condition $\text{End}(J_{C_{\overline{\eta}}}) = \mathbb{Z}$ is known to hold for several general families of curves, e.g.
- the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ over an arbitrary field $k = \overline{k}$ by [25];
 - the universal family of hyperelliptic curves over an arbitrary field $k = \overline{k}$ by [36, Thm. 6.5];
 - the family of smooth curves belonging to a linear system on a regular smooth projective surface if $k = \mathbb{C}$ by [45].
- (2) The condition $\text{End}(J_{C_{\overline{\eta}}}) = \mathbb{Z}$ implies that the Néron–Severi of $J_{C_{\overline{\eta}}}$ is equal to

$$\text{NS}(J_{C_{\overline{\eta}}}) = \mathbb{Z} \cdot [\Theta_{C_{\overline{\eta}}}], \tag{2.4}$$

where $[\Theta_C]$ is the class of a theta divisor $J_{C_{\overline{\eta}}}$ (using the fact that $\text{NS}(J_{C_{\overline{\eta}}})$ is the subgroup of $\text{End}(J_{C_{\overline{\eta}}})$ consisting of elements that are invariant under the Rosari involution). But the viceversa is false, e.g. for an elliptic curve with complex multiplication.

Pirola has shown in [38] that condition (2.4) holds true for any family $C \rightarrow S$ of genus g curves over $k = \mathbb{C}$ such that the image of S in \mathcal{M}_g via the modular map has codimension less than or equal to $g - 2$ (e.g. the universal families of k -gonal curves of genus $g \geq 2$ for any $k \geq 2$). It would be interesting to know if for such families also the stronger statement $\text{End}(J_{C_{\overline{\eta}}}) = \mathbb{Z}$ holds true.

2.2. Reductive groups

A *reductive group* (over $k = \overline{k}$) is a smooth and connected linear algebraic group over k (i.e. a closed algebraic subgroup of $(\text{GL}_n)_k$, or equivalently an affine group scheme of finite type over k), which does not contain non-trivial connected normal unipotent algebraic subgroups. To any reductive group G , we can associate a cross-like diagram of reductive groups

$$\begin{array}{ccccc}
 & & \mathcal{D}(G) & & \\
 & & \downarrow \mathcal{D} & \searrow & \\
 \mathcal{R}(G) & \xrightarrow{\mathcal{R}} & G & \xrightarrow{ss} & G^{ss} \\
 & \searrow & \downarrow ab & & \\
 & & G^{ab} & &
 \end{array} \tag{2.5}$$

where

- $\mathcal{D}(G) := [G, G]$ is the derived subgroup of G ;
- $G^{ab} := G/\mathcal{D}(G)$ is called the abelianization of G ;
- $\mathcal{R}(G)$ is the radical subgroup of G , which is equal (since G is reductive) to the connected component $\mathcal{Z}(G)^o$ of the centre $\mathcal{Z}(G)$ and
- $G^{ss} := G/\mathcal{R}(G)$ is called the semisimplification of G .

In the above diagram, the horizontal and vertical lines are short exact sequences of reductive groups, the morphisms $\mathcal{D}(G) \rightarrow G^{ss}$ and $\mathcal{R}(G) \rightarrow G^{ab}$ are central isogenies of, respectively, semisimple groups and tori with the same kernel. Since the two semisimple groups $\mathcal{D}(G)$ and G^{ss} are isogenous, they share the same simply connected cover, which we will denote by G^{sc} , and the same adjoint quotient, which we will denote by G^{ad} .

Given a reductive group G over k , we will denote by T_G a maximal torus of G and by B_G the positive Borel subgroup of G containing T_G . As usual, we denote by $\mathcal{W}_G := N(T_G)/T_G$ its Weyl group.

Given a torus T , we will denote by $\Lambda(T) := \text{Hom}(\mathbb{G}_m, T)$ its cocharacter lattice and by $\Lambda^*(T) := \text{Hom}(T, \mathbb{G}_m)$ its character lattice. Note that these lattices are dual to each other via the bilinear map

$$\begin{aligned}
 \Lambda^*(T) \times \Lambda(T) &\longrightarrow \text{Hom}(\mathbb{G}_m, \mathbb{G}_m) = \mathbb{Z}, \\
 (\chi, d) &\mapsto \chi \circ d := \chi(d).
 \end{aligned}$$

The *fundamental group* of G is $\pi_1(G) := \frac{\Lambda(T_G)}{\Lambda_{\text{coroots}}}$, where Λ_{coroots} is the sublattice of $\Lambda(T_G)$, which is generated by integral linear combinations of coroots. The vertical exact sequence in Equation (2.5) induces an exact sequence of finitely generated abelian groups

$$0 \rightarrow \pi_1(\mathcal{D}(G)) \rightarrow \pi_1(G) \xrightarrow{\pi_1(ab)} \pi_1(G^{ab}) \rightarrow 0, \tag{2.6}$$

which identifies $\pi_1(\mathcal{D}(G))$ with the torsion subgroup of $\pi_1(G)$ and $\pi_1(G^{ab})$ with the torsion-free quotient of $\pi_1(G)$.

2.3. The stack of G -bundles on a family of curves

In this subsection, G will be a connected and smooth linear algebraic group over $k = \bar{k}$. Further restrictions on G , like reductiveness, will be specified when needed.

For any family of curves $\pi : C \rightarrow S$, we denote by $\text{Bun}_G(C/S)$ the *moduli stack of G -bundles on $C \rightarrow S$* , i.e. the algebraic stack over S whose fibre over $V \rightarrow S$ is the groupoid of G -bundles $E \rightarrow C_V := C \times_S V$. By definition, we have a forgetful surjective morphism

$$\Phi_G = \Phi_G(C/S) : \text{Bun}_G(C/S) \longrightarrow S. \tag{2.7}$$

The stack $\text{Bun}_G(C/S)$ comes equipped with a universal G -torsor \mathcal{P} on the universal family $\widehat{\pi} : \text{Bun}_G(C/S) \times_S C \rightarrow \text{Bun}_G(C/S)$.

The stack $\text{Bun}_G(C/S)$ is functorial in S , i.e. for any $V \rightarrow S$, we have that

$$\text{Bun}_G(C_V/V) = \text{Bun}_G(C/S) \times_S V.$$

In particular, the fibre of $\Phi_G(C/S) : \text{Bun}_G(C/S) \rightarrow S$ over a k -point $s \in S$ is the moduli stack $\text{Bun}_G(C_s)$ of G -bundles on the curve C_s .

In the special case where the family of curves is the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ over the moduli stack of n -pointed curves of genus g , we set

$$\text{Bun}_{G,g,n} := \text{Bun}_G(\mathcal{C}_{g,n}/\mathcal{M}_{g,n}).$$

In particular, if the family $C \rightarrow S$ has constant relative genus $g = g(C/S)$, then we have that

$$\text{Bun}_G(C/S) = S \times_{\mathcal{M}_g} \text{Bun}_{G,g},$$

with respect to the modular morphism $S \rightarrow \mathcal{M}_g$ associated to the family $C \rightarrow S$.

Any morphism of connected and smooth linear algebraic groups $\phi : G \rightarrow H$ determines a morphism of stacks over S

$$\begin{aligned} \phi_{\#} = \phi_{\#}(C/S) : \text{Bun}_G(C/S) &\longrightarrow \text{Bun}_H(C/S) \\ (E \rightarrow C_V) &\longmapsto \left(\phi_{\#}(E) := E \times^{\phi, G} H = (E \times H)/G \rightarrow C_V \right), \end{aligned} \tag{2.8}$$

where the (right) action of G on $E \times H$ is $(p, h).g := (p.g, \phi(g)^{-1}h)$.

We collect in the following fact the geometric properties of $\text{Bun}_G(C/S)$ and of the forgetful morphism $\Phi_G(C/S)$.

Fact 2.6. *Let G be a connected and smooth linear algebraic group and let $\pi : C \rightarrow S$ be a family of curves.*

- (1) *The morphism $\Phi_G(C/S)$ is locally of finite presentation, smooth, with affine and finitely presented relative diagonal.*
- (2) *There is a functorial decomposition into a disjoint union*

$$\Phi_G(C/S) : \coprod_{\delta \in \pi_1(G)} \text{Bun}_G^{\delta}(C/S) \xrightarrow{\Phi_G^{\delta}(C/S)} S \tag{2.9}$$

such that the morphism $\Phi_G^\delta(C/S) : \text{Bun}_G^\delta(C/S) \rightarrow S$ has geometric integral fibres for every $\delta \in \pi_1(G)$.

- (3) If G is reductive, then for any $\delta \in \pi_1(G)$, the morphism $\Phi_G^\delta(C/S) : \text{Bun}_G^\delta(C/S) \rightarrow S$ is Stein, i.e. it is fpqc and cohomologically flat in degree zero (which means that the natural morphism $(\Phi_G^\delta(C/S))^\sharp : \mathcal{O}_S \rightarrow \Phi_G^\delta(C/S)_*(\mathcal{O}_{\text{Bun}_G^\delta(C/S)})$ is a universal isomorphism), and of relative dimension equal to $(g - 1) \dim G$.

Proof. Part (1): see [21, Thm. 3.1] and the references therein. Part (2): see [21, Thm. 3.1.1, Cor. 3.1.2] and the references therein. Part (3): see [21, Thm. 3.1.3] and the proof of [21, Prop. 3.3.2]. □

Since the centre $\mathcal{Z}(G)$ of a reductive group G acts functorially on any G -bundle, we have that $\mathcal{Z}(G)$ sits functorially inside the automorphism group of any T -point $(C_T \rightarrow T, E)$ of $\text{Bun}_G^\delta(C/S)$ for any $\delta \in \pi_1(G)$. Hence, we can form the rigidification

$$\nu_G^\delta = \nu_G^\delta(C/S) : \text{Bun}_G^\delta(C/S) \rightarrow \text{Bun}_G^\delta(C/S) // \mathcal{Z}(G) := \mathfrak{Bun}_G^\delta(C/S), \tag{2.10}$$

which turns out to be a $\mathcal{Z}(G)$ -gerbe, i.e. a gerbe banded by $\mathcal{Z}(G)$.

3. The Picard group of $\text{Bun}_G^\delta(C/S)$

We will assume throughout this section that:

- ★ $\pi : C \rightarrow S$ is a family of smooth curves of genus $g = g(C/S)$ over an integral and regular quotient stack S (locally of finite type over k). We will denote by $\bar{\eta} \rightarrow S$ a geometric generic point of S and by $C_{\bar{\eta}} := C \times_S \bar{\eta}$ the corresponding geometric generic fibre.

The aim of this section is to study the Picard group of $\text{Bun}_G^\delta(C/S)$ for a reductive group G .

3.1. Recollection on $\text{Pic Bun}_G^\delta(C/S)$

In this subsection, we recall some facts from [21, 22] on the Picard group of $\text{Bun}_G^\delta(C/S)$, for a reductive group G and a fixed $\delta \in \pi_1(G)$. The case $g = 0$ is slightly different from the other cases, and for unifying the statements, we need the following definition:

Definition-Lemma 3.1. *Let G be a reductive group and pick a maximal torus $\iota : T_G \hookrightarrow G$.*

- (1) *We say that a cocharacter $d \in \Lambda(T_G)$ is generic if its image $d^{\text{ss}} \in \Lambda(T_{G^{\text{ss}}}) \subset \Lambda(T_{G^{\text{ad}}})$, into the cocharacter lattice of the adjoint group, satisfies one of the following equivalent conditions (see [21, Lemma 2.2.3]):*

(i) *The contraction homomorphism*

$$(d^{ss}, -) : (\text{Sym}^2 \Lambda^*(T_{G^{sc}}))^{\mathscr{W}G} = \text{Bil}^{s, \text{ev}}(\Lambda(T_{G^{sc}}))^{\mathscr{W}G} \rightarrow \Lambda^*(T_{G^{sc}})$$

$$b \mapsto b(d^{ss}, -)$$

is injective.

(ii) *Let $G^{\text{ad}} = G_1^{\text{ad}} \times \dots \times G_s^{\text{ad}}$ be the decomposition of G^{ad} into almost simple factors (which are then automatically semisimple adjoint groups) and choose maximal tori in such a way that $T_{G^{\text{ad}}} = T_{G_1^{\text{ad}}} \times \dots \times T_{G_s^{\text{ad}}}$. Then, d^{ss} satisfies the following condition:*

$$d^{ss} = d_1 + \dots + d_s \in \Lambda(T_{G^{\text{ad}}}) = \Lambda(T_{G_1^{\text{ad}}}) \oplus \dots \oplus \Lambda(T_{G_s^{\text{ad}}}),$$

with $d_i \neq 0$ for every $1 \leq i \leq s$.

(2) *Any $\delta \in \pi_1(G)$ admits a lift $d \in \pi_1(T_G) = \Lambda(T_G)$, which is generic.*

We collect in the following Fact the main results proved in [21] for the Picard group of $\text{Bun}_G^\delta(C/S)$.

Fact 3.2. *Let G be a reductive group and fix $\delta \in \pi_1(G)$.*

(1) *There exists an exact sequence*

$$0 \rightarrow \text{Pic}(S) \rightarrow \text{Pic Bun}_G^\delta(C/S) \rightarrow \text{Pic Bun}_G^\delta(C_{\bar{\eta}}),$$

where the first map is the pull-back map and the second one is the restriction to the fibre of $C \rightarrow S$ over a geometric generic point $\bar{\eta}$ of S .

(2) *For any lift $d \in \pi_1(B_G) = \pi_1(T_G) = \Lambda(T_G)$ of δ (which is moreover generic if $g=0$), the pull-back maps along the inclusions $\iota : T_G \hookrightarrow B_G \hookrightarrow G$*

$$\iota_{\#}^* : \text{Pic Bun}_G^\delta(C/S) \hookrightarrow \text{Pic Bun}_{B_G}^d(C/S) \xrightarrow{\cong} \text{Pic Bun}_{T_G}^d(C/S)$$

are such that the first map is injective and the second one is an isomorphism.

(3) *There exists a unique homomorphism (called transgression map)*

$$\tau_G^\delta = \tau_G^\delta(C/S) : (\text{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}G} \hookrightarrow \text{Pic Bun}_G^\delta(C/S) \tag{3.1}$$

that is functorial in S and G and fits in the following commutative diagram for any lift $d \in \pi_1(T_G)$ (which is generic if $g=0$) of $\delta \in \pi_1(G)$:

$$\begin{array}{ccc}
 (\mathrm{Sym}^2 \Lambda^*(T_G))^{\#_G} & \xrightarrow{\tau_G^\delta} & \mathrm{Pic} \mathrm{Bun}_G^\delta(C/S), \\
 \downarrow & & \downarrow t_\#^* \\
 \mathrm{Sym}^2 \Lambda^*(T_G) & \xrightarrow{\tau_{T_G}^d} & \mathrm{Pic} \mathrm{Bun}_{T_G}^d(C/S),
 \end{array} \tag{3.2}$$

$$\chi \cdot \chi' \xrightarrow{\quad\quad\quad} \langle \mathcal{L}_\chi, \mathcal{L}_{\chi'} \rangle_{\hat{\pi}}$$

11

where $\langle -, - \rangle_\pi$ is the Deligne pairing of the family $\hat{\pi} : C \times_S \mathrm{Bun}_{T_G}^d(C/S) \times \mathrm{Bun}_{T_G}^d(C/S)$ and \mathcal{L}_χ is the pull-back of the universal \mathbb{G}_m -bundle (i.e. line bundle) over $C \times_S \mathrm{Bun}_{\mathbb{G}_m}(C/S)$ via the morphism $\mathrm{id} \times \chi_\# : C \times_S \mathrm{Bun}_G^\delta(C/S) \rightarrow C \times_S \mathrm{Bun}_{\mathbb{G}_m}^{(\delta, \chi)}(C/S)$.

Proof. Part (1) follows from [21, Prop. 5.1.1]. Part (2) follows from [21, Cor. 5.1.2, Thm. 6.0.1]. Part (3) follows from [21, Thm. 5.0.1, Rmk. 5.2.2] (whose proof works also for $g = 0$). □

Using Fact 3.2(1), we will often restrict to study the *relative Picard group*

$$\mathrm{RPic} \mathrm{Bun}_G^\delta(C/S) := \frac{\mathrm{Pic} \mathrm{Bun}_G^\delta(C/S)}{\mathrm{Pic}(S)}. \tag{3.3}$$

3.2. The Picard group of $\mathrm{Bun}_T^d(C/S)$ for a torus T

The goal of this subsection is to determine the structure of the Picard group of $\mathrm{Bun}_T^d(C/S)$, where

- ★ T is a torus and $d \in \Lambda(T)$.

We first introduce some natural line bundles on $\mathrm{Bun}_T^d(C/S)$. For any character $\chi \in \Lambda^*(T)$ and $M \in \mathrm{Pic}(C)$, we consider the following line bundle on $C \times_S \mathrm{Bun}_T^d(C/S)$:

$$\mathcal{L}_\chi(M) := \mathcal{L}_\chi \otimes p_1^* M,$$

where p_1 is the projection onto the first factor and \mathcal{L}_χ is the pull-back of the universal \mathbb{G}_m -bundle (i.e. line bundle) over $C \times_S \mathrm{Bun}_{\mathbb{G}_m}(C/S)$ via the morphism $\mathrm{id} \times \chi_\# : C \times_S \mathrm{Bun}_T^d(C/S) \rightarrow C \times_S \mathrm{Bun}_{\mathbb{G}_m}^{(d, \chi)}(C/S)$ induced by χ .

Definition 3.3. *The tautological Picard group of $\mathrm{Bun}_T^d(C/S)$ is the subgroup $\mathrm{Pic}^{\mathrm{taut}} \mathrm{Bun}_T^d(C/S)$ of $\mathrm{Pic} \mathrm{Bun}_T^d(C/S)$ generated by the pull-back of the line bundles on S and the following line bundles (called tautological line bundles):*

$$\{d_{\widehat{\pi}}(\mathcal{L}_{\chi}(M)) : \chi \in \Lambda^*(T), M \in \text{Pic}(C)\},$$

where $d_{\widehat{\pi}}$ denotes the determinant of cohomology with respect to the family $\widehat{\pi} : C \times_S \text{Bun}_T^d(C/S) \rightarrow \text{Bun}_T^d(C/S)$.

The relative tautological Picard group of $\text{Bun}_T^d(C/S)$ is the subgroup $\text{RPic}^{\text{taut}} \text{Bun}_T^d(C/S) \subseteq \text{RPic} \text{Bun}_T^d(C/S)$, which is the image of $\text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S)$ into the relative Picard group.

Remark 3.4. The tautological Picard group $\text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S)$ contains all the line bundles

$$\{\langle \mathcal{L}_{\chi}(M), \mathcal{L}_{\mu}(N) \rangle_{\widehat{\pi}} : \chi, \mu \in \Lambda^*(T), M, N \in \text{Pic}(C)\},$$

where $\langle -, - \rangle_{\widehat{\pi}}$ is the Deligne pairing, since we have the following isomorphism:

$$\langle \mathcal{L}_{\chi}(M), \mathcal{L}_{\mu}(N) \rangle_{\widehat{\pi}} \cong d_{\widehat{\pi}}(\mathcal{L}_{\chi+\mu}(M \otimes N)) \otimes d_{\widehat{\pi}}(\mathcal{L}_{\chi}(M))^{-1} \otimes d_{\widehat{\pi}}(\mathcal{L}_{\mu}(N))^{-1} \otimes d_{\widehat{\pi}}(\mathcal{O}). \tag{3.4}$$

The Picard group of $\text{Bun}_T^d(C/S)$ is endowed with a *weight function*

$$\text{Pic} \text{Bun}_T^d(C/S) \xrightarrow{w_T^d} \Lambda^*(T), \tag{3.5}$$

which is defined as it follows. Let \mathcal{F} be a line bundle on $\text{Bun}_T^d(C/S)$ and fix a k -point $\xi := (C_s, E)$ of $\text{Bun}_T^d(C/S)$ over $s \in S$. The automorphism group of ξ acts on the fibre \mathcal{F}_{ξ} of \mathcal{F} over ξ . Since the torus T is contained in the automorphism group of ξ , this defines an action of T on $\mathcal{F}_{\xi} \cong k$, which is given by a character of T . This character, which is independent of the chosen k -point ξ and on the chosen isomorphism $\mathcal{F}_{\xi} \cong k$, coincides with $w_T^d(\mathcal{F})$. Note that w_T^d factors through the relative Picard group.

The weight function on the tautological Picard group is easily described.

Lemma 3.5. *The weight function w_T^d is given on tautological line bundles by*

$$w_T^d(d_{\widehat{\pi}}(\mathcal{L}_{\chi}(M))) = [\chi(d) + \text{deg}_{\pi}(M) + 1 - g]\chi. \tag{3.6}$$

In particular, we have that

$$w_T^d(\langle \mathcal{L}_{\chi}(M), \mathcal{L}_{\mu}(N) \rangle_{\widehat{\pi}}) = [\mu(d) + \text{deg}_{\pi}(N)]\chi + [\chi(d) + \text{deg}_{\pi}(M)]\mu. \tag{3.7}$$

Proof. This is proved as in [21, Prop. 4.1.2(1)]. □

Using the weight function, we can now describe the structure of the Picard group of $\text{Bun}_T^d(C/S)$ for a family of curves of genus 0.

Theorem 3.6 *Assume that $g(C/S) = 0$. Then,*

$$(1) \text{ Pic} \text{Bun}_T^d(C/S) = \text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S).$$

(2) the weight function

$$w_T^d : \text{RPic Bun}_T^d(C/S) \hookrightarrow \Lambda^*(T)$$

is injective on the relative Picard group.

(3) The image of w_T^d is equal to

$$\text{Im}(w_T^d) = \begin{cases} \Lambda^*(T) & \text{if } C/S \text{ is Zariski-locally trivial,} \\ \{\chi \in \Lambda^*(T) : \chi(d) \text{ is even}\} & \text{if } C/S \text{ is not Zariski-locally trivial.} \end{cases}$$

The above Theorem was proved for the universal family $\mathcal{C}_{0,n} \rightarrow \mathcal{M}_{0,n}$ (which is Zariski-locally trivial if and only if $n > 0$, see Remark 2.2) in [21, Thm. 4.2].

Proof. Part (2): since $g(C/S) = 0$, the rigidification $\text{Bun}_T^d(C/S) // T$ is isomorphic to S . Hence, the Leray spectral sequence for \mathbb{G}_m associated to the T -gerbe $\text{Bun}_T^d(C/S) \rightarrow \text{Bun}_T^d(C/S) // T$ gives the desired exact sequence

$$0 \rightarrow \text{Pic}(S) = \text{Pic}(\text{Bun}_T^d(C/S) // T) \rightarrow \text{Pic Bun}_T^d(C/S) \xrightarrow{w_T^d} \text{Pic}(BT) = \Lambda^*(T).$$

Note that $\delta(C/S) = 1$ or 2 according to whether C/S is Zariski-locally trivial or not (see Remark 2.2(4)). Hence, using the above exact sequence, in order to prove parts (1) and (3), it is enough to prove the two equalities

$$w_T^d(\text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S)) = \text{Im}(w_T^d) = \begin{cases} \Lambda^*(T) & \text{if } \delta(C/S) = 1, \\ \{\chi \in \Lambda^*(T) : \chi(d) \text{ is even}\} & \text{if } \delta(C/S) = 2. \end{cases} \tag{3.8}$$

We will distinguish two cases.

Case 1: $\delta(C/S) = 1$.

In this case, the relative degree map (2.1) is an isomorphism. Hence, given any $\chi \in \Lambda^*(T)$, we can choose a line bundle $M_\chi \in \text{Pic}(C)$ having relative degree equal to $g - \chi(d)$, and then formula (3.6) implies that

$$w_T^d(d_{\tilde{\pi}}(\mathcal{L}_\chi(M_\chi))) = \chi.$$

This shows that

$$w_T^d(\text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S)) = \Lambda^*(T),$$

which gives the desired two equalities in Equation (3.8).

Case 2: $\delta(C/S) = 2$.

In this case, the image of the relative degree map (2.1) is the subgroup $2\mathbb{Z} \subset \mathbb{Z}$. We will divide the proof of the equalities in Equation (3.8) into three steps.

- $w_T^d(\text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S)) \subseteq \{\chi \in \Lambda^*(T) : \chi(d) \text{ is even}\}.$

To prove this, since $\text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S)$ is generated the tautological line bundles $d_{\widehat{\pi}}(\mathcal{L}_\chi(M))$, it is enough to show, using formula 3.6, that

$$d_{\widehat{\pi}}(\mathcal{L}_\chi(M))(d) = [\chi(d) + \text{deg}_\pi(M) + 1]\chi(d) \text{ is even}$$

for any $\chi \in \Lambda^*(T)$ and any $M \in \text{Pic}(C)$. This is obvious if $\chi(d)$ is even. Otherwise, if $\chi(d)$ is odd, then it follows from the fact that $\chi(d) + \text{deg}_\pi(M) + 1$ is even because $\text{deg}_\pi(M)$ is even.

- $w_T^d(\text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S)) \supseteq \{\chi \in \Lambda^*(T) : \chi(d) \text{ is even}\}$.

Let $\chi \in \Lambda^*(T)$ such that $\chi(d)$ is even. Then, we can choice a line bundle $M_\chi \in \text{Pic}(C)$ such that $\text{deg}_\pi(M_\chi) = -\chi(d)$. We then have, using formula (3.6), that

$$d_{\widehat{\pi}}(\mathcal{L}_\chi(M_\chi)) = [\chi(d) + \text{deg}_\pi(M_\chi) + 1]\chi = \chi,$$

which shows that $\chi \in w_T^d(\text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S))$.

- $\text{Im}(w_T^d) \subseteq \{\chi \in \Lambda^*(T) : \chi(d) \text{ is even}\}$.

Let L be a line bundle on $\text{Bun}_T^d(C/S)$ and set $\chi := w_T^d(L) \in \Lambda^*(T)$. We want to show that $\chi(d)$ is even. Consider the following Cartesian diagram

$$\begin{array}{ccccc}
 & & \text{Bun}_T^d(C/S) \times_S C & \longleftarrow & \text{Bun}_T^d(C \times_S C/C) \times_C (C \times_S C) \\
 & \swarrow & \downarrow \widehat{\pi} & \square & \downarrow \widehat{\pi}_2 \uparrow \widetilde{\Delta} \\
 C & \longleftarrow & \text{Bun}_T^d(C/S) & \xleftarrow{\pi_T} & \text{Bun}_T^d(C \times_S C/C) & \square & \longrightarrow & C \times_S C \\
 & \searrow \pi & \downarrow \Phi_S & \square & \downarrow \Phi_C & \nearrow \pi_2 & \nearrow \Delta \\
 & & S & \xleftarrow{\pi} & C & & &
 \end{array} \tag{3.9}$$

The pull-back $\pi_T^*(L)$ on $\text{Bun}_T^d(C \times_S C/C)$ verifies $w_T^d(\pi_T^*(L)) = \chi$, since the weight function is functorial with respect to base change. Consider now the line bundle $\langle \mathcal{L}_\chi, \mathcal{O}(\mathcal{O}(\widetilde{\Delta})) \rangle_{\widehat{\pi}_2} = \langle \mathcal{L}_\chi, \mathcal{O}(\widetilde{\Delta}) \rangle_{\widehat{\pi}_2}$ on $\text{Bun}_T^d(C \times_S C/C)$. By Equation (3.7) and the fact that $\mathcal{O}(\Delta)$ has π_2 -relative degree equal to 1, we have that

$$w_T^d(\langle \mathcal{L}_\chi, \mathcal{O}(\widetilde{\Delta}) \rangle_{\widehat{\pi}_2}) = \chi.$$

Therefore, since $\pi_T^*(L)$ and $\langle \mathcal{L}_\chi, \mathcal{O}(\widetilde{\Delta}) \rangle_{\widehat{\pi}_2}$ have the same weight, part (2) implies that

$$\pi_T^*(L) = \langle \mathcal{L}_\chi, \mathcal{O}(\widetilde{\Delta}) \rangle_{\widehat{\pi}_2} \otimes \Phi_C^*(M)$$

for some $M \in \text{Pic}(C)$. Taking the relative degree with respect to π_T , we obtain that

$$\begin{aligned}
 0 &= \text{deg}_{\pi_T}(\pi_T^*(L)) = \text{deg}_{\pi_T}(\langle \mathcal{L}_\chi, \mathcal{O}(\widetilde{\Delta}) \rangle_{\widehat{\pi}_2}) + \text{deg}_{\pi_T}(\Phi_C^*(M)) = \\
 &= \text{deg}_{\pi_T} \widetilde{\Delta}^*(\mathcal{L}_\chi) + \text{deg}_\pi(M) = \chi(d) + \text{deg}_\pi(M),
 \end{aligned}$$

where we have used the isomorphism $\langle \mathcal{L}_\chi, \mathcal{O}(\widetilde{\Delta}) \rangle_{\widehat{\pi}_2} \cong \widetilde{\Delta}^*(\mathcal{L}_\chi)$ has π_T -relative degree equal to $\chi(d)$. This implies that $\chi(d)$ is even since $\text{deg}_\pi(M)$ is even because of the hypothesis that $\delta(C/S) = 2$. □

We will now focus on families of curves of positive genus. In this case, the relative tautological Picard group is endowed with the following homomorphism:

Lemma 3.7. *Assume that $g(C/S) > 0$. There exists a homomorphism*

$$\begin{aligned} \gamma_T^d : \text{RPic}^{\text{taut}} \text{Bun}_T^d(C/S) &\longrightarrow \text{Bil}^s \Lambda(T) = (\Lambda^*(T) \otimes \Lambda^*(T))^s, \\ d_{\widehat{\pi}}(\mathcal{L}_\chi(M)) &\mapsto \chi \otimes \chi. \end{aligned}$$

In particular, we have that

$$\gamma_T^d(\langle \mathcal{L}_\chi(M), \mathcal{L}_\mu(N) \rangle_{\widehat{\pi}}) = \chi \otimes \mu + \mu \otimes \chi.$$

Proof. This is proved as in [21, Prop. 4.1.2(2)] (see also [22, Def./Lemma 3.5(2)]). \square

Using the above homomorphism γ_T^d , we can now describe the (relative) tautological Picard group of $\text{Bun}_T^d(C/S)$ for families of curves of positive genus.

Theorem 3.8. *Assume that $g = g(C/S) > 0$.*

(1) *Then there exists a commutative diagram with exact rows, functorial in S and T*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Lambda^*(T) \otimes \text{RPic}^0(C/S) & \xrightarrow{j_T^d} & \text{RPic}^{\text{taut}} \text{Bun}_T^d(C/S) & \xrightarrow{w_T^d \oplus \gamma_T^d} & \Lambda^*(T) \oplus \text{Bil}^s(\Lambda(T)) \\ & & \downarrow & & \parallel & & \downarrow \\ 0 & \longrightarrow & \Lambda^*(T) \otimes \text{RPic}(C/S) & \xrightarrow{i_T^d} & \text{RPic}^{\text{taut}} \text{Bun}_T^d(C/S) & \xrightarrow{\gamma_T^d} & \text{Bil}^s(\Lambda(T)) \longrightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow \epsilon \\ 0 & \longrightarrow & \Lambda^*(T) \otimes \text{RPic}(C/S) \oplus \text{Sym}^2 \Lambda^*(T) & \xrightarrow{i_T^d \oplus \tau_T^d} & \text{RPic}^{\text{taut}} \text{Bun}_T^d(C/S) & \xrightarrow{\rho_T^d} & \text{Hom}(\Lambda(T), \mathbb{Z}/2\mathbb{Z}) \longrightarrow 0 \end{array} \quad (3.10)$$

where

- $\text{RPic}(C/S)$ is the relative Picard group of C/S and $\text{RPic}^0(C/S)$ is the subgroup consisting of line bundles of relative degree 0;
- i_T^d is defined by

$$i_T^d(\chi \otimes M) = \langle \mathcal{L}_\chi, M \rangle_{\widehat{\pi}};$$

- j_T^d is the restriction of i_T^d to $\Lambda^*(T) \otimes \text{RPic}^0(C/S)$;
- τ_T^d is the transgression map (see Fact 3.2(3));
- γ_T^d is the map of Lemma 3.7;
- ρ_T^d is defined by

$$\rho_T^d(d_{\widehat{\pi}}(\mathcal{L}_\chi(M)))(x) = \chi(x)^2 \pmod{2};$$

- ϵ is defined by

$$\epsilon(b)(x) := b(x, x) \pmod{2}.$$

(2) The image of $w_T^d \oplus \gamma_T^d$ is equal to

$$\{(\chi, b) \in \Lambda^*(T) \oplus \text{Bil}^s(\Lambda(T)) : \delta(C/S)|\chi(x) - b(d, x) + (g - 1)b(x, x) \text{ for every } x \in \Lambda(T)\}.$$

Most of the results of the previous theorem were known for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ with $g \geq 1$: the exactness of the first line in Equation (3.10) and part (2) were proved in [21, Prop. 4.3.1]; the exactness of the second line in Equation (3.10) was proved in [22, Thm. 3.6] (where it is also proved for an arbitrary reductive group); the exactness of the third line in Equation (3.10) was proved for $g = 1$ in [21, Thm. B] (while for genus $g \geq 2$, a different presentation is obtained in loc. cit.).

Proof. Let us first prove part (1). Observe that the diagram is commutative: the commutativity of the two left squares and the upper right square are obvious; the commutativity of the lower right square follows from the fact that $(\chi \otimes \chi)(x, y) = \chi(x)\chi(y)$.

We now divide the rest of the proof of part (1) into three steps.

Step 1: The middle row is exact.

First of all, observe that i_T^d is well-defined since the Deligne pairing $\langle -, - \rangle_{\hat{\pi}}$ is bilinear and it is trivial on the pull-back of line bundle on $\text{Bun}_T^d(C/S)$, and hence in particular on those coming from S . Consider now the following diagram

$$\begin{CD} \Lambda^*(T) \otimes \text{RPic}(C/S) @>i_T^d>> \text{RPic}^{\text{taut}} \text{Bun}_T^d(C/S) \\ @VVV @VVV \\ \Lambda^*(T) \otimes \text{Pic}(C_{\bar{\eta}}) @>i_{C_{\bar{\eta}}}^d>> \text{RPic}^{\text{taut}} \text{Bun}_T^d(C_{\bar{\eta}}) \end{CD}$$

where the vertical maps are restriction maps to the geometric generic fibre $C_{\bar{\eta}}$ of C/S and the map $i_{C_{\bar{\eta}}}^d$ is the restriction of i_T^d to the the geometric generic fibre. Since the restriction maps are injective by Fact 3.2(1) and [21, Prop. 2.3.2] and the map $i_T^d(C_{\bar{\eta}})$ is injective by [7, § 3], we deduce that i_T^d is also injective.

Lemma 3.7 implies that $\text{Im}(i_T^d) \subseteq \ker(\gamma_T^d)$ and that γ_T^d is surjective, using that $(\Lambda^*(T) \otimes \Lambda^*(T))^s$ is generated by the elements $\{\chi \otimes \chi : \chi \in \Lambda^*(T)\}$ (see [21, § 2.2]).

It remains to prove that $\ker(\gamma_T^d) \subseteq \text{Im}(i_T^d)$. To this aim, we fix a basis $\{\chi_1, \dots, \chi_r\}$ of $\Lambda^*(T)$. Then, $\text{RPic}^{\text{taut}}(\text{Bun}_T^d(C/S))$ is generated by the line bundles

$$\{d_{\hat{\pi}}(\mathcal{L}_{\chi_i}), \langle \mathcal{L}_{\chi_j}, \mathcal{L}_{\chi_k} \rangle_{\hat{\pi}}, \langle \mathcal{L}_{\chi_h}, M \rangle_{\hat{\pi}}\},$$

with $1 \leq i, h \leq r, 1 \leq j \leq k \leq r$ and $M \in \text{RPic}(C/S)$. The line bundles $\langle \mathcal{L}_{\chi_h}, M \rangle_{\hat{\pi}}$ belong to $\text{Im}(i_T^d) \subseteq \ker(\gamma_T^d)$ as proved above. Suppose that we have a line bundle on $\text{Bun}_T^d(C/S)$

$$K := \bigotimes_i d_{\hat{\pi}}(\mathcal{L}_{\chi_i})^{a_i} \bigotimes_{j \leq k} \langle \mathcal{L}_{\chi_j}, \mathcal{L}_{\chi_k} \rangle_{\hat{\pi}}^{b_{jk}} \text{ for some } a_i, b_{jk} \in \mathbb{Z}$$

whose class in the relative tautological Picard group belongs to $\ker(\gamma_T^d)$. Then, Lemma 3.7 implies that

$$0 = \gamma_T^d \left(\bigotimes_i d_{\hat{\pi}}(\mathcal{L}_{\chi_i})^{a_i} \bigotimes_{j \leq k} \langle \mathcal{L}_{\chi_j}, \mathcal{L}_{\chi_k} \rangle_{\hat{\pi}}^{b_{jk}} \right) = \sum_i a_i \chi_i \otimes \chi_i + \sum_{j \leq k} b_{jk} (\chi_j \otimes \chi_k + \chi_k \otimes \chi_j). \tag{3.11}$$

Since a basis of $\text{Bil}^s(\Lambda(T)) = (\Lambda^*(T) \otimes \Lambda^*(T))^s$ is given by $\{\{\chi_i \otimes \chi_i\}_i \cup \{\chi_j \otimes \chi_k + \chi_k \otimes \chi_j\}_{j < k}\}$, the relation Equation (3.11) is equivalent to

$$\begin{cases} b_{jk} = 0 & \text{for any } j < k, \\ a_i + 2b_{ii} = 0 & \text{for any } i. \end{cases}$$

Hence, the line bundle K is equal to

$$K = \bigotimes_i [d_{\hat{\pi}}(\mathcal{L}_{\chi_i})^{-2} \bigotimes \langle \mathcal{L}_{\chi_i}, \mathcal{L}_{\chi_i} \rangle_{\hat{\pi}}^{b_{ii}}] = \bigotimes_i [\langle \mathcal{L}_{\chi_i}, \omega_{\hat{\pi}} \rangle_{\hat{\pi}} \otimes d_{\hat{\pi}}(\mathcal{O})^{-2}]^{b_{ii}},$$

where the last equality follows from [21, Rmk. 3.5.1]. This implies that the class of K in the relative Picard group belongs to $\text{Im}(i_T^d)$, and we are done.

Step 2: The first row is exact.

Indeed, the map j_T^d is injective since it is the restriction of the injective map i_T^d . Using the exactness of the middle row, the kernel of $w_T^d \oplus \gamma_T^d$ is equal to

$$\ker(w_T^d \oplus \gamma_T^d) = \{i_T^d(\chi \otimes M) : \text{wt}_T^d(i_T^d(\chi \otimes M)) = 0\}.$$

By the definition of i_T^d and Equation (3.7), we compute

$$\text{wt}_T^d(i_T^d(\chi \otimes M)) = w_T^d(\langle \mathcal{L}_{\chi}, M \rangle_{\hat{\pi}}) = \text{deg}_{\pi}(M)\chi.$$

Hence, the kernel of $w_T^d \oplus \gamma_T^d$ coincides with the image via i_T^d (or equivalently via j_T^d) of $\Lambda^*(T) \otimes \text{RPic}^0(C/S)$, and we are done.

Step 3: The last row is exact.

Indeed, the composition $\gamma_T^d \circ \tau_T^d$ is given by

$$(\gamma_T^d \circ \tau_T^d)(\gamma \cdot \gamma') = \gamma_T^d(\langle \mathcal{L}_{\gamma}, \mathcal{L}_{\gamma'} \rangle_{\hat{\pi}}) = \gamma \otimes \gamma' + \gamma' \otimes \gamma.$$

Hence, $\gamma_T^d \circ \tau_T^d$ is injective, and its image is the subgroup $\text{Bil}^{s,\text{ev}}(\Lambda(T)) \subset \text{Bil}^s(\Lambda(T))$ of even symmetric bilinear forms (see [21, § 2.2]). We now deduce the exactness of the last row from the exactness of the second row and the fact that $\text{Bil}^{s,\text{ev}}(\Lambda(T))$ is the kernel of the surjective map ϵ .

Let us prove now part (2). We first observe that the image of $w_T^d \oplus \gamma_T^d$ is contained in

$$I := \{(\chi, b) \in \Lambda^*(T) \oplus \text{Bil}^s(\Lambda(T)) : \delta(C/S)|\chi(x) - b(d, x) + (g - 1)b(x, x) \text{ for every } x \in \Lambda(T)\}.$$

Indeed, for the generators of $\text{RPic}^{\text{taut}} \text{Bun}_T^d(C/S)$, we compute (using Equation (3.6) and Lemma 3.7) that

$$(w_T^d, \gamma_T^d)(d_{\bar{\pi}}(\mathcal{L}_\chi(M))) = (\chi(d) + \text{deg}_\pi(M) + 1 - g)\chi, \chi \otimes \chi).$$

Hence, for any $x \in \Lambda(T)$, we get

$$\begin{aligned} [\chi(d) + \text{deg}_\pi(M) + 1 - g]\chi(x) - \chi(d)\chi(x) + (g - 1)\chi(x)^2 &= [\text{deg}_\pi(M) + 1 - g]\chi(x) \\ &\quad + (g - 1)\chi(x)^2 = \\ &= \text{deg}_\pi(M)\chi(x) + (g - 1)[\chi(x)^2 - \chi(x)] \equiv \text{deg}_\pi(M)\chi(x) \pmod{2g - 2} \\ &\equiv 0 \pmod{\delta(C/S)}, \end{aligned}$$

where in the last congruence relation we used the fact that $\text{deg}_\pi(M)$ is a multiple of $\delta(C/S)$ (by definition of $\delta(C/S)$) and that $\delta(C/S)$ is a multiple of $2g - 2$ (by Remark 2.2(1)).

Next, in order to show that the inclusion $\text{Im}(w_T^d \oplus \gamma_T^d) \subseteq I$ is an equality, we consider the following commutative diagram with exact vertical columns:

$$\begin{array}{ccccc} \text{Im}(w_T^d \oplus \gamma_T^d) \cap (\Lambda^*(T) \times \{0\}) & \hookrightarrow & I \cap (\Lambda^*(T) \times \{0\}) & \hookrightarrow & \Lambda^*(T) \\ \downarrow & & \downarrow & & \downarrow \\ \text{Im}(w_T^d \oplus \gamma_T^d) & \hookrightarrow & I & \hookrightarrow & \Lambda^*(T) \oplus \text{Bil}^s(\Lambda(T)) \\ \downarrow & & \downarrow & & \downarrow p_2 \\ p_2(\text{Im}(w_T^d \oplus \gamma_T^d)) & \hookrightarrow & p_2(I) & \hookrightarrow & \text{Bil}^s(\Lambda(T)) \end{array}$$

We now conclude using that

- $p_2(\text{Im}(w_T^d \oplus \gamma_T^d)) = \text{Bil}^s(\Lambda(T))$ as it follows from the right upper square of the diagram (3.10);
- the inclusion

$$\text{Im}(w_T^d \oplus \gamma_T^d) \cap (\Lambda^*(T) \times \{0\}) \subseteq I \cap (\Lambda^*(T) \times \{0\}) = \delta(C/S) \cdot \Lambda^*(T)$$

is an equality since it coincides with the boundary homomorphism coming from the snake lemma applied to the first two horizontal lines of the diagram (3.10). □

After having obtained in Theorem 3.8, a complete description of the tautological Picard group of $\text{Bun}_T^d(C/S)$ for families of curves of positive genus, the next natural question is to determine when the entire Picard group coincides with the tautological subgroup. A sufficient condition is provided by the following:

Theorem 3.9 Assume that $g = g(C/S) > 0$. If $\text{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$ and $\text{RPic}(C/S) = \text{Pic}_{C/S}(S)$, then

$$\text{Pic Bun}_T^d(C/S) = \text{Pic}^{\text{taut}} \text{Bun}_T^d(C/S).$$

The above theorem was shown in [7] (see also [21, Prop. 4.1.1]) for a curve C over $S = \text{Spec } k$, with k an algebraically closed field, in which case the second assumption is automatic and the first assumption $\text{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$ is also necessary for the theorem to hold. Theorem 3.9 recovers the equality $\text{Pic Bun}_T^d(\mathcal{C}_{g,n}/\mathcal{M}_{g,n}) = \text{Pic}^{\text{taut}} \text{Bun}_T^d(\mathcal{C}_{g,n}/\mathcal{M}_{g,n})$, which was shown in [21, Thm. 4.1] for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ (with $g \geq 1$), since the assumption of the Theorem is known to hold for $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ (see Remarks 2.3 and 2.5(1)).

Proof. We will distinguish three cases.

Case I: $d = 0$ and $\pi : C \rightarrow S$ admits a section σ .

Since $d = 0$, we have the canonical identifications

$$\begin{cases} \text{Bun}_T^0(C/S) = \mathcal{J}^0(C/S) \otimes \Lambda(T), \\ \text{Bun}_T^0(C/S) // T = J^0(C/S) \otimes \Lambda(T), \end{cases}$$

where $\mathcal{J}^0(C/S)$ is the Jacobian stack of C/S and $J^0(C/S) = \mathcal{J}^0(C/S) // T$ is the Jacobian scheme of C/S . The theorem is implied in this case by the following commutative diagram with exact rows:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Lambda^*(T) \otimes \text{RPic}^0(C/S) & \xrightarrow{j_T^0} & \text{RPic}^{\text{taut}} \text{Bun}_T^0(C/S) & \xrightarrow{w_T^0 \oplus \gamma_T^0} & \Lambda^*(T) \oplus \text{Bil}^s(\Lambda(T)) \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & (J^0(C/S) \otimes \Lambda(T))^\vee(S) & \longrightarrow & \text{RPic} \text{Bun}_T^0(C/S) & \xrightarrow{\text{res}_T^0(C_{\bar{\eta}})} & \text{NS}(\text{Bun}_T^0(C_{\bar{\eta}})) \end{array} \tag{3.12}$$

where

- the first line is exact by Theorem 3.8 using that $\delta(C/S) = 1$ since $\pi : C \rightarrow S$ has a section by assumption;
- the second exact line comes from the fact that the rigidification $\text{Bun}_T^0(C/S) // T = J^0(C/S) \otimes \Lambda(T) \rightarrow S$ is an abelian scheme (see e.g. the proof of [19, Prop. 3.6]), where $(J^0(C/S) \otimes \Lambda(T))^\vee = J^0(C/S)^\vee \otimes \Lambda^*(T)$ is the dual abelian scheme and the last map is the restriction homomorphism onto the geometric generic fibre;
- the left vertical isomorphism follows from the fact that $\text{RPic}^0(C/S) = J^0(C/S)^\vee(S)$;
- the right vertical isomorphism follows from the fact that (see [7, § 3.2])

$$\text{NS}(\text{Bun}_T^0(C_{\bar{\eta}})) = \Lambda^*(T) \oplus \text{Hom}^s(\Lambda(T) \otimes \Lambda(T), \text{End}(J_{C_{\bar{\eta}}}))$$

where $\text{Hom}^s(\Lambda(T) \otimes \Lambda(T), \text{End}(J_{C_{\overline{\eta}}}))$ is the set of bilinear forms $\text{Hom}(\Lambda(T) \otimes \Lambda(T), \text{End}(J_{C_{\overline{\eta}}}))$ that are symmetric with respect to the Rosati involution, together with the assumption that $\text{End}(J_{C_{\overline{\eta}}}) = \mathbb{Z}$.

- the commutativity of the left square follows from the proof of [21, Lemma 4.2.1]¹ together with formula (3.4);
- the commutativity of the right square follows from [21, Prop. 4.1.2]².

Case II: $\pi : C \rightarrow S$ admits a section σ .

Fix an isomorphism $T \cong \mathbb{G}_m^r$, which induces a canonical isomorphism $\Lambda(T) \cong \mathbb{Z}^r$ under which d corresponds to (d_1, \dots, d_r) . Then, we have an isomorphism

$$\text{Bun}_T^d(C/S) \cong \text{Bun}_{\mathbb{G}_m}^{d_1}(C/S) \times_S \dots \times_S \text{Bun}_{\mathbb{G}_m}^{d_r}(C/S).$$

Using the section σ , we define an isomorphism

$$\begin{aligned} \mathfrak{t} : \text{Bun}_{\mathbb{G}_m}^{d_1}(C/S) \times_S \dots \times_S \text{Bun}_{\mathbb{G}_m}^{d_r}(C/S) &\xrightarrow{\cong} \text{Bun}_{\mathbb{G}_m}^0(C/S) \times_S \dots \times_S \text{Bun}_{\mathbb{G}_m}^0(C/S) \\ &\cong \text{Bun}_T^0(C/S) \\ (L_1, \dots, L_r) &\mapsto (L_1(-d_1\sigma), \dots, L_r(-d_r\sigma)). \end{aligned}$$

By construction, we have that $(\mathfrak{t} \times \text{id})^*(\mathcal{L}_{e_i}) = \mathcal{L}_{e_i}(-d_i\sigma)$, where $\{e_i\}$ is the canonical basis of $\mathbb{Z}^r \cong \Lambda^*(T)$. Hence, using the functoriality of the determinant of cohomology, we deduce that the pull-back isomorphism

$$\mathfrak{t}^* : \text{Pic Bun}_T^0(C/S) \xrightarrow{\cong} \text{Pic Bun}_T^d(C/S)$$

preserves the tautological subgroups. Hence, the conclusion is implied by Case I.

Case III: General case.

Let $\pi : C \rightarrow S$ be an arbitrary family of curves (i.e. possibly without a section). Fix a smooth cover $S' \rightarrow S$ such that the family $C' := C \times_S S' \rightarrow S'$ admits a section (e.g. $C \rightarrow S$). By Case II, we have the following morphism of exact sequences:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Lambda^*(T) \otimes \text{RPic}(C'/S') & \longrightarrow & \text{RPic}(\text{Bun}_T^d(C'/S')) & \longrightarrow & \text{Bil}^s(\Lambda(T)) \longrightarrow 0 \\ & & \downarrow \text{Id}_{\Lambda^*(T)} \otimes (p_1^* - p_2^*) & & \downarrow p_1^* - p_2^* & & \downarrow 0 \\ 0 & \longrightarrow & \Lambda^*(T) \otimes \text{RPic}(C''/S'') & \longrightarrow & \text{RPic}(\text{Bun}_T^d(C''/S'')) & \longrightarrow & \text{Bil}^s(\Lambda(T)) \longrightarrow 0 \end{array} \tag{3.13}$$

where $(C'' \rightarrow S'') := (C' \times_{S'} S' \rightarrow S' \times_{S'} S')$ and $p_1, p_2 : C'' \rightarrow C'$ denotes the natural projections.

Consider the following commutative diagram of abelian groups:

¹the proof of loc. cit. for the universal family $\mathcal{C}_{g,n}/\mathcal{M}_{g,n}$ extends mutatis mutandis to an arbitrary family C/S .

²the proof of loc. cit. for the universal family $\mathcal{C}_{g,n}/\mathcal{M}_{g,n}$ extends mutatis mutandis to an arbitrary family C/S .

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \Lambda^*(T) \otimes \mathrm{RPic}(C/S) & \xrightarrow{i_T^d} & \mathrm{RPic}^{\mathrm{taut}}(\mathrm{Bun}_T^d(C/S)) & \longrightarrow & \mathrm{Bil}^s(\Lambda(T)) \longrightarrow 0 \\
 & & \parallel & & \downarrow \iota^{\mathrm{taut}} & & \downarrow \\
 0 & \longrightarrow & \Lambda^*(T) \otimes \mathrm{RPic}(C/S) & \xrightarrow{\iota^{\mathrm{taut}} \circ i_T^d} & \mathrm{RPic}(\mathrm{Bun}_T^d(C/S)) & \longrightarrow & K := \mathrm{coker}\{\iota^{\mathrm{taut}} \circ i_T^d\} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & \Lambda^*(T) \otimes \mathrm{Pic}_{C/S}(S) & \longrightarrow & \ker\{p_1^* - p_2^*\} & \longrightarrow & \mathrm{Bil}^s(\Lambda(T))
 \end{array}$$

where all the rows are exact: the first one by Theorem 3.8, the second one by definition and the third one is obtained by taking the kernels of the vertical arrows in Diagram (3.13).

By applying the snake lemma at the first and the third rows, and using the assumption $\mathrm{RPic}(C/S) = \mathrm{Pic}_{C/S}(S)$, we deduce that the central vertical arrows in the middle must be equalities, and thus, we have the assertion. \square

Remark. If we remove the hypothesis $\mathrm{RPic}(C/S) = \mathrm{Pic}_{C/S}(S)$, Theorem 3.9 is false. Indeed, from the proof of Case III, we may deduce the following exact sequence of abstract groups:

$$0 \rightarrow \Lambda^*(T) \otimes \mathrm{Pic}_{C/S}(C) \rightarrow \mathrm{Pic}_{\mathrm{Bun}_T^d(C/S)/S}(S) \rightarrow \mathrm{Bil}^s(\Lambda(T)) \rightarrow 0,$$

where $\mathrm{Pic}_{\mathrm{Bun}_T^d(C/S)/S}$ denotes the relative Picard sheaf of the relative moduli stack $\mathrm{Bun}_T^d(C/S) \rightarrow S$. If we assume $\mathrm{RPic}(C/S) \subsetneq \mathrm{Pic}_{C/S}(S)$ and $\mathrm{RPic}(\mathrm{Bun}_T^d(C/S)) = \mathrm{Pic}_{\mathrm{Bun}_T^d(C/S)/S}(S)$ (e.g. if $d=0$, because $\mathrm{Bun}_T^0(C/S) \rightarrow S$ has sections), we get $\mathrm{RPic}^{\mathrm{taut}}(\mathrm{Bun}_T^d(C/S)) \subsetneq \mathrm{RPic}(\mathrm{Bun}_T^d(C/S))$.

3.3. The Picard group of $\mathrm{Bun}_G^\delta(C/S)$ for G reductive

In this subsection, we study when the Picard group of $\mathrm{Bun}_G^\delta(C/S)$ is generated by the image of the pull-back along the abelianization map

$$\mathrm{ab}_\# : \mathrm{Bun}_G^\delta(C/S) \rightarrow \mathrm{Bun}_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}}(C/S),$$

where $\delta^{\mathrm{ab}} := \pi_1(\mathrm{ab})(\delta)$, and the image of the transgression map (3.1).

More precisely, we investigate when the following commutative diagram, which is functorial in G and S :

$$\begin{array}{ccc}
 \mathrm{Sym}^2 \Lambda^*(G^{\mathrm{ab}}) \xrightarrow{\mathrm{Sym}^2 \Lambda_{\mathrm{ab}}^*} & \mathrm{Sym}^2 \Lambda^*(T_G) & \\
 \downarrow \tau_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}} & & \downarrow \tau_G^\delta \\
 \mathrm{Pic} \mathrm{Bun}_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}}(C/S) & \xrightarrow{\mathrm{ab}_\#} & \mathrm{Pic} \mathrm{Bun}_G^\delta(C/S)
 \end{array} \tag{3.14}$$

is a push-out diagram, where τ_G^δ and $\tau_{G^{\mathrm{ab}}}^{\delta^{\mathrm{ab}}}$ are, respectively, the transgression maps for G and G^{ab} (see Fact 3.2(3)), $\mathrm{Sym}^2 \Lambda_{\mathrm{ab}}^*$ is the injective homomorphism induced by the

inclusion $\Lambda_{ab}^* : \Lambda^*(G^{ab}) \hookrightarrow \Lambda^*(T_G)$, which is invariant under the \mathscr{W}_G -action on $\Lambda^*(T_G)$, and ab_{\sharp}^* is injective by the following:

Lemma 3.11. *The pull-back map*

$$ab_{\sharp}^* : \text{Pic Bun}_{G^{ab}}^{\delta^{ab}}(C/S) \rightarrow \text{Pic Bun}_G^{\delta}(C/S)$$

is injective.

Proof. The composition

$$\phi : T_G \xrightarrow{\iota} G \xrightarrow{\text{ab}} G^{ab}$$

is a surjective homomorphism of tori, hence it admits a section. This implies that, for any lift $d \in \pi_1(T_G)$ of $\delta \in \pi_1(G)$, the homomorphism

$$\phi_{\sharp}^* : \text{Pic Bun}_{G^{ab}}^{\delta^{ab}}(C/S) \xrightarrow{ab_{\sharp}^*} \text{Pic Bun}_G^{\delta}(C/S) \xrightarrow{\iota_{\sharp}^*} \text{Pic Bun}_{T_G}^d(C/S)$$

is injective. Hence, also the homomorphism ab_{\sharp}^* must be injective. □

We will use the following easy lemma on push-out diagrams of abelian groups.

Lemma 3.12. *Consider a commutative diagram of abelian groups*

$$\begin{array}{ccc} A & \xrightarrow{i} & B \\ \downarrow & & \downarrow \\ C & \xrightarrow{j} & D \end{array} \tag{3.15}$$

where i and j are injective homomorphisms. Then Equation (3.15) is a push-out diagram if and only if the natural map $\text{coker } i \rightarrow \text{coker } j$ is an isomorphism.

Proof. This is easy and left to the reader. □

Hence, the diagram (3.14) is a push-out diagram if and only if the induced homomorphism

$$\text{coker}(\text{Sym}^2 \Lambda_{ab}^*) \rightarrow \text{coker}(ab_{\sharp}^*)$$

is an isomorphism.

The cokernel of $\text{Sym}^2 \Lambda_{ab}^*$ is described explicitly as follows:

Lemma 3.13.

- (1) *We have an exact sequence of lattices, functorial in G ,*

$$\begin{aligned}
 0 \rightarrow \mathrm{Sym}^2 \Lambda^*(G^{\mathrm{ab}}) \xrightarrow{\mathrm{Sym}^2 \Lambda_{\mathrm{ab}}^*} \mathrm{Sym}^2 \Lambda^*(T_G) \mathscr{W}_G \xrightarrow{\mathrm{res}_{\mathscr{D}}} \mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_{\mathscr{D}(G)})|\Lambda(T_{G^{\mathrm{ss}}})) \mathscr{W}_G \rightarrow 0, \\
 b \mapsto b|_{\Lambda(T_{\mathscr{D}(G)}) \otimes \Lambda(T_{\mathscr{D}(G)})}
 \end{aligned}
 \tag{3.16}$$

where we identify $\mathrm{Sym}^2 \Lambda^*(T_G)$ with the lattice $\mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_G))$ of even symmetric bilinear forms and the last group is defined by

$$\begin{aligned}
 & \mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_{\mathscr{D}(G)})|\Lambda(T_{G^{\mathrm{ss}}})) \mathscr{W}_G \\
 & := \left\{ b \in \mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_{\mathscr{D}(G)})) \mathscr{W}_G : b|_{\Lambda(T_{\mathscr{D}(G)}) \otimes \Lambda(T_{G^{\mathrm{ss}}})} \text{ is integral} \right\},
 \end{aligned}$$

where $b^{\mathbb{Q}}$ is the extension of b to a rational bilinear symmetric form on $\Lambda(T_{G^{\mathrm{ss}}})$.

- (2) If $G^{\mathrm{sc}} \cong G_1 \times \dots \times G_s$ is the decomposition of the universal cover G^{sc} of $\mathscr{D}(G)$ into simply connected quasi-simple groups, then $\mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_{\mathscr{D}(G)})|\Lambda(T_{G^{\mathrm{ss}}})) \mathscr{W}_G$ is a finite index sub-lattice of

$$(\mathrm{Sym}^2 \Lambda^*(T_{G^{\mathrm{sc}}})) \mathscr{W}_G = \mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_{G^{\mathrm{sc}}})) \mathscr{W}_G = \mathbb{Z}\langle b_1 \rangle \oplus \dots \oplus \mathbb{Z}\langle b_s \rangle,$$

where b_i is the basic inner product of G_i .

- (3) If $\mathscr{D}(G)$ is simply connected (i.e. $\mathscr{D}(G) = G^{\mathrm{sc}}$), then

$$\mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_{\mathscr{D}(G)})|\Lambda(T_{G^{\mathrm{ss}}})) \mathscr{W}_G = \mathrm{Bil}^{\mathrm{s},\mathrm{ev}}(\Lambda(T_{G^{\mathrm{sc}}})) \mathscr{W}_G.$$

Proof. Part (1) is proved in [22, Prop. 2.4]. Part (2) follows from the fact that we have finite index inclusions of lattices

$$\Lambda(T_{G^{\mathrm{sc}}}) \hookrightarrow \Lambda(T_{\mathscr{D}(G)}) \hookrightarrow \Lambda(T_{G^{\mathrm{ss}}})$$

together with [21, Lemma 2.2.1]. Part (3) follows from [7, Lemma 4.3.4]. □

A first case where Equation (3.15) is a push-out diagram is provided by the following:

Theorem 3.14. *Assume that G is a reductive group such that $\mathscr{D}(G)$ is simply connected and fix $\delta \in \pi_1(G)$. There exists an exact sequence functorial in S and G*

$$0 \rightarrow \mathrm{Pic} \mathrm{Bun}_{G^{\mathrm{ab}}}^{\delta,\mathrm{ab}}(C/S) \xrightarrow{\mathrm{ab}^*_{\#}} \mathrm{Pic} \mathrm{Bun}_G^{\delta}(C/S) \rightarrow (\mathrm{Sym}^2 \Lambda^*(T_{G^{\mathrm{sc}}})) \mathscr{W}_G \rightarrow 0 \tag{3.17}$$

together with a (non-canonical) splitting functorial in S .

In particular, the commutative diagram (3.15) is a push-out diagram.

This theorem was proved: for G semisimple by Faltings [17], for a fixed curve C over $k = \bar{k}$ by Biswas–Hoffmann [7] and for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ for $g \geq 1$ by the authors in [21, Thm. C(2)] and [22, Cor. 3.4].

Proof. The second assertion follows from the first one using Lemmas 3.12 and 3.13, and the assumption that $\mathscr{D}(G)$ is simply connected. Hence, it is enough to prove the first assertion.

Let $\bar{\eta}$ be a geometric generic point of S . The family $\pi : C \rightarrow S$ sits in the following Cartesian diagram of families of curves:

$$\begin{array}{ccccc}
 C_{\bar{\eta}} & \longrightarrow & C & \longrightarrow & C_g \\
 \downarrow & & \downarrow \pi & & \downarrow \\
 \bar{\eta} & \longrightarrow & S & \longrightarrow & \mathcal{M}_g
 \end{array}$$

The above diagram gives rise to the following commutative diagram with exact rows:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \text{RPic Bun}_{G^{\text{ab}},g}^{\delta^{\text{ab}}} & \xrightarrow{\text{ab}_{\sharp}^*} & \text{RPic Bun}_{G,g}^{\delta} & \longrightarrow & (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\#G} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \rho \\
 0 & \longrightarrow & \text{RPic Bun}_{G^{\text{ab}}(C/S)}^{\delta^{\text{ab}}} & \xrightarrow{\text{ab}_{\sharp}^*(C/S)} & \text{RPic Bun}_{G(C/S)}^{\delta} & \longrightarrow & \mathcal{Q} \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \sigma \\
 0 & \longrightarrow & \text{Pic Bun}_{G^{\text{ab}}(C/\bar{\eta})}^{\delta^{\text{ab}}} & \xrightarrow{\text{ab}_{\sharp}^*(C/\bar{\eta})} & \text{Pic Bun}_{G(C/\bar{\eta})}^{\delta} & \longrightarrow & (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\#G} \longrightarrow 0
 \end{array} \tag{3.18}$$

where

- the first line is exact by [22, Cor. 3.4] for $g \geq 1$ and by Lemma 3.15 for $g = 0$, where $\text{Bun}_{G,g}^{\delta} := \text{Bun}_{G}(C_g/\mathcal{M}_g)$ and similarly for G^{ab} ;
- \mathcal{Q} is the cokernel of $\text{ab}_{\sharp}^*(C/S)$, so that the second line is exact by definition;
- the third line is exact as it follows by applying [7, Thm. 5.3.1] to the morphism ab and using [7, Prop. 5.2.11] together with our hypothesis that $\mathcal{D}(G)$ is simple connected;
- the vertical morphisms in the left and central columns are injective by Fact 3.2(1);
- the composition $\sigma \circ \rho$ is the identity, which implies that ρ is injective and σ is surjective.

It remains to show that σ is injective (or, equivalently, that ρ is surjective), and that the exact sequence Equation (3.17) admits a functorial splitting.

To this aim, consider the surjective morphisms of tori $T_G \hookrightarrow G \xrightarrow{\text{ab}} G^{\text{ab}}$ and pick a section $\tilde{\iota} : G^{\text{ab}} \hookrightarrow T_G$. The composition $\iota : G^{\text{ab}} \xrightarrow{\tilde{\iota}} T_G \hookrightarrow G$ is a section of ab . The induced morphism

$$\iota_{\sharp}(C/S) : \text{Bun}_{G^{\text{ab}}(C/S)}^{\delta^{\text{ab}}} \hookrightarrow \text{Bun}_{G(C/S)}^{\delta}$$

is a section of the morphism $\text{ab}_{\sharp}(C/S)$. By taking the pull-back at the level of the Picard groups, we deduce that the inclusion

$$\text{ab}_{\sharp}^*(C/S) : \text{Pic Bun}_{G^{\text{ab}}(C/S)}^{\delta^{\text{ab}}} \hookrightarrow \text{Pic Bun}_{G(C/S)}^{\delta}$$

admits a splitting, which is functorial in S , namely

$$\iota_{\sharp}(C/S)^* : \text{Pic Bun}_{G(C/S)}^{\delta} \twoheadrightarrow \text{Pic Bun}_{G^{\text{ab}}(C/S)}^{\delta^{\text{ab}}}$$

This shows that the exact rows in Equation (3.15) admit compatible splittings, which implies that σ is injective, and we are done. \square

The following lemma extends [22, Cor. 3.4] to genus zero, for the reductive groups with simply connected derived subgroup.

Lemma 3.15. *Let G be a reductive group such that $\mathcal{D}(G)$ is simply connected. For every $n \geq 0$, there exists an exact sequence, functorial in G*

$$0 \rightarrow \text{Pic Bun}_{G^{\text{ab}},0,n}^{\delta^{\text{ab}}} \xrightarrow{\text{ab}_{\#}^*} \text{Pic Bun}_{G,0,n}^{\delta} \rightarrow (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\mathcal{W}G} \rightarrow 0. \tag{3.19}$$

Proof. We will use the notation and results from [21, § 5]. Since the map $\pi_1(\text{ab}) : \pi_1(G) \xrightarrow{\cong} \pi_1(G^{\text{ab}}) = \Lambda(G^{\text{ab}})$ is an isomorphism by the hypothesis that $\mathcal{D}(G)$ is simply connected (see Equation (2.6)), we can pick a lift $d \in \Lambda(T_G)$ of $\delta \in \pi_1(G)$, which is generic (see Definition/Lemma 3.1) and such that the following condition holds:

$$\delta^{\text{ab}} \text{ is 2-divisible in } \Lambda(G^{\text{ab}}) \implies d \text{ is 2-divisible in } \Lambda(T_G). \tag{*}$$

By [21, Thm. 5.2], there exists a commutative diagram

$$\begin{CD} \text{RPic}(\text{Bun}_{G^{\text{ab}},0,n}^{\delta^{\text{ab}}}) @<{\text{ab}_{\#}^*}<< \text{RPic}(\text{Bun}_{G,0,n}^{\delta}) @<{\iota_{\#}^d}<< \text{RPic}(\text{Bun}_{T_G,0,n}^d) \\ @V{\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}}}VV @V{\omega_G^{\delta}}VV @V{\omega_{T_G}^d}VV \\ \Lambda^*(G^{\text{ab}}) @<{\iota}<< \Omega_d^*(T_G) @<{\iota}<< \Lambda^*(T_G) \end{CD} \tag{3.20}$$

where $\iota : T_G \hookrightarrow G$.

By the definition of $\Omega_d(T_G)$ and the fact that d is generic (see Definition/Lemma 3.1), the contraction homomorphism defines an isomorphism

$$\begin{aligned} (d^{\text{ss}}, -) : (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\mathcal{W}G} &\xrightarrow{\cong} \frac{\Omega_d(T_G)}{\Lambda^*(G^{\text{ab}})} \subseteq \frac{\Lambda^*(T_G)}{\Lambda^*(G^{\text{ab}})} = \Lambda^*(T_{G^{\text{sc}}}), \\ b &\mapsto b(d^{\text{ss}}, -). \end{aligned} \tag{3.21}$$

Now, if $n \geq 1$, then the maps $\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ and ω_G^{δ} are isomorphisms by [21, Thm. 5.2(2)], and we conclude by putting together Equations (3.20) and (3.21). On the other hand, if $n = 0$, then [21, Thm. 5.2(2)] gives that

$$\begin{cases} \text{Im}(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}}) = \{\chi \in \Lambda^*(G^{\text{ab}}) : \chi(\delta^{\text{ab}}) \in 2\mathbb{Z}\}, \\ \text{Im}(\omega_G^{\delta}) = \{\chi \in \Omega_d^*(T_G) : \chi(d) \in 2\mathbb{Z}\}. \end{cases} \tag{3.22}$$

This implies that the commutative diagram of abelian groups is a pull-back diagram

$$\begin{array}{ccc}
 \Lambda^*(G^{\text{rab}}) & \hookrightarrow & \Omega_d^*(T_G) \\
 \uparrow & & \uparrow \\
 \text{Im}(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}}) & \hookrightarrow & \text{Im}(\omega_G^\delta)
 \end{array}$$

since if $\chi \in \Lambda^*(G^{\text{ab}})$, then $\chi(d) = \chi(\delta^{\text{ab}})$. Hence, we get an injective homomorphism

$$\alpha : \frac{\Lambda^*(G^{\text{rab}})}{\text{Im}(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}})} \hookrightarrow \frac{\Omega_d^*(T_G)}{\text{Im}(\omega_G^\delta)}. \tag{3.23}$$

Now if δ^{ab} is not 2-divisible, then

$$\frac{\Lambda^*(G^{\text{rab}})}{\text{Im}(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}})} \cong \mathbb{Z}/2\mathbb{Z} \cong \frac{\Omega_d^*(T_G)}{\text{Im}(\omega_G^\delta)}.$$

If, instead, δ^{ab} is 2-divisible, then condition (*) implies that

$$\frac{\Lambda^*(G^{\text{rab}})}{\text{Im}(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}})} = 0 = \frac{\Omega_d^*(T_G)}{\text{Im}(\omega_G^\delta)}.$$

In any case, the morphism α of Equation (3.23) is an isomorphism, which implies, by the snake lemma applied to Equation (3.22), that

$$\frac{\text{Im}(\omega_G^\delta)}{\text{Im}(\omega_{G^{\text{ab}}}^{\delta^{\text{ab}}})} \cong \frac{\Omega_d^*(T_G)}{\Lambda^*(G^{\text{ab}})}.$$

Hence, we get the conclusion using again Equations (3.20) and (3.21). □

A second case where Equation (3.15) is a push-out diagram is described in the following:

Theorem 3.16. *Let G be a reductive group and fix $\delta \in \pi_1(G)$. If the family C/S has genus $g(C/S) > 0$ and satisfies the following properties:*

- (1) $\text{End}(J_{C/\bar{\eta}}) = \mathbb{Z}$ and $\text{RPic}(C/S) = \text{Pic}_{C/S}(S)$;
- (2) $\text{RPic}^0(C/S)$ is torsion-free;

then the commutative diagram (3.15) is a push-out diagram.

This theorem was proved for the universal family $C_{g,n} \rightarrow \mathcal{M}_{g,n}$ with $g \geq 1$ (which satisfy the assumptions, by Remarks 2.3 and 2.5(1) and the Franchetta’s conjecture) in [21, Thm. C(2)].

Proof. The proof follows the same lines of the proof of [21, Thm. C(2)], as we now indicate.

First of all, because of the assumptions, the Picard group of $\text{Bun}_{T_G}^d(C/S)$ (where $d \in \pi_1(T_G)$ is a fixed lift of $\delta \in \pi_1(G)$) is generated by tautological classes by Theorem 3.9. Hence, using Theorem 3.8, we can define an action of the Weyl group \mathscr{W}_G on $\text{Pic}^d(\text{Bun}_{T_G}^d(C/S))$ by setting

$$w \cdot d_{\tilde{\pi}}(\mathcal{L}_\chi(M)) := d_{\tilde{\pi}}(\mathcal{L}_{w \cdot \chi}(M)) \quad \text{for any } \chi \in \Lambda^*(T) \text{ and any } M \in \text{Pic}(C/S),$$

in such a way that the diagram (3.10) for $\text{RPic}^d(\text{Bun}_{T_G}^d(C/S))$ is \mathscr{W}_G -equivariant.

Now, using the bottom exact sequence of Equation (3.10) and arguing as in the proof of [21, Lemma 5.3.3] for the case $m=2$, we get that the subgroup of invariants of the above action of \mathscr{W}_G sit in the following push-out diagram:

$$\begin{CD} \text{Sym}^2 \Lambda^*(G^{\text{ab}}) @<\text{Sym}^2 \Lambda_{\text{ab}}^*<< (\text{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}_G} \\ @VV\tau_{G^{\text{ab}}}^{\delta^{\text{ab}}}\downarrow V @VV\tau_{T_G}^d\downarrow V \\ \text{Pic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S) @<(\text{ab}\circ\iota)_{\#}^*<< (\text{Pic Bun}_{T_G}^d(C/S))^{\mathscr{W}_G} \end{CD} \tag{3.24}$$

Therefore, it remains to show that the natural inclusion

$$\psi : \left(\text{Pic Bun}_{T_G}^d(C/S) \right)^{\mathscr{W}_G} \hookrightarrow \text{Pic Bun}_G^\delta(C/S) \tag{3.25}$$

induced by the push-out diagram (3.24) and the commutative diagram (3.14) is an isomorphism. This will be achieved in two steps:

Step 1: $\text{coker } \psi$ is torsion.

Indeed, consider the two inclusions

$$\text{ab}_{\#}^* : \text{Pic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S) \xrightarrow{(\text{ab}\circ\iota)_{\#}^*} \left(\text{Pic Bun}_{T_G}^d(C/S) \right)^{\mathscr{W}_G} \xrightarrow{\psi} \text{Pic Bun}_G^\delta(C/S).$$

We get an induced exact sequence

$$0 \rightarrow \frac{\left(\text{Pic Bun}_{T_G}^d(C/S) \right)^{\mathscr{W}_G}}{\text{Pic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S)} \rightarrow \frac{\text{Pic Bun}_G^\delta(C/S)}{\text{Pic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S)} \rightarrow \text{coker } \psi \rightarrow 0.$$

By the push-out diagram (3.24) and using Lemmas 3.12 and 3.13, we get that

$$\frac{\left(\text{Pic Bun}_{T_G}^d(C/S) \right)^{\mathscr{W}_G}}{\text{Pic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S)} \cong \mathbb{Z}^s,$$

where s is the number of quasi-simple factors of G^{sc} . Moreover, arguing as in the proof of [21, Lemma 5.3.4], we deduce that

$$\text{rk} \frac{\text{Pic Bun}_G^\delta(C/S)}{\text{Pic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S)} \leq s.$$

By putting everything together, we deduce that $\text{coker } \psi$ is a finite group.

Step 2: $\text{coker } \psi$ is torsion-free.

Indeed, the natural inclusion of the subgroup of \mathscr{W}_G -invariants of $\text{Pic Bun}_{T_G}^d(C/S)$ factors is

$$\left(\text{Pic Bun}_{T_G}^d(C/S)\right)^{\mathscr{W}_G} \xrightarrow{\psi} \text{Pic Bun}_G^\delta(C/S) \xrightarrow{\iota_{\sharp}^*} \text{Pic Bun}_{T_G}^d(C/S),$$

where ι_{\sharp}^* is injective by Fact 3.2(2). Therefore, $\text{coker } \psi$ injects into the quotient

$$Q := \frac{\text{Pic Bun}_{T_G}^d(C/S)}{\left(\text{Pic Bun}_{T_G}^d(C/S)\right)^{\mathscr{W}_G}} = \frac{\text{RPic Bun}_{T_G}^d(C/S)}{\left(\text{RPic Bun}_{T_G}^d(C/S)\right)^{\mathscr{W}_G}}$$

Because of our three assumptions on the family C/S , Theorems 3.9 and 3.8 imply that $\text{RPic Bun}_{T_G}^d(C/S)$ is torsion-free. Hence, the quotient Q is torsion-free by [21, Lemma 5.3.6], which implies that $\text{coker } \psi$ is torsion-free. \square

We now explain how to obtain two alternative presentations of the relative Picard group of $\text{Bun}_G^\delta(C/S)$, for a family of curves C/S of positive genus and such that $\text{End}(J_{C/\bar{\eta}}) = \mathbb{Z}$ and $\text{RPic}(C/S) = \text{Pic}_{C/S}(S)$, assuming that Equation (3.14) is a push-out diagram.

We first need the following:

Definition 3.17. (see [22, Cor. 2.6, Def. 3.9]) *Let G be a reductive group with maximal torus T_G and Weyl group \mathscr{W}_G , and fix $\delta \in \pi_1(G)$.*

- (1) $\text{Bil}^{s, \mathscr{D}^{-\text{ev}}}(\Lambda(T_G))^{\mathscr{W}_G} := \left\{ b \in \text{Bil}^s(\Lambda(T_G))^{\mathscr{W}_G} : b|_{\Lambda(T_{\mathscr{D}(G)}) \times \Lambda(T_{\mathscr{D}(G)})} \text{ is even} \right\}$.
- (2) *The Néron–Severi group of Bun_G^δ is the subgroup*

$$\text{NS}(\text{Bun}_G^\delta) \subseteq \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})} \oplus \text{Bil}^{s, \mathscr{D}^{-\text{ev}}}(\Lambda(T_G))^{\mathscr{W}_G}$$

consisting of all the elements $([\chi], b)$ such that

$$\left[\chi|_{\Lambda(T_{\mathscr{D}(G)})} \right] = b(\delta \otimes -)|_{\Lambda(T_{\mathscr{D}(G)})} := \left[b(d \otimes -)|_{\Lambda(T_{\mathscr{D}(G)})} \right] \in \frac{\Lambda^*(T_{\mathscr{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})}, \tag{3.26}$$

where $d \in \Lambda(T_G)$ is any lift of δ .

For a more detailed study of the Néron–Severi group $\text{NS}(\text{Bun}_G^\delta)$, see [22, Prop. 3.11].

Theorem 3.18. *Assume that the family C/S has genus $g(C/S) > 0$ and satisfies the following conditions:*

- (1) $\text{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$ and $\text{RPic}(C/S) = \text{Pic}_{C/S}(S)$;
- (2) either $\text{RPic}^0(C/S)$ is torsion-free or $\mathcal{D}(G)$ is simply connected.

Then, there exists a commutative diagram, functorial in S and G ,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \Lambda^*(G^{\text{ab}}) \otimes \text{RPic}^0(C/S) & \xrightarrow{j_G^\delta} & \text{RPic Bun}_G^\delta(C/S) & \xrightarrow{\omega_G^\delta \oplus \gamma_G^\delta} & \text{NS}(\text{Bun}_G^\delta) \\
 & & \downarrow & & \parallel & & \downarrow \text{res}_G^{\text{NS}} \\
 0 & \longrightarrow & \Lambda^*(G^{\text{ab}}) \otimes \text{RPic}(C/S) & \xrightarrow{i_G^\delta} & \text{RPic Bun}_G^\delta(C/S) & \xrightarrow{\gamma_G^\delta} & \text{Bil}^{s, \mathcal{D}\text{-ev}}(\Lambda(T_G))^{\#G} \longrightarrow 0
 \end{array} \tag{3.27}$$

where the rows are exact, the left vertical morphism is injective and the right vertical morphism is surjective.

Moreover, the image of $\omega_G^\delta \oplus \gamma_G^\delta$ is equal to

$$\text{Im}(\omega_G^\delta \oplus \gamma_G^\delta) = \left\{ ([\chi], b) \in \text{NS}(\text{Bun}_G^\delta) : \begin{array}{l} [\chi(x) - b(\delta \otimes x)] + (g - 1)b(x \otimes x) \\ \text{is divisible by } \delta(C/S), \text{ for any } x \in \Lambda(T_G) \end{array} \right\}.$$

This theorem was proved for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ for $g \geq 1$ (and an arbitrary reductive group) in [22, Thm. 1.1], and it can be deduced for a fixed curve C over $k = \bar{k}$ (in the case $g(C) > 0$, $\text{End}(J_{C_{\bar{\eta}}}) = \mathbb{Z}$ and $\mathcal{D}(G)$ simply connected) from [7, Thm. 5.3.1].

Proof. Observe that our assumptions imply that Equation (3.14) is a push-out diagram (by Theorems 3.14 and 3.16), and that $\text{Pic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S) = \text{Pic}^{\text{taut}} \text{Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S)$ by Theorem 3.9.

Let us first describe the maps appearing in the above diagram (3.27):

- the map i_G^δ is defined by

$$i_G^\delta(\chi \otimes M) := \langle \mathcal{L}_\chi, M \rangle_{\hat{\pi}}$$

and j_G^δ is the restriction of i_G^δ .

- the morphism ω_G^δ is induced by the weight homomorphism

$$\text{wt}_G^\delta : \text{Pic Bun}_G^\delta(C/S) \rightarrow \Lambda^*(\mathcal{Z}(G)) = \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}})}$$

coming from the exact sequence Equation (4.1).

- the morphism γ_G^δ is uniquely determined, using that the diagram (3.14) is a push-out, by

(1) The composition $\gamma_G^\delta \circ \tau_G^\delta$ is the inclusion

$$\alpha : (\text{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}G} \cong \text{Bil}^{s,\text{ev}}(\Lambda(T_G))^{\mathscr{W}G} \hookrightarrow \text{Bil}^{s,\mathscr{D}-\text{ev}}(\Lambda(T_G))^{\mathscr{W}G}.$$

(2) The composition $\gamma_G^\delta \circ \text{ab}_\#^*$ is equal to the following composition:

$$\begin{aligned} \text{RPic}^{\text{taut}} \text{Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S) &\xrightarrow{\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}} (\Lambda^*(G^{\text{ab}}) \otimes \Lambda^*(G^{\text{ab}}))^s \cong \text{Bil}^s(\Lambda(G^{\text{ab}})) \\ &\xrightarrow{B_{\text{ab}}^*} \text{Bil}^{s,\mathscr{D}-\text{ev}}(\Lambda(T_G))^{\mathscr{W}G}, \end{aligned}$$

where $\gamma_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ is the homomorphism of Lemma 3.7 and B_{ab}^* is obtained by pull-back.

Now the proof is deduced from Theorems 3.8, 3.9 and 3.14, arguing as in the proofs of [22, Thm. 3.6, Thm. 3.12]. □

We end this subsection by describing the relative Picard group of $\text{Bun}_G^\delta(C/S)$, for a family of curves C/S of genus zero and an arbitrary reductive group G . We first need the following:

Definition 3.19. (see [7, Def. 5.2.1]) Let G be a reductive group and let $\delta \in \pi_1(G)$. Define

$$\text{NSBun}_G^\delta(\mathbb{P}^1) := \{(\chi, b) : l + b(d^{\text{ss}}, -) \in \Lambda^*(T_G)\} \subseteq \Lambda^*(\mathscr{Z}(G)) \times (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\mathscr{W}G},$$

where d^{ss} is the image in $\Lambda(T_{G^{\text{ss}}})$ of any lift $d \in \Lambda(T_G)$ of δ and $l + b(d^{\text{ss}}, -)$ is interpreted as an element of

$$\Lambda^*(\mathscr{Z}(G))_{\mathbb{Q}} \oplus \Lambda^*(T_{G^{\text{sc}}})_{\mathbb{Q}} = \Lambda^*(T_G)_{\mathbb{Q}}.$$

It is easily checked that the above definition does not depend on the choice of the lifting d of δ (see [7, Lemma 5.2.2]) and that it is functorial in G (see [7, Def. 5.2.7]).

Theorem 3.20. Let $C \rightarrow S$ be a family of curves of genus $g=0$ on a regular and integral quotient stack. Let G be a reductive group, fix $\delta \in \pi_1(G)$ and set $\delta^{\text{ab}} := \pi_1(\text{ab})(\delta) \in \pi_1(G^{\text{ab}})$. Fix a lift $d \in \Lambda(T_G)$ of δ and denote by d^{ss} its image in $\Lambda(T_{G^{\text{ss}}})$.

There exists an injective homomorphism, functorial in S and G :

$$c_G^\delta(C/S) = c_G^\delta : \text{RPic Bun}_G^\delta(C/S) \rightarrow \text{NSBun}_G^\delta(\mathbb{P}^1)$$

such that

(1) if $G = T$ is a torus, then c_G^δ coincides with the weight homomorphism;

(2) the image of c_G^δ is equal to

$$\text{Im}(c_G^\delta) = \begin{cases} \text{NS Bun}_G^\delta(\mathbb{P}^1) & \text{if } C/S \text{ is Zariski locally trivial,} \\ \left\{ (l, b) \in \text{NS Bun}_G^\delta(\mathbb{P}^1) : \right. & \text{if } C/S \text{ is not Zariski locally trivial.} \\ \left. \begin{array}{l} l(\delta^{\text{ab}}) + b(d^{\text{ss}}, d^{\text{ss}}) \text{ is even} \end{array} \right\} & \end{cases} \tag{3.28}$$

In particular, if C/S is not Zariski locally trivial, then $\text{Im}(c_G^\delta)$ is an index two subgroup of $\text{NS Bun}_G^\delta(\mathbb{P}^1)$.

Note that the second subgroup in Equation (3.28) is well-defined since if $\tilde{d} \in \Lambda(T_G)$ is another lift of δ , then its image \tilde{d}^{ss} in $\Lambda(T_{G^{\text{ss}}})$ is such that $\tilde{d}^{\text{ss}} = d^{\text{ss}} + \epsilon$ for some $\epsilon \in \Lambda(T_{G^{\text{sc}}})$, and we have

$$b(\tilde{d}^{\text{ss}}, \tilde{d}^{\text{ss}}) = b(d^{\text{ss}}, d^{\text{ss}}) + 2b(d^{\text{ss}}, \epsilon) + b(\epsilon, \epsilon) \in b(d^{\text{ss}}, d^{\text{ss}}) + 2\mathbb{Z},$$

using that $b(\epsilon, \epsilon) \in 2\mathbb{Z}$ since b is an even symmetric bilinear form on $\Lambda(T_{G^{\text{sc}}})$, and that $b(d^{\text{ss}}, \epsilon) \in \mathbb{Z}$ by [7, Lemma 4.3.4].

Proof. Note that C/S is Zariski locally trivial if and only if C/S admits a section. Therefore, for a choice of a geometric generic point $\bar{\eta}$ of S , we get the following diagram:

$$\begin{array}{ccccc}
 C_{\bar{\eta}} & \longrightarrow & C & \longrightarrow & \mathcal{C}_{0,n} \\
 \downarrow & & \square & & \downarrow \\
 \bar{\eta} & \longrightarrow & S & \longrightarrow & \mathcal{M}_{0,n}
 \end{array} \tag{3.29}$$

with Cartesian squares, where

$$n = \begin{cases} 1 & \text{if } C/S \text{ is Zariski locally trivial,} \\ 0 & \text{if } C/S \text{ is not Zariski locally trivial.} \end{cases}$$

The above diagram induces the following homomorphisms, functorial in S and in G :

$$\text{RPic Bun}_{G,0,n}^\delta \hookrightarrow \text{RPic Bun}_G^\delta(C/S) \hookrightarrow \text{Pic Bun}_G^\delta(\mathbb{P}^1_{\bar{\eta}}) \xrightarrow[\cong]{c_G^\delta(\mathbb{P}^1_{\bar{\eta}})} \text{NS Bun}_G^\delta(\mathbb{P}^1), \tag{3.30}$$

where the first two pullback homomorphisms are injective by Fact 3.2(1) (note that $\bar{\eta}$ is also a geometric generic point of $\mathcal{M}_{0,n}$) and the last isomorphism is the one of [7, Thm. 5.3.1(iii)]. Therefore, we get our desired injective homomorphism, functorial in S and G

$$c_G^\delta(C/S) : \text{RPic Bun}_G^\delta(C/S) \hookrightarrow \text{Pic Bun}_G^\delta(\mathbb{P}^1_\eta) \xrightarrow[\cong]{c_G^\delta(\mathbb{P}^1_\eta)} \text{NS Bun}_G^\delta(\mathbb{P}^1),$$

which coincides with the weight homomorphism if G is a torus, because this is the case for the isomorphism $c_G^\delta(\mathbb{P}^1_\eta)$ (see the proof of [7, Thm. 5.3.1(iii)]).

It remains to prove part (2) by distinguishing the two cases C/S Zariski locally trivial or not.

If C/S is Zariski locally trivial (and hence $n = 1$ in the above diagram), then we get that $c_G^\delta(C/S)$ is surjective since the composite homomorphism in Equation (3.30)

$$c_G^\delta(\mathcal{C}_{0,1}/\mathcal{M}_{0,1}) : \text{RPic Bun}_{G,0,1}^\delta \rightarrow \text{NS Bun}_G^\delta(\mathbb{P}^1)$$

is surjective by [21, Thm. 5.0.2].

If C/S is not Zariski locally trivial (and hence $n = 0$ in the above diagram), then we get that

$$\{(l, b) \in \text{NS Bun}_G^\delta(\mathbb{P}^1) : l(\delta^{\text{ab}}) + b(d^{\text{ss}}, d^{\text{ss}}) \text{ is even}\} \subseteq \text{Im}(c_G^\delta(C/S)),$$

since the same is true for the image of $c_G^\delta(\mathcal{C}_0/\mathcal{M}_0)$ by [21, Thm. 5.0.2]. In order to prove the other inclusion, consider the following commutative diagram (for a chosen lift $d \in \Lambda(T_G)$ of δ):

$$\begin{CD} \text{RPic Bun}_G^\delta(C/S) @>{\iota_\#^*}>> \text{RPic Bun}_{T_G}^d(C/S) \\ @VV{c_G^\delta(C/S)}V @VV{c_{T_G}^d(C/S)}V \\ \text{NS Bun}_G^\delta(\mathbb{P}^1) @>{\iota_\#^{\text{NS},*}}>> \text{NS Bun}_{T_G}^d(\mathbb{P}^1) = \Lambda^*(T_G) \end{CD}$$

where the bottom map is defined by (see [7, Def. 5.2.5])

$$\begin{aligned} \iota_\#^{\text{NS},*} : \text{NS Bun}_G^\delta(\mathbb{P}^1) &\rightarrow \text{NS Bun}_{T_G}^d(\mathbb{P}^1) = \Lambda^*(T_G) \\ (l, b) &\mapsto l + b(d^{\text{ss}}, -). \end{aligned}$$

The above commutative diagram, together with Theorem 3.6(2), implies that

$$\begin{aligned} \text{Im}(c_G^\delta(C/S)) &\subseteq (\iota_\#^{\text{NS},*})^{-1}(\text{Im}(c_{T_G}^d(C/S))) \\ &= \{(l, b) \in \text{NS Bun}_G^\delta(\mathbb{P}^1) : l(\delta^{\text{ab}}) + b(d^{\text{ss}}, d^{\text{ss}}) \text{ is even}\}, \end{aligned}$$

and we are done. □

Corollary 3.21. *Same assumptions of Theorem 3.20. There exists an exact sequence functorial in S and G*

$$0 \rightarrow \text{Pic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S) \xrightarrow{\text{ab}_\#^*} \text{Pic Bun}_G^\delta(C/S) \xrightarrow{p_G^\delta} \left\{ \begin{array}{l} b \in (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\mathscr{W}_G} \\ b^{\mathbb{Q}}(d^{\text{ss}}, -)|_{\Lambda(T_{\mathcal{D}(G)})} \text{ is integral} \end{array} \right\}, \tag{3.31}$$

where the last homomorphism p_G^δ has image of index at most two and it is surjective if either $\mathcal{D}(G)$ is simply connected or C/S is Zariski locally trivial.

Note that if $\mathcal{D}(G)$ is simply connected, then the last group of Equation (3.31) is equal to $(\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\mathcal{W}G}$ by Lemma 3.13(3).

Proof. Theorem 3.20 implies that the following diagram

$$\begin{CD} \text{RPic Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S) @<{\text{ab}_\#^\delta}<< \text{RPic Bun}_G^\delta(C/S) \\ @V{\text{wt}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S)}VV @VV{c_G^\delta(C/S)}V \\ \text{NS Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(\mathbb{P}^1) = \Lambda^*(G^{\text{ab}}) @<{\text{ab}_\#^{\text{NS},*}}<< \text{NS Bun}_G^\delta(\mathbb{P}^1) \end{CD}$$

is Cartesian, where the map $\text{ab}_\#^{\text{NS},*}$ sends χ into $(\chi, 0)$. Therefore, we get that the cokernel of $\text{ab}_\#^\delta$ injects into the cokernel of $\text{ab}_\#^{\text{NS},*}$, which is equal (by [7, Prop. 5.2.11]) to

$$\text{coker}(\text{ab}_\#^{\text{NS},*}) = \left\{ b \in (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\mathcal{W}G} : b^\mathbb{Q}(d^{\text{ss}}, -)|_{\Lambda(T_{\mathcal{D}(G)})} \text{ is integral} \right\}.$$

This proves the existence of the exact sequence Equation (3.31), as well as of the following exact sequence:

$$0 \rightarrow \text{coker wt}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S) \rightarrow \text{coker } c_G^\delta(C/S) \rightarrow \text{coker } p_G^\delta \rightarrow 0.$$

Since $\text{coker } c_G^\delta(C/S)$ is a finite group of order at most two by Theorem 3.20 and it is trivial if C/S is Zariski locally trivial, we deduce that the same is true for $\text{coker } p_G^\delta$. Moreover, $\text{coker } p_G^\delta$ is trivial if $\mathcal{D}(G)$ is simply connected by Theorem 3.14. \square

4. The rigidification of $\text{Bun}_G^\delta(C/S)$ by the centre

The aim of this section is to study, for a reductive group G , the $\mathcal{Z}(G)$ -gerbe (see Equation (2.10))

$$\nu_G^\delta = \nu_G^\delta(C/S) : \text{Bun}_G^\delta(C/S) \rightarrow \text{Bun}_G^\delta(C/S) // \mathcal{Z}(G) := \mathfrak{Bun}_G^\delta(C/S),$$

obtained by rigidifying the stack $\text{Bun}_G^\delta(C/S)$ by the centre $\mathcal{Z}(G)$ of the reductive group G .

From the Leray spectral sequence

$$E_{p,q}^2 = H^p(\mathfrak{Bun}_G^\delta(C/S), R^q(\nu_G^\delta)_*(\mathbb{G}_m)) \Rightarrow H^{p+q}(\text{Bun}_G^\delta(C/S), \mathbb{G}_m),$$

and using that $(\nu_G^\delta)_*(\mathbb{G}_m) = \mathbb{G}_m$ and that $R^1(\nu_G^\delta)_*(\mathbb{G}_m)$ is the constant sheaf $\Lambda^*(\mathcal{Z}(G)) = \text{Hom}(\mathcal{Z}(G), \mathbb{G}_m)$, we get the exact sequence

$$\begin{aligned} \text{Pic}(\mathfrak{Bun}_G^\delta(C/S)) \xrightarrow{(\nu_G^\delta)^*} \text{Pic}(\text{Bun}_G^\delta(C/S)) \xrightarrow{\text{wt}_G^\delta} \Lambda^*(\mathcal{Z}(G)) := \text{Hom}(\mathcal{Z}(G), \mathbb{G}_m) \\ \xrightarrow{\text{obs}_G^\delta} H^2(\mathfrak{Bun}_G^\delta(C/S), \mathbb{G}_m) \xrightarrow{(\nu_G^\delta)^*} H^2(\text{Bun}_G^\delta(C/S), \mathbb{G}_m), \end{aligned} \tag{4.1}$$

and an analogue one in which the Picard groups Pic are replaced by the relative Picard groups RPic.

Recall (see [22, § 5]) the following geometric descriptions of the *weight homomorphism* wt_G^δ and the *obstruction homomorphism* obs_G^δ :

- given a line bundle \mathcal{L} on $\text{Bun}_G^\delta(C/S)$, the character $\text{wt}_G^\delta(\mathcal{L}) \in \Lambda^*(\mathcal{Z}(G))$ is such that, for any $\mathcal{E} := (E \rightarrow C_V) \in \text{Bun}_G^\delta(C/S)(V)$, we have the factorization

$$\text{wt}_G^\delta(V) : \mathcal{Z}(G)(V) \hookrightarrow \text{Aut}_{\text{Bun}_G^\delta(C/S)(V)}(\mathcal{E}) \xrightarrow{\text{Aut}(\mathcal{L}_V)} \text{Aut}_{\mathcal{O}_V}(\mathcal{L}_V(\mathcal{E})) = \mathbb{G}_m(V),$$

where the first homomorphism is given by the canonical action of $\mathcal{Z}(G)$ on every G -gerbe and the second homomorphism is induced by the functor of groupoids

$$\mathcal{L}_V : \text{Bun}_G^\delta(C/S)(V) \rightarrow \{\text{Line bundles on } V\}$$

determined by \mathcal{L} .

- given any character $\lambda \in \text{Hom}(\mathcal{Z}(G), \mathbb{G}_m)$, the element $\text{obs}_G^\delta(\lambda)$ is the class in $H^2(\mathfrak{Bun}_G^\delta(C/S), \mathbb{G}_m)$ of the \mathbb{G}_m -gerbe $\lambda_*(\nu_G^\delta)$ obtained by push-forwarding the $\mathcal{Z}(G)$ -gerbe ν_G^δ along λ .

Recall also (see [22, § 2.1]) that the character group of the centre $\mathcal{Z}(G)$ is equal to

$$\Lambda^*(\mathcal{Z}(G)) = \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}}},$$

and it sits into the following exact sequence induced by the vertical exact sequence of Equation (2.5):

$$0 \rightarrow \Lambda^*(G^{\text{ab}}) \xrightarrow{\overline{\Lambda^*_{\text{ab}}}} \Lambda^*(\mathcal{Z}(G)) = \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}}} \xrightarrow{\overline{\Lambda^*_{\mathcal{Z}}}} \Lambda^*(\mathcal{Z}(\mathcal{Z}(G))) = \frac{\Lambda^*(T_{\mathcal{Z}(G)})}{\Lambda^*(T_{G^{\text{ad}}}} \rightarrow 0. \tag{4.2}$$

Our goal is to determine the (relative) Picard group of the rigidification $\mathfrak{Bun}_G^\delta(C/S)$ and the group

$$\text{coker}(\text{wt}_G^\delta) \cong \text{Im}(\text{obs}_G^\delta) = \ker(\nu_G^\delta)^*, \tag{4.3}$$

which is an obstruction to the triviality of the $\mathcal{Z}(G)$ -gerbe ν_G^δ .

With this aim, we need to recall two crucial homomorphisms from [22].

Definition 4.1. (see [22, Definition/Lemma 2.8, Def. 5.3]) Let G be a reductive group with fixed maximal torus T_G and Weyl group \mathscr{W}_G , and fix $\delta \in \pi_1(G)$.

- (1) There is a well-defined homomorphism (called evaluation homomorphism)

$$\begin{aligned} \text{ev}_{\mathscr{D}(G)}^\delta : \text{Bil}^{s,\text{ev}}(\Lambda(T_{\mathscr{D}(G)})|\Lambda(T_{G^{\text{ss}}}))^{\mathscr{W}_G} &\longrightarrow \frac{\Lambda^*(T_{\mathscr{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})}, \\ b \mapsto b(\delta^{\text{ss}} \otimes -) &:= [b^{\mathbb{Q}}(d^{\text{ss}} \otimes -)], \end{aligned} \tag{4.4}$$

where $d^{\text{ss}} \in \Lambda(T_{G^{\text{ss}}})$ is any lift of $\delta^{\text{ss}} := \pi_1(\text{ss})(\delta) \in \pi_1(G^{\text{ss}}) = \frac{\Lambda(T_{G^{\text{ss}}})}{\Lambda(T_{G^{\text{sc}}})}$ and $d^{\mathbb{Q}}$ is the rational extension of b to $\Lambda(T_{G^{\text{ss}}})$.

- (2) The Néron–Severi group of \mathfrak{Bun}_G^δ is the subgroup

$$\text{NS}(\mathfrak{Bun}_G^\delta) \subset \text{Bil}^{s,\mathscr{D}-\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G}$$

consisting of those $b \in \text{Bil}^{s,\mathscr{D}-\text{ev}}(\Lambda(T_G))^{\mathscr{W}_G}$ such that

$$0 = b(\delta \otimes -)|_{\Lambda(T_{\mathscr{D}(G)})} := \left[b(d \otimes -)|_{\Lambda(T_{\mathscr{D}(G)})} \right] \in \frac{\Lambda^*(T_{\mathscr{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}})}, \tag{4.5}$$

where $d \in \Lambda(T_G)$ is any lift of δ .

Note that there is an injective homomorphism

$$\begin{aligned} \nu^{\delta,\text{NS}} : \text{NS}(\mathfrak{Bun}_G^\delta) &\hookrightarrow \text{NS}(\text{Bun}_G^\delta) \\ b &\mapsto ([0], b). \end{aligned}$$

For a more detailed description of $\text{NS}(\mathfrak{Bun}_G^\delta)$, see [22, Prop. 5.4].

Example 1. If $\mathscr{D}(G)$ is simply connected, i.e. $\mathscr{D}(G) = G^{\text{sc}}$, then the cokernel of $\text{ev}_{G^{\text{sc}}}^\delta$ can be computed explicitly as it follows:

- (1) If $G^{\text{sc}} = G_1^{\text{sc}} \times \dots \times G_r^{\text{sc}}$ is the decomposition of G^{sc} into simply connected and almost simple algebraic groups and $\delta = (\delta_1, \dots, \delta_r) \in \pi_1(G^{\text{ad}}) = \pi_1(G_1^{\text{ad}}) \times \dots \times \pi_1(G_r^{\text{ad}})$, then

$$\text{coker ev}_{G^{\text{sc}}}^\delta \cong \text{coker ev}_{G_1^{\text{sc}}}^{\delta_1} \times \dots \times \text{coker ev}_{G_r^{\text{sc}}}^{\delta_r}.$$

- (2) If G^{sc} is simply connected and almost simple, then the cokernel of $\text{ev}_{G^{\text{sc}}}^\delta$ is equal to (see [22, § 7]):

- (a) If $G^{\text{sc}} = \text{SL}_n$ (and hence $G^{\text{ad}} = \text{PSL}_n$), then, for any $\delta \in \pi_1(\text{PSL}_n) = \mathbb{Z}/n\mathbb{Z}$, we have that

$$\text{coker ev}_{\text{SL}_n}^\delta = \frac{\mathbb{Z}}{\text{gcd}(n, \delta)\mathbb{Z}}.$$

- (b) If $G^{\text{sc}} = \text{Spin}_{2n+1}$ (and hence $G^{\text{ad}} = \text{SO}_{2n+1}$), then, for any $\delta \in \pi_1(\text{SO}_{2n+1}) = \mathbb{Z}/2\mathbb{Z}$, we have that

$$\text{coker ev}_{\text{Spin}_{2n+1}}^\delta = \mathbb{Z}/2\mathbb{Z}.$$

- (c) If $G^{\text{sc}} = \text{Sp}_{2n}$ (and hence $G^{\text{ad}} = \text{PSp}_{2n}$), then, for any $\delta \in \pi_1(\text{PSp}_{2n}) = \mathbb{Z}/2\mathbb{Z}$, we have that

$$\text{coker ev}_{\text{Sp}_{2n}}^\delta = \begin{cases} \mathbb{Z}/2\mathbb{Z} & \text{if either } \delta = 0 \text{ or } n \text{ is even,} \\ 0 & \text{otherwise.} \end{cases}$$

- (d) If $G^{\text{sc}} = \text{Spin}_{2n}$ (and hence $G^{\text{ad}} = \text{PSO}_{2n}$), then, for any

$$\delta \in \pi_1(\text{PSO}_{2n}) = \begin{cases} \mathbb{Z}/4\mathbb{Z} & \text{if } n \text{ is odd,} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} & \text{if } n \text{ is even,} \end{cases}$$

we have that

$$\text{coker ev}_{\text{Spin}_{2n}}^\delta = \begin{cases} \mathbb{Z}/4\mathbb{Z} & \text{if } \delta = 0 \text{ and } n \text{ is odd,} \\ \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} & \text{if } \delta = 0 \text{ and } n \text{ is even,} \\ \mathbb{Z}/2\mathbb{Z} & \text{if } \text{ord}(\delta) = 2, \\ 0 & \text{if } \text{ord}(\delta) = 4. \end{cases}$$

- (e) If $G^{\text{sc}} = \mathbb{E}_6^{\text{sc}}$, then, for any $\delta \in \pi_1(\mathbb{E}_6^{\text{ad}}) = \mathbb{Z}/3\mathbb{Z}$, we have that

$$\text{coker ev}_{\mathbb{E}_6^{\text{sc}}}^\delta = \begin{cases} \mathbb{Z}/3\mathbb{Z} & \text{if } \delta = 0, \\ 0 & \text{if } \delta \neq 0. \end{cases}$$

- (f) If $G^{\text{sc}} = \mathbb{E}_7^{\text{sc}}$, then, for any $\delta \in \pi_1(\mathbb{E}_7^{\text{ad}}) = \mathbb{Z}/2\mathbb{Z}$, we have that

$$\text{coker ev}_{\mathbb{E}_7^{\text{sc}}}^\delta = \begin{cases} \mathbb{Z}/2\mathbb{Z} & \text{if } \delta = 0, \\ 0 & \text{if } \delta \neq 0. \end{cases}$$

- (g) If G^{sc} is of type \mathbb{E}_8 or \mathbb{F}_4 or \mathbb{G}_2 , then

$$\text{coker ev}_{G^{\text{sc}}}^\delta = 0.$$

We first describe the (relative) Picard group of the rigidification $\mathfrak{Bun}_G^\delta(C/S)$ for families of positive genus, under certain assumptions.

Theorem 4.2. *Let $C \rightarrow S$ be a family of curves of genus $g > 0$ on a regular and integral quotient stack with geometric generic fibre $C_{\overline{\eta}}$. Assume that the following two conditions hold true:*

- (1) $\text{End}(J_{C_{\overline{\eta}}}) = \mathbb{Z}$ and $\text{RPic}(C/S) = \text{Pic}_{C/S}(S)$;
- (2) either $\text{RPic}^0(C/S)$ is torsion-free or $\mathcal{D}(G)$ is simply connected.

Then, for any fixed $\delta \in \pi_1(G)$, we have the following exact sequence, functorial in S and G :

$$0 \rightarrow \Lambda^*(G^{\text{ab}}) \otimes \text{RPic}^0(C/S) \xrightarrow{j_G^\delta} \text{RPic}(\mathfrak{Bun}_G^\delta(C/S)) \xrightarrow{\gamma_G^\delta} \text{NS}(\mathfrak{Bun}_G^\delta), \tag{4.6}$$

where $\overline{j_G^\delta}$ and $\overline{\gamma_G^\delta}$ are uniquely determined by

$$(\nu_G^\delta)^* \circ \overline{j_G^\delta} = j_G^\delta \quad \text{and} \quad \nu_G^{\delta, \text{NS}} \circ \overline{\gamma_G^\delta} = (\omega_G^\delta \oplus \gamma_G^\delta) \circ (\nu_G^\delta)^*,$$

see Theorem 3.18.

Moreover, the image of $\overline{\gamma_G^\delta}$ is equal to

$$\text{Im}(\overline{\gamma_G^\delta}) = \left\{ b \in \text{NS}(\mathfrak{Bun}_G^\delta) : \begin{array}{l} b(\delta \otimes x) + (g-1)b(x \otimes x) \text{ is a multiple of } \delta(C/S) \\ \text{for any } x \in \Lambda(T_G) \end{array} \right\}. \tag{4.7}$$

This theorem was proved for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ for $g \geq 1$ in [22, Thm. 5.5].

Proof. This is deduced from Theorem 3.18, arguing as in the proofs of [22, Thm. 5.5]. □

We now describe the cokernel of the weight homomorphism.

Theorem 4.3. *Let $C \rightarrow S$ be a family of curves of genus $g > 0$ on a regular and integral quotient stack with geometric generic fibre $C_{\overline{\eta}}$. Assume that the following two conditions hold true:*

- (1) $\text{End}(J_{C_{\overline{\eta}}}) = \mathbb{Z}$ and $\text{RPic}(C/S) = \text{Pic}_{C/S}(S)$;
- (2) either $\text{RPic}^0(C/S)$ is torsion-free or $\mathcal{D}(G)$ is simply connected.

Then, for any fixed $\delta \in \pi_1(G)$, we have the following exact sequence, There exists an exact sequence

$$0 \rightarrow \text{coker}(\overline{\gamma_G^\delta}) \xrightarrow{\partial_G^\delta} \text{Hom} \left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{\delta(C/S)\mathbb{Z}} \right) \xrightarrow{\widetilde{\Lambda_{\text{ab}}^*}} \text{coker}(\text{wt}_G^\delta) \xrightarrow{\widetilde{\Lambda_{\mathcal{D}}^*}} \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta) \rightarrow 0, \tag{4.8}$$

where $\widetilde{\Lambda}_{\text{ab}}^*$ and $\widetilde{\Lambda}_{\mathcal{D}}^*$ are the homomorphisms induced by, respectively, $\overline{\Lambda}_{\text{ab}}^*$ and $\overline{\Lambda}_{\mathcal{D}}^*$ of Equation (4.2) and ∂_G^δ is defined as follows:

$$\begin{aligned} \partial_G^\delta : \text{coker}(\overline{\gamma_G^\delta}) &\longrightarrow \text{Hom}\left(\Lambda(G^{\text{ab}}), \frac{\mathbb{Z}}{\delta(C/S)\mathbb{Z}}\right) \\ [b] &\mapsto \{x \mapsto [b(\delta \otimes \tilde{x}) + (1-g)b(\tilde{x} \otimes \tilde{x})]\}, \end{aligned} \tag{4.9}$$

where $b \in \text{NS}(\mathfrak{Bun}_{G,g,n}^\delta) \subset \text{Bil}^{s,\mathcal{D}\text{-ev}}(\Lambda(T_G))^{\mathcal{W}G}$ and $\tilde{x} \in \Lambda(T_G)$ is any lift of $x \in \Lambda(G^{\text{ab}})$.

This theorem was proved for the universal family $\mathcal{C}_{g,n} \rightarrow \mathcal{M}_{g,n}$ for $g \geq 1$ in [22, Thm. 5.7] and for a curve C over $k = \bar{k}$ of genus $g(C) \geq 3$ in [8, Thm. 6.8] (and in both results the general case of an arbitrary reductive group is also treated).

Proof. This is deduced from Theorem 3.18, arguing as in the proofs of [22, Thm. 5.7]. □

Corollary 4.4. *Same assumptions as in Theorem 4.3.*

(1) *If $\delta(C/S) = 1$, then there is an isomorphism*

$$\widetilde{\Lambda}_{\mathcal{D}}^* : \text{coker}(\text{wt}_G^\delta) \xrightarrow{\cong} \text{coker}(\text{ev}_{\mathcal{D}(G)}^\delta).$$

(2) *If $G = T$ is a torus and $d \in \Lambda(T)$, then there is an exact sequence*

$$0 \rightarrow \text{coker}(\overline{\gamma_T^d}) \xrightarrow{\partial_T^d} \text{Hom}\left(\Lambda(T), \frac{\mathbb{Z}}{\delta(C/S)\mathbb{Z}}\right) \rightarrow \text{coker}(\text{wt}_T^d) \rightarrow 0, \tag{4.10}$$

Moreover, we have that

$$\begin{aligned} \text{coker}(\overline{\gamma_T^d}) &\cong \frac{\mathbb{Z}}{\frac{\delta(C/S)}{\text{gcd}(\delta(C/S), \text{div}(d)+1-g)}\mathbb{Z}} \oplus \left[\frac{\mathbb{Z}}{\frac{\delta(C/S)}{\text{gcd}(\delta(C/S), g-1, \text{div}(d))}\mathbb{Z}} \right]^{\oplus(\dim(T)-1)} \\ \text{coker}(\text{wt}_T^d) &\cong \frac{\mathbb{Z}}{\text{gcd}(\delta(C/S), \text{div}(d) + 1 - g)\mathbb{Z}} \oplus \left[\frac{\mathbb{Z}}{\text{gcd}(\delta(C/S), g - 1, \text{div}(d))\mathbb{Z}} \right]^{\oplus(\dim(T)-1)}, \end{aligned}$$

where $\text{div}(d)$ is the divisibility of d in the lattice $\Lambda(T)$, with the convention that $\text{coker}(\overline{\gamma_T^d}) = \{0\}$ if $\delta(C/S) = 0$ (when the above expression for $\text{coker}(\overline{\gamma_T^d})$ is not well-defined).

The above Corollary for $G = \mathbb{G}_m$ recovers the main result of [35]: a Poincaré line bundle exists on $\mathfrak{Bun}_{\mathbb{G}_m}^d(C/S) \times_S C$ (which is equivalent to the triviality of the \mathbb{G}_m -gerbe $\nu_{\mathbb{G}_m}^d$) if and only if $\text{gcd}(\delta(C/S), d + 1 - g) = 1$, where $d \in \Lambda(\mathbb{G}_m) = \mathbb{Z}$.

Proof. Part (1) and the first part of (2) follow from Theorem 4.2.

The second part of (2) follows by choosing a basis $\{e_1, \dots, e_r\}$ of $\Lambda(T)$ (with $r = \dim T$) such that $d = \text{div}(d)e_1$ and computing the composition

$$\overline{\partial}_T^d : \text{Bil}^s(\Lambda(T)) \rightarrow \text{coker}(\overline{\gamma}_T^d) \xrightarrow{\partial_T^d} \text{Hom}\left(\Lambda(T), \frac{\mathbb{Z}}{\delta(C/S)\mathbb{Z}}\right)$$

on the basis $\{e_i^* \otimes e_i^*\}_i, \{e_i^* \otimes e_j^* + e_j^* \otimes e_i^*\}_{i>j}$ of $\text{Bil}^s(\Lambda(T))$:

$$\begin{cases} \overline{\partial}_T^d(e_1^* \otimes e_1^*) = [\text{div}(d) + 1 - g]e_1^*, \\ \overline{\partial}_T^d(e_i^* \otimes e_i^*) = [1 - g]e_i^* \text{ for any } i \geq 2, \\ \overline{\partial}_T^d(e_1^* \otimes e_i^* + e_i^* \otimes e_1^*) = \text{div}(d)e_i^* \text{ for any } i \geq 2, \\ \overline{\partial}_T^d(e_i^* \otimes e_j^* + e_j^* \otimes e_i^*) = 0 \text{ for any } 2 \leq i < j. \end{cases}$$

□

We end this subsection by computing the relative Picard group of $\mathfrak{Bun}_G^\delta(C/S)$ and the group $\text{coker}(\text{wt}_G^\delta)$ for families of genus zero, under certain assumptions.

Theorem 4.5. *Let $\pi : C \rightarrow S$ is a family of smooth curves of genus $g(C/S) = 0$ over an integral and regular quotient stack S . Assume that at least one of the following two assumptions is satisfied:*

- (1) *the family C/S is Zariski locally trivial;*
- (2) *the derived subgroup $\mathcal{D}(G)$ is simply connected.*

For a given $\delta \in \pi_1(G)$, consider the homomorphism

$$\widehat{\text{ev}}_{\mathcal{D}(G)}^\delta : \left\{ \begin{array}{l} b \in (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\mathcal{W}G} : \\ b^\mathbb{Q}(d^{\text{ss}}, -)|_{\Lambda(T_{\mathcal{D}(G)})} \text{ is integral} \end{array} \right\} \rightarrow \frac{\Lambda^*(T_{\mathcal{D}(G)})}{\Lambda^*(T_{G^{\text{ad}}}} \tag{4.11}$$

$$b \mapsto b(\delta^{\text{ss}} \otimes -) := [b^\mathbb{Q}(d^{\text{ss}} \otimes -)],$$

where $d^{\text{ss}} \in \Lambda(T_{G^{\text{ss}}})$ is any lift of $\delta^{\text{ss}} := \pi_1(\text{ss})(\delta) \in \pi_1(G^{\text{ss}}) = \frac{\Lambda(T_{G^{\text{ss}}})}{\Lambda(T_{G^{\text{sc}}})}$ and $d^\mathbb{Q}$ is the rational extension of b to $\Lambda(T_{G^{\text{ss}}})$.

- (1) *We have an isomorphism*

$$\text{RPic}(\mathfrak{Bun}_G^\delta(C/S)) \cong \ker(\widehat{\text{ev}}_{\mathcal{D}(G)}^\delta),$$

which is functorial in S and G .

(2) The homomorphism $\overline{\Lambda}_{\mathcal{D}}^*$ of Equation (4.2) induces a surjection

$$\widetilde{\Lambda}_{\mathcal{D}}^* : \text{coker}(\text{wt}_G^\delta) \rightarrow \text{coker}(\widehat{\text{ev}}_{\mathcal{D}(G)}^\delta), \tag{4.12}$$

whose kernel is equal to

$$\text{ker}(\widetilde{\Lambda}_{\mathcal{D}}^*) = \begin{cases} 0 & \text{if either } \delta(C/S) = 1 \text{ or } \delta^{\text{ab}} \text{ is 2-divisible in } \Lambda^*(G^{\text{ab}}), \\ \mathbb{Z}/2\mathbb{Z} & \text{if } \delta(C/S) = 2 \text{ and } \delta^{\text{ab}} \text{ is not 2-divisible in } \Lambda^*(G^{\text{ab}}). \end{cases} \tag{4.13}$$

This theorem can be deduced for the universal family $\mathcal{C}_{0,n} \rightarrow \mathcal{M}_{0,n}$ with $n \geq 1$ (which is Zariski locally trivial) from [22, Rmk. 5.10]. Note that if $\mathcal{D}(G)$ is simply connected, then the homomorphism $\widehat{\text{ev}}_{\mathcal{D}(G)}^\delta$ coincides with the homomorphism $\text{ev}_{G^{\text{sc}}}^\delta$ by Lemma 3.13(3).

Proof. Set

$$Q := \left\{ b \in (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\mathscr{W}G} : b^{\mathbb{Q}}(d^{\text{ss}}, -)|_{\Lambda(T_{\mathcal{D}(G)})} \text{ is integral} \right\}.$$

Consider the following diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{RPic}(\text{Bun}_{G^{\text{ab}}}^{\delta^{\text{ab}}}(C/S)) & \xrightarrow{\text{ab}_G^{\delta^{\text{ab}}}} & \text{RPic}(\text{Bun}_G^\delta(C/S)) & \longrightarrow & Q \longrightarrow 0 \\ & & \downarrow \text{wt}_{G^{\text{ab}}}^{\delta^{\text{ab}}} & & \downarrow \text{wt}_G^\delta & & \downarrow \widehat{\text{ev}}_{\mathcal{D}(G)}^\delta \\ 0 & \longrightarrow & \Lambda^*(G^{\text{ab}}) & \xrightarrow{\overline{\Lambda}_{\text{ab}}^*} & \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{nd}}})} & \xrightarrow{\overline{\Lambda}_{\mathcal{D}}^*} & \frac{\Lambda^*(T_{G^{\text{sc}}})}{\Lambda^*(T_{G^{\text{nd}}})} \longrightarrow 0. \end{array} \tag{4.14}$$

where the first row is exact by Corollary 3.21, the second row is exact by Equation (4.2) and the diagram is commutative by the same proof of [22, Prop. 5.2, Prop. 5.4].

By applying the snake lemma and using that $\text{wt}_{G^{\text{ab}}}^{\delta^{\text{ab}}}$ is injective by Theorem 3.6(2), we get an exact sequence

$$0 \rightarrow \text{RPic}(\mathfrak{Bun}_G^\delta(C/S)) \rightarrow \text{ker}(\widehat{\text{ev}}_{\mathcal{D}(G)}^\delta) \xrightarrow{\partial} \text{coker}(\text{wt}_{G^{\text{ab}}}^{\delta^{\text{ab}}}) \rightarrow \text{coker}(\text{wt}_G^\delta) \xrightarrow{\widetilde{\Lambda}_{\mathcal{D}}^*} \text{coker}(\widehat{\text{ev}}_{\mathcal{D}(G)}^\delta) \rightarrow 0.$$

Observe now that Theorem 3.6(3) implies that $\text{coker}(\text{wt}_{G^{\text{ab}}}^{\delta^{\text{ab}}})$ is equal to the group in the right-hand side of Equation (4.13). Hence, we conclude by the previous exact sequence and the following:

Claim: the coboundary map ∂ is zero.

Indeed, assume that $\text{coker}(\text{wt}_{G^{\text{ab}}}^{\delta^{\text{ab}}}) \cong \mathbb{Z}/2\mathbb{Z}$; otherwise, the statement is trivially true. In particular, C/S is not Zariski locally trivial, and hence $\mathcal{D}(G) = G^{\text{sc}}$ due to our assumptions. This implies that $\widehat{\text{ev}}_{\mathcal{D}(G)}^\delta = \text{ev}_{G^{\text{sc}}}^\delta$ as observed above. Let $b \in \text{ker}(\text{ev}_{G^{\text{sc}}}^\delta) \subset (\text{Sym}^2 \Lambda^*(T_{G^{\text{sc}}}))^{\mathscr{W}G}$. By Lemma 3.13(1), we can choose $\widetilde{b} \in (\text{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}G}$ such that

$\tilde{b}|_{\Lambda(T_{G^{sc}}) \times \Lambda(T_{G^{sc}})} = b$, which implies that the class of $\tau_G^\delta(\tilde{b}) \in \text{RPic}(\text{Bun}_G^\delta(C/S))$ is a lift of b . By the same argument of [22, Def./Lemma 3.7, Prop. 5.2], we have that

$$\text{wt}_G^\delta(\tau_G^\delta(\tilde{b})) = \tilde{b}(\delta \otimes -) := [\tilde{b}(d \otimes -)] \in \frac{\Lambda^*(T_G)}{\Lambda^*(T_{G^{\text{ad}}}},$$

where $d \in \Lambda(T_G)$ is any lift of δ . Since b is in the kernel of $\text{ev}_{G^{sc}}^\delta$, the element $\text{wt}_G^\delta(\tau_G^\delta(\tilde{b}))$ lies in the image of $\overline{\Lambda_{\text{ab}}^*}$, i.e. there exists $\chi \in \Lambda^*(G^{\text{ab}})$ such that

$$\tilde{b}(\delta \otimes -) = \chi(\Lambda_{\text{ab}}(-)). \tag{4.15}$$

By the definition of the boundary homomorphism, we have that

$$\partial(b) = [\chi(\delta^{\text{ab}})] \in \mathbb{Z}/2\mathbb{Z} \cong \text{coker}(\text{wt}_{G^{\text{ab}}}^{\delta^{\text{ab}}}).$$

Since $\delta^{\text{ab}} = \Lambda_{\text{ab}}(d)$ for any lift $d \in \Lambda(T_G)$ of δ , Equation (4.15) implies that

$$\chi(\delta^{\text{ab}}) = \tilde{b}(\delta \otimes b) = \tilde{b}(d \otimes d).$$

But $\tilde{b}(d \otimes d)$ is even since $\tilde{b} \in (\text{Sym}^2 \Lambda^*(T_G))^{\mathscr{W}G} \cong \text{Bil}^{\text{s, ev}}(\Lambda(T_G))^{\mathscr{W}G}$, which forces $\partial(b)$ to be zero, q.e.d. □

Acknowledgement. We thank Guido Lido for useful conversations concerning Remark 2.5.

Funding Statement. F.V. is funded by the MIUR ‘Excellence Department Project’ MATH@TOV, awarded to the Department of Mathematics, University of Rome Tor Vergata, CUP E83C18000100006, by the PRIN 2022 ‘Moduli Spaces and Birational Geometry’ and the PRIN-2020 ‘Curves, Ricci flat Varieties and their Interactions’ funded by MIUR (Ministero dell’Istruzione e del Merito), by the project EXPL/MAT-PUR/1162/2021 (DOI:10.54499/EXPL/MAT-PUR/1162/2021) funded by FCT (Fundação para a Ciência e a Tecnologia), by the CMUC (Centro de Matemática da Universidade de Coimbra).

References

- (1) E. Ambrosi, Specialization of Néron-Severi groups in positive characteristic, *Ann. Sci. Éc. Norm. Supér. (4)* **56**(3) (2023), 665–711.
- (2) E. Arbarello and M. Cornalba, The Picard groups of the moduli spaces of curves, *Topology* **26**(2) (1987), 153–171.
- (3) A. Beauville, Vector bundles on Riemann surfaces and conformal field theory, In *R.C.P. 25, Vol. 46 (French) (Strasbourg, 1992/1994)*, Prépubl. Inst. Rech. Math. Av., Volume 1994/29, pp. 127–147 (Universit Louis-Pasteur, Strasbourg, 1994).
- (4) A. Beauville, Conformal blocks, fusion rules and the Verlinde formula, In *Proceedings of the Hirzebruch 65 conference on algebraic geometry (Ramat Gan, 1993)*, Israel Math. Conf. Proc., Volume 9, pp. 75–96 (Bar-Ilan University, Ramat Gan, 1996).

- (5) A. Beauville and Y. Laszlo, Conformal blocks and generalized theta functions, *Comm. Math. Phys.* **164**(2) (1994), 385–419.
- (6) A. Beauville, Y. Laszlo and C. Sorger, The Picard group of the moduli of G -bundles on a curve, *Compos. Math.* **112**(2) (1998), 183–216.
- (7) I. Biswas and N. Hoffmann, The line bundles on moduli stacks of principal bundles on a curve, *Doc. Math.* **15** (2010), 35–72.
- (8) I. Biswas and N. Hoffmann, Poincaré families of G -bundles on a curve, *Math. Ann.* **352**(1) (2012), 133–154.
- (9) I. Biswas and Y. I. Holla, Brauer group of moduli of principal bundles over a curve, *J. Reine Angew. Math.* **677** (2013), 225–249.
- (10) S. Bosch, W. Lütkebohmert and M. Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], Volume 21, (Springer-Verlag, Berlin, 1990).
- (11) A. Boysal and S. Kumar, Explicit determination of the Picard group of moduli spaces of semistable G -bundles on curves, *Math. Ann.* **332**(4) (2005), 823–842.
- (12) A. Christensen, Specialization of Néron-Severi groups in characteristic p , Preprint arXiv:1810.06550, 2018.
- (13) C. Ciliberto, Rationally determined line bundles on families of curves, In *The curves seminar at Queen's, Vol. IV (Kingston, ON, 1985–1986)*, Queen's Papers in Pure and Appl. Math., Volume 76, ExNo. D, 48 (Queen's University, Kingston, ON, 1986).
- (14) C. Ciliberto, On rationally determined line bundles on a family of projective curves with general moduli, *Duke Math. J.* **55**(4) (1987), 909–917.
- (15) J.-M. Drezet and M. S. Narasimhan, Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques, *Invent. Math.* **97**(1) (1989), 53–94.
- (16) G. Faltings, A proof for the Verlinde formula, *J. Algebr. Geom.* **3**(2) (1994), 347–374.
- (17) G. Faltings, Algebraic loop groups and moduli spaces of bundles, *J. Eur. Math. Soc. (JEMS)* **5**(1) (2003), 41–68.
- (18) R. Fringuelli, The Picard group of the universal moduli space of vector bundles on stable curves, *Adv. Math.* **336** (2018), 477–557.
- (19) R. Fringuelli and R. Pirisi, The Picard group of the universal abelian variety and the Franchetta conjecture for abelian varieties, *Michigan Math. J.* **68**(3) (2019), 651–671.
- (20) R. Fringuelli and R. Pirisi, The Brauer group of the universal moduli space of vector bundles over smooth curves, *Int. Math. Res. Not. IMRN* **2021**(18) (2019), 13609–13644.
- (21) R. Fringuelli and F. Viviani, The Picard group of the universal moduli stack of principal bundles on pointed smooth curves, *J. Topol.* **15**(4) (2022), 2065–2142.
- (22) R. Fringuelli and F. Viviani, The Picard group of the universal moduli stack of principal bundles on pointed smooth curves II, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **24**(1) (2023), 367–447.
- (23) R. Fringuelli and F. Viviani, On the Brauer group of the moduli stack of principal bundles over families of curves. In preparation.
- (24) R. Fringuelli F. Viviani, *On the Picard group scheme of the moduli stack of stable pointed curves*, Preprint arXiv:2005.06920, 2020.
- (25) S. Koizumi, The ring of algebraic correspondences on a generic curve of genus g , *Nagoya Math. J.* **60** (1976), 173–180.
- (26) A. Kouvidakis, The Picard group of the universal Picard varieties over the moduli space of curves, *J. Differential Geom.* **34**(3) (1991), 839–850.
- (27) A. Kouvidakis, On the moduli space of vector bundles on the fibers of the universal curve, *J. Differential Geom.* **37**(3) (1993), 505–522.
- (28) S. Kumar, M. S. Narasimhan and A. Ramanathan, Infinite Grassmannians and moduli spaces of G -bundles, *Math. Ann.* **300**(1) (1994), 41–75.

- (29) S. Kumar and M. S. Narasimhan., Picard group of the moduli spaces of G -bundles, *Math. Ann.* **308**(1) (1997), 155–173.
- (30) Y. Laszlo, Linearization of group stack actions and the Picard group of the moduli of SL_r/μ_s -bundles on a curve, *Bull. Soc. Math. France* **125**(4) (1997), 529–545.
- (31) Y. Laszlo and C. Sorger, The line bundles on the moduli of parabolic G -bundles over curves and their sections, *Ann. Sci. École Norm. Sup. (4)* **30**(4) (1997), 499–525.
- (32) D. Maulik and B. Poonen, Néron-Severi groups under specialization, *Duke Math. J.* **161**(11) (2012), 2167–2206.
- (33) M. Melo and F. Viviani, The Picard group of the compactified universal Jacobian, *Doc. Math.* **19** (2014), 457–507.
- (34) N. Mestrano, Conjecture de Franchetta forte, *Invent. Math.* **87**(2) (1987), 365–376.
- (35) N. Mestrano and S. Ramanan, Poincaré bundles for families of curves, *J. Reine Angew. Math.* **362** (1985), 169–178.
- (36) S. Mori, The endomorphism rings of some Abelian varieties, *Jpn. J. Math. (N.S.)* **2**(1) (1976), 109–130.
- (37) C. Pauly, Espaces de modules de fibrés paraboliques et blocs conformes, *Duke Math. J.* **84**(1) (1996), 217–235.
- (38) G. Pietro Pirola, Base number theorem for abelian varieties. An infinitesimal approach, *Math. Ann.* **282**(3) (1988), 361–368.
- (39) M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Mathematics, Volume 119, (Springer-Verlag, Berlin-New York, 1970).
- (40) S. Schröer, The strong Franchetta conjecture in arbitrary characteristics, *Internat. J. Math.* **14**(4) (2003), 371–396.
- (41) C. Sorger, La formule de Verlinde, *Astérisque* **237** (1996), ExNo. 794, 3, 87–114 Séminaire Bourbaki, Vol. 1994/95.
- (42) C. Sorger, On moduli of G -bundles of a curve for exceptional G , *Ann. Sci. École Norm. Sup. (4)* **32**(1) (1999), 127–133.
- (43) C. Teleman, Borel-Weil-Bott theory on the moduli stack of G -bundles over a curve, *Invent. Math.* **134**(1) (1998), 1–57.
- (44) The Stacks Project Authors, *Stacks project*, <https://stacks.math.columbia.edu>, 2018.
- (45) O. Zariski, On a theorem of Severi, *Amer. J. Math.* **50**(1) (1928), 87–92.
- (46) W. Zomervrucht, *Ineffective descent of genus one curves*, <http://arxiv.org/abs/1501.04304>, 2015.