

The municipalities offers a home visit to the population at the ages of 75 and 80 years to assess the need of care and prevent sickness. These home visits are well established and might offer an unused opportunity to detect cognitive impairment and dementia.

Objectives: To assess impaired cognition at home visits in order to initiate clinical examination for dementia.

Methods: A feasibility study with the use of Brief Assessment of Impaired Cognition Questionnaire (BASIC-Q) (sensitivity 0.92, specificity 0.97) at home visits. It is expected to include 1000 participants without a dementia diagnosis at the ages of 75 and 80 years. Participants will be included in a period of 12 months (in the year of 2022), in a number of municipalities.

If the screening for cognitive impairment is positive, the participant is motivated for clinical examination at the general practitioner. Follow-up through registers and general practitioners.

Results: Preliminary results will be presented at the conference.

Conclusions: Assessment of cognition might give an opportunity to start medication and social support early in the elderly with impaired cognition and undiagnosed dementia.

Disclosure: No significant relationships.

Keywords: screening; Dementia; cognitive impairment

EPV0946

The good compliance is an opportunity to avoid pathological brain aging

A. Sidenkova

Ural State Medical University, Psychiatry, Yekaterinburg, Russian Federation

doi: 10.1192/j.eurpsy.2022.1678

Introduction: Preservation of health, increase in life expectancy determine the need to improve the effectiveness of medical recommendations, which, despite the success of pharmacology, are insufficient for reasons related to the low level of compliance with these recommendations by patients.

Objectives: Participants of the study-148 employees of medical institutions: 12 men, 136 women, their age ranged from 27 to 74 years.

Methods: Despite the absence of signs of decompensation of concomitant pathology, representatives of the subgroups took a different amount of concomitant therapy. Using the scale of assessment of drug compliance, it was found that compliance is most reduced in the subgroup of 41-50 years. In this subgroup, a comprehensive decrease in compliance across the "behavioral", "emotional", and "cognitive" domains was detected in 87.8% of cases, while in the younger subgroup partial non-compliance was 32.4%, in the older subgroup - 74.5%

Results: An analysis of the states of cognitive functions in 52 representatives of the middle age subgroup with low compliance rates showed that, unlike other representatives of the same subgroup, their indices for a number of neuropsychological tests are close to the results of more adult participants in the study. Individuals demonstrating low compliance with quite favorable CNS resources are at risk for the formation of pathological aging.

Conclusions: Compliance is considered as Compliance is considered as a control mechanism for preventing normal aging into pathological by regulating risk factors that are dangerous for the brain and associated with the formation of dementia

Disclosure: No significant relationships.

Keywords: compliance; pathological brain aging

EPV0948

Electronic smart-hub based intervention during COVID-19 in a rural Psychiatry of Old Age service in North-West Ireland.

S. Patel^{1,2}, A. Gannon³, M. Cryan³, C. Dolan^{2,4}, C. McDonald^{1,2}, C. Houghton⁵ and G. Mccarthy^{2,4*}

¹University Hospital Galway, Psychiatry, Galway, Ireland; ²NUI Galway, School Of Medicine, Galway, Ireland; ³Sligo Leitrim Mental Health Services, St. Columba's Hospital,, Psychiatry, Sligo, Ireland;

⁴Sligo Leitrim Mental Health Services, St. Columba's Hospital,, Psychiatry, Sligo, Ireland and ⁵NUI Galway, School Of Nursing And Midwifery, Galway, Ireland

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1679

Introduction: The COVID-19 pandemic caused significant disruptions in services and necessitated innovation to continue care provision to the vulnerable population of older adults with psychiatric needs.

Objectives: The objective of this study was to examine the experiences of staff and patients using a hands-free electronic smart-hub (eSMART hub) intervention to keep patients connected with psychiatry of old age following COVID-19 restrictions.

Methods: A risk stratification register was created of all patients known to the Psychiatry of Old Age service in the North-West of Ireland to identify those at highest risk of relapse. These patients were offered a smart-hub with remote communication and personal assistant technology to be installed into their homes. Smart-hubs were also installed in the team base to facilitate direct device to device communication. Semi-structured qualitative interviews were conducted with 10 staff members and 15 patients at 6-12 months following the installation of the smart-hubs.

Results: The smart-hubs were utilized by the POA team to offer remote interventions over video including clinician reviews, regular contact with key workers and day-hospital based therapeutic interventions such as anxiety management groups and OT led physical exercises. Patients also used the personal assistant aspect of the hub to attend to personal hobbies such as accessing music and radio. Positive feedback related to companionship during isolation and connectivity to services. Negative feedback was mainly related to technology, particularly internet access and narrow scope of communication abilities.

Conclusions: Electronic smart-hub devices may offer an acceptable avenue for remote intervention and communication for isolated high-risk older persons.

Disclosure: The smart hub devices used in this study were donated by Amazon. However, the company was not involved in any other aspect of the study and the researchers have no significant financial interest, consultancy or other relationship with products, manufacturer

Keywords: Covid-19; smart hub; Older Adults; technology

EPV0949

The occurrence of symptoms of fear of COVID-19 among participants of the Polish University of the Third Age

M. Cybulski^{1*}, U. Cwalina² and E. Krajewska-Kulak¹

¹Medical University of Białystok, Department Of Integrated Medical Care, Białystok, Poland and ²Medical University of Białystok, Department Of Statistics And Medical Informatics, Białystok, Poland

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1680

Introduction: The older adults have been considered one of the groups at highest risk of SARS-CoV-2 infection and death due to COVID-19. Fear of SARS-CoV-2 infections has become widespread. It's constantly being enhanced by the media reports and social distancing principle.

Objectives: The aim of the study was to assessment the occurrence of symptoms of fear of COVID-19 among participants of the Polish University of the Third Age (UTA).

Methods: The study included 296 participants of the UTA in Poland, including 258 women and 38 men. The study conducted with the use of the following validated psychometric scales: General Anxiety Disorder-7 (GAD-7), Short Health Anxiety Inventory (SHAI) and State-Trait Anxiety Inventory (STAI).

Results: The mean scores in STAI and SHAI demonstrated mild symptoms indicative of anxiety disorders in the older adults. Women and men did differ significantly in terms of the scores obtained in STAI(X-1) ($p=0.002$) and STAI(X-2) ($p=0.020$). There were no statistically significant differences between respondents with higher education and those with a different level of education. The single respondents differed significantly from divorced ones in terms of STAI(X-1) ($p=0.046$). Moreover, widows/widowers differed significantly from divorced ones in terms of STAI(X-2) ($p=0.045$) and GAD-7 ($p=0.032$).

Conclusions: The subjective experience of anxiety symptoms associated with fear of contracting COVID-19 was increased due to the ongoing pandemic, but was not significantly high in the analysed population of older people. COVID-19-related anxiety was significantly more common in lonely individuals. Women and men differed significantly in terms of perceived state anxiety and trait anxiety measured by STAI.

Disclosure: No significant relationships.

Keywords: fear; Covid-19; Older Adults; Anxiety

EPV0950

An old way of forgetting

J.P. Rebeca Coelho*

Centre Neuchâtois de Psychiatrie, Cnpa1, Neuchâtel, Switzerland

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1681

Introduction: During the last decades, the incidence of syphilis is on the rising, particularly in the United States of America and Europe. Neurosyphilis is a disease that has a vast differential diagnostic. With that in mind, clinicians have some difficulties to identify it rapidly. A case of a 57-year-old man is presented, with a brutal change in his behavior, associated with a dementia-like syndrome. He is diagnosed with neurosyphilis.

Objectives: The main goal is to present his clinical psychiatric symptoms and diagnosis procedure, the treatment that he received and his clinical outcome in the psychogeriatric department.

Methods: The treatment was based in an integrated framework of pharmacology and psychotherapy.

Results: The patient was able to slowly recover and to get back home, we a solid structure to make the follow up.

Conclusions: This clinical vignette represents a growing number of adult patients that present themselves for the first time with dementia-like symptoms. It is important to remember, that many diseases are capable of mimicking dementia and their exclusion before admitting a diagnosis of dementia is mandatory.

Disclosure: No significant relationships.

Keywords: Syphilis; Dementia

EPV0951

What we know about Auditory Charles Bonnet Syndrome?

C. Moreno Menguiano^{1*}, F. Garcia Sánchez², J.J. Vazquez Vazquez³, M.M. Gutiérrez Rodríguez² and M.D.L.A. Corral Y Alonso²

¹Hospital Universitario de Mostoles, Psychiatry, Mostoles (Madrid), Spain; ²HOSPITAL UNIVERSITARIO DE MÓSTOLES, Psychiatry, MÓSTOLES (MADRID), Spain and ³Centro de Salud Mental de Móstoles / Hospital Universitario de Móstoles, Psychiatry, Móstoles, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1682

Introduction: Charles Bonnet syndrome (CBS) is defined like visual hallucinations found in individuals who are not necessarily mentally ill, who have visual impairment and no cognitive deficits. Although CBS make reference to visual hallucinations, in this case we are going to deal about Auditory Charles Bonnet Syndrome (aCBS), a very infrequent condition that consists in the presentation of musical hallucinations in patients with sensorineural hearing loss and which etiology is not clearly due to a psychiatric condition.

Objectives: Review the scientific literature available on aCBS to see how much we know about this syndrome.

Methods: Review of available literature sources were obtained through electronic search in PubMed database.

Results: Musical hallucination is a complex form of auditory hallucinations. The most common of these are idiopathic and they present in elderly patients with deafness or impaired audition, which suggests a deterioration of cerebral function. The pathophysiological mechanism is not understood. These patients tend to have intact reality tests. The time course is variable. In those cases in which it is possible, treating the hearing loss can lead to a significant improvement of the symptom. However, when every this strategies are insufficient, pharmacological treatments can be considered.

Conclusions: - aCBS is an uncommon condition characterized by the presence of complex auditory hallucinations that mainly affect elderly patients with hearing loss. - In most cases there is no previous psychiatric history. - The etiology and pathophysiological are not well defined. - There is no etiological treatment. We can use pharmacological and no pharmacological methods of treatment.

Disclosure: No significant relationships.

Keywords: old age; Charles-Bonnet Syndrome; auditory