


Article

Seasonal paleoecological records from antler collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$

Rachel Schwartz-Narbonne* , Tessa Plint , Elizabeth Hall, Grant Zazula,
and Fred J. Longstaffe

Abstract.—Cervids living in high latitudes have evolved to thrive in ecosystems that experience dramatic seasonal changes. Understanding these seasonal adaptations is important for reconstructing cervid life histories, ecosystem dynamics, and responses in the distant and not-so-distant past to changing seasonality caused by climate change. Cervid antlers provide a rare opportunity for insight into faunal seasonal ecology, as they are grown and shed each year. Stable isotopes of carbon and nitrogen measured directly from antlers have the potential to provide seasonal dietary data for individuals. If the isotopic signals in bone and antler are controlled by the same metabolic processes, then the stable carbon and nitrogen isotope compositions of collagen ($\delta^{13}\text{C}_{\text{Coll}}$ and $\delta^{15}\text{N}_{\text{Coll}}$) from incrementally grown antler tissue provide time-constrained dietary signals from the spring and summer growth season. Bone, by comparison, provides an average signal over several years. The amino acid (glutamate and phenylalanine) $\delta^{15}\text{N}$ in antlers from modern captive caribou showed similar trophic discrimination factors to earlier results for other collagenous tissues (bone, tooth dentin, and cementum). Hence, growth rate was not the primary control on the stable isotope composition of antler collagen. We applied this knowledge to assess seasonal shifts in Quaternary fossils of three Cervidae species: elk (*Cervus elaphus*), moose (*Alces alces*), and caribou (*Rangifer tarandus*). Paired antler–bone $\delta^{13}\text{C}_{\text{Coll}}$ and $\delta^{15}\text{N}_{\text{Coll}}$ from the same individual were used to identify differences between summer and annual diet and ecology. Intra-antler isotopic variability from serially sampled antlers was used to examine seasonal dietary shifts and specialization.

Rachel Schwartz-Narbonne[†], Tessa Plint[‡], and Fred J. Longstaffe. Department of Earth Sciences, University of Western Ontario, London, Ontario, Canada N6A 5B7. E-mail: flongsta@uwo.ca. Present addresses:

[†]Biomolecular Sciences Research Centre, Sheffield Hallam University, Sheffield S1 1WB, U.K.

E-mail: r.schwartz-narbonne@shu.ac.uk. [‡]Lyell Centre, School of Energy, Geoscience, Infrastructure and Society, Heriot-Watt University, Edinburgh EH14 4AS, U.K. E-mail: tp46@hw.ac.uk

Elizabeth Hall and Grant Zazula[§], Government of Yukon, Department of Tourism & Culture (Palaeontology Program), Whitehorse, Yukon Territory Y1A 2C6, Canada. Email: elizabeth.hall@gov.yk.ca, and grant.zazula@gov.yk.ca. [§]Additional address: Collections and Research, Canadian Museum of Nature, P.O. Box 3443, Station D, Ottawa, Ontario, Canada, K1P 6P4.

Accepted: 31 December 2020

*Corresponding author.

Introduction

Seasonal Ecological Variation

High-latitude ecosystems experience extreme seasonality, with the timing of their short growing season dependent on a combination of temperature, snowmelt, and photoperiod (Ernakovich et al. 2014). Animals have adapted to this seasonality, with annual metabolic events (i.e., pregnancy, weaning, use of fat reserves, hibernation) coordinated based on seasonal resource availability (Gilg et al. 2012). Temperature and precipitation, and thus the growing season, at high latitudes were strongly affected by climatic

variability during the Pleistocene (e.g., Guthrie 2001) and are expected to change again dramatically with contemporary climate change (Gilg et al. 2012; Ernakovich et al. 2014). Large terrestrial herbivores such as cervids (family Cervidae) may experience “trophic mismatch” if they are unable to adapt the timing of annual metabolic events to changing seasonality (Gilg et al. 2012; Ernakovich et al. 2014). Methods to track cervid responses to environmental change, such as isotope-based tracking of seasonal diets, are therefore important to study high-latitude ecosystem shifts caused by climate change (Zhao et al. 2019). These methods are also

© The Author(s), 2021. Published by Cambridge University Press on behalf of The Paleontological Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

0094-8373/21

important to elucidate Pleistocene ecology. The diversity of Pleistocene herbivore megafauna coexisting in high-latitude environments may have involved seasonally based resource partitioning, with dietary specialization during periods of winter resource scarcity and generalist diets during the growing season (Schwartz-Narbonne et al. 2019). Seasonality may thus have played a key role in Pleistocene herbivore megafauna interactions.

Cervid antler tissue grows rapidly, and the antler sets are fully developed after 4 months (Chapman 1975). Despite this intense energy investment, cervids that live in high-latitude environments shed and regrow their antlers annually. Antler is structurally analogous to bone and is composed of hydroxyapatite mineral deposited within a collagen protein matrix. The stable carbon and nitrogen isotope compositions ($\delta^{13}\text{C}_{\text{Coll}}$ and $\delta^{15}\text{N}_{\text{Coll}}$) of bone collagen closely correlate with those of the individual's diet (Kelly 2000). In this study, we assess whether seasonal changes in diet or nutritional stress are recorded in cervid antler $\delta^{13}\text{C}_{\text{Coll}}$ and $\delta^{15}\text{N}_{\text{Coll}}$. In addition, we compare the spring–summer isotopic signal in antler with the multi-year signal in bone from the same individual.

The $\delta^{13}\text{C}_{\text{Coll}}$ and $\delta^{15}\text{N}_{\text{Coll}}$ of three cervid species were examined in this study: elk (*Cervus elaphus*, also known as red deer or wapiti), moose (*Alces alces*, also known as Eurasian elk), and caribou (*Rangifer tarandus*, also known as reindeer). We tested for metabolic controls on antler $\delta^{15}\text{N}_{\text{Coll}}$ using antler amino acid (glutamate and phenylalanine) $\delta^{15}\text{N}$ ($\delta^{15}\text{N}_{\text{Glu}}$ and $\delta^{15}\text{N}_{\text{Phe}}$) from modern caribou living in captivity. The study includes 8 paleo-antlers measured for intra-antler isotopic variability (2 elk; 2 moose; 4 caribou), 2 paired antler–bone paleo-samples (1 moose; 1 caribou), and 6 modern samples (5 caribou; 1 Ankole bull). All paleontological specimens originate from Yukon Territory, Canada, and a subset were confirmed by radiocarbon dating to be of Pleistocene or Holocene age.

Temporal changes in an animal's diet can be assessed through stable isotope analysis of (1) a single tissue repeatedly sampled over time, (2) multiple tissues with varying turnover rates, or (3) a single tissue with progressive growth (Dalerum and Angerbjörn 2005; Davis and

Pineda-Munoz 2016). Previous studies of present-day cervids examined short-term ecological changes by comparing the isotopic signal of bone collagen, which provides a multi-year average, to that of tooth collagen, which represents only the brief period of time during youth when the tooth formed (Drucker et al. 2001, 2012). Seasonal changes in diet have also been identified by serial sampling of keratin-based tissues such as hooves (Barnett 1994; Kielland 2001; Walter and Leslie 2009) and hair (Drucker et al. 2010; Zhao et al. 2019). Keratinous tissues, however, do not typically preserve well over archeological or paleontological timescales, whereas bone, tooth, and antler are more likely to survive. In some contexts, antler represents a large proportion of the available sample material in paleontological archives, surpassing the number of other skeletal elements from the same species (Jay and Richards 2007; Kuitens et al. 2015; Miller et al. 2016). Thus, understanding the temporal scale of dietary information that stable isotope analysis of antler can—and cannot—provide is highly relevant (Davis and Pineda-Munoz 2016).

Antler and Bone Growth

Antler growth is rapid, with continuous remodeling occurring spatially at the cellular level within the tissue during its formation. Antler development begins at the pedicle, or attachment point to the skull, with new tissue deposited at the tip of the growing antler (Fig. 1) (Banks and Newbrey 1983). The upper portion of the antler grows faster than the lower portion, correlated with the greater availability of high-quality forage during late spring and summer (Van Ballenberghe 1983; Finstad and Kielland 2011; Gomez et al. 2013). The outer circumference of an antler is composed of layers of woven bone that are systematically remodeled into dense lamellar cortical bone. In contrast, the interior of an antler is composed of spongy cancellous bone deposited toward the very end of the growth period (Fig. 1) (Gomez et al. 2013; Kierdorf et al. 2013). The final stage of tissue growth consists of an extremely thin layer of woven bone deposited over the pedicle and antler base (Fig. 1) (Gomez et al. 2013; Kierdorf et al. 2013). All growth takes




FIGURE 1. Sampling strategy: A, an example of the sampling pattern (arrows) for a caribou skull and antler (specimen YG 361.52; bone sample removed from right occipital condyle); B, conceptual model of the timing of antler tissue formation in cranial bone versus antler; and C, cross-section of an antler shaft differentiating cortical and cancellous bone. This study measured variability along the length of the antler in cortical bone.

place while the antler is covered in a layer of highly vascular skin (called velvet) that supplies nutrients and oxygen to the living bone tissue of the antler. Once antler growth is complete, the velvet is shed. The finished antler is effectively dead avascular tissue in which no further remodeling takes place (Gomez et al. 2013).

The overall growth pattern makes it possible to record seasonal changes in tissue $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ across both the length and width of an antler (Fig. 1). In contrast, the bone tissue that comprises the rest of the skeletal elements in the body is vascular and remodels continuously over the course of the animal's lifetime, yielding an isotopic signal that is an average of multiple years. In most cervid species, only males take on the calcium burden of growing antlers, with the exception of caribou. Both male and female caribou grow antlers. Males use antlers primarily to attract females for mating and for fighting with other males during the rut; the antlers are commonly shed before winter. Antler growth in males generally occurs between the late spring and summer months, and the velvet is shed during late summer or early autumn before the rutting season (Chapman 1975; Van Ballenberghe 1983). Resorption of mineralized tissue between the base of the antler and pedicle occurs after the

mating season, and the antler is eventually shed. Antler growth in caribou females begins about a month later than male antler growth (Chapman 1975; Leader-Williams 1988). Female caribou typically retain their antlers over the winter months; females primarily use antlers to access forage underlying snow to obtain sufficient resources to sustain themselves and their unborn calves.

Stable Isotopes

Stable isotope compositions are reported using δ -notation, which compares sample isotope ratios (e.g., $^{13}\text{C}/^{12}\text{C}$, $^{15}\text{N}/^{14}\text{N}$) to internationally accepted standard isotope ratios, in per mille (‰). The reference standards are VPDB for carbon isotope ratios (Coplen et al. 2006) and AIR for nitrogen isotope ratios (Mariotti 1983).

$\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of Plants.—Cervids consume a herbivorous diet. The isotopic composition of photosynthetic organisms is highly variable. The main variability in terrestrial plant isotopic compositions depends on the method of carbon fixation. Modern plant species that employ the C_3 photosynthetic pathway have an average $\delta^{13}\text{C}$ of $\sim -27\text{‰}$. Plants that utilize the C_4 photosynthetic pathway comprised the vast majority (and at times, the entirety) of the

vegetation in North American high-latitude regions during the late Pleistocene and Holocene (Blinnikov et al. 2011; Gaglioti et al. 2011; Kristensen et al. 2011). Other factors that can influence C_3 vegetation stable carbon isotope composition include extensive canopy cover, which typically leads to more negative plant $\delta^{13}C$; warmer temperatures, which lead to more positive plant $\delta^{15}N$; and increased aridity, which results in more positive plant $\delta^{13}C$ and $\delta^{15}N$ (O'Leary 1981; Tieszen 1991; Koch et al. 1994; Kohn 2010). While plant isotopic baselines can vary among sites and over time (Schwartz-Narbonne et al. 2019), the average $\delta^{13}C$ and $\delta^{15}N$ of aquatic macrophytes, C_3 vascular terrestrial plants (shrubs and herbs), lichens, mosses, and fungi in high-latitude ecosystems tend to follow a predictable pattern (Fig. 2). The observed isotopic patterns for these plants and fungi from high-latitude sites is: $\delta^{13}C$ of shrubs < herbs < fungi < mosses < lichens < freshwater macrophytes. For $\delta^{15}N$, the pattern is: moss < lichens < shrubs < herbs < freshwater macrophytes < fungi (Fig. 2) (Barnett 1994; Nadelhoffer et al. 1996; Ben-David et al. 2001; Kielland 2001; Tischler 2004; Wang and Wooller 2006; Milligan 2008; Drucker et al. 2010; Finstad and Kielland 2011; Kristensen et al. 2011; Tahmasebi et al. 2017; Plint et al. 2019). The extensive overlap in the data compiled from a range of high-latitude sites is expected, given the range of $\delta^{13}C$ and $\delta^{15}N$ among plant types within a single site.

$\delta^{13}C$ and $\delta^{15}N$ of Herbivores.—The stable carbon and nitrogen isotope compositions of a consumer's bodily tissues reflect the protein component of diet, after accounting for trophic discrimination. For large herbivores, there is a well-documented trophic increase of $\sim +5\%$ for $\delta^{13}C$, and $+3$ to $+5\%$ for $\delta^{15}N$ from diet to collagenous tissues (bone, tooth dentin and cementum) (Koch et al. 1994; Bocherens 2015). Nutritional stress is thought to produce higher $\delta^{15}N$ in cervids (Parker et al. 2005), although this effect is disputed (see Drucker et al. 2012). We use $\Delta^{15}N_{\text{Glu-Phe}}$ ($\delta^{15}N_{\text{glutamate}} - \delta^{15}N_{\text{phenylalanine}}$) to evaluate whether rapid tissue growth, such as occurs within antlers, has affected the $\delta^{15}N$ of collagenous tissues, including antlers, as has been suggested in the past

(Trueman et al. 2005; Waters-Rist and Katzenberg 2010; Madgwick et al. 2013; Osborne 2017).

The $\delta^{15}N$ of individual amino acids in bone collagen can be used to assess the extent of metabolic enrichment between diet and tissue. The difference between the $\delta^{15}N$ of phenylalanine (Phe: a "source" amino acid whose $\delta^{15}N$ derives from diet) and of glutamate (Glu: a "trophic" amino acid whose $\delta^{15}N$ increases with each subsequent metabolic cycling event) is a measure of metabolic enrichment in an animal. Higher $\Delta^{15}N_{\text{Glu-Phe}}$ occurs in animals higher up in the food chain (Chikaraishi et al. 2007, 2009, 2015; McMahon and McCarthy 2016). Values of $\Delta^{15}N_{\text{Glu-Phe}}$ can change in animals fed a low-quality diet (McMahon and McCarthy 2016) or experiencing starvation (Barreto-Curiel et al. 2017) (although the latter is debated; see Ishikawa et al. 2017). Thus, metabolic controls, in addition to trophic level, have the potential to change an animal's $\Delta^{15}N_{\text{Glu-Phe}}$. If bone, tooth, and antler $\delta^{15}N_{\text{Coll}}$ are controlled by the same metabolic processes, we expect the $\Delta^{15}N_{\text{Glu-Phe}}$ in these three tissues to be the same for ruminant herbivores, in the absence of dietary stress. If rapid tissue growth is a metabolic control on antler $\delta^{15}N_{\text{Coll}}$, we expect that to be reflected in distinct antler $\Delta^{15}N_{\text{Glu-Phe}}$.

Materials and Methods

Specimen Collection

The Pleistocene and Holocene caribou, elk, and moose fossil specimens originate from Qikiqtaruk (Herschel Island), southern Beaufort Sea, and the Klondike region of west-central Yukon Territory (Fig. 3). These specimens are curated in the fossil collections of the Yukon Palaeontology Program of the Yukon Government in Whitehorse, Yukon (specimen prefix YG = Yukon Government). Antlers naturally shed by modern caribou were obtained over multiple years from the collections of the Bowmanville Zoo, located in Bowmanville, Ontario, Canada (specimen BZ = Bowmanville Zoo). Coeval skeletal material from these caribou was not available, as none had died recently. As such, the only bone tissue sample available from Bowmanville Zoo during the time of our sampling was from a recently deceased Anko-




FIGURE 2. Stable carbon and nitrogen isotope compositions of common, modern, high-latitude vegetation based on previous studies of Arctic and subarctic ecosystems (Barnett 1994; Wang and Wooller 2006; Milligan 2008; Tahmasebi et al. 2017; Plint et al. 2019) and presented as: A, data points; and B, ellipses. Herbs include annual and perennial grasses, forbs, and sedges (Tahmasebi et al. 2017). Woody plant angiosperms (AS) include angiosperm shrubs, shrubs, and trees. Woody plant gymnosperms (GS) include gymnosperm trees. Ellipses encompass 40% of the data for each forage type, corrected for small sample size (Parnell and Jackson 2013). Forage $\delta^{13}\text{C}$ was corrected for the Suess effect to the last glacial maximum $\delta^{13}\text{C}_{\text{atmosphere}}$ using the date of sample collection (Long et al. 2005; Tahmasebi et al. 2018). Isotopic results for all plant tissues were included unless the study identified them as roots or as outliers. Despite substantial overlap, different plant types tend to occupy different positions in $\delta^{15}\text{N}$ versus $\delta^{13}\text{C}$ space on this isotopic cross-plot.

Watusi bull. For ease of handling, the zoo-keepers sampled the easily removed dewclaw bone, which is a vestigial digit in cattle.

Collagen Extraction

Individual bone and antler samples were removed using a Dremel drill fit with a cutting wheel blade. Serial sampling along the length of antlers was performed using a 0.625 cm drill core attached to a drill press. Serial samples were taken every ~10 cm along one side of the main antler beam, with the base of the antler

designated as 0 cm (Fig. 1). Cancellous bone was removed from bone samples to allow for sampling of purely cortical bone. Outer antler tissue was preferentially sampled when antler thickness permitted. All antlers were fully mineralized, and no velvet remained.

Collagen extraction was performed at room temperature following the modified Longin method (Longin 1971; Schwartz-Narbonne et al. 2019). Samples were dissolved for 24 h in 0.25 M HCl, and subsequently in 0.5 M HCl, during which time the acid was refreshed every 1–3 days until the tissues were fully demineralized. Samples were then rinsed three times with deionized water. Humic substances were removed using 0.1 M NaOH treatment for 20 min at room temperature, which was repeated until the liquid remained colorless. Samples were then rinsed seven times with deionized water, and the pH was adjusted to <3. Samples were then placed in a 90°C oven for ~16 h to solubilize the collagen. The solubilized collagen was decanted and dried at 90°C and then weighed for stable isotope analysis.




FIGURE 3. Map of sample collection locations, with inset detail of paleo-sample collection locations. QI, Qikiqtaruk (Herschel Island); KD, the Klondike; BZ, Bowmanville Zoo. The modern samples were collected at a site distant from the paleo-samples.

Stable Isotope Measurements

The collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ were measured using a Costech elemental analysis (EA)

combustion system (ECS 4010) attached to a Thermo Scientific Delta V isotope ratio mass spectrometer (IRMS) operated in continuous-flow mode with helium as the carrier gas. Collagen samples were measured over a total of 11 analytical sessions. The stable carbon isotope data were calibrated to VPDB using a two-point scale anchored by either NBS-22 (accepted $\delta^{13}\text{C} = -30.03\text{‰}$; $\pm 0.0\text{‰}$ 1 SD, $n = 3$) (Coplen et al. 2006) and IAEA-CH-6 (accepted $\delta^{13}\text{C} = -10.45\text{‰}$; $\pm 0.2\text{‰}$ SD, $n = 18$) (Coplen et al. 2006), or USGS-40 (accepted $\delta^{13}\text{C} = -26.39\text{‰}$; $\pm 0.0\text{‰}$ SD, $n = 21$) (Coplen et al. 2006) and USGS-41 (accepted $\delta^{13}\text{C} = +37.63\text{‰}$; $\pm 0.1\text{‰}$ SD, $n = 24$) (Coplen et al. 2006). The nitrogen isotope data were calibrated to AIR using a two-point scale anchored by USGS-40 (accepted $\delta^{15}\text{N} = -4.52\text{‰}$; $\pm 0.1\text{‰}$ SD, $n = 21$) (Qi et al. 2003) and either IAEA-N2 (accepted $\delta^{15}\text{N} = +20.39\text{‰}$; $\pm 0.2\text{‰}$ SD, $n = 15$) (Qi et al. 2003) or USGS-41 (accepted $\delta^{15}\text{N} = +47.57\text{‰}$; $\pm 0.2\text{‰}$ SD, $n = 24$) (Qi et al. 2003). These standards were also used to calibrate the carbon and nitrogen contents and to calculate the atomic C/N ratio of each sample. Standards not used in the calibration curve were measured as unknowns to monitor instrument accuracy and precision. Every analytical session also included a keratin in-house standard (MP Biomedicals, cat. no. 90211, lot no. 9966H) for which the following average results (SD) were obtained ($n = 38$): $\delta^{13}\text{C} = -24.1 \pm 0.1\text{‰}$, $\delta^{15}\text{N} = +6.3 \pm 0.1\text{‰}$, and atomic C/N ratio = 3.7 ± 0.1 . These results compare well with the accepted values of $\delta^{13}\text{C} = -24.0\text{‰}$, $\delta^{15}\text{N} = +6.4\text{‰}$, and atomic C/N ratio of 3.7. A subset of samples ($n = 14$) were analyzed in duplicate or triplicate; reproducibility (SD) ranged from ± 0.0 to $\pm 0.2\text{‰}$ for $\delta^{13}\text{C}$, and from ± 0.0 to $\pm 0.3\text{‰}$ for $\delta^{15}\text{N}$.

Amino Acid $\delta^{15}\text{N}$ Measurements

In preparation for isotopic analysis, collagen was hydrolyzed into its constituent amino acids, which were then derivatized following the *N*-acetyl-methyl ester derivatives (NACME) procedure (Corr et al. 2007a,b; Styring et al. 2010). Approximately 6 mg of collagen was dissolved in 2 ml of double-distilled 6 N HCl and vortexed, and the air in the sample tube was replaced with $\text{N}_{2(\text{g})}$. Samples were then heated

at $100\text{--}110\text{°C}$ for 24 h, followed by drying at 40°C under a gentle stream of $\text{N}_{2(\text{g})}$. The hydrolyzed amino acids were then dissolved in 2 ml of methanol and stored at -25°C until derivatization. At this point, a 0.5 ml aliquot was taken, and 0.5 ml of 0.2 mg norleucine/ml methanol was added as an internal standard. Methanol was removed by drying under a gentle stream of $\text{N}_{2(\text{g})}$ at room temperature. A 1.85 M acidified methanol solution was prepared by dropwise addition of 0.8 ml of acetyl chloride to 5 ml of anhydrous methanol in an ice or dry ice bath. One milliliter of acidified methanol was added to each sample, and samples were vortexed and then heated at 75°C for 1 h. Samples were then dried in an ice bath using a gentle stream of $\text{N}_{2(\text{g})}$. One milliliter of dichloromethane (DCM) was added to each sample, and the resulting liquid was dried in an ice bath using a gentle stream of $\text{N}_{2(\text{g})}$. Addition of DCM and subsequent drying of the sample were repeated three times, after which the samples were stored at -25°C overnight.

A solution of acetic anhydride, triethylamine, and acetone (1 ml:2 ml:5 ml) was then prepared, of which 1 ml was added to each sample. Samples were then vortexed and heated at 60°C for 10 min, followed by drying in an ice bath under $\text{N}_{2(\text{g})}$. To extract the organic phase, 2 ml of ethyl acetate and 1 ml of saturated NaCl in MilliQ water were added to each sample, and the solution was shaken. After the inorganic and organic phases had separated, the organic phase was transferred by pipette to another vial. An additional 1 ml of ethyl acetate was then added to the inorganic phase to ensure complete removal of the organic phase. The extracted organic phases were combined and evaporated to dryness in an ice bath under a gentle stream of nitrogen. One milliliter of DCM was then added to each sample, and the solution was evaporated in an ice bath using a gentle stream of $\text{N}_{2(\text{g})}$. This step was repeated three times. Each sample was then dissolved in $50\ \mu\text{l}$ of ethyl acetate and vortexed, following which a $25\ \mu\text{l}$ aliquot was removed for nitrogen isotope analysis. Samples were stored at -25°C until the isotopic analysis was performed.

Amino acid $\delta^{15}\text{N}$ was measured using an Aliigent 6890N-Thermo Scientific Gas Chromatograph-Combustion 3-Thermo Scientific

DeltaPlus XL (GC-C-IRMS). The GC ramping program used follows Styring et al. (2010) with minor adjustments, beginning at 60°C (1 min), ramping up at 15°C/min to 120°C, 3°C/min to 190°C, and then at 5°C/min to a final temperature of 250°C (15 min). An Agilent Technologies VF-23MS column was used in the GC. Samples were injected in splitless mode at an injection temperature of 200°C. Samples were oxidized over CuO/NiO/Pt at 940°C and reduced over Cu at 650°C. The nitrogen gas from the sample was isolated by passing the gas through a liquid nitrogen trap using helium flowing at a rate of 2ml/min. Three reference gas pulses were introduced into the IRMS at the beginning of each analytical session, and one pulse was introduced at the end of each session. The isotopic composition of the reference gas was calibrated using four amino acid standards. Three of these standards, alanine, leucine, and phenylalanine, were purchased as their NACME derivatives from Sigma Aldrich. The fourth, proline, was purchased as an amino acid and derivatized in-house. The nitrogen isotope compositions of the derivatized standards were established by multiple measurements of their isotope ratios by EA-IRMS, as described earlier for bulk collagen.

All samples were analyzed in triplicate. An internal standard, norleucine, was analyzed as an unknown, along with the samples. Norleucine $\delta^{15}\text{N}$ was offset from the expected values by +1.0 to +4.3‰, with an average offset of +2.5‰ and an average reproducibility of $\pm 1.3\%$ for triplicate measurements. A graph of proline versus hydroxyproline $\delta^{15}\text{N}$ produces a linear relationship ($R^2 = 0.95$; $y = 1.2x - 0.9$) within the expected 1:1 relationship (O’Connell and Collins 2018), accounting for instrumental error.

Radiocarbon Dating and Suess Effect Correction of $\delta^{13}\text{C}$

Radiocarbon (^{14}C) dates were obtained for a subset of antler and bone samples. Collagen was extracted, combusted, graphitized, and dated at the University of Arizona Accelerator Mass Spectrometry (AMS) Laboratory. Dates are presented as uncalibrated radiocarbon years before present (1950 AD) and are listed

TABLE 1. Sample collection information.

Common Name	Species	Lab ID	Institution ID	Locality	Collection site	Latitude °N	Longitude °W	Classification	^{14}C date	^{14}C lab no.	^{14}C date reference	Tissue	Skeletal element
Caribou	<i>Rangifer tarandus</i>	YT126	YG 381.52	Qikiqtaruk	Last Chance Creek	69.6	139.0	Female	Post-bomb	AA103889	This work	Bone	Right occipital condyle
Caribou	<i>Rangifer tarandus</i>	YT126	YG 381.52	Qikiqtaruk	Last Chance Creek	69.6	139.0	Female	Post-bomb	AA103889	This work	Antler	
Caribou	<i>Rangifer tarandus</i>	YT125	YG 109.9	Klondike	Quartz Creek	63.8	139.0	Female	>41,100	AA103824	This work	Antler	
Caribou	<i>Rangifer tarandus</i>	YT127	YG 404.657	Klondike	Hunker Creek	64.0	139.2	Male	29,570 \pm 970	AA103825	This work	Antler	
Caribou	<i>Rangifer tarandus</i>	YT128	YG 306.468	Klondike	Last Chance Creek	64.0	139.2	Male				Antler	
Elk	<i>Cervus elaphus</i>	YT85/123	YG 104.04	Klondike	Last Chance Creek	64.0	139.2	Male	11,675 \pm 45; 11,785 \pm 50	OxA20918; OxA20919	Meiri et al. 2014	Antler	
Elk	<i>Cervus elaphus</i>	YT124	YG 420.1	Klondike	Dawson City	64.1	139.4	Male	9064 \pm 41	WK-32828	Kristensen and Hefner 2011	Antler	
Moose	<i>Alces alces</i>	YT101B	YG 193.1	Klondike								Bone	Skull
Moose	<i>Alces alces</i>	YT101A	YG 193.1	Klondike	Lucky Lady Sulphur Creek	63.7	138.9	Male				Antler	
Moose	<i>Alces alces</i>	YT121	YG 190.1	Klondike	Laskey Creek	63.7	138.7	Male				Antler	
Moose	<i>Alces alces</i>	YT122	YG 162.46	Klondike	Thistle Creek	63.1	139.3	Male	1363 \pm 35; 1197 \pm 27	AA103820; OxA22267	This work; Meiri et al. 2014	Antler	

TABLE 2. Cervid antler collagen bulk stable carbon and nitrogen isotope measurements, C and N elemental data (%), and preservation parameters. Average values for duplicate or triplicate analyses are indicated in bold font.

Species	Institution ID	Locality	¹⁴ C date	Base circumference (cm)	Length of sampling edge (cm)	Complete?	Semi-major axis (cm)	Semi-minor axis (cm)	Log (area)	Location	$\delta^{13}\text{C}_{\text{Coll}}$ ‰; VPDB	$\delta^{15}\text{N}_{\text{Coll}}$ ‰; AIR	%C	%N	Atomic C/N	% Yield
Caribou	YG 381.52	Qikiqtaruk	Post-bomb	9	39	Yes	1.5	1.3	0.8	0 cm	-22.3	+0.9	40.8	14.4	3.3	26.3
										10 cm	-22.5	+1.1	40.7	14.5	3.3	29.6
										20 cm	-22.4	+1.5	39.6	14.2	3.3	31.9
										30 cm	-22.3	+1.7	41.5	14.8	3.3	27.8
										Mean	-22.4	+1.3	40.7	14.5	3.3	28.9
Tip – Base	-0.1	+0.8														
Caribou	YG 109.9	Klondike		7	34	Near complete	1.0	0.9	0.5	0 cm	-18.0	+3.7	38.7	13.6	3.3	7.2
										10 cm	-18.6	+3.7	34.5	12.0	3.3	4.5
										20 cm	-18.5	+3.6	38.3	13.5	3.3	21.6
										Mean	-18.4	+3.7	37.2	13.0	3.3	11.1
										Tip – Base	-0.5	-0.1				
Caribou	YG 404.657	Klondike	>41,100	12	70	Near complete	1.8	1.6	1.0	0 cm	-18.5	+1.8	39.1	13.8	3.3	20.6
										10 cm	-18.7	+1.8	36.3	12.7	3.3	15.2
										20 cm	-18.9	+1.8	38.1	13.3	3.3	19.2
										30 cm	-19.1	+1.8	36.4	12.7	3.3	15.1
										40 cm	-19.2	+1.8	37.0	12.9	3.3	13.2
										50 cm	-19.0	+1.9	38.3	13.5	3.3	22.1
										60 cm	-19.2	+1.8	37.6	13.2	3.3	17.1
										70 cm	-19.5	+1.9	37.8	13.4	3.3	22.8
										Mean	-19.0	+1.8	37.6	13.2	3.3	18.2
										Tip – Base	-1.1	0.0				
Caribou	YG 306.468	Klondike	29,570 ± 970	13	53	No	2.3	1.9	1.1	0 cm	-18.4	+4.5	36.6	13.0	3.3	44.2
										10 cm	-18.7	+4.2	39.4	13.9	3.3	12.0
										20 cm	-18.9	+4.2	35.7	12.3	3.4	5.4
										30 cm	-18.8	+4.1	42.3	15.0	3.3	17.8
										40 cm	-18.8	+4.0	37.6	13.2	3.3	10.6
										50 cm	-18.8	+4.1	39.4	13.9	3.3	15.8
										Mean	-18.7	+4.2	38.5	13.5	3.3	17.7
Tip – Base	-0.4	-0.4														
Elk	YG 104.4	Klondike	11,675 ± 45; 11,785 ± 50	22	71	No				0 cm	-19.2	+2.6	37.2	13.4	3.2	15.7
										10 cm	-19.1	+2.4	37.3	13.3	3.3	9.1
										20 cm	-19.2	+2.5	37.2	13.5	3.2	17.1
										30 cm	-19.2	+2.5	38.7	14.0	3.2	15.9
										40 cm	-19.1	+2.5	37.9	13.8	3.2	13.4
										50 cm	-19.1	+2.6	37.3	13.6	3.2	14.1
										60 cm	-19.1	+2.8	37.7	13.7	3.2	15.6
										Mean	-19.2	+2.6	37.6	13.6	3.2	14.4
Tip – Base	0.1	+0.2														

Elk	YG 420.1	Klondike	9064 ± 41	22	43	No	0 cm	-21.1	+2.4	40.1	14.2	3.3	25.3
							10 cm	-21.0	+2.2	39.6	13.8	3.3	12.9
							20 cm	-21.1	+2.1	40.3	14.1	3.3	12.3
							30 cm	-21.2	+2.0	38.7	13.5	3.3	13.4
							Mean	-21.1	+2.2	39.7	13.9	3.3	16.0
Moose	YG 190.1	Klondike		18	60	Yes	Tip – Base	-0.1	-0.4				
							0 cm	-19.9	+1.8	34.8	12.1	3.3	3.9
							10 cm	-19.8	+2.1	39.9	14.0	3.3	17.5
							20 cm	-19.9	+1.5	40.3	14.0	3.3	13.5
							30 cm	-20.2	+1.9	37.0	12.9	3.3	10.8
							40 cm	-20.4	+2.3	36.5	12.7	3.3	7.4
							50 cm	-20.4	+2.5	34.4	11.8	3.4	7.7
Moose	YG 162.46	Klondike	1363 ± 35; 1197 ± 27	20	44	Near complete	Mean	-20.1	+2.0	37.2	12.9	3.4	10.1
							Tip – Base	-0.5	+0.8				
							0 cm	-20.5	+2.4	36.1	12.5	3.4	3.2
							10 cm	-20.7	+2.2	34.6	12.0	3.4	1.4
							20 cm	-20.5	+2.2	36.8	12.9	3.3	10.6
							30 cm	-20.6	+2.2	38.6	13.5	3.3	15.2
							40 cm	-20.8	+2.4	39.5	13.9	3.3	15.5
							Mean	-20.6	+2.3	37.1	13.0	3.3	9.2
Tip – Base	-0.2	0.0											

TABLE 3. Paired bone and antler collagen bulk stable carbon and nitrogen isotope measurements, offset between tissues ($\Delta\text{B} - \text{A}$), C and N elemental data (%), and preservation parameters. “Single” refers to specimens from which only a single sample was taken. Averages for samples analyzed in duplicate are shown in bold font. *Average isotopic and elemental compositions are listed for serially sampled antler YG 381.52.

Species	Institution ID	Stable isotope reference	^{14}C date	Locality	Tissue	Sampling method	$\delta^{13}\text{C}_{\text{Coll}}$ ‰; VPDB	$\delta^{15}\text{N}_{\text{Coll}}$ ‰; AIR	%C	%N	Atomic C/N	% Yield
Caribou	YG 381.52	This work	Post-bomb	Qikiqtaruk	Bone	Single	-21.0	+2.6	34.6	11.9	3.4	18.0
				Suess effect corrected to 1987			-20.2					
Caribou	YG 381.52	This work	Post-bomb	Qikiqtaruk	Antler	Serial*	-22.4	+1.3	40.7	14.5	3.3	28.9
				Suess effect corrected to 1987			-21.6					
						$\Delta(\text{B}-\text{A})$	+1.4	+1.3				
Moose	YG 193.1	Schwartz-Narbonne et al. 2019		Klondike	Bone	Single	-20.0	+3.0	39.9	15.3	3.0	18.4
Moose	YG 193.1	This work		Klondike	Antler	Single	-19.9	+2.1	39.4	14.9	3.1	17.1
						$\Delta(\text{B} - \text{A})$	-0.1	+0.9				

TABLE 4. Bowmanville Zoo animal bone and antler collagen bulk stable carbon and nitrogen isotope measurements, C and N elemental data (%), and preservation parameters. Averages for samples analyzed in duplicate are shown in bold font.

Lab ID	Species	Tissue	$\delta^{13}\text{C}_{\text{Coll}}$ ‰; VPDB	$\delta^{15}\text{N}_{\text{Coll}}$ ‰; VPDB	%C	%N	Atomic C/N	% Yield
BZ13	Caribou	Antler	-19.8	+4.3	36.7	13.1	3.3	7.9
BZ14	Caribou	Antler	-16.1	+5.7	41.3	14.4	3.3	14.7
BZ15	Caribou	Antler	-15.1	+6.7	40.0	13.8	3.4	27.2
BZ16	Caribou	Antler	-21.2	+4.6	41.1	14.2	3.4	13.3
BZ18	Caribou	Antler	-16.8	+6.3	36.5	12.6	3.4	19.3
BZ19a	Ankole bull	Bone	-21.5	+5.9	36.0	12.7	3.3	10.4

in Tables 1–3 alongside previously published results (Kristensen and Heffner 2011; Meiri et al. 2014; Schwartz-Narbonne et al. 2019). Dated samples covered a wide range, from non-finite dates (YG 404.657) to post-Industrial Revolution/post-bomb (YG 381.52).

While the $\delta^{13}\text{C}$ of atmospheric CO_2 varied somewhat during the Pleistocene and pre-Industrial Revolution Holocene, it has decreased substantially since the advent of fossil fuel use, a change commonly referred to as the Suess effect (Long et al. 2005). The $\delta^{13}\text{C}$ of the post-bomb sample (YG 381.52) was therefore corrected, following the formula of Long et al. (2005), to the value for atmospheric $\delta^{13}\text{C}$ before the Industrial Revolution. As samples covered a wide range of Pleistocene and Holocene ages, we followed previous work and corrected to the last glacial maximum as a known point for comparison (Tahmasebi et al. 2018). Because an exact date for this specimen is not known, the midpoint (1987) between the start of the post-bomb time period (1964) and the year of collection (2009) was used to make the correction. In the absence of radiocarbon dates, a Suess correction cannot be made for the smaller shifts in atmospheric CO_2 $\delta^{13}\text{C}$ during the Pleistocene and Holocene. Hence, no

direct comparisons were made among the $\delta^{13}\text{C}$ of the Quaternary samples. Plant $\delta^{15}\text{N}$ baselines are hypothesized to have shifted during the Quaternary (Tahmasebi et al. 2018; Schwartz-Narbonne et al. 2019), and so direct comparisons were also not made among samples for $\delta^{15}\text{N}$.

Results

Information about the origin and age of the paleontological specimens is summarized in Table 1. Bulk antler measurements are listed in Table 2. The surface area of the pedicle (Miller et al. 2013) and the total size of the caribou antlers were used to identify male versus female animals. Bulk stable isotope measurements for paired antler and bone collagen samples from Yukon Territory are listed in Table 3, and bulk measurements for antler (caribou) and bone (Ankole-Watusi bull) collagen from the Bowmanville Zoo are listed in Table 4. Table 5 provides the nitrogen isotope results for amino acid phenylalanine and glutamate from the Bowmanville Zoo collagen samples.

Preservation

All collagen samples were considered well preserved based on preservation parameters

TABLE 5. $\delta^{15}\text{N}_{\text{Phe}}$, $\delta^{15}\text{N}_{\text{Glu}}$ and $\Delta^{15}\text{N}_{\text{Glu-Phe}}$ data. Results for amino acids (Glu and Phe) are averages of triplicate measurements.

Lab ID	Species	Tissue	$\delta^{15}\text{N}_{\text{Phe}}$ ‰; AIR	$\delta^{15}\text{N}_{\text{Glu}}$ ‰; AIR	$\Delta^{15}\text{N}_{\text{Glu-Phe}}$ ‰; AIR	Bulk $\delta^{15}\text{N}$ ‰; AIR
BZ13	Caribou	Antler	+7.6 ± 1.0	+6.5 ± 1.0	-1.2 ± 2.0	+4.3
BZ14	Caribou	Antler	+7.7 ± 0.9	+8.3 ± 1.1	+0.5 ± 2.0	+5.7
BZ15	Caribou	Antler	+7.9 ± 1.3	+7.5 ± 1.1	-0.4 ± 2.4	+6.7
BZ16	Caribou	Antler	+8.0 ± 2.4	+7.5 ± 0.4	-0.5 ± 2.8	+4.6
BZ18	Caribou	Antler	+7.9 ± 0.4	+7.6 ± 0.4	-0.3 ± 0.8	+6.3
BZ19a	Ankole bull	Bone	+6.4 ± 0.7	+9.1 ± 0.6	+2.7 ± 1.3	+5.9

(collagen yield and atomic C/N) (Tables 2–4). In all cases, collagen extraction yields were >1%, atomic C/N ratios were between 2.9 and 3.6, and carbon and nitrogen contents were ≥ 13 wt.% for carbon and ≥ 4.8 wt.% for nitrogen (DeNiro 1985; Ambrose 1990; van Klinken 1999).

Collagen Stable Carbon and Nitrogen Isotope Compositions

Changes in Antler $\delta^{13}\text{C}_{\text{Coll}}$ and $\delta^{15}\text{N}_{\text{Coll}}$.—A small but measurable change in carbon and/or nitrogen isotope compositions was observed along the length of four out of the eight serially sampled antlers (Fig. 4, Table 3). Length of antler was not a determining factor for the presence or absence of isotopic variation. Antlers from two species (moose and caribou) exhibited changes in isotopic composition that were greater than twice instrumental analytical error ($2 \times \pm 0.2\%$ for $\delta^{13}\text{C}$, $2 \times \pm 0.3\%$ for $\delta^{15}\text{N}$). Values of $\delta^{13}\text{C}_{\text{Coll}}$ decreased with antler growth in YG 109.9 (−0.5%; Fig. 4B) and YG 404.657 (−1.1%; Fig. 4C). Values of $\delta^{15}\text{N}_{\text{Coll}}$ increased in YG 381.52 (+0.8%; Fig. 4A) and YG 190.1 (+0.8%; Fig. 4G) with progressive growth of the antler. There were no significant variations in the “along-antler” isotopic compositions for the other four samples (two elk, one caribou, and one moose). Paired bone and antler collagen samples from individual specimens had different isotopic compositions ($n = 2$; *Alces alces* and *Rangifer tarandus*) (Table 3).

$\Delta^{15}\text{N}_{\text{Glu-Phe}}$

The $\delta^{15}\text{N}$ of amino acids Phe and Glu were measured in several samples of caribou antler and one sample of dewclaw bone from an Ankole-Watusi bull. Both species are ruminants and originated from the Bowmanville Zoo (Table 5). The turnover rate of dewclaw bone, a vestigial digit, is unknown. These results are illustrated in Figure 5, together with data from the literature for bone, tooth dentin, and tooth cement collagen for mammalian herbivore species (Naito et al. 2010, 2013; Styring et al. 2010; Schwartz-Narbonne et al. 2015; Kendall et al. 2017). The expanded dataset includes wild cervid bone collagen (Styring et al. 2010; Naito et al. 2013) and tooth dentin collagen from pastured adult cattle (Kendall et al. 2017).

Discussion

Are Antler and Bone Metabolically Equivalent?

The impact of metabolic processes on antler tissue formation must be understood before interpreting antler collagen carbon and nitrogen isotope signals. We first needed to establish if the isotopic offset between diet and collagen is the same for both bone and antler or the offset for antler includes an additional shift caused by its faster growth rate than bone. If the offset is the same, it can be reasonably assumed that the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of both bone and antler tissues correlate with diet or seasonally determined physiological changes (i.e., starvation), rather than additional tissue-specific metabolic processes. Prior studies discuss rapid growth rate as a potential control on the $\delta^{15}\text{N}$ of collagenous tissues (Trueman et al. 2005; Waters-Rist and Katzenberg 2010; Madgwick et al. 2013; Osborne 2017); accordingly, nitrogen isotopes were the focus of our investigation.

Our examination of caribou antler collagen $\Delta^{15}\text{N}_{\text{Glu-Phe}}$ used animals from a zoo population that consumed grass present in their enclosure as well as feed (e.g., grain) provided as a supplement. The antler bulk collagen $\delta^{13}\text{C}$ suggests a mixed C_3/C_4 diet (unpublished data), which is consistent with both C_3 and C_4 plants growing in Ontario. Use of a captive population allowed us to control for physiological sources of ^{15}N enrichment in antlers other than rapid growth, as these animals did not experience dietary stress or nutritional deficiencies. Only one bone specimen was available from Bowmanville Zoo, and it was from a non-cervid species not indigenous to North America (Ankole-Watusi bull). We therefore supplemented our results with published values for bone and tooth collagen $\Delta^{15}\text{N}_{\text{Glu-Phe}}$, which included ruminant herbivores (cow, buffalo, sheep, steenbok, deer, elk, and auroch), as well as terrestrial carnivores and high-marine-protein consumers (Naito et al. 2010, 2013; Styring et al. 2010; Schwartz-Narbonne et al. 2015; Kendall et al. 2017) (Fig. 5). The use of $\Delta^{15}\text{N}_{\text{Glu-Phe}}$ largely corrects for the variability in isotopic composition among diets, thus allowing for more direct comparisons to be made. The Bowmanville Zoo caribou antler collagen $\Delta^{15}\text{N}_{\text{Glu-Phe}}$ values lie within the range




FIGURE 4. Antler $\delta^{13}\text{C}_{\text{Coll}}$ (left) and $\delta^{15}\text{N}_{\text{Coll}}$ (right): A, female caribou antler, post-bomb, YG 381.52; B, female caribou antler, undated, YG 109.9; C, male caribou antler, >41,100 ^{14}C BP, YG 404.657; D, male caribou antler, 29,570 \pm 970 ^{14}C BP, YG 306.468; E, male elk antler, 11,675 \pm 45 ^{14}C BP, YG 104.4; F, male elk antler, 9064 ^{14}C BP, YG 420.1; G, male moose antler, undated, YG 190.1; and H, moose antler, 1363 \pm 35/1197 \pm 27 ^{14}C BP, YG 162.46. Black stars indicate specimens from Qikiqtaruk (Herschel Island). Variations greater than twice experimental error were measured for $\delta^{13}\text{C}_{\text{Coll}}$ in YG 109.9 and YG 404.657 and for $\delta^{15}\text{N}_{\text{Coll}}$ in YG 381.52 and YG 190.1.

previously reported for ruminant collagenous tissues (Fig. 5). While variation in herbivore diets may have increased the range of

$\Delta^{15}\text{N}_{\text{Glu-Phe}}$ (Kendall et al. 2019), all the herbivore collagenous tissues (antler, bone, and teeth) have lower $\Delta^{15}\text{N}_{\text{Glu-Phe}}$ than carnivorous

Do Intra-antler Isotopic Differences Reflect Changes in Diet?

The small sample set, covering a range of species, locations, and time intervals within the Quaternary renders it difficult to discern a consistent pattern in the stable carbon and nitrogen isotope variations. Consideration of these results plus prior isotopic studies of antlers and other temporally variable tissues, however, suggests some general observations:

1. Antler collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ can be indicative of trophic niche width, where less isotopic variation equates to less available dietary choice or participation in specialist feeding behavior. Both Darr and Hewitt's (2008) controlled feeding study of white-tailed deer and Finstad and Kielland's (2011) study of captive versus free-ranging caribou populations observed small, or no, isotopic changes within antlers from individuals with reduced dietary variability. Based on this concept, the four antlers that displayed no notable change in $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ (Pleistocene male caribou, Fig. 4D, YG 306.468; Pleistocene–Holocene boundary elk, Fig. 4E, YG 104.4; early Holocene elk, Fig. 4F, YG 420.1; an undated moose antler, Fig. 4H, YG 162.46) may reflect dietary specialists. Dietary specialization may be caused by a number of factors such as interspecies or intraspecies competition for resources, individual dietary preferences, small range sizes, or lack of varied forage types within a large habitat. Contemporary climate change is causing terrestrial herbivores to migrate northward to track their optimal habitat, with adverse effects for species unable to colonize new habitats (Gilg et al. 2012). Small range sizes and dependence on specialized diets may therefore be risk factors for cervid populations adapting to climate change.
2. Changes in antler collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ can produce results generally consistent with expected dietary shifts in $\delta^{13}\text{C}$ and/or $\delta^{15}\text{N}$. Finstad and Kielland (2011) observed an increase in antler $\delta^{15}\text{N}$ over time consistent with increased consumption of graminoids and forbs during the transition of spring to summer, and a similar increase in antler $\delta^{15}\text{N}$ was observed for one female

caribou antler in this study (post-bomb, Fig. 4A, YG 381.52). In a modern Alaskan caribou herd, low $\delta^{15}\text{N}_{\text{Coll}}$ in antler deposited in early spring correlated with winter dietary stress, likely because these individuals sought out high-protein shrubs rather than grasses to consume (Finstad and Kielland 2011). Unfortunately, without a larger dataset with strong temporal control, similar comparisons between Pleistocene antler $\delta^{15}\text{N}_{\text{Coll}}$ cannot be made here. Only limited research has been done correlating antler $\delta^{13}\text{C}$ to diet; however, temporally based shifts in caribou $\delta^{13}\text{C}$ of hoof keratin and tooth dentin collagen (Barnett 1994; Drucker et al. 2001, 2012) have been observed and attributed in part to a reduction in consumption of ^{13}C -rich lichen as other, more protein-rich forage became available in spring and summer. This shift is consistent with one female caribou antler (undated, Fig. 4B, YG 109.9), and one male caribou antler (non-finite date, Fig. 4C, YG 404.657) measured here. Differences in caribou $\delta^{13}\text{C}$ of hair keratin grown in summer were related to differing summer availability of nonmycorrhizal vascular plants between sites (Zhao et al. 2019); again, a larger sample set with strong temporal control is needed for similar comparisons within a Pleistocene dataset. Similarly, studies of serially sampled moose hooves found higher $\delta^{15}\text{N}$ in layers grown during the summer, related to increased consumption of aquatic plants (see Fig. 2) (Kielland 2001; Tischler et al. 2019). This is consistent with the pattern observed here for the undated moose antler (Fig. 4G, YG 190.1). Seasonal shifts in Pleistocene caribou diet, as observed in specimen YG 404.657, would be consistent with the hypothesis of Schwartz-Narbonne et al. (2019) that the Pleistocene dietary niche of caribou changed seasonally. However, a seasonal dietary shift was not observed in Pleistocene caribou antler specimen YG 306.468. Further study of dated caribou antlers is necessary to test this hypothesis.

Results of previous studies are currently insufficient to explain the entirety of our intra-antler variability, particularly where the

expected signal response in one isotope was observed, but the corresponding signal in the other isotope was not. For example, an increasing $\delta^{15}\text{N}$ consistent with summer consumption of ^{15}N -rich aquatic plants was observed for the undated moose antler (Fig. 4G, YG 190.1); however, the expected corresponding increase in $\delta^{13}\text{C}$ from consumption of ^{13}C -rich aquatic plants did not occur. Such decoupling of the stable carbon and nitrogen isotope signals has been observed previously for cervid tissues (Drucker et al. 2001, 2010; Walter and Leslie 2009) and merits further study. A more comprehensive model incorporating a range of controls, including forage isotopic composition, protein content, and nutritional stress, may be needed to fully deconvolute antler isotopic signals. Our work is insufficient to resolve these questions, but it highlights the value of such an investigative approach to paleoecological and modern ecological studies.

Conclusion

Antler collagen $\Delta^{15}\text{N}_{\text{Glu-Phe}}$ is consistent with ruminant bone and tooth collagen $\Delta^{15}\text{N}_{\text{Glu-Phe}}$, suggesting that rapid growth is not a primary isotopic control on antler $\delta^{15}\text{N}$ and that the metabolic processes affecting antlers are similar to those occurring in other collagenous tissues. We therefore suggest two applications of antler $\delta^{13}\text{C}_{\text{Coll}}$ and $\delta^{15}\text{N}_{\text{Coll}}$ measurements. First, antlers can be used to assess changes in diet during the spring and early summer. Antlers with lower isotopic variability likely represent dietary/habitat specializations or limited dietary choice during the spring and summer months. Second, antler $\delta^{13}\text{C}_{\text{Coll}}$ and $\delta^{15}\text{N}_{\text{Coll}}$ reflect environmental and dietary signals acquired during spring and early summer, while bone collagen records an average multi-year signal. Hence, antler isotopic signals can be used to compare spring and early summer conditions to yearly conditions in ancient ecosystems. Additional studies employing larger sample sizes of both wild and captive cervid populations are needed to better understand the causes of variability in antler stable carbon and nitrogen isotope compositions before this tissue can be fully integrated in the isotopic toolbox for paleoenvironmental studies.

Acknowledgments

We thank K. Law and L. Huang for technical advice. Samples were collected with the permission of the Yukon Government Palaeontology Program and the Bowmanville Zoo. We are grateful to the placer gold mining community and the Tr'ondëk Hwëch'in First Nation for their continued support and partnership with our research in the Klondike goldfields region. We thank the people of the Inuvialuit Settlement Region, along with the Herschel/Qikiqtaruk Territorial Park Management Committee, which includes Yukon Parks, the Wildlife Management Advisory Committee, the Aklavik Hunters and Trappers Committee, the Aklavik Community Corporation, and the Yukon Government Cultural Services Branch, for support of research at Qikiqtaruk/Herschel Island. We thank Bowmanville Zoo staff for information on animal diets. G. Hodgins and the staff at the National Science Foundation Arizona AMS Laboratory assisted with the radiocarbon dating. K. Allan drew the artwork for Figures 1 and 3. We thank the editor and two anonymous reviewers for constructive comments that improved the article. Funding was provided by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant (F.J.L.), and an NSERC Canada Graduate Scholarship, an NSERC Postgraduate Scholarship, and an Ontario Graduate Scholarship (all to R.S.-N.). Laboratory infrastructure was funded by the Canada Foundation for Innovation (F.J.L.) and the Ontario Research Fund (F.J.L.). Additional time for research was funded through the Canada Research Chairs program (F.J.L.). The Western Graduate Research Fund (R.S.-N.) funded one radiocarbon date. This is Laboratory for Stable Isotope Science Contribution no. 340.

Literature Cited

- Ambrose, S. H. 1990. Preparation and characterization of bone and tooth collagen for isotopic analysis. *Journal of Archaeological Science* 17:431–451.
- Banks, W. J., and J. W. Newbrey. 1983. Light microscopic studies of the ossification process in developing antlers. Pp. 231–260 in R. D. Brown, ed. *Antler development in Cervidae*. Caesar Kleburg Wildlife Research Institute, Kingsville, Tex.
- Barnett, B. A. 1994. Carbon and nitrogen isotope ratios of caribou tissues, vascular plants, and lichens from northern Alaska. M.Sc. thesis, University of Alaska, Fairbanks.

- Barreto-Curiel, F., U. Focken, L. R. D'Abramo, and M. T. Viana. 2017. Metabolism of *Seriola lalandi* during starvation as revealed by fatty acid analysis and compound-specific analysis of stable isotopes within amino acids. *PLoS ONE* 12:e0170124.
- Ben-David, M., E. Shochat, and L. G. Adams. 2001. Utility of stable isotope analysis in studying foraging ecology of herbivores: examples from moose and caribou. *Alces* 37:421–434.
- Blinnikov, M. S., B. V. Gaglioti, D. A. Walker, M. J. Wooller, and G. D. Zazula. 2011. Pleistocene graminoid-dominated ecosystems in the Arctic. *Quaternary Science Reviews* 30:2906–2929.
- Bocherens, H. 2015. Isotopic tracking of large carnivore palaeoecology in the Mammoth Steppe. *Quaternary Science Reviews* 117:42–71.
- Chapman, D. I. 1975. Antlers—bones of contention. *Mammal Review* 5:121–172.
- Chikaraishi, Y., Y. Kashiyama, N. O. Ogawa, H. Kitazato, and N. Ohkouchi. 2007. Metabolic control of nitrogen isotope composition of amino acids in macroalgae and gastropods: implications for aquatic food web studies. *Marine Ecology Progress Series* 342:85–90.
- Chikaraishi, Y., N. O. Ogawa, Y. Kashiyama, Y. Takano, H. Suga, A. Tomitani, H. Miyashita, H. Kitazato, and N. Ohkouchi. 2009. Determination of aquatic food-web structure based on compound-specific nitrogen isotope composition of amino acids. *Limnology and Oceanography: Methods* 2003:740–750.
- Chikaraishi, Y., S. A. Steffan, Y. Takano, and N. Ohkouchi. 2015. Diet quality influences isotopic discrimination among amino acids in an aquatic vertebrate. *Ecology and Evolution* 5:2048–2059.
- Clarke, M. 2012. Flat Creek Wetlands: preliminary habitat assessment. Yukon Fish and Wildlife Branch Report SR-12-02. <http://www.env.gov.yk.ca>.
- Coplen, T. B., W. A. Brand, M. Gehre, M. Gröning, and H. Meijer. 2006. New guidelines for $\delta^{13}\text{C}$ measurements. *Analytical Chemistry* 78:2439–2441. <http://pubs.acs.org/doi/abs/10.1021/ac052027c>.
- Cormie, A. B., and H. P. Schwarcz. 1994. Stable isotopes of nitrogen and carbon of North American white-tailed deer and implications for paleodietary and other food web studies. *Palaeogeography, Palaeoclimatology, Palaeoecology* 107:227–241.
- Corr, L. T., R. Berstan, and R. Evershed. 2007a. Development of N-acetyl methyl ester derivatives for the determination of $\delta^{13}\text{C}$ values of amino acids using gas chromatography-combustion-isotope ratio mass spectrometry. *Analytical Chemistry* 79:9082–9090.
- Corr, L. T., R. Berstan, and R. P. Evershed. 2007b. Optimisation of derivatisation procedures for the determination of $\delta^{13}\text{C}$ values of amino acids by gas chromatography/combustion/isotope ratio mass spectrometry. *Rapid Communications in Mass Spectrometry* 21:3759–3771.
- Dalerum, F., and A. Angerbjörn. 2005. Resolving temporal variation in vertebrate diets using naturally occurring stable isotopes. *Oecologia* 144:647–658.
- Darr, R. L., and D. G. Hewitt. 2008. Stable isotope trophic shifts in white-tailed deer. *Journal of Wildlife Management* 72:1525–1531.
- Davis, M., and S. Pineda-Munoz. 2016. The temporal scale of diet and dietary proxies. *Ecology and Evolution* 6:1883–1897.
- DeNiro, M. J. 1985. Postmortem preservation and alteration of in vivo bone collagen isotope ratios in relation to palaeodietary reconstruction. *Nature* 317:806–809.
- Drucker, D. G., H. Bocherens, A. Pike-Tay, and A. Mariotti. 2001. Isotopic tracking of seasonal dietary change in dentine collagen: preliminary data from modern caribou. *Comptes Rendus de l'Académie Des Sciences, series IIA* 333:303–309.
- Drucker, D. G., K. A. Hobson, J. P. Ouellet, and R. Courtois. 2010. Influence of forage preferences and habitat use on ^{13}C and ^{15}N abundance in wild caribou (*Rangifer tarandus caribou*) and moose (*Alces alces*) from Canada. *Isotopes in Environmental and Health Studies* 46:107–121.
- Drucker, D. G., K. A. Hobson, S. C. Münzel, and A. Pike-Tay. 2012. Intra-individual variation in stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotopes in mandibles of modern caribou of Qamanirjuaq (*Rangifer tarandus groenlandicus*) and Banks Island (*Rangifer tarandus pearyi*): implications for tracing seasonal and temporal changes in diet. *International Journal of Osteoarchaeology* 22:494–504.
- Ernakovich, J. G., K. A. Hopping, A. B. Berdanier, R. T. Simpson, E. J. Kachergis, H. Steltzer, and M. D. Wallenstein. 2014. Predicted responses of Arctic and alpine ecosystems to altered seasonality under climate change. *Global Change Biology* 20:3256–3269.
- Finstad, G. L., and K. Kielland. 2011. Landscape variation in the diet and productivity of reindeer in Alaska based on stable isotope analyses. *Arctic, Antarctic, and Alpine Research* 43:543–554.
- Gaglioti, B. V., B. M. Barnes, G. D. Zazula, A. B. Beaudoin, and M. J. Wooller. 2011. Late Pleistocene paleoecology of Arctic ground squirrel (*Urocitellus parryi*) caches and nests from interior Alaska's Mammoth Steppe ecosystem, USA. *Quaternary Research* 76:373–382.
- Gilg, O., K. M. Kovacs, J. Aars, J. Fort, G. Gauthier, D. Grémillet, R. A. Ims, H. Meltøfte, J. Moreau, E. Post, and N. M. Schmidt. 2012. Climate change and the ecology and evolution of Arctic vertebrates. *Annals of the New York Academy of Sciences* 1249:166–190.
- Gomez, S., A. J. Garcia, S. Luna, U. Kierdorf, H. Kierdorf, L. Gallego, and T. Landete-Castillejos. 2013. Labeling studies on cortical bone formation in the antlers of red deer (*Cervus elaphus*). *Bone* 52:506–515.
- Guthrie, R. D. 2001. Origin and causes of the Mammoth Steppe: a story of cloud cover, woolly mammal tooth pits, buckles, and inside-out Beringia. *Quaternary Science Reviews* 20:549–574.
- Ishikawa, N. F., F. Hayashi, Y. Sasaki, Y. Chikaraishi, and N. Ohkouchi. 2017. Trophic discrimination factor of nitrogen isotopes within amino acids in the dobsonfly *Prothermes grandis* (Megaloptera: Corydalidae) larvae in a controlled feeding experiment. *Ecology and Evolution* 7:1674–1679.
- Jay, M., and M. P. Richards. 2007. British Iron Age diet: stable isotopes and other evidence. *Proceedings of the Prehistoric Society* 73:169–190.
- Kelly, J. F. 2000. Stable isotopes of carbon and nitrogen in the study of avian and mammalian trophic ecology. *Canadian Journal of Zoology* 78:1–27.
- Kendall, I. P., M. R. Lee, and R. P. Evershed. 2017. The effect of trophic level on individual amino acid $\delta^{15}\text{N}$ values in a terrestrial ruminant food web. *STAR: Science & Technology of Archaeological Research* 3:135–145.
- Kendall, I. P., P. Woodward, J. P. Clark, A. K. Styring, J. V. Hanna, and R. P. Evershed. 2019. Compound-specific $\delta^{15}\text{N}$ values express differences in amino acid metabolism in plants of varying lignin content. *Phytochemistry* 161:130–138.
- Kielland, K. 2001. Stable isotope signatures of moose in relation to seasonal forage composition: a hypothesis. *Alces* 37:329–337.
- Kierdorf, U., S. Flohr, S. Gomez, T. Landete-Castillejos, and H. Kierdorf. 2013. The structure of pedicle and hard antler bone in the European roe deer (*Capreolus capreolus*): a light microscope and backscattered electron imaging study. *Journal of Anatomy* 223:364–384.
- Koch, P. L., M. L. Fogel, and N. Tuross. 1994. Tracing the diets of fossil animals using stable isotopes. Pp. 63–92 in K. Lajtha and R. H. Michener, eds. *Stable isotopes in ecology and environmental science*. Blackwell Scientific, Malden, Mass.
- Kohn, M. J. 2010. Carbon isotope compositions of terrestrial C_3 plants as indicators of (paleo)ecology and (paleo)climate. *Proceedings of the National Academy of Sciences USA* 107:19691–19695.
- Kristensen, D. K., E. Kristensen, M. C. Forchhammer, A. Michelsen, and N. M. Schmidt. 2011. Arctic herbivore diet can be inferred from stable carbon and nitrogen isotopes in C_3 plants, faeces, and wool. *Canadian Journal of Zoology* 89:892–899.
- Kristensen, T., and T. Heffner. 2011. Heritage resources impact assessment of the Yukon Hospital Corporation proposed Dawson City Multi-Level Care Facility conducted under permit 10-02ASR. Government of Yukon, Heritage Resources Unit, Whitehorse, YT.

- Kuitemans, M., J. van der Plicht, D. G. Drucker, T. Van Kolfshoten, S. W. Palstra, and H. Bocherens. 2015. Carbon and nitrogen stable isotopes of well-preserved middle Pleistocene bone collagen from Schöningen (Germany) and their paleoecological implications. *Journal of Human Evolution* 89:105–113.
- Leader-Williams, N. 1988. *Reindeer on south Georgia*. Cambridge: Cambridge University Press.
- Long, E. S., R. A. Sweitzer, D. R. Diefenbach, and M. Ben-David. 2005. Controlling for anthropogenically induced atmospheric variation in stable carbon isotope studies. *Oecologia* 146:148–156.
- Longin, R. 1971. New method of collagen extraction for radiocarbon dating. *Nature* 230:241–242.
- Madgwick, R., N. Sykes, H. Miller, R. Symmons, J. Morris, and A., Lamb. 2013. Fallow deer (*Dama dama dama*) management in Roman south-east Britain. *Archaeological and Anthropological Sciences* 5:111–122.
- Malcolm, D. 2013. Climate change impacts, adaptation, and the technology interface. Pp. 113–127 in G. H. Gjørv, D. Bazely, M. Goloviznina, and A. Tanentzap, eds. *Environmental and human security in the Arctic*. Routledge, London.
- Mariotti, A. 1983. Atmospheric nitrogen is a reliable standard for natural ^{15}N abundance. *Measurements*. *Nature* 303:685–687.
- McMahon, K. W., and M. D. McCarthy. 2016. Embracing variability in amino acid $\delta^{15}\text{N}$ fractionation: mechanisms, implications, and applications for trophic ecology. *Ecosphere* 7:e01511.
- McMillan, J. F. 1953. Some feeding habits of moose in Yellowstone Park. *Ecology* 34:102–110.
- Meiri, M., A. M. Lister, M. J. Collins, N. Tuross, T. Goebel, S. Blockley, G. D. Zazula, N. van Doorn, R. D. Guthrie, G. G. Boeskorov, and G. F. Baryshnikov. 2014. Faunal record identifies Bering Isthmus conditions as constraint to end-Pleistocene migration to the New World. *Proceedings of the Royal Society of London B* 281:20132167.
- Miller, H., R. F. Carden, J. Evans, A. Lamb, R. Madgwick, D. Osborne, R. Symmons, and N. Sykes. 2016. Dead or alive? Investigating long-distance transport of live fallow deer and their body parts in antiquity. *Environmental Archaeology* 21:246–259.
- Miller, J. H., P. Druckenmiller, and V. Bahn. 2013. Antlers on the Arctic refuge: capturing multi-generational patterns of calving ground use from bones on the landscape. *Proceedings of the Royal Society of London B* 280:20130275.
- Milligan, H. 2008. *Aquatic and terrestrial foraging by a subarctic herbivore: the beaver*. M.Sc. thesis. McGill University, Montreal.
- Nadelhoffer, K., G. Shaver, B. Fry, A. Giblin, L. Johnson, and R. McKane. 1996. ^{15}N Natural abundances and N use by tundra plants. *Oecologia* 107:386–394.
- Naito, Y. I., N. V. Honch, Y. Chikaraishi, N. Ohkouchi, and M. Yoneda. 2010. Quantitative evaluation of marine protein contribution in ancient diets based on nitrogen isotope ratios of individual amino acids in bone collagen: an investigation at the Kitakogane Jomon site. *American Journal of Physical Anthropology* 143:31–40.
- Naito, Y. I., Y. Chikaraishi, N. Ohkouchi, D. G. Drucker, and H. Bocherens. 2013. Nitrogen isotopic composition of collagen amino acids as an indicator of aquatic resource consumption: insights from Mesolithic and Epipalaeolithic archaeological sites in France. *World Archaeology* 45:338–359.
- O'Connell, T. C., and M. J. Collins. 2018. Comment on “Ecological Niche of Neanderthals from Spy Cave Revealed by Nitrogen Isotopes of Individual Amino Acids in Collagen” [J. Hum. Evol. 93 (2016) 82–90]. *Journal of Human Evolution* 117:53–55.
- O'Leary, M. H. 1981. Carbon isotope fractionation in plants. *Phytochemistry* 20:553–567.
- Osborne, D. 2017. Imports and isotopes: a modern baseline study for interpreting Iron Age and Roman trade in fallow deer antlers. *Papers from the Institute of Archaeology* 27. <https://doi.org/10.5334/pia-482>.
- Parker, K. L., P. S. Barboza, and T. R. Stephenson. 2005. Protein conservation in female caribou (*Rangifer tarandus*): effects of decreasing diet quality during winter. *Journal of Mammalogy* 86:610–622.
- Parnell, A., and A. Jackson. 2013. SIAR: stable isotope analysis in R. <https://cran.r-project.org/package=siar>.
- Plint, T., F. J. Longstaffe, and G. Zazula. 2019. Giant beaver palaeoecology inferred from stable isotopes. *Scientific Reports* 9:7179.
- Qi, H., T. B. Coplen, H. Geilmann, W. A. Brand, and J. K. Böhlke. 2003. Two new organic reference materials for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ measurements and a new value for the $\delta^{13}\text{C}$ of NBS 22 oil. *Rapid Communications in Mass Spectrometry* 17:2483–2487.
- Schwartz-Narbonne, R., F. J. Longstaffe, J. Z. Metcalfe, and G. Zazula. 2015. Solving the woolly mammoth conundrum: amino acid ^{15}N -enrichment suggests a distinct forage or habitat. *Scientific Reports* 5:9791.
- Schwartz-Narbonne, R., F. J. Longstaffe, K. J. Kardynal, P. Druckenmiller, K. A. Hobson, C. N. Jass, J. Z. Metcalfe, and G. Zazula. 2019. Reframing the mammoth steppe: insights from analysis of isotopic niches. *Quaternary Science Reviews* 215:1–21.
- Stevens, R. E., and T. C. O'Connell. 2016. Red deer bone and antler collagen are not isotopically equivalent in carbon and nitrogen. *Rapid Communications in Mass Spectrometry* 30:1969–1984.
- Styring, A. K., J. C. Sealy, and Evershed, R. P. 2010. Resolving the bulk $\delta^{15}\text{N}$ values of ancient human and animal bone collagen via compound-specific nitrogen isotope analysis of constituent amino acids. *Geochimica et Cosmochimica Acta* 74:241–251.
- Tahmasebi, F., F. J. Longstaffe, G. Zazula, and B. Bennett. 2017. Nitrogen and carbon isotopic dynamics of subarctic soils and plants in southern Yukon Territory and its implications for paleoecological and paleodietary studies. *PLoS ONE* 12:e0183016.
- Tahmasebi, F., F. J. Longstaffe, and G. Zazula. 2018. Nitrogen isotopes suggest a change in nitrogen dynamics between the late Pleistocene and modern time in Yukon, Canada. *PLoS ONE* 13:e0192713.
- Tieszen, L. L. 1991. Natural variations in the carbon isotope values of plants: implications for archaeology, ecology, and paleoecology. *Journal of Archaeological Science* 18:227–248.
- Tischler, K. B. 2004. *Aquatic plant nutritional quality and contribution to moose diet at Isle Royale National Park*. M.Sc. thesis. Michigan Technological University, Houghton, Mich.
- Tischler, K. B., W. J. Severud, R. O. Peterson, and J. K. Bump. 2019. Aquatic macrophytes are seasonally important dietary resources for moose. *Diversity* 11:209.
- Trueman, C. N., R. A. McGill, and P. H. Guyard. 2005. The effect of growth rate on tissue-diet isotopic spacing in rapidly growing animals. An experimental study with Atlantic salmon (*Salmo salar*). *Rapid Communications in Mass Spectrometry* 19:3239–3247.
- Van Ballenberghe, V. 1983. Growth and development of moose antlers in Alaska. Pp. 37–48 in R.D. Brown, ed. *Antler development in Cervidae*. Caesar Kleberg Wildlife Research Institute, Kingsville, Tex.
- Van Klinken, G. J. 1999. Bone collagen quality indicators for palaeodietary and radiocarbon measurements. *Journal of Archaeological Science* 26:687–695.
- Walter, W. D., and D. M. Leslie. 2009. Stable isotope ratio analysis to differentiate temporal diets of a free-ranging herbivore. *Rapid Communications in Mass Spectrometry* 23:2190–2194. <https://doi.org/10.1002/rcm.4135>.
- Wang, Y., and M. J. Wooller. 2006. The stable isotopic (C and N) composition of modern plants and lichens from northern Iceland: with ecological and paleoenvironmental implications. *Jökull* 56:27–38.
- Waters-Rist, A. L., and M. A. Katzenberg. 2010. The effect of growth on stable nitrogen isotope ratios in subadult bone collagen. *International Journal of Osteoarchaeology* 20:172–191.
- Zhao, L. Z., A. S. Colman, R. J. Irvine, S. R. Karlsen, G. Olack, and E. A. Hobbie. 2019. Isotope ecology detects fine-scale variation in Svalbard reindeer diet: implications for monitoring herbivory in the changing Arctic. *Polar Biology* 42:793–805.