

Violence, mental health and nutritional status in pregnant women: the Araraquara Cohort Study

Leonardo Domingos Biagio¹, Delanjathan Devakumar², Paula Louro da Silva¹, Rossana Verónica Mendoza López³, Perla Pizzi Argentato¹, Liania Alves Luzia¹ and Patrícia Helen Carvalho Rondó¹

¹Nutrition Department, School of Public Health, University of São Paulo (USP), Brazil;

²Institute for Global Health, University College London (UCL), United Kingdom; ³São Paulo State Cancer Institute (ICESP), Brazil

Correspondence to: Patrícia Helen Carvalho Rondó, Nutrition Department, School of Public Health, University of São Paulo, Avenida Dr Arnaldo 715, São Paulo, CEP-01246-904, Brazil. Phone: + 55 11 3061 7867; e-mail: phcrondo@usp.br

Short title: Violence and nutritional status in pregnancy

Acknowledgments: We would like to acknowledge Prof. Walter Manso Figueiredo and Prof. Angela Aparecida Costa for assistance in data collection, and the Health Secretary of Araraquara for allowing us to carry out the research in the city



This is an Accepted Manuscript for Public Health Nutrition. This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI 10.1017/S1368980024002295

Public Health Nutrition is published by Cambridge University Press on behalf of The Nutrition Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Financial Support: This study was supported by the São Paulo Research Foundation (FAPESP) (grant number 2015/03333-6) and Medical Research Council (MRC-UK). LDB received an MPH scholarship from the Coordination for the Improvement of Higher Education Personnel (CAPES) and PPA had a scholarship from FAPESP (2018/17824-0).

Conflict of Interest: The authors have no conflict of interest to declare.

Authorship: PHCR designed the study protocol, secured funding, and facilitated data collection; LDB, PLS and PPA collected the data; PHCR and LAL coordinated the fieldwork; PHCR, RVML, and LDB were responsible for data management and performing analyses; PHCR, RVML, DD, and LDB interpreted the data; LDB, PHCR, RVML and PPA revised each draft for important intellectual content. All authors read and approved the final version of the manuscript.

Ethical Standards Disclosure: Ethical Standards Disclosure: The study was conducted in accordance with the guidelines laid down in the Declaration of Helsinki. All procedures were approved by the Ethics Committee of the School of Public Health, University of São Paulo, (No. 59787216.2.0000.5421). Written informed consent was obtained from all participants.

Abstract

Objective: To investigate the relationship between violence and the nutritional status of pregnant women, and whether mental health could be a mediator in this relationship.

Design: Cross-sectional study. Violence and mental health status were investigated using the following questionnaires: World Health Organization Violence Against Women (WHO-VAW), Abuse Assessment Screen (AAS), Patient Health Questionnaire (PHQ-9), and General Health Questionnaire (GHQ). Demographic, socioeconomic, obstetric, and lifestyle factors (smoking/alcohol consumption) were also investigated. The nutritional status of the women was assessed by the body mass index.

Setting: Data were collected from February 2021 to August 2022 in Araraquara city, Brazil.

Participants: Four hundred pregnant women recruited at 34 health units and the municipal maternity hospital.

Results: Experience of violence was reported by 52.2% of the women and psychological violence in the last 12 months was the most prevalent type of domestic violence (19.5%). Approximately 43% of the women showed mental health changes and 59.7% had a risk of major depression. Women with mental health changes had an increased risk (OR=2.34) of obesity. Psychological violence in the last 12 months was associated with obesity ($p=0.01$) when mediated by mental health changes. The mediation effect was significant ($\beta=0.708$; 95%BCa CI=0.004-1.460), with mental health changes mediating 46.1% of the relationship between psychological violence and obesity.

Conclusions: The relationship between psychological violence and obesity during pregnancy was mediated by changes in mental health. This original study shows that nutritional status is not limited to biological factors and highlights the importance of social, mental, and psychological factors.

Keywords: nutritional status, pregnant women, violence, mental health.

Introduction

According to the World Health Organization (WHO), 30% of women worldwide suffered physical and sexual violence between 2000 and 2018⁽¹⁾. Pregnancy can be a particularly vulnerable period for domestic violence (DV) because of the physical and emotional changes that occur during this period. Moreover, social and economic status may increase this risk in minority groups. A review conducted in 2021 that involved 7 systematic reviews and 5 meta-analyses reported a prevalence of psychological and physical DV during pregnancy ranging from 1.8 to 67.4% and from 1.6 to 78%, respectively. The reviews involved high, but mainly low-and middle-income countries and showed a higher risk of violence in the latter countries due to social factors⁽²⁾. In a meta-analysis involving data from Brazil, James *et al.*⁽³⁾ found an overall prevalence of violence of 63.40% (40.40% of psychological DV, 12.35% of physical DV, and 4.55% of sexual DV) in pregnant women.

Pregnant women exposed to DV are more susceptible to mental health changes such as depression, anxiety, emotional distress, suicidal ideation, and lack of self-efficacy and self-esteem^(4,5). A recent systematic review and metanalysis⁽⁶⁾ assessed the prevalence of DV and the magnitude of the association between violence and mental health outcomes in 250,599 women who were primarily from high-income countries. Lifetime psychological violence was reported by 32.8% of the women, while physical and sexual violence was reported by 18.3% and 9.6%, respectively. Perinatal women were more likely to have experienced lifetime physical DV (28.8%), than the other women investigated (women in the community and help-seeking women). The metanalysis suggested an association between DV and an increased risk of all mental health outcomes including depression, posttraumatic stress disorder, and suicidality.

Violence against women has a negative impact not only on mental health, but also on physical health, considering that stress can increase the concentrations of glucocorticoids and insulin influencing eating behavior by either increasing or decreasing food intake, and affecting metabolism⁽⁷⁻⁹⁾. Therefore, some mental health changes, particularly stress, have a bidirectional effect on food intake, and women may thus be either under- or overweight.

Chronic stress has been found to cause weight loss by increasing the metabolic rate and energy expenditure^(7,8). Conversely, long-term chronic stress can trigger prolonged hyperactivation of the hypothalamus-pituitary-adrenal axis, with further increases in abdominal fat retention and triglyceride concentrations, that lead to overweight and obesity^(7,10,11).

In a cohort of 206 pregnant women, Farias *et al.*⁽¹²⁾ observed that persistent depressive symptoms resulted in a 2.40 times higher risk of insufficient gestational weight gain. However, the authors found no significant association of generalized anxiety disorder or suicide risk with weight gain. In a Thai study⁽¹³⁾, pregnant women with high scores on depression and anxiety scales had a higher body mass index (BMI) during pregnancy. The results are similar to the findings of two meta-analyses in which severe symptoms of depression were associated with excessive gestational weight gain^(14,15).

Studies on non-pregnant women suggest that DV can influence their nutritional status in different ways, but the results are limited and inconsistent⁽¹⁶⁻¹⁹⁾. To our knowledge, there is only one large epidemiological study that has evaluated the association between prenatal depressive symptoms, maternal nutrition, DV and social support among pregnant women in rural Ethiopia. The authors observed that several factors were associated with depression, including DV, mid-upper arm circumference less than 23 cm, and anemia⁽²⁰⁾. Furthermore, maternal mental health changes, either directly or as a result of DV, may compromise their offspring earlier in utero, predisposing to low birth weight or premature labor^(21,22), and even to later malnutrition^(23,24).

Apparently, the number of studies evaluating the impact of DV and mental health changes on maternal nutritional status is much smaller than the number of studies evaluating the impact of DV and mental health changes on adverse outcomes in offspring.

Therefore, this study investigated the relationship between violence and the nutritional status of pregnant women, and whether mental health could be a mediator in this relationship.

Materials and methods

This cross-sectional study investigated violence and mental health in 400 pregnant women. The study is part of a large epidemiological prospective cohort - the Araraquara Cohort Study - that assessed the relationship between maternal and offspring adiposity during the fetal, neonatal, and infant periods. Pregnant women with a gestational age ≤ 26 weeks were included in this study.

Data were collected from February 2021 to August 2022. The women were selected by trained interviewers at the 34 health units and at the municipal maternity hospital of Araraquara city, São Paulo, Brazil. The participants answered a questionnaire previously used in pregnant women⁽²⁵⁾, which contained data on demographic, socioeconomic, obstetric, and

lifestyle factors (smoking/alcohol consumption). The women attended the Serviço Especial de Saúde de Araraquara (SESA) for assessment of their nutritional status and their gestational age by ultrasonography. They were also asked about violence and mental health using standardized questionnaires that were applied online.

Before the start of any conversation about violence, the interviewers were instructed to ask whether the participant felt safe to speak and she could answer simply with “yes” or “no”. If she did not feel safe, a better time for the interview was suggested. In addition, the interviewers asked the participant if she was alone, ensuring that the abuser was not in the same room with her. The WHO ethical and safety recommendations for research on violence against women were followed (WHO, 2001).

Questionnaires for assessing violence

World Health Organization Violence Against Women (WHO-VAW)⁽²⁶⁾

The instrument consists of 13 items that evaluate marital violence (psychological, physical, and sexual) in the last 12 months. This questionnaire has been used in several international and national studies, including those involving Brazilian pregnant women^(27,28).

Abuse Assessment Screen (AAS)⁽²⁹⁾

The AAS is a specific instrument used to assess violence during pregnancy and has already been applied to Brazilian pregnant women⁽²⁹⁾. The instrument contains five questions designed to identify the frequency and severity of events, the locations of bodily harm suffered over a given period, and the profile of the perpetrator. The questions address lifetime experiences of abuse, physical violence in the past year, physical violence during pregnancy, sexual abuse in the past 12 months, and fear of a current partner or someone close.

Questionnaires for assessing mental health

General Health Questionnaire (GHQ)⁽³⁰⁾

The GHQ measures psychological well-being in the adult population, including pregnant women⁽²⁵⁾. The questionnaire addresses common mental health problems/domains of depression, anxiety, somatic symptoms, and social withdrawal. It is simple to administer, easy to complete and score, and widely used in many studies⁽³¹⁾. The version used consisted of 12 questions. The GHQ score was classified as low (0-3) and high (≥ 4)⁽³⁰⁾.

Patient Health Questionnaire (PHQ-9)⁽³²⁾

The PHQ-9 aims to assess the risk of major depression and has been validated for the Brazilian population and applied to pregnant women^(33,34). It consists of nine symptoms and the frequency of each symptom is rated over the past two weeks on a Likert scale from 0 to 3. The total score ranges from 0 to 27 and a score ≥ 10 is defined as the presence of depression.

Assessment of nutritional status

The nutritional status of the pregnant women was assessed based on BMI at gestational age ≤ 26 weeks. Weight was evaluated by electrical bioimpedance using the Inbody 370 equipment (Biospace®, Seoul, Korea). Height was measured with a Seca 206 stadiometer (Seca®, Hamburg, Germany). The Atalah reference curve was used to classify the pregnant women according to their BMI⁽³⁵⁾.

Statistical analysis

Descriptive analysis was performed to obtain the frequency and percentage of the independent, dependent, and confounding variables. The Shapiro-Wilk test was applied to analyze the normality of the data. The chi-squared test and Fisher's exact test were used to evaluate the associations among the variables. Multivariate logistic regression models were built to estimate the relative risk ratios between violence and the nutritional status of the pregnant women. The models were adjusted for the following confounding variables: age, parity, and gestational age. The results were reported as odds ratio (OR) and 95% confidence interval (CI) and a p-value < 0.05 was considered statistically significant. To assess the effect of mental health as a mediator in the relationship between violence (X) and nutritional status (Y), we used the PROCESS macro for SPSS. This tool is widely employed to estimate direct and indirect effects in single and multiple mediator (parallel and serial) models. In this study, the mediation process involved only one mediating variable (mental health changes) and was therefore simple mediation- a type of mediation that occurs when one variable (M) mediates the effect of X on Y⁽³⁶⁻³⁸⁾. Statistical analysis was performed using the SPSS® software (version 20; Chicago, USA).

Results

A total of 471 pregnant women were recruited for the study. Seventy-one (15.1%) women refused to participate due to the following reasons: lack of interest and lack of time. Therefore, 400 women completed the study.

Table 1 shows the demographic, socioeconomic, obstetric, nutritional, and lifestyle characteristics, violence at any point in their lifetime, DV in the last 12 months, and mental health of the pregnant women studied. Most pregnant women (87%) were 20-30 years of age and were non-white (52%), were married or had a common-law union (87.5%), had ≤ 12 years of schooling (81%), and had a per capita income of 0.5-1 Brazilian minimum wage (56.3%). Less than half of the women had an intimate partner as head of the household (40.5%), and most of them did not consume alcohol, did not smoke and did not use illicit drugs during pregnancy. Almost 43% of the women were primiparous and 78.7% were at 14-26 weeks of gestation. The majority of pregnant women had a BMI above the adequate range (56.6%), with similar percentages of overweight and obesity, while a minority were underweight (12.8%).

Experience of violence at any point in their lifetime was reported by 52.2% of the pregnant women. Psychological violence was the most prevalent type of DV in the last 12 months (19.5%) followed by physical (7.5%) and sexual violence (2.7%). The GHQ scores revealed mental health changes in 42.7% of the pregnant women and 59.7% had a risk of major depression based on PHQ scores.

Table 2 shows the associations of the nutritional status of pregnant women with demographic, socioeconomic, obstetric, and lifestyle characteristics, DV at any point in their lifetime, type of violence in the last 12 months, and mental health changes. There were statistically significant associations between the current nutritional status of the pregnant women and mental health assessed by the GHQ ($p=0.021$), age ($p=0.001$), and parity ($p\leq 0.001$).

Table 3 shows the results obtained with four multivariate regression models that assessed the associations of the nutritional status category of pregnant women with mental health and DV in the last 12 months, controlling for the confounders age, parity and gestational age. Mental health changes assessed by the GHQ were associated with an increased risk of the pregnant woman being obese (OR=2.34) compared to women with a normal BMI ($p=0.002$). Similarly, women with altered PHQ scores were more likely to be obese (OR=1.81). Psychological and physical violence in the last 12 months were not

associated with any of the nutritional status categories. As expected, the confounder age <20 years was positively associated with the nutritional status of the pregnant women (undernutrition) in all models.

Table 4 shows the results obtained with multivariate regression models that assessed the associations of the nutritional status category of pregnant women with DV in the last 12 months, mediated by mental health and controlling for the confounders age, parity and gestational age. In model 5, psychological violence in the last 12 months was associated with obesity ($p=0.01$). We sought to investigate the extent to which mental health changes mediated the relationship between psychological violence and obesity. The mediation effect (indirect effect) was significant ($\beta=0.708$; 95% BCa CI=0.004-1.460). Obese pregnant women had a 2.82 times higher chance of having mental health changes than women with a normal BMI ($p\leq 0.001$). As shown in Figure 1, mental health changes mediated 46.1% of the relationship between psychological violence and obesity. According to model 6, physical violence was not associated with any of the nutritional status categories. There was a 2.36 times higher chance of a pregnant woman with obesity having mental health changes compared to women with a normal BMI ($p=0.002$).

Discussion

The objective of this study was to assess the impact of DV and mental health on the nutritional status of pregnant women. Moreover, we sought to investigate the extent to which mental health changes mediated the relationship between psychological violence and nutritional status of the women.

Psychological violence, assessed by the WHO-VAW, was more prevalent (19.5%) than physical (7.5%) and sexual (2.7%) violence among the pregnant women investigated, in agreement with previous studies^(3,39,40). However, studies that assess DV are sensitive to methodological factors. In addition, the perception of abuse as violence is low, especially sexual violence⁽⁴¹⁾.

According to GHQ scores, 42.7% of the women had mental health changes, a higher percentage than that reported in a cohort study conducted by our group, with a similar population, in which the frequency of high GHQ scores (≥ 4) was 30.4% among adult pregnant women in the first trimester of gestation⁽⁴²⁾. A recent study conducted in Nigeria⁽⁴³⁾ detected high GHQ scores among 51.8% of the 991 pregnant women investigated compared to 28.6% of non-pregnant women. The PHQ-9 identified that 59.7% of the women included

in our study had depression. A Chinese study reported a lower prevalence of depression (36.12%) among 681 pregnant women who attended various obstetrics and gynecology clinics in Changzhou city, located in the economically developed eastern part of China⁽⁴⁴⁾.

More than half of the pregnant women (56.6%) were overweight/obese. Nutritional status was significantly associated with GHQ and PHQ-9 scores. Multivariate regression models, controlling for the confounders age, parity and gestational age, showed that overweight/obese pregnant women had a 2.32 times higher chance of having mental changes and a 1.81 times higher chance of having major depression than women with a normal BMI. Similarly, studies conducted in different regions of the world^(13–15,45) found associations of excessive BMI during pregnancy with depression and anxiety. There was no association between undernutrition and GHQ or PHQ-9 scores.

Psychological and physical violence in the last 12 months were not associated with the nutritional status of pregnant women, in multivariate regression analysis controlling for the confounders age, parity and gestational age. However, in the final multivariate regression model that included mental health (GHQ scores), psychological violence was significantly associated with overweight/obesity. The inclusion of GHQ in the model showed that mental health changes acted as a mediator between psychological violence and the nutritional status of pregnant women. Obese pregnant women had a 2.82 times higher chance of having mental health changes than women with a normal BMI. Mental health changes mediated 46.1% of the relationship between psychological violence and obesity. To our knowledge there are no studies that assessed the relationship between DV and the nutritional status of pregnant women, considering mental health as a mediator between these two factors.

Physical violence in the last 12 months, on the other hand, was not associated with nutritional status of the pregnant women, in contrast to the findings of another Brazilian study⁽¹⁸⁾ involving non-pregnant women, that revealed an inverse association of physical violence with BMI. The results showed that physical intimate partner violence was negatively associated with BMI between the 25th and 85th percentiles, corresponding to 22.9 and 31.2kg/m², respectively. Adhikari et al.⁽¹⁹⁾ did not find any association between underweight and DV among non-pregnant Nepalese women, but overweight was associated with physical violence, including severe physical violence.

We found no association between sexual violence and the nutritional status of pregnant women, probably due to the low prevalence of sexual violence (2.7%). It was therefore not possible to include the data in the regression analysis. The experience of

physical and sexual violence was a marker for increased risk of chronic undernutrition in women of reproductive age in Bangladesh⁽⁴⁶⁾, which was more strongly accentuated among the poorest women. In a study of Nepalese women, emotional and physical violence was significantly associated with the risk of being overweight and obese, while sexual violence alone was associated with a BMI compatible with underweight⁽⁴⁷⁾.

One limitation of our study is its cross-sectional design. Due to the COVID-19 pandemic, we were unable to follow up the women throughout pregnancy. In addition, some studies reported an increase in violence and mental health changes during the COVID-19 pandemic^(48,49), a fact that may have influenced the results of this study. Another limitation of the study is that we did not use a specific questionnaire to assess anxiety, only the GHQ, that measures mental health changes in general. A strength of the study is its originality in evaluating the relationship between DV, mental health changes, and nutritional status in pregnant women since the studies available so far only involve non-pregnant women. Furthermore, there are no studies assessing the role of mental changes as a mediator in the relationship between violence and the nutritional status of pregnant women.

Prospective cohort studies are necessary to assess violence and mental health during pregnancy. Moreover, it would be important to evaluate the dietary pattern of pregnant women exposed to violence in view of its impact not only for the mothers but also for their offspring. A recent study showed that physical violence by an intimate partner, was associated with greater adherence to a dietary pattern of lower nutritional quality among pregnant women from a Brazilian cohort⁽⁵⁰⁾.

Conclusion

The relationship between psychological violence in the last 12 months and overweight/obesity, during pregnancy was mediated by mental health changes. Mental health changes mediated 46.1% of the relationship between psychological violence and overweight/obesity. The findings of this original study show that nutritional status is not limited to biological factors and highlights the importance of mental and psychological factors.

It is very important to detect situations of violence and mental health changes in pregnancy, because of the intense physiological changes during this period of life, which probably render the pregnant women more vulnerable to these events. Furthermore, we

emphasize the need for multiple antenatal screenings to detect DV and mental health changes at an early stage of pregnancy, in order to develop safe and appropriate interventions that will prevent complications not only for the mother, but also for their offspring. Finally, cohort studies assessing violence and mental health status of women throughout pregnancy are needed to evaluate the impact of these events during the different trimesters of pregnancy on maternal and child health.

References

1. WHO. Violence against women prevalence estimates, 2018: global, regional and national prevalence estimates for intimate partner violence against women and global and regional prevalence estimates for non-partner sexual violence against women 2021. 1–112 p. Available from: <https://www.who.int/publications/i/item/9789240022256> (accessed April 2024).
2. Román-Gálvez RM, Martín-Peláez S, Martínez-Galiano JM *et al.* (2021) Prevalence of intimate partner violence in pregnancy: An umbrella review. *Int J Environ Res Public Health* **18**, 1–13.
3. James L, Brody D & Hamilton Z (2013) Risk factors for domestic violence during pregnancy: A meta-analytic review. *Violence Vict* **28**, 359–380.
4. Halim N, Beard J, Mesic A *et al.* (2018) Intimate partner violence during pregnancy and perinatal mental disorders in low and lower middle income countries: A systematic review of literature, 1990–2017. *Clin Psychol Rev* **66**, 117–135.
5. Alipour Z, Kheirabadi GR, Kazemi A *et al.* (2018) The most important risk factors affecting mental health during pregnancy: A systematic review. *East Mediterr Health J* **24**, 549–559.
6. White SJ, Sin J, Sweeney A *et al.* (2024) Global prevalence and mental health outcomes of intimate partner violence among women: A systematic review and meta-analysis. *Trauma Violence Abuse* **25**, 494–511.
7. Björntorp P (2001) Do stress reactions cause abdominal obesity and comorbidities? *Obes Rev* **2**, 73–86.
8. Singh M (2014) Mood, food and obesity. *Front Psychol* **5**, 1–35.
9. Tank AW & Wong DL (2015) Peripheral and central effects of circulating catecholamines. *Compr Physiol* **5**, 1–15.

10. Kyrou I, Chrousos GP & Tsigos C (2006) Stress, visceral obesity, and metabolic complications. *Ann N Y Acad Sci* **1083**, 77–110.
11. Avila C, Holloway AC, Hahn MK *et al.* (2015) An Overview of Links Between Obesity and Mental Health. *Curr Obes Rep* **4**, 303–310.
12. Farias DR, Carrilho TRB, Freitas-Costa NC *et al.* (2021) Maternal mental health and gestational weight gain in a Brazilian Cohort. *Sci Rep* **11**, 1–11.
13. Roomruangwong C, Kanchanatawan B, Sirivichayakul S *et al.* (2017) High incidence of body image dissatisfaction in pregnancy and the postnatal period: Associations with depression, anxiety, body mass index and weight gain during pregnancy. *Sex Reprod Healthc* **13**, 103–109.
14. Kapadia MZ, Gaston A, Van Blyderveen S *et al.* (2015) Psychological antecedents of excess gestational weight gain: A systematic review. *BMC Pregnancy Childbirth* **15**, 1–30.
15. Hartley E, McPhie S, Skouteris H *et al.* (2015) Psychosocial risk factors for excessive gestational weight gain: A systematic review. *Women Birth* **28**, e99–e109.
16. Issah AN, Yeboah D, Kpordoxah MR *et al.* (2022) Association between exposure to intimate partner violence and the nutritional status of women and children in Nigeria. *PLoS One* **17**, 1–17.
17. Ferdos J & Rahman MM (2018) Exposure to intimate partner violence and malnutrition among young adult Bangladeshi women: Cross-sectional study of a nationally representative sample. *Cad Saude Publica* **34**, 1–11.
18. Ferreira M de F, Moraes CL de, Reichenheim ME *et al.* (2015) Effect of physical intimate partner violence on body mass index in low-income adult women. *Cad Saude Publica* **31**, 161–172.
19. Adhikari RP, Yogi S, Acharya A *et al.* (2020) Intimate partner violence and nutritional status among nepalese women: An investigation of associations. *BMC Womens Health* **20**, 1–11.
20. Woldetensay YK, Belachew T, Biesalski HK *et al.* (2018) The role of nutrition, intimate partner violence and social support in prenatal depressive symptoms in rural Ethiopia: Community based birth cohort study. *BMC Pregnancy Childbirth* **18**, 1–10.
21. Voit FAC, Kajantie E, Lemola S *et al.* (2022) Maternal mental health and adverse birth outcomes. *PLoS One* **17**, 1–18.
22. Kpordoxah MR, Adiak AA, Issah AN *et al.* (2024) Magnitude of self-reported intimate

- partner violence against pregnant women in Ghana's northern region and its association with low birth weight. *BMC Pregnancy Childbirth* **24**, 1–10.
23. Hassan BK, Werneck GL & Hasselmann MH (2016) Maternal mental health and nutritional status of six-month-old infants. *Rev Saude Publica* **50**, 1–9.
 24. Memiah P, Bond T, Opanga Y *et al.* (2020) Neonatal, infant, and child mortality among women exposed to intimate partner violence in East Africa: A multi-country analysis. *BMC Womens Health* **20**, 1–16.
 25. Rondó PHC, Ferreira RF, Nogueira F *et al.* (2003) Maternal psychological stress and distress as predictors of low birth weight, prematurity and intrauterine growth retardation. *Eur J Clin Nutr* **57**, 266–272.
 26. Garcia-Moreno C, Jansen HAFM, Ellsberg M *et al.* (2006) Prevalence of intimate partner violence: Findings from the WHO multi-country study on women's health and domestic violence. *Lancet* **368**, 1260–1269.
 27. Ribeiro MRC, Silva AAM, De Britto E Alves MTSS *et al.* (2017) Effects of socioeconomic status and social support on violence against pregnant women: A structural equation modeling analysis. *PLoS One* **12**, 1–16.
 28. Ribeiro MRC, De Britto E Alves MTSS, Batista RFL *et al.* (2014) Confirmatory factor analysis of the WHO violence against women instrument in pregnant women: Results from the BRISA prenatal cohort. *PLoS One* **9**, 1–16.
 29. Reichenheim ME, Moraes CL & Hasselmann MH (2000) Equivalência semântica da versão em português do instrumento Abuse Assessment Screen para rastrear a violência contra a mulher grávida. *Rev Saude Publica* **34**, 610–616.
 30. Jesus Mari J De & Williams P (1985) A Comparison of the Validity of Two Psychiatric Screening Questionnaires (GHQ-12 and SRQ-20) In Brazil, Using Relative Operating Characteristic (ROC) Analysis. *Psychol Med* **15**, 651–659.
 31. Jackson C (2006) The General Health Questionnaire. *Occup. Med. (Chic. Ill)*. **57**, 79–79.
 32. Kroenke K, Spitzer RL & Williams JBW (2001) The PHQ-9: Validity of a brief depression severity measure. *J Gen Intern Med* **16**, 606–613.
 33. Faisal-Cury A, Levy RB, Azeredo CM *et al.* (2021) Prevalence and associated risk factors of prenatal depression underdiagnosis: A population-based study. *Int J Gynaecol Obstet* **153**, 469–475.
 34. de Pádua Borges R, de Azevedo Jacob Reichelt A, de Brito A *et al.* (2021) Impact of

- the COVID-19 pandemic on mental health of pregnant women with diabetes mellitus and hypertension. *Rev Assoc Med Bras* **67**, 1268–1273.
35. Atalah SE, Castillo LC, Castro SR *et al.* (1997) Propuesta de un nuevo estándar de evaluación nutricional en embarazadas. *Rev Med Chil* **125**, 1429–1436.
 36. Preacher KJ & Hayes AF (2008) Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods* **40**, 879–891.
 37. Preacher KJ & Hayes AF (2004) SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behav Res Methods Instrum Comput* **36**, 717–731.
 38. Kristofferzon ML, Engström M & Nilsson A (2018) Coping mediates the relationship between sense of coherence and mental quality of life in patients with chronic illness: a cross-sectional study. *Qual Life Res* **27**, 1855–1863.
 39. Shafiei S, Chegeni M, Afrashteh S *et al.* (2022) Prevalence of violence in Iranian pregnant women: A systematic review and metanalysis. *Matern Child Health J* **26**, 1983-2019.
 40. Devries KM, Kishor S, Johnson H *et al.* (2010) Intimate partner violence during pregnancy: Analysis of prevalence data from 19 countries. *Reprod Health Matters* **18**, 158–170.
 41. Ellsberg M, Heise L, Penã R *et al.* (2001) Researching domestic violence against women: methodological and ethical considerations. *Stud Fam Plann* **32**, 1–16.
 42. Rondó PHC, Souza MR, Moraes F *et al.* (2004) Relationship between nutritional and psychological status of pregnant adolescents and non-adolescents in Brazil. *J Health Popul Nutr* **22**, 34–45.
 43. Bello OO, Bella-Awusah TT, Adebayo AM *et al.* (2023) Psychiatric morbidity among pregnant and non pregnant women in Ibadan, Nigeria. *J Obstet Gynaecol* **43**(1), 2205503.
 44. Wang L, Yang N, Zhou H *et al.* (2022) Pregnant Women’s Anxiety and Depression Symptoms and Influence Factors in the COVID-19 Pandemic in Changzhou, China. *Front Psychol* **13**, 1–8.
 45. Holton S, Fisher J, Nguyen H *et al.* (2019) Pre-pregnancy body mass index and the risk of antenatal depression and anxiety. *Women Birth* **32**, e508–e514.

46. Rahman M, Nakamura K, Seino K *et al.* (2013) Intimate partner violence and chronic undernutrition among married Bangladeshi women of reproductive age: Are the poor uniquely disadvantaged? *Eur J Clin Nutr* **67**, 301–307.
47. Chaudhary A, Nakarmi J & Goodman A (2022) Association between intimate partner violence and nutritional status of married Nepalese women. *Glob Health Res Policy* **7**, 1–14.
48. Hildersley R, Easter A, Bakolis I *et al.* (2022) Changes in the identification and management of mental health and domestic abuse among pregnant women during the COVID-19 lockdown: regression discontinuity study. *BJPsych Open* **8**, 1–12.
49. Botteman H, Vahdat B, Jouault C *et al.* (2022) Becoming a mother during COVID-19 pandemic: how to protect maternal mental health against stress factors. *Front Psychiatry* **12**, 1–11.
50. Vaz JDS, Souza MEMDC, Valério ID, Silva MTD *et al.* (2022) Physical intimate partner violence and dietary patterns in pregnancy: a Brazilian cohort. *Cien Saude Colet* **27**, 1317–1326.

Table 1. Demographic, socioeconomic, obstetric, nutritional and lifestyle characteristics, violence at any point in their lifetime, DV in the last 12 months, and mental health of the pregnant women.

Characteristics	n (%)
Age (years)	
<20	30 (7.5)
20-30	348 (87)
>30	22 (5.5)
Race	
White	192 (48)
Non-white	208 (52)
Marital status	
Married/common-law union	350 (87.5)
Single/divorced/widow	50 (12.5)
Education (years of schooling)	
<12	324 (81)
≥12	76 (19)
Per capita income(BMW)*	
<0.5	148 (37)
0.5-1	225 (56.3)
>1	27 (6.8)
Head of household	
Intimate partner	162 (40.5)
Pregnant woman	73 (18.2)
Intimate partner and pregnant woman	88 (22)
Other	77 (19.2)
Alcohol consumption	
Yes	53 (13.2)
No	347 (86.7)
Smoking	
Yes	26 (6.5)
No	374 (93.5)

Use of illicit drugs	
Yes	8 (2)
No	392 (98)
Parity	
0	171 (42,8)
1-2	190 (47,5)
≥3	39 (9,8)
Gestational age (weeks)	
≤13	85 (21.6)
14-26	315 (78.7)
BMI (kg/m ²)	
Undernutrition	51 (12.8)
Normal	123 (30.8)
Overweight	113 (28.3)
Obesity	113 (28,3)
AAS (violence at any point in their lifetime)	209 (52.2)
WHO-VAW (DV in the last 12 months)	
Psychological	78 (19.5)
Physical	30 (7.5)
Sexual	11 (2.7)
GHQ	
0-3	229 (57.3)
≥4	171 (42.7)
PHQ	
0-9	161 (40.3)
≥10	239 (59.7)

*One BMW (Brazilian Minimum Wage) = US\$ 267.00; DV = domestic violence; BMI = body mass index; AAS=Assessment Abuse Screen; WHO-VAW=World Health Organization-Violence Against Women; GHQ = General Health Questionnaire; PHQ = Patient Health Questionnaire.

Table 2. Associations of the nutritional status of pregnant women with demographic, socioeconomic, obstetric and lifestyle characteristics, DV at any point in their lifetime, type of violence in the last 12 months, and mental health changes.

Variable	Undernutrition	Normal	Overweight	Obesity	p-value*
Age (years)					
<20	9 (17.6%)	8 (6.5%)	6 (5.3%)	1 (0,9%)	<0.001
20-30	34 (66.6%)	73 (59.3%)	64 (56.6%)	59 (52.3%)	
>30	8 (15.7%)	42 (34.1%)	43 (13.3%)	53 (46.9%)	
Per capita income(BMW)***					
<0.5	12 (23.5%)	45 (36.6%)	53 (46.9%)	38 (33.6%)	0.265
0.5-1	35 (68.6%)	70 (56.9%)	50 (44.2%)	70 (61.9%)	
>1	4 (7.8%)	8 (6.5%)	10 (8.8%)	5 (6.8%)	
Race					
White	26 (51%)	62 (50.4%)	51 (45.1%)	53 (46.9%)	0.827
Non-white [†]	25 (49%)	61 (49.6%)	62 (54.9%)	60 (53.1%)	
Marital status					
Married/common-law union	43 (84.3%)	107 (87%)	99 (87.6%)	101 (89.4%)	0.833
Single/divorced/widow	8 (15.7%)	16 (13%)	14 (12.4%)	12 (10.6%)	
Head of household					
Intimate partner	18 (35.3%)	48 (39%)	48 (42.5%)	48 (42.5%)	0.354

Pregnant woman	11 (21.6%)	24 (19.5%)	22 (19.5%)	16 (14.2%)	
Intimate partner and pregnant woman	6 (11.8%)	29 (23.6%)	24 (21.2%)	29 (25.7%)	
Other	16 (31.4%)	22 (17.9%)	19 (16.8%)	20 (17.7%)	
Education (years of schooling)					
<12	39 (76.5%)	99 (80.5%)	94(83.2%)	92 (81.4%)	0.786
≥12	12 (23.4%)	24 (19.5%)	19 (16.8%)	21 (18.6%)	
Smoking					
Yes	45 (88.2%)	119 (96.7%)	107 (92.9%)	105 (92.9%)	0.205
No	6 (11.8%)	4 (3.3%)	8 (7.1%)	8 (7.1%)	
Use of illicit drugs					
Yes	49 (96.1%)	121 (98.4%)	109 (96.5%)	113 (100%)	0.125**
No	2 (3.9%)	2 (1.6%)	4 (3.5%)	0 (0%)	
Alcohol consumption					
Yes	40 (78.4%)	102 (82.9%)	84 (74.3%)	90 (79.6%)	0.447
No	11 (21.6%)	21 (17.1%)	29 (25.7%)	23 (20.4%)	
Gestational age (weeks)					
≤13	10 (19.6%)	31 (25.2%)	24 (21.2%)	20 (17.7%)	0.556
14-26	41 (80.4%)	92 (74.8%)	89 (78.8%)	93 (82.3%)	

Parity					
0	33 (64.7%)	58 (47.2%)	37 (32.7%)	43 (38.1%)	<0.001
1-2	14 (27.5%)	55 (44.7%)	64 (56.6%)	57 (50.4%)	
≥3	4 (7.8%)	10 (8.1%)	12 (10.6%)	13 (11.5%)	
GHQ					
0-3	30 (58.8%)	83 (67.5%)	62 (54.9%)	54 (47.8%)	0.021
≥4	21 (41.2%)	40 (32.5%)	51 (45.1%)	59 (52.2%)	
PHQ					
0-9	17 (33.3%)	61 (49.6%)	42 (37.2%)	41 (46.3%)	0.083
≥10	34 (66.7%)	62 (50.4%)	71 (62.8%)	72 (63.7%)	
WHO-VAW- Psychological violence					
No	39 (76.5%)	97 (78.9%)	90 (79.6%)	96 (85%)	0.528
Yes	12 (23.5%)	26 (21.1%)	23 (20.4%)	17 (15%)	
WHO-VAW- Physical violence					
No	45 (88.2%)	114 (92.7%)	106 (93.8%)	105 (92.9%)	0.648
Yes	6 (11.8%)	9 (7.3%)	7 (6.2%)	8 (7.1%)	
WHO-VAW- Sexual violence					
No	49 (96.1%)	118	113 (100%)	109	0.107**

		(95.9%)		(96.5%)	
Yes	2 (3.9%)	5 (4.1%)	0 (0%)	4 (3.5%)	
AAS (violence at any point in their lifetime)					
No	24 (47.1%)	59 (48%)	53 (46.9%)	55 (48.7%)	0.994
Yes	27 (52.9%)	64 (52%)	60 (53.1%)	58 (51.3%)	

*Chi-squared test. **Fisher's exact test. ***One BMW (Brazilian Minimum Wage) = US\$ 267.00. [†]Non-white = black, mixed race, yellow, and indigenous people. Abbreviations: AAS = Assessment Abuse Screen; WHO-VAW = World Health Organization-Violence Against Women; GHQ = General Health Questionnaire; PHQ = Patient Health Questionnaire.

Table 3. Associations of nutritional status category of the pregnant women with mental health and DV in the last 12 months.

Variable	Undernutrition			Overweight			Obesity		
	OR (95% CI)	P		OR (95% CI)	p		OR (95% CI)	P	
MODEL 1									
GHQ									
≥4	1.50 (0.748 – 3.022)	0.253		1.61 (0.936 – 2.767)	0.085		2.34 (1.359 – 4.051)	0.002	
0-3	Ref.			Ref.			Ref.		
Age (years)									
>30	2.12 (0.858 – 5.287)	0.103		0.94 (0.530- 1.690)	0.851		0.58 (0.327- 1.041)	0.068	
<20	4.89 (1.318 – 18.189)	0.018		1.09 (0.327- 3.661)	0.884		0.10 (0.012 - 0.929)	0.043	
20-30	Ref.			Ref.			Ref.		
Parity									
0	0.56 (0.150 – 2.133)	0.400		1.03 (0.406- 2.614)	0.950		0.98 (0.388- 2.511)	0.979	
≥ 3	0.97 (0.261 – 3.634)	0.968		0.59 (0.220- 1.599)	0.302		0.97 (0.366- 2.612)	0.964	
1-2	Ref.			Ref.			Ref.		
Gestational age (weeks)									
0-13	1.36 (0.600 – 3.121)	0.455		1.25 (0.677- 2.338)	0.468		1.53 (0.801- 2.946)	0.197	
14-26	Ref.			Ref.			Ref.		
MODEL 2									
PHQ									
>9	1.78 (0.887-3.599)	0.105		1.63 (0.959 – 2.773)	0.071		1.81 (1.062- 3.103)	0.029	

0-9	Ref.		Ref.		Ref.			
Age (years)								
>30	2.07 (0.832-5.148)	0.117	0.94 (1.684)	(0.527- 0.842	0.61	0.61	(0.343- 0.093	
<20	4.52 (1.206-16.97)	0.025	1.03 (3.470)	(0.307- 0.960	0.10	0.10	(0.012- 0.038	
20-30	Ref.		Ref.		Ref.			
Parity								
0	0.57 (0.152-2.180)	0.417	1.04 (2.654)	(0.412- 0.926	0.985	0.985	(0.390- 0.974	
1-2	0.96 (0.260-3.615)	0.962	0.578 (1.55)	(0.215- 0.278	0.886	0.886	(0.335- 0.807	
≥3	Ref.		Ref.		Ref.			
Gestational age (weeks)								
0-13	1.29 (0.568-2.69)	0.536	1.21 (2.265)	(0.654- 0.535	1.46	1.46	(0.767- 0.247	
14-26	Ref.		Ref.		Ref.			
MODEL 3								
WHO-VAW-psychological violence								
Yes	0.77 (0.347-1.747)	0.544	1.14 (2.186)	(0.602- 0.676	1.58	1.58	(0.798- 0.187	
No	Ref.		Ref.		Ref.			
Age (years)								
>30	2.22 (0.899-5.523)	0.084	1.03 (0.581-1.83)	0.913	0.70	0.70	(0.399- 0.222	

							1.238)		
<20	5.081 (1.369- 18.848)	0.015	1.11 (0.332- 3.710)	0.865	0.11 (0.013- 0.047	0.969)			
20-30	Ref.		Ref.		Ref.				
Parity									
0	0.55 (0.146-2.089)	0.382	0.99 (0.392- 2.514)	0.987	0.90 (0.357- 0.824	2.270)			
1-2	0.92 (0.246- 3.465)	0.906	0.52 (0.195- 1.413)	0.202	0.74 (0.281- 0.548	1.962)			
≥3	Ref.		Ref.		Ref.				
Gestational age (weeks)									
0-13	1.38 (0.608-3.155)	0.439	1.26 (0.680- 2.338)	0.461	0.74 (0.795- 0.207	2.882)			
14-26	Ref.		Ref.		Ref.				
MODEL 4									
WHO- VAW- physical violence									
Yes	1.60 (0.524-4.907)	0.408	0.82 (0.291- 2.309)	0.707	1.04 (0.383- 0.936	2.836)			
No	Ref.		Ref.		Ref.				
Age (years)									
>30	2.22 (0.897-5.493)	0.085	1.03 (0.581- 1.834)	0.914	0.67 (0.383- 0.170	1.184)			
<20	4.91 (1.32-18.246)	0.017	1.12 (0.337- 3.765)	0.847	0.11 (0.013- 0.048	0.977)			
20-30	Ref.		Ref.		Ref.				
Parity									

0	0.53 (0.141-2.022)	0.355	1.00	0.399-	0.986	0.93	(0.373-	0.889	
			2.549)			2.351)			
1-2	0.88 (0.237-3.278)		0.53	(0.201-	0.215	0.80	(0.308-	0.662	
		0.851	1.436)			2.113)			
≥3	Ref.		Ref.			Ref.			
Gestational									
age									
(weeks)									
0-13	1.353	(0.594-	0.472	1.26	(0.683-	0.454	1.52	(0.801-	0.200
	3.082)			2.347)			2.894)		
14-26	Ref.		Ref.			Ref.			

Multivariate regression models. Confounding factors included in the models: age, parity, and gestational age. Abbreviations: OR= odds ratio; CI= confidence interval; Ref.= Reference; AAS= Assessment Abuse Screen; WHO-VAW= World Health Organization-Violence Against Women; GHQ= General Health Questionnaire; PHQ= Patient Health Questionnaire.

Table 4. Associations of nutritional status category of the pregnant women with DV in the last 12 months, mediated by mental health.

Variable	Undernutrition		Overweight			Obesity	
	OR (95% CI)	P	OR (95% CI)	p	OR (95% CI)	P	
WHO-VAW- psychological violence							
Yes	0.90 (0.380)	0.825	1.45 (0.730-2.915)	0.286	2.42 (1.160-5.08)	<0.001	
No	Ref.		Ref.		Ref.		
GHQ							
≥4	1.42 (0.672 - 3.002)	0.358	1.75 (0.980 - 3.115)	0.058	2.82 (1.586 - 5.022)	<0.001	
0-3	Ref.		Ref.		Ref.		
Age (years)							
>30	2.12 (0.852-5.289)	0.106	0.95 (0.536-1.718)	0.889	0.61 (0.340-1.093)	0.096	
<20	4.93 (1.326-18.385)	0.017	1.06 (0.319-3.586)	0.914	0.10 (0.012-0.889)	0.039	
20-30	Ref.		Ref.		Ref.		
Parity							
0	0.55 (0.148-2.114)	0.391	1.00 (0.396-2.558)	0.990	0.92 (0.362-2.374)	0.873	
1-2	0.97 (0.259-3.678)	0.972	0.56 (0.209-1.540)	0.266	0.87 (0.323-2.354)	0.787	
≥3	Ref.		Ref.		Ref.		
Gestational age (weeks)							
0-13	1.37 (0.604-3.148)	0.445	1.25 (0.673-2.328)	0.479	1.52 (0.792-2.948)	0.206	
14-26	Ref.		Ref.		Ref.		

MODEL 6								
WHO-VAW-								
physical violence								
Yes	1.39 (0.438-4.440)	0.574	0.66 (0.230-1.930)	0.45	0.74 (0.263-2.084)	0.570		
No	Ref.		Ref.		Ref.			
GHQ								
≥4	1.40 (0.683 - 2.859)	0.360	1.64 (0.944 - 2.850)	0.07	2.36 (1.355 - 4.097)	0.002		
0-3	Ref.		Ref.		Ref.			
Age (years)								
>30	2.11 (0.849-5.256)	0.108	0.96 (0.536-1.719)	0.89	0.59 (0.330-1.054)	0.075		
<20	4.83 (1.300-17.975)	0.019	1.01 (0.330-3.709)	0.87	0.10 (0.012-0.937)	0.043		
20-30	Ref.		Ref.		Ref.			
Parity								
0	0.54 (0.144-2.073)	0.374	1.04 (0.411-2.647)	0.93	0.99 (0.391-2.529)	0.990		
1-2	0.95 (0.254-3.563)	0.940	0.59 (0.222-1.615)	0.31	0.98 (0.368-2.628)	0.974		
≥3	Ref.		Ref.		Ref.			
Gestational age (weeks)								
0-13	1.35 (0.592-3.081)	0.474	1.26 (0.678-2.36)	0.46	1.53 (0.799-2.943)	0.198		
14-26	Ref.		Ref.		Ref.			

Multivariate regression models, considering mental health as mediator. Confounding factors included in the models: age, parity, and, gestational age. Abbreviations: OR= odds ratio; CI= confidence interval; Ref= Reference; WHO-VAW= World Health Organization-Violence Against Women; GHQ= General Health Questionnaire; PHQ= Patient Health Questionnaire.




Figure 1. Model of psychological violence as a predictor of obesity, mediated by mental health changes. The bias-corrected and accelerated confidence interval (BCa CI) was estimated using the bootstrapping technique (5,000, re-sampling).