

NOTES FOR AUTHORS

Papers submitted for publication in the *Bulletin of Entomological Research* should be sent, in duplicate, to the Director, C·A·B International Institute of Entomology, 56 Queen's Gate, London SW7 5JR. They must deal with original research concerning insects, mites or ticks of economic importance in the agricultural, medical or veterinary fields; they must be of more than purely local interest, and taxonomic papers will not be accepted unless the organisms dealt with are of definite importance in the field of applied entomology. Review articles may also be submitted, but will normally be by invitation. Papers must not have been published or accepted for publication elsewhere. They will normally be published in order of their receipt in definitive form.

For general guidance on presentation and style, including the preparation of illustrations, contributors are urged to consult *Writing Scientific Papers in English. An ELSE-Ciba Foundation Guide for Authors* (by Maeve O'Connor & E. Peter Woodford). 116 pp., Amsterdam, Associated Scientific Publishers, 1975, ISBN 90-219-4035-3.

Text

Papers should be typewritten (double spaced, with ample margins and on one side of the paper only) and a current number of the *Bulletin* should be consulted for details of style and lay-out. Titles should be as brief as is consistent with clarity. Abbreviations should conform to *British Standard 1991, Part I, 1967*, and are alike in singular and plural; stops will be used only where ambiguity might occur, e.g. 'in.' not 'in' for inch, but 'in²' for square inch. Metric units are preferred; if non-metric units are used, metric equivalents should be given at least once in each article.

Tables

Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must on no account be of a size or form that will not conveniently fit into the page of the *Bulletin*.

Illustrations

Line drawings and graphs must be in jet-black indian ink, preferably on Bristol-board or tracing cloth. Graphs may be drawn on paper ruled in pale blue or light grey; paper having green, red, yellow or brown lines should not be used. Any stippling or hatching, as well as any numerals or lettering, on the text-figures must be completed, by the author, in a form suitable for reproduction; this work is *not* undertaken by the printer. Text-figures, when reproduced, cannot exceed 12·7 cm in width; if, therefore, the original drawings are larger, the lettering on them must be in proportion, so that it is still legible when reduced; special attention should be paid to the relative scales of lettering and line thicknesses in relation to the figure as a whole. Photographs for reproduction as figures are admissible when they are a real contribution to the text. Captions to figures should be typed on a separate sheet.

Abstract

Each paper must contain a carefully prepared and accurate informative abstract that shall be complete in itself and intelligible without reference to the text or figures. It should not exceed 250 words, and should be included at the beginning immediately after the title, author and author's affiliation.

References

References must be based on the name and year system and conform to the present usages of the *Bulletin*. Titles of journals should be abbreviated as in the 4th edition of *The World List of Scientific Periodicals*. References to unpublished information should normally be included only in the text—(A. B. Jones, pers. comm.) or (A. B. Jones, Control of sandflies by insecticides. Unpublished report. Ross Institute, London, 1950).

Offprints

Free copies of offprints are allowed as follows: One author—20 copies, two or more authors—30 copies. Further copies may be obtained on payment, and the number required should be specified when the paper is submitted.

BULLETIN OF ENTOMOLOGICAL RESEARCH

(World List Abbreviation: *Bull. ent. Res.*)

ISSN 0007-4853

Original articles

PAGE

LUTWAMA, J. J. & MATANMI, B. A. Efficacy of <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> and <i>Baculovirus heliothis</i> foliar applications for suppression of <i>Helicoverpa armigera</i> (Hübner) (Noctuidae) and other lepidopterous larvae on tomato in south-western Nigeria	173
PAGE, W. W. Varying durations of arrested oocyte development in relation to migration in the African armyworm moth, <i>Spodoptera exempta</i> (Walker) (Lepidoptera: Noctuidae)	181
VAN DRIESCHE, R. G. Field measurement of population recruitment of <i>Apanteles glomeratus</i> (L.) (Hymenoptera: Braconidae), a parasitoid of <i>Pieris rapae</i> (L.) (Lepidoptera: Pieridae), and factors influencing adult parasitoid foraging success in kale	199
TAYLOR, B. The impact of yellow stem-borer, <i>Scirpophaga incertulas</i> (Walker) (Lepidoptera: Pyralidae), on deepwater rice, with special reference to Bangladesh	209
KELLY, P. M., STERLING, P. H., SPEIGHT, M. R. & ENTWHISTLE, P. F. Preliminary spray trials of a nuclear polyhedrosis virus as a control agent for the brown-tail moth, <i>Euproctis chryorrhoea</i> (L.) (Lepidoptera: Lymantriidae) .	227
MATTHIESSEN, J. N. & PALMER, M. J. Prediction of temperatures in cattle dung for estimating development times of coprophilous organisms	235
LOXDALE, H. D. & BROOKES, C. P. Electrophoretic study of enzymes from cereal aphid populations. V. Spatial and temporal genetic similarity of holocyclic populations of the bird-cherry oat aphid, <i>Rhopalosiphum padi</i> (L.) (Hemiptera: Aphididae), in Britain	241
REYNOLDS, D. R. & RILEY, J. R. A migration of grasshoppers, particularly <i>Diabolocatantops axillaris</i> (Thunberg) (Orthoptera: Acrididae), in the West African Sahel	251
EISEMANN, C. H. Upwind flight by gravid Australian sheep blowflies, <i>Lucilia cuprina</i> (Wiedemann) (Diptera: Calliphoridae), in response to stimuli from sheep	273
BURSELL, E., GOUGH, A. J. E., BEEVOR, P. S., CORK, A., HALL, D. R. & VALE, G. A. Identification of components of cattle urine attractive to tsetse flies, <i>Glossina</i> spp. (Diptera: Glossinidae)	281
VALE, G. A., HALL, D. R. & GOUGH, A. J. E. The olfactory responses of tsetse flies, <i>Glossina</i> spp. (Diptera: Glossinidae), to phenols and urine in the field	293
BRADY, J. & SHERENI, W. Landing responses of the tsetse fly <i>Glossina morsitans morsitans</i> Westwood and the stable fly <i>Stomoxys calcitrans</i> (L.) (Diptera: Glossinidae & Muscidae) to black-and-white patterns: a laboratory study ...	301
NOYES, J. S. <i>Gyranusoidea tebygi</i> sp. n. (Hymenoptera: Encyrtidae), a parasitoid of <i>Rastrococcus</i> (Hemiptera: Pseudococcidae) on mango in India	313
LANGTON, P. H., CRANSTON, P. S. & ARMITAGE, P. The parthenogenetic midge of water supply systems, <i>Paratanytarsus grimmii</i> (Schneider) (Diptera: Chironomidae)	317
RAHBÉ, Y., FEBVAY, G. & KERMARREC, A. Foraging activity of the attine ant <i>Acromyrmex octospinosus</i> (Reich) (Hymenoptera: Formicidae) on resistant and susceptible yam varieties	329
PADGHAM, D. E. & WOODHEAD, S. Variety-related feeding patterns in the brown planthopper, <i>Nilaparvata lugens</i> (Stål) (Hemiptera: Delphacidae), on its host, the rice plant	339
BROWN, P. A. & BLACKMAN, R. L. Karyotype variation in the corn leaf aphid, <i>Rhopalosiphum maidis</i> (Fitch), species complex (Hemiptera: Aphididae) in relation to host-plant and morphology	351

© C.A.B International, 1988

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without prior permission of the copyright owner.