

Research Article

*These authors contributed equally to this work and should be considered co-first authors.

Cite this article: Zhang L, Liu K, Deng J, Xu H, Wang J, Wang K, Qu W, Liu G and Gu X (2024). Occurrence of antimicrobial resistance and class 1 integrons in *Staphylococcus aureus* isolated from bovine mastitis in China. *Journal of Dairy Research* **91**, 298–304. <https://doi.org/10.1017/S0022029924000566>

Received: 21 May 2024
Revised: 23 May 2024
Accepted: 7 June 2024

Keywords:

antimicrobial resistance; bovine mastitis; class 1 integrons; *Staphylococcus aureus*

Corresponding author:

Xiaolong Gu;
Email: xuewu24588@126.com

Occurrence of antimicrobial resistance and class 1 integrons in *Staphylococcus aureus* isolated from bovine mastitis in China

Limei Zhang^{1,*}, Kai Liu^{1,*}, Jianming Deng¹, Hui Xu¹, Juyu Wang¹, Kuan Wang¹, Weijie Qu¹, Gang Liu² and Xiaolong Gu¹

¹College of Veterinary Medicine, Yunnan Agricultural University, Kunming, Yunnan Province 650201, PR China and ²College of Veterinary Medicine, China Agricultural University, Beijing 100193, PR China

Abstract

Integrons are important genetic elements that allow easy acquisition and dissemination of antimicrobial resistance genes. Studies reporting occurrence of integrons in *Staphylococcus aureus* (*S. aureus*) isolated from bovine mastitis in large dairy farms across China are scarce. The aim of this study was to investigate the occurrence of class 1 integrons (intI1), antimicrobial resistance (AMR) and associated genes in *S. aureus* isolated from bovine mastitis and their associations. Minimum inhibitory concentrations (MICs) were determined to evaluate the AMR phenotypes, whereas PCR was carried out to assess the occurrence of AMR genes and intI1. In addition, index cluster analysis was used to estimate associations between AMR phenotype, genotype and intI1 in 103 isolates. Overall, 83% of *S. aureus* were intI1-positive and 5 types of gene cassettes were detected. Susceptibility against single antimicrobial agents ranged from 0% (erythromycin), 12% (ampicillin) and 16% (penicillin G) to 96% (gentamicin). Most isolates (64%) were intermediate-resistant against erythromycin, whereas resistance against ceftriaxone (22%), clindamycin (4%), cefotaxime (2%), tetracycline (1%) and ciprofloxacin (1%) were relatively uncommon. The predominant resistant gene was *blaZ* gene ($n = 88$, 85%) followed by *tetD* gene ($n = 85$, 83%). With an estimated prevalence of 12% of the *meCA* gene, methicillin-resistant *S. aureus* isolates had higher MIC₅₀ and MIC₉₀ for majority of antimicrobials than methicillin-susceptible *S. aureus* isolates. Presence of the *ermC* gene was associated with erythromycin resistance. Ampicillin, erythromycin and penicillin G resistance were associated with intI1. The data presented in our study indicated that class 1 integron-mediated resistance possibly plays an important role in dissemination of AMR in *S. aureus* isolated from bovine mastitis.

Staphylococcus aureus is one of the major contagious udder pathogens and causes great economic losses in the dairy industry by way of compromising milk quality, decreasing milk production and causing recurrent infections (Wang *et al.*, 2016; Gao *et al.*, 2017). Antimicrobial agents have been used to treat bovine mastitis of bacterial origin worldwide. However, antimicrobial resistance (AMR) in bacteria challenges efficacy, making treatment costly, time consuming and increasing the risk of therapeutic failure (Sharma *et al.*, 2017). Indeed, AMR can disseminate without any geographical boundaries and affect diverse animal populations including humans (Littmann and Viens, 2015). It has been widely recognized as one of the most serious public health challenges of the twenty-first century (Hernando-Amado *et al.*, 2019). It is, therefore, important to survey and study the occurrence of AMR in pathogenic flora of dairy cows, to understand the possible mechanism of AMR and to develop alternative strategies and detect novel target sites for combating AMR.

Bacteria can develop AMR either vertically (gene mutation) or horizontally (obtaining new genes from other species or strains: Martinez and Baquero, 2000; Palmer *et al.* 2010). It has been reported that integrons play a key role in dissemination of AMR through horizontal gene transfer, particularly in Gram-negative pathogens (Xia *et al.*, 2016). Nandi *et al.* (2004) reported that in poultry litter, the major reservoir for class 1 integrons is Gram-positive bacteria instead of Gram-negative Enterobacteriaceae. For *S. aureus*, presence of class 1 integrons was reported in samples from humans (Deng *et al.*, 2015a), however, studies on presence of integrons in *S. aureus* isolates obtained from dairy cows are still scarce. Therefore, the objectives of this study were to determine the phenotypic and genotypic AMR as well as the occurrence of class 1 integrons in *S. aureus* strains isolated from bovine clinical mastitis in China, and to investigate the associations between class 1 integrons and AMR genotypes and phenotypes.

Materials and methods

The study was conducted in compliance with guidelines of the Beijing Municipality on the Review of Welfare and Ethics of Laboratory Animals, approved by the Beijing Municipality

Administration Office of Laboratory Animals (BAOLA) and approved by the China Agricultural University Animal Ethics Committee (protocol CAU-AEC-2010-0603).

Herds and sampling

A total of 103 *S. aureus* isolates were collected over a period of four years (2013–2016) from quarter milk samples of Holstein cows with clinical mastitis in 19 large (> 500 cows) dairy herds in 9 provinces which are the major producers of milk in China (online Supplementary Fig. S1). Briefly, 1021 milk samples were collected following NMC guidelines (2017) from lactating cows (one sample per cow) suffering from clinical mastitis (NMC, 2017). Clinical mastitis was diagnosed based on visual abnormality /inflammation of the udder or its milk secretion. Samples were collected in sterile bottles (50 mL) and transported on ice to the mastitis laboratory at the China Agricultural University (Beijing, China).

Isolation and identification of *S. aureus*

Milk samples (10 µl out of 50 ml) were spread-plated on TSA (trypticase soya agar; Sigma, India) supplemented with 5% defibrinated sheep blood and incubated at 37 °C for 16–24 h. *S. aureus* isolates were identified by colony morphology, hemolysis, Gram staining, catalase test, tube coagulase test, oxidation and fermentation of mannitol salt agar (AOBOX, Beijing, China), as well as 16S rRNA, *coa* and *nuc* gene sequence analysis, as described by Wang *et al.* (2016).

Antimicrobial susceptibility testing

To analyze the antimicrobial susceptibility profiles of all isolates, minimum inhibitory concentrations (MICs) were determined against a panel of 11 different antimicrobials using the broth microdilution method following the Clinical Laboratory and Standard Institutes (CLSI) guidelines (CLSI, 2018). Antimicrobial agents were selected either according to their availability in commercial products or working as representative of an antimicrobial family: penicillin, ampicillin, oxacillin, erythromycin, tetracycline, ciprofloxacin, ceftriaxone, cefotaxime, gentamicin, clindamycin. A concentration of 0.125–64 µg/ml was used for *S. aureus*. *Staphylococcus aureus* ATCC 29213 was used as the quality control strain. Breakpoints were defined as described by the CLSI (CLSI, 2018). Isolates were classified as susceptible, intermediate or resistant to each antimicrobial. Multidrug resistance was defined as resistant to at least 3 antimicrobials. As defined by CLSI, MRSA (methicillin resistant *S. aureus*) are those strains that possess *mecA* gene. Frequency distribution of MIC₅₀ and MIC₉₀ were calculated for each antimicrobial.

Presence of antimicrobial resistance genes

Template DNA of the 103 *S. aureus* isolates were prepared using bacterial genomic DNA extraction kit (CW, Beijing, China) according to the manufacturer instructions (Zhang *et al.*, 2018). The 13 acquired resistance genes were analyzed using PCR (primers identified in online Supplementary Table S1).

Detection of three classes of intergrons

Presence of class 1, 2 and 3 intergrons was analyzed in all isolates using PCR as per Goldstein *et al.* (2001). The following program was employed: 5 min of initial denaturation at 94 °C, followed by

30 cycles (30 s of denaturation at 94 °C, 30 s of annealing at 55 °C and 1 min of extension at 72 °C) and a final extension step. The PCR products were randomly selected for sequencing to ensure specificity and accuracy.

Gene cassettes inserted in the variable regions of class 1 intergrons were amplified as described by Lévesque *et al.* (1995). To determine whether each cassette gene PCR amplicon with the same size had the same content, PCR products were digested with *AluI* and *RsaI* (New England Biolabs, Beijing, China). PCR products with the same restriction fragment length polymorphism (RFLP) pattern were considered to contain the same gene cassettes (Lévesque *et al.*, 1995; Li *et al.*, 2015).

Statistical analysis

SPSS version 22.0 (IBM, Chicago, IL, USA) was used for statistical analysis. Index hierarchical cluster analysis was performed using between groups linkage method with measure of binary squared Euclidean distance to explore the similarity among antimicrobial resistance phenotypes, resistances genes and int1 (Zhang *et al.*, 2017). Also, the association between phenotypic and genotypic resistance was explored using χ^2 or Fisher's exact test on contingency tables (positive and negative for phenotype or genotype of isolates).

Results

Phenotypic antimicrobial susceptibility

Overall, for antimicrobials tested, MIC₅₀ ranged from 0.25 to 4 µg/ml and MIC₉₀ from 0.5 to 64 µg/ml (Table 1). Susceptibility for single antimicrobial agents ranged from 0% (for erythromycin) to 96% (for gentamicin). All *S. aureus* isolates demonstrated a low sensitivity to ampicillin (12%) and penicillin (16%). Most isolates (64%) were of intermediate susceptibility for erythromycin, followed by ceftriaxone (22%), clindamycin (4%), cefotaxime (2%), cefotaxime (1%) and ciprofloxacin (1%). In total, 84% of *S. aureus* isolates were susceptible to oxacillin (Table 2).

A total of 12 isolates (12%) were identified as MRSA (methicillin-resistant *Staphylococcus aureus*, *mecA* gene positive), which were distributed in 8 regions except Inner Mongolia. Compared to MSSA (methicillin-sensitive *Staphylococcus aureus*, *mecA* gene negative) isolates, MRSA isolates demonstrated higher resistance rate to all the 10 antimicrobial agents. In addition, the MIC ranges of gentamicin, ciprofloxacin, penicillin, ampicillin, oxacillin, cefotaxime and ceftriaxone were larger in MRSA isolates. Similarly, the MIC₅₀ results of MRSA isolates were higher than MSSA for all antimicrobial agents except for tetracycline, gentamicin, penicillin, ampicillin and ceftriaxone; MIC₉₀ values of MRSA isolates were also higher than MSSA isolates except for penicillin and ampicillin (Table 3). In addition, the multidrug resistance phenotype was much more frequent in MRSA isolates (12, 100%) than in MSSA isolates (51, 56%; Fig. 1). MSSA isolates demonstrated different resistance patterns: 24 MSSA isolates (47%) showed resistance to 3 antimicrobials, 15 isolates (10%) to 5 antimicrobials and 10 isolates to 4 antimicrobials. While for MRSA isolates, resistance to 4, 5, 6, 9 and 10 antimicrobials were identified with 4, 4, 2, 1 and 1 isolates, respectively.

Occurrence of resistance genes

Of the 13 resistance genes, *blaZ* was detected most frequently (85%), followed by *tetD* (83%). *tetK*, *ermT* and *ermC* were also detected in

Table 1. Minimal inhibitory concentration (MIC) distribution for the 103 *Staphylococcus aureus* isolates

Antimicrobial	Distribution (%) of MIC (ug/ml)												
	%Susc. ^a	0.125	0.25	0.5	1	2	4	8	16	32	64	MIC ₅₀ ^b	MIC ₉₀ ^c
Tetracycline	93	21	63	6	2	0	1	1	0	6	0	0.25	0.5
Gentamicin	96	8	52	27	3	5	2	0	1	2	1	0.25	1
Erythromycin	0	0	0	0	0	18	46	8	2	0	26	4	64
Ciprofloxacin	84	7	68	4	5	1	5	5	6	0	1	0.25	8
Penicillin	16	16	15	4	5	7	11	14	15	11	6	4	32
Oxacillin	84	0	79	6	0	0	13	1	2	0	0	0.25	4
Ampicillin	12	8	4	4	11	21	14	13	13	3	1	4	16
Clindamycin	80	0	60	11	1	8	4	0	3	3	11	0.25	64
Cefotaxime	95	0	2	22	40	31	2	0	0	2	1	1	2
Ceftriaxone	75	0	0	0	18	56	22	6	1	2	0	2	4

^aPercent of susceptible isolates according to CLSI (2018).

^bThe MIC value that inhibits growth of 50% of the isolates.

^cThe MIC value that inhibits growth of 90% of the isolates. indicates the CLSI epidemiological cut off values.

Table 2. Antimicrobial activity and potency of 11 antimicrobials against MRSA and MSSA isolated from clinical mastitis in Chinese dairy herds

Antimicrobial	MRSA/MSSA			
	Resistance (%)	MIC ₅₀	MIC ₉₀	MIC range
	25/3	0.25/0.25	32/0.5	0.125–32/0.125–32
Gentamicin	17/2	0.25/0.25	2/1	0.125–64/0.125–32
Erythromycin	75/28	64/4	64/64	2–64/2–64
Ciprofloxacin	58/10	8/0.25	16/4	0.25–64/0.125–16
Penicillin	100/73	4/4	16/16	0.25–64/0.125–64
Oxacillin	100/4	4/0.25	16/0.5	4–16/0.25–4
Ampicillin	100/77	2/2	16/16	0.5–64/0.125–16
Clindamycin	33/13	0.5/0.25	64/32	0.25–64/0.25–64
Cefotaxime	17/1	2/1	32/2	0.5–32/0.25–64
Ceftriaxone	33/13	4/4	64/8	2–64/2–16

considerable proportion (Table 3). Furthermore, we analyzed the frequency of drug-specific resistance on presence or absence of a particular antimicrobial resistance gene, results indicated that presence of some resistance genes (*ermC*, *blaZ* and *mecA*) was associated with drug-specific resistance phenotype (Table 3).

Gene patterns were commonly observed for resistant isolates included the presence of single *tetL* (17% of tetracycline-resistant isolates), *tetD-tetK* (33% of tetracycline-resistant isolates), *tetD-tetL* (17% of tetracycline-resistant isolates), *ermT* (24% of erythromycin-resistant isolates), *ermC* (27% of erythromycin-resistant isolates), *ermB* (5% of erythromycin-resistant isolates) and *ermT-ermC* (3% of erythromycin-resistant isolates) (Fig. 2).

Class 1 integrons

Of the 103 isolates, 84 (83%) were positive for *intI1*. From 76 of 84 isolates (89%), the class 1 integrons variable regions were

amplified by PCR, yielding amplicons of 3 distinct sizes: 1600 bp (74 isolates), 1000 bp (1 isolate) and 750 bp (1 isolate). Digested with *AluI* and *RsaI*, a total of 5 different patterns were detected (Fig. 3).

Associations among resistance genes, AMR and *intI*.

Index cluster analysis indicated that GM-resistance, CTX-resistance, TET-resistance, CIP-resistance, OX-resistance, CL-resistance, CRO-resistance and ERY-resistance in *S. aureus* isolates were highly associated (Fig. 4a). Resistance to AMP and PEN was associated with *intI1*. Correspondingly, aminoglycosides resistance genes (*aphA3*, *aadA1/A2*), macrolides resistance genes (*ermB* and *ermT*), tetracycline resistance genes (*tetL* and *tetK*), β -lactams resistance gene *mecA* and lincosamides resistance gene *lunA* were also highly associated. A similar association was present for *intI1* and *blaZ* (Fig. 4b).

Table 3. The MIC_{50/90} of *Staphylococcus aureus* isolates with or without a particular antimicrobial resistance gene and association between resistance phenotype and genotype

Antimicrobial	MIC _{50/90} (ug/mL)		Positive (%)	Resistance (%)		P value
	Positive	Negative		Gene absent	Gene present	
Tetracycline	0.25/0.5	0.25/32	<i>tetD</i> (83%)	11	6	0.292
	0.25/8	0.25/0.5	<i>tetK</i> (25%)	7	0	0.359
	0.25/0.5	0.25/0.5	<i>tetL</i> (8%)	6	0	0.662
Erythromycin	-	-	<i>ermB</i> (4%)	35	100	0.359
	4/64	4/64	<i>ermT</i> (21%)	38	13	0.15
	4/64	4/64	<i>ermC</i> (20%)	20	49	0.002
Penicillin	4/16	0.125/16	<i>blaZ</i> (85%)	53	90	0.00
Oxacillin	4/16	0.25/0.5	<i>mecA</i> (12%)	4	92	0.00
Gentamicin	0.25/1	0.25/2	<i>aadA1/A2</i> (5%)	31	0	1.00
	0.5/32	0.25/1	<i>aacA-aphD</i> (17%)	1	20	0.34
	0.25/0.5	0.25/2	<i>aphA3</i> (4%)	3	0	1.00
	0.25/1	0.25/2	<i>aadD</i> (3%)	4	0	0.575
Clindamycin	0.5/16	0.25/64	<i>lnuA</i> (16%)	14	36	0.06




Figure 1. Multi-resistance patterns of MRSA and MSSA isolates from clinical mastitis in China. Different numbers above bars indicate the number of isolates corresponding to the resistance pattern.

Discussion

Despite considerable progress in alternative treatment strategies such as vaccines, AMR remains a major problem in the treatment of bovine mastitis, and poses a serious threat to public health (Wang *et al.*, 2016; Cheng *et al.*, 2019). In this study, AMR of *S. aureus* was exhibited against antimicrobials frequently used in dairy herds, such as erythromycin, penicillin, and ampicillin. Some AMR genes were involved in drug-specific AMR and vice versa. Class 1 integrons were associated with resistance to several drugs (penicillin and ampicillin) and antimicrobial resistance genes *blaZ* and *tetD*. Of note, the MIC₉₀ of *S. aureus* against erythromycin was very high. Despite the lack of availability of an erythromycin clinical breakpoint for *S. aureus* isolated from bovine mastitis, concern should be raised for this, since it suggests that macrolides such as erythromycin might not be effective

for treatment of mastitis caused by *S. aureus* in China. Furthermore, the MIC₅₀/MIC₉₀ (4/64) of erythromycin was higher than in both MSSA (< 0.25/4) and MRSA (> 4/> 4) isolates from human hospitals in previous report (Pfaller *et al.*, 2018). Similar results were present for CL (0.25/64 vs. < 0.25/2), which indicates that erythromycin and clindamycin resistance in *S. aureus* is developing regardless of bacterial origin. We observed a lower resistance rate to tetracycline (6 vs. 19%), ciprofloxacin (16 vs. 53%), ampicillin (88 vs. 91%) and penicillin (83 vs. 91%) compared to another study in the mid-east of China, in which 200 CM milk samples were included (Zhang *et al.*, 2016). Regarding the comparison between different studies, differences in resistance prevalence were observed among countries (Wang *et al.*, 2016; Beyene *et al.*, 2017; Freitas *et al.*, 2018). We inferred these differences may be related to the source of samples,




Figure 2. Gene patterns in *Staphylococcus aureus* from clinical mastitis resistant to antimicrobials of TET (a) and ERY (b).




Figure 3. Representative profiles identified among *Staphylococcus aureus* from clinical mastitis by *AluI* and *RsaI* digestion in class 1 integrons gene cassette. M, DNA ladder; 1, SA1 (Pattern 1); 2, SA2 (Pattern 2); 3, SA3 (Pattern 2); 4, SA4 (Pattern 3); 5, SA5 (Pattern 4); 6, SA6 (Pattern 1); 7, SA7 (Pattern 4); 8, SA8 (Pattern 1); 9, SA9 (Pattern 4); 10, SA10 (Pattern 1); 11, SA11 (Pattern 3); 12, SA12 (Pattern 5); 13, SA13 (Pattern 5); 14, SA14 (Pattern 5).

geographical distribution and bacterial genotype. The high and intermediate resistance against erythromycin (65%) and ceftriaxone (20%) indicated that veterinarians need to be more prudent when considering these antimicrobials.

MRSA has been reported in livestock in recent years all over the world with different prevalence. In our study, 12% of isolates harbored the *mecA* gene, much higher than the 0.05% reported by Saini *et al.* (2012) in Canada, and the 1% by Ahangari *et al.* (2017) in Iran. In total, the prevalence of MRSA strains reported from 7% to 19% in China was much higher than that in other

countries, perhaps resulting from a misuse of various antibiotics (Zhang *et al.*, 2016; Li *et al.*, 2017; Qu *et al.*, 2019). Compared with MSSA, MRSA in this study showed larger MIC range, higher MIC_{50/90} values for most antimicrobials and more multi-drug resistance, which was consistent with the previous study (Wang *et al.*, 2016).

In this study, *blaZ* gene was widespread in the herds, which was consistent with the prominent resistance to penicillin. However, the presence of other AMR genes did not always correspond with the resistance phenotype. For instance, *tetD* was the




Figure 4. Index cluster analysis of *Staphylococcus aureus* from clinical mastitis in antimicrobial resistance phenotypes (a), resistances genes and mobile genetic element *int1* (b). When branch point was less than 5, these branches were determined as related to each other significantly.

most prevalent gene whereas clinical resistance to tetracycline was uncommon. In addition, an association was observed between detection of gene *ermC*, *blaZ* and *mecA* encoding resistance to erythromycin, penicillin and oxacillin and their resistance phenotype. The discordance between phenotypic and genotypic resistance could be due to lack of expression of resistance genes or MIC increases below the adopted thresholds.

Integrations are considered an important source of AMR genes, which facilitate the horizontal spread of resistance genes within microbial populations (Deng *et al.*, 2015b). Class 1 integrons were detected in 83% of isolates, lower than another study conducted in northwestern China, in which all 121 *S. aureus* isolates from clinical mastitis samples were positive for the *int1* gene (Li *et al.*, 2018). The proportion of class 1 integrons in *S. aureus* isolates can be different among farm animals, hospitals and environment. For example, they were present in 73% of wounds, blood, urines, nasal and throat *S. aureus* isolates from hospitals in Tehran (Mostafa *et al.*, 2015), absent from all the MRSA isolates from a Chinese municipal wastewater treatment plant and a swine slaughterhouse wastewater treatment plant but present in all of the other wastewater samples tested (Wan and Chou, 2015). High occurrence of *int1* gene suggests that *S. aureus* isolated from bovine mastitis can acquire and easily disseminate resistance genes.

Amplification and restriction digest of the variable region of class 1 integrons from 76 isolates allowed clustering into 5 types. The reasons why some *int1*-positive isolates did not harbor gene cassettes may be that mutations happened at the 3'CS; or the variable region was too long to be amplified. Many studies reported different gene cassette in *int1*-positive *S. aureus* isolates, including *aadA2* (59%), *dfrA12-orfF-aadA2* (49%), *aacA4-cmlA1* (3%), *dfrA17-aadA5* (1%), *dfrA1-aadA1* (45%), and *aadA1* (26%) (Xia *et al.*, 2016; Li *et al.*, 2018). Sometimes, the gene cassettes in class 1 integrons did not account for the total resistance phenotypes for erythromycin, tetracycline and penicillin (Li *et al.*, 2018). We did not sequence the gene cassettes in this study but used index cluster analysis to estimate the association between *int1* and antimicrobial resistance phenotype and genotype. Results showed that, consistent with the penicillin resistance

phenotype, incidence of *int1* was also related to *blaZ* gene. Similarly, a strong association was shown between *int1* and penicillin-resistant phenotypes in *E. coli* isolated from swine (Zhang *et al.*, 2020). This association suggests that these resistance mechanisms are co-selected, and warrants further investigations on how these determinants disseminating in dairy herds.

In conclusion, a high occurrence of AMR against erythromycin, ampicillin and penicillin in *S. aureus* isolated from clinical bovine mastitis in large dairy farms was detected, and the most frequent resistance genes were *blaZ* and *tetD* genes. Twelve percent of *S. aureus* isolates were *mecA*-positive. In addition, class 1 integrons were commonly detected in *S. aureus* and associated with penicillin resistance. This suggests that Class 1 integrons and penicillin resistance are genetically linked in *S. aureus* isolated from bovine mastitis in China.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0022029924000566>

Acknowledgments. This research was financially supported by the Basic Research Project-Youth Project (202001AU070098); National Natural Science Foundation of China (31902335); Yunnan Fundamental Research Projects (grant no. 202301AS070081), Rural Vitalization Science and technology Project -Gejiu Cattle Industry Science and Technology Mission in Yunnan Province (grant no. 202304BI090007).

References

- Ahangari Z, Ghorbanpoor M, Shapouri MRS, Gharibi D and Ghazvini K (2017) Methicillin resistance and selective genetic determinants of *Staphylococcus aureus* isolates with bovine mastitis milk origin. *Iranian Journal of Microbiology* **9**, 152–159.
- Beyene T, Hayishe H, Gizaw F, Beyi AF, Abunna F, Mammo B, Ayana D, Waktole H and Abdi RD (2017) Prevalence and antimicrobial resistance profile of *Staphylococcus* in dairy farms, abattoir and humans in Addis Ababa, Ethiopia. *BMC Research Notes* **10**, 171.
- Cheng J, Qu W, Barkema HW, Nobrega DB, Gao J, Liu G, De Buck J, Kastelic JP, Sun H and Han B (2019) Antimicrobial resistance profiles of 5 common bovine mastitis pathogens in large Chinese dairy herds. *Journal of Dairy Science* **102**, 2416–2426.

- CLSI (2018) *Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard*. Wayne, PA, USA: Clinical and Laboratory Standards Institute.
- Deng Y, Bao X, Ji L, Chen L, Liu J, Miao J, Chen D, Bian H, Li Y and Yu G (2015a) Resistance integrons: class 1, 2 and 3 integrons. *Annals of Clinical Microbiology and Antimicrobials* **14**, 45.
- Deng Y, Liu J, Peters BM, Chen L, Miao J, Li B, Li L, Chen D, Yu G, Xu Z and Shirliff ME (2015b) Antimicrobial resistance investigation on *Staphylococcus* strains in a local hospital in Guangzhou, China, 2001–2010. *Microbial Drug Resistance* **21**, 102–104.
- Freitas CH, Mendes JF, Villarreal PV, Santos PR, Gonçalves CL, Gonzales HL and Nascente PS (2018) Identification and antimicrobial susceptibility profile of bacteria causing bovine mastitis from dairy farms in Pelotas, Rio Grande do Sul. *Brazil Journal of Biology* **78**, 661–666.
- Gao J, Barkema HW, Zhang L, Liu G, Deng Z, Cai L, Shan R, Zhang S, Zou J, Kastelic JP and Han B (2017) Incidence of clinical mastitis and distribution of pathogens on large Chinese dairy farms. *Journal of Dairy Science* **100**, 4797–4806.
- Goldstein C, Lee MD, Sanchez S, Hudson C, Phillips B, Register B, Grady M, Liebert C, Summers AO, White DG and Maurer JJ (2001) Incidence of class 1 and 2 integrons in clinical and commensal bacteria from livestock, companion animals, and exotics. *Antimicrobial Agents and Chemotherapy* **45**, 723–726.
- Hernando-Amado S, Coque TM, Baquero F and Martínez JL (2019) Defining and combating antibiotic resistance from one health and global health perspectives. *Nature Microbiology* **4**, 1432–1442.
- Lévesque C, Piche L, Larose C and Roy P (1995) PCR mapping of integrons reveals several novel combinations of resistance genes. *Journal of Antimicrobial Chemotherapy* **39**, 185–191.
- Li T, Lu H, Wang X, Gao Q, Dai Y, Shang J and Li M (2017) Molecular characteristics of *Staphylococcus aureus* causing bovine mastitis between 2014 and 2015. *Frontiers in Cellular and Infection Microbiology* **7**, 127.
- Littmann J and Viens AM (2015) The ethical significance of antimicrobial resistance. *Public Health Ethics* **8**, 209–224.
- Martínez JL and Baquero F (2000) Mutation frequencies and antibiotic resistance. *Antimicrobial Agents and Chemotherapy* **44**, 1771–1777.
- Mostafa M, Siadat SD, Shahcheraghi F, Vaziri F, Japoni-Nejad A, Vand Yousefi J, Rajaei B, Harifi Mood E, Ebrahim Zadeh N, Moshiri A, Seyed Siamdoust SA and Rahbar M (2015) Variability in gene cassette patterns of class 1 and 2 integrons associated with multi drug resistance patterns in *Staphylococcus aureus* clinical isolates in Tehran-Iran. *BMC Microbiology* **15**, 152.
- Nandi S, Maurer JJ, Hofacre C and Summers AO (2004) Gram-positive bacteria are a major reservoir of Class 1 antibiotic resistance integrons in poultry litter. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 7118–7122.
- NMC (2017) *Laboratory Handbook on Bovine Mastitis*. National Mastitis Council. Madison, WI: NMC Inc.
- Palmer KL, Veronica K and Gilmore MS (2010) Horizontal gene transfer and the genomics of enterococcal antibiotic resistance. *Current Opinion in Microbiology* **13**, 632–639.
- Pfaller MA, Sader HS, Flamm RK, Castanheira M and Mendes RE (2018) Oritavancin in vitro activity against Gram-positive organisms from European and United States medical centers: results from the SENTRY antimicrobial surveillance program for 2010–2014. *Diagnostic Microbiology and Infectious Disease* **91**, 199–204.
- Qu Y, Zhao H, Nobrega DB, Cobo ER, Han B, Zhao Z, Li S, Li M, Barkema HW and Gao J (2019) Molecular epidemiology and distribution of antimicrobial resistance genes of *Staphylococcus* species isolated from Chinese cows with clinical mastitis. *Journal of Dairy Science* **102**, 1571–1583.
- Saini V, McClure JT, Léger D, Keefe GP, Scholl DT, Morck DW and Barkema HW (2012) Antimicrobial resistance profiles of common mastitis pathogens on Canadian dairy farms. *Journal of Dairy Science* **95**, 4319–4332.
- Sharma C, Rokana N, Chandra M, Singh BP, Gulhane RD, Gill JPS, Ray P, Puniya AK and Panwar H (2017) Antimicrobial resistance: its surveillance, impact, and alternative management strategies in dairy animals. *Frontiers in Veterinary Science* **4**, 237.
- Wan MT and Chou CC (2015) Class 1 Integrons and the antiseptic resistance gene (*qacδ1*) in municipal and swine slaughterhouse wastewater treatment plants and wastewater-associated methicillin-resistant *Staphylococcus aureus*. *International Journal of Environmental Research and Public Health* **12**, 6249–6260.
- Wang D, Zhang L, Zhou X, He Y, Yong C, Shen M, Szenci O and Han B (2016) Antimicrobial susceptibility, virulence genes, and randomly amplified polymorphic DNA analysis of *Staphylococcus aureus* recovered from bovine mastitis in Ningxia, China. *Journal of Dairy Science* **99**, 9560–9569.
- Xia W, Xu T, Qin T, Li P, Liu Y, Kang H, Gu B and Ma P (2016) Characterization of integrons and novel cassette arrays in bacteria from clinical isolates in China, 2000–2014. *Journal of Biomedical Research* **30**, 292–303.
- Zhang L, Li Y, Bao H, Wei R, Zhou Y, Zhang H and Wang R (2016) Population structure and antimicrobial profile of *Staphylococcus aureus* strains associated with bovine mastitis in China. *Microbial Pathogenesis* **97**, 103–109.
- Zhang W, Meng C, Chen H, Tu L and Zhang J (2017) Index cluster analysis of drug resistance-related genes in multidrug-resistant *Shigella flexneri*. *Chinese Journal of Laboratory Medicine* **40**, 970–979.
- Zhang L, Gao J, Barkema HW, Ali T, Liu G, Deng Y, Naushad S, Kastelic JP and Han B (2018) Virulence gene profiles, alpha-hemolysin and clonal diversity in *Staphylococcus aureus* isolates from bovine clinical mastitis in China. *BMC Veterinary Research* **14**, 63.
- Zhang X, Li X, Wang W, Qi J, Wang D, Xu L, Liu Y, Zhang Y and Guo K (2020) Diverse gene cassette arrays prevail in commensal *Escherichia coli* from intensive farming swine in four provinces of China. *Frontiers in Microbiology* **11**, 565349.