

This is an Accepted Manuscript for *International Psychogeriatrics*.

This version may be subject to change during the production process.

DOI: 10.1017/S1041610223000820

## **Predicting and promoting resilience in later life**

Lauren R Miller-Lewis<sup>1,2</sup>

<sup>1</sup>Cluster for Resilience and Wellbeing, Appleton Institute, College of Psychology, School of Health Medical and Applied Sciences, CQUniversity, Adelaide Campus, 44 Greenhill Road, Wayville, 5034 Adelaide, South Australia, Australia.

<sup>2</sup>Research Centre for Palliative Care, Death and Dying, College of Nursing and Health Sciences, Flinders University, Adelaide, South Australia, Australia.

A commentary on "Contributors to mental health resilience in middle-aged and older adults: An analysis of the Canadian Longitudinal Study on Aging" by Hopper et al., (2023).

There is something almost spellbinding about a person who flourishes despite the challenges life throws at them – what is their ‘secret’? The concept of ‘resilience’ has intrigued researchers for decades, with the seminal work with children by investigators such as Emmy Werner and Michael Rutter leading the way in helping us to understand why some people manage to survive and thrive despite being exposed to the most challenging of life circumstances (Werner & Smith, 1989; Rutter, 1979). Resilience is most-commonly defined as the occurrence of positive adaptation when exposed to significant adversity (Cosco, Howse & Brayne, 2017; Cosco, Kok et al., 2019; Hopper et al., 2023; Miller-Lewis et al., 2013). The definition specifies adversity as an inherent precondition of resilience, i.e., ‘resilience’ can only exist when there is experience of adversity. This multi-part nature of the phenomenon makes investigating resilience challenging, because these adversity-adaptation dyads (of which there can be many permutations) need to be operationalized (Cosco, Kok et al., 2019).

As eloquently put by Michael Ungar (2019, p. 2), it is “important that resilience researchers answer the question: ‘Which *promotive and protective factors or processes* are best for which people in which *contexts* at what *level of risk exposure* and for which *outcomes*?’ ”. One can see from this statement the complex nature of resilience as a concept – to truly capture ‘resilience’, a number of components need to be taken into account simultaneously. This has led to a range of

different strategies for operationalizing the construct of resilience, and there is still no gold-standard method for measuring resilience (Cosco, Kaushal et al., 2017; Costenoble et al., 2022; Chmitorz et al., 2018; Ungar et al., 2019; Wister et al., 2022). As pointed out by Costenoble et al., (2022) and others (Chmitorz et al., 2018; Cosco, Kok et al., 2019), the different approaches to operationalizing resilience might yield different results, thus clouding research findings with inconsistent conclusions. We continue to deal with the same challenging issues we were facing a decade ago when I first published in this field of resilience research (Miller-Lewis et al., 2013). But if we can get this right, the pay-off is huge in terms of the knowledge we can gain about the best tactics to increase the likelihood of resilient outcomes for people.

A considerable body of research has emerged about resilience in younger people, but there has been comparatively less attention to investigating resilience in older adults (e.g., Costenoble et al., 2022; Hopper et al., 2023; Ong et al., 2009; Wister et al., 2022). Yet understanding the way we as human beings adapt to adverse changes as we age is of vital importance, especially given the ageing population growth worldwide. Ageing comes with a set of inherent adversities humans are faced with, such as decline in physical functioning, multi-morbidity, grief and bereavement, and even acceptance of one's own eventual mortality (e.g., Wister et al., 2016; 2022). These adversities experienced in later life, in addition to the other adversities we accumulate during life, deserve consideration.

A gerontological research group based in Canada has recently been making considerable headway in furthering our understanding of resilience in older people (Cosco, Cooper et al., 2018; Cosco, Hardy et al., 2019; Wister et al., 2020). Based on older adult participants in population-based cohort studies in Canada and Britain, these studies used different strategies to investigate resources associated with resilience, yet a common thread in their findings was that stronger social support networks played a role in resilience to adversity.

The recently published study by Hopper, Best, Wister and Cosco (2023), in this issue of *International Psychogeriatrics*, adds to the growing set of investigations by this research group that is delving into predictors of resilient outcomes in older people. Hopper and colleagues' (2023) study takes advantage of data gathered from a very large population-based sample of over 30,000 Canadian adults aged over 45, to identify factors contributing to mental health resilience. They conceptualize resilient outcomes in their study as better-than-expected mental health (fewer

depressive symptoms) based on their level of objectively-measured physical performance (a composite of grip strength, sit-to-stand movements, and standing balance). They provide a good example of a clearly outlined adversity-adaptation dyad for operationalizing resilience. Hopper et al., (2023) implemented the ‘resilience residuals’ approach, which provides a quantified indicator of the extent to which a person’s health outcomes are, in a statistical sense, better-than-expected or worse-than-expected given their level of exposure to adversities (i.e., ensuring that key ‘adversity component’ within the definition of resilience is taken into account). The resilience residuals approach originated from studies of resilience in early childhood (e.g., Kim-Cohen et al., 2004; Miller-Lewis et al., 2013). In demonstration of the value of exploring research methods used in other domains for strategies that can be usefully applied in new disciplines, the residuals approach is now becoming more frequently used for studies in older adults (Cosco, Cooper et al., 2018; Hopper et al., 2023; Yu et al., 2022). The resilience residuals approach involves statistically regressing the variable representing adaptation on levels of the variable(s) representing adversity, and calculating the discrepancy between a person’s actual adaptation score and the adaptation score predicted by their level of adversity. This residual variance score can then be used as a continuous vulnerability-to-resilience score, with the size of the residual quantifying the extent of their resilience (Hopper et al., 2023; Miller-Lewis et al., 2013). Cosco, Kok et al., (2019) and Hölzge & Ungar (2022) provide useful visual depictions of resilience residuals. Recent research indicates that the residuals method is a valid and reliable approach to operationalizing resilience (Cahill et al., 2022). It is dynamic, process-oriented, and does not assume that a person demonstrates resilience at all times, during all stages of life, or in all life-circumstances. This is where the residuals approach to operationalizing resilience is fundamentally different from measurement scales that assess resilience as a stable trait-oriented characteristic.

Though studies which use measurement tools designed to capture resilience at a dispositional level may offer useful insights into resilience in older people (e.g., Costenoble et al., 2022), there are three key advantages of the residuals approach to resilience, all of which are clearly demonstrated by Hopper et al.’s (2023) study. First, the residuals approach provides, for every participant in a study, a quantified continuous resilience outcome variable that accounts for the adversity-adaptation dyadic element inherent in the concept. Second, this continuous resilience outcome variable opens the door for the examination of potential resource factors that can predict

resilient outcomes, including the exploration of more complex mechanisms through which these resource factors are interrelated and interact with each other to influence resilience. For example, the impact of distal resource factors on resilient outcomes might be mediated by more proximal resource factors. This information on the mediational processes and temporal pathways by which resources exert their effects on resilience provide valuable details about what resources should be prioritized for intervention and how they should be targeted (Cahill et al., 2022; Miller-Lewis et al., 2013). Third, the resilience residuals approach enables investigation into resilience in older individuals without the need to include an explicit resilience measurement tool in the assessment battery for the study (Cosco, Kok et al., 2019). This is particularly useful in large population cohort studies, where a number of constructs might be assessed, and a separate measure of resilience is not feasible for inclusion in the test-battery. Combining adversity-adaptation dyads to conceptualize resilience outcomes based on the existing data in cohort studies is an opportunity of untapped potential. There are numerous existing secondary datasets from population cohorts in across the world where the resilience residuals approach could be applied to available data in order to investigate promotive resources that predict resilience in older people. Hopper and colleagues (2023) have taken advantage of this opportunity using data from the population-based Canadian Longitudinal Study on Aging.

Hopper et al., (2023) hypothesized that in older people, socio-economic position, leisure-time physical activity, and social networks would be related to mental health resilience (operationalized using the residuals approach as described above). They also hypothesized that both physical activity and social network would mediate the relationship between socio-economic position and mental health resilience in middle- and older-age Canadians. In line with their hypotheses, Hopper et al.'s (2023) results demonstrated that indicators of holding a higher socio-economic position were associated with greater mental health resilience, and that participants reporting a greater frequency of leisure-based physical activity and larger social networks had greater mental health resilience. These modifiable resource factors, especially the extent of an individual's social network, were found to partially mediate the influence of socio-economic position on mental health resilience. This implies that when socio-economic position is low, one may be able to foster mental health resilience via increasing social networks and physical activity.

As the number of studies like Hopper et al.'s (2023) grow, and additional positive resources predictive of resilient outcomes in older people are discovered, they can provide valuable insights for developing intervention strategies. Hopper et al.'s (2023) findings offer us such insights regarding not just *what* to target in interventions, but also *who* to target. They conclude that interventions encouraging leisure-based physical activities and activities that bolster social connections may help build mental health resilience in older adults, and such interventions may be particularly beneficial for older people from a lower socio-economic position. Because the study is observational and cross-sectional, the authors rightly note that these recommendations are tentative and that future research is needed to infer causality in the direction of effects between variables. Yet given the paucity of resilience interventions designed for older adults (three systematic reviews of resilience interventions found none targeted at older people; Chmitorz et al., 2018; Macedo et al., 2014; MacLeod et al., 2016), these knowledge contributions are sorely needed. One recently published pragmatic trial of a group intervention to increase resilience in seniors (Treichler et al., 2020) is an important step forward, finding that a brief positive psychological intervention focused on gratitude, savoring, positive emotions, and values-based activities led to increased resilience. Also needed is an expansion beyond cross-sectional studies like Hopper et al., (2023) into prospective longitudinal studies (Cosco, Kaushal et al., 2017) capturing a greater period of the life-course and which have the capacity to demonstrate temporal precedence of resource factors prior to resilient outcomes. Capitalizing further on existing population cohort studies tracking individuals over time will be worthwhile.

The flexibility of the resilience residuals approach for conceptualizing adversity-adaptation dyads representative of resilience also opens up opportunities to further expand on Hopper et al.'s (2023) research. Different combinations of adversities and adaptations can be investigated. For example, Cosco, Kaushal and colleagues (2017) point out that most longitudinal studies measure the 'positive adaptation' component of resilience as the absence of psychopathology, with few studies measuring 'positive adaptation' with positively-oriented constructs such as mental wellbeing and life-satisfaction. There is a need for resilience research to move beyond the deficit-based conceptualizations into asset-based understandings of resilient outcomes. There are also many resource factors that could be investigated as potential contributors to mental health resilience in older people, adding to the groundwork on the socio-economic, physical activity, and social network resources considered in this initial study by Hopper et al., (2023). This might

include external resources within one's wider socio-ecological environments, and resources internal within the individual, as commonly proposed in models of resilience (e.g., Wister et al., 2016; 2022). Potential examples of internal resources worthy of further investigation are the capacity for emotional regulation and a positive outlook (Kiosses & Sachs-Ericsson, 2020; Treichler et al., 2020).

In conclusion, while the debates continue about the best way to operationalize resilience in research (Cosco, Kok et al., 2019), like Ungar's (2019) recommendations for designing childhood resilience research, I suggest that future studies with older adults ensure they clearly and explicitly outline how they have operationalized and measured the key components of resilience: (a) adversity; (b) desired adaptive outcomes; and (c) the promotive resources under investigation. Consideration should also be given to using more than one method to operationally define resilience in the same study where there is adequate power, in order to determine if findings are convergent across methods of analysis. If similar resource factors emerge as predictive of resilience from different methodological approaches, this can increase our confidence that targeting these resources with concerted intervention efforts to foster resilient outcomes in the later years of life is a worthwhile endeavor.

## ***References***

- Cahill, S., Hager, R., & Chandola, T. (2022). The validity of the residuals approach to measuring resilience to adverse childhood experiences. *Child and Adolescent Psychiatry and Mental Health, 16*(1), 18. <https://doi.org/10.1186/s13034-022-00449-y>
- Chmitorz, A., Kunzler, A., Helmreich, I., Tüscher, O., Kalisch, R., Kubiak, T., Wessa, M., & Lieb, K. (2018). Intervention studies to foster resilience – A systematic review and proposal for a resilience framework in future intervention studies. *Clinical Psychology Review, 59*, 78-100. <https://doi.org/10.1016/j.cpr.2017.11.002>
- Cosco, T., Cooper, R., Kuh, D., & Stafford, M. (2018). Socioeconomic inequalities in resilience and vulnerability among older adults: A population-based birth cohort analysis. *International Psychogeriatrics, 30*(5), 695-703. doi:10.1017/S1041610217002198

- Cosco, T., Hardy, R., Howe, L., & Richards, M. (2019). Early-life adversity, later-life mental health, and resilience resources: A longitudinal population-based birth cohort analysis. *International Psychogeriatrics*, *31*(9), 1249-1258. doi:10.1017/S1041610218001795
- Cosco, T., Howse, K., & Brayne, C. (2017). Healthy ageing, resilience and wellbeing. *Epidemiology and Psychiatric Sciences*, *26*(6), 579-583. doi:10.1017/S2045796017000324
- Cosco, T. D., Kok, A., Wister, A., & Howse, K. (2019). Conceptualising and operationalising resilience in older adults. *Health Psychology and Behavioral Medicine*, *7*(1), 90-104. <https://doi.org/10.1080/21642850.2019.1593845>
- Cosco, T. D., Kaushal, A., Hardy, R., Richards, M., Kuh, D., & Stafford, M. (2017). Operationalising resilience in longitudinal studies: A systematic review of methodological approaches. *Journal of Epidemiology and Community Health*, *71*(1), 98. <http://dx.doi.org/10.1136/jech-2015-206980>
- Costenoble, A., Rossi, G., Knoop, V., Debain, A., Smeys, C., Bautmans, I., . . . Gorus, E. (2022). Does psychological resilience mediate the relation between daily functioning and prefrailty status? *International Psychogeriatrics*, *34*(3), 253-262. doi:10.1017/S1041610221001058
- Höltge, J., & Ungar, M. (2022). Quantifying resilience as an outcome: Advancing the residual approach with influence statistics to derive more adequate thresholds of resilience. *Adversity and Resilience Science*, *3*, 381–390. <https://doi.org/10.1007/s42844-022-00078-6>
- Hopper, S., Best, J., Wister, A., & Cosco, T. (2023). Contributors to mental health resilience in middle-aged and older adults: An analysis of the Canadian Longitudinal Study on Aging. *International Psychogeriatrics*, 1-10. doi:10.1017/S1041610223000224
- Kim-Cohen, J., Moffitt, T. E., Caspi, A., & Taylor, A. (2004). Genetic and environmental processes in young children's resilience and vulnerability to socioeconomic deprivation. *Child Development*, *75*(3), 651-668. <https://doi.org/10.1111/j.1467-8624.2004.00699.x>
- Kiosses, D., & Sachs-Ericsson, N. (2020). Increasing resilience in older adults. *International Psychogeriatrics*, *32*(2), 157-159. doi:10.1017/S1041610220000046
- Macedo, T., Wilhelm, L., Gonçalves, R., Coutinho, E. S. F., Vilete, L., Figueira, I., & Ventura, P. (2014). Building resilience for future adversity: a systematic review of interventions in

non-clinical samples of adults. *BMC Psychiatry*, 14(1), 227.  
<https://doi.org/10.1186/s12888-014-0227-6>

MacLeod, S., Musich, S., Hawkins, K., Alsgaard, K., & Wicker, E. R. (2016). The impact of resilience among older adults. *Geriatric Nursing*, 37(4), 266-272.  
<https://doi.org/10.1016/j.gerinurse.2016.02.014>

Miller-Lewis, L. R., Searle, A. K., Sawyer, M. G., Baghurst, P. A., & Hedley, D. (2013). Resource factors for mental health resilience in early childhood: An analysis with multiple methodologies. *Child and Adolescent Psychiatry and Mental Health* 7, 6.  
<https://doi.org/10.1186/1753-2000-7-6>

Ong, A. D., Bergeman, C. S., & Boker, S. M. (2009). Resilience comes of age: Defining features in later adulthood. *Journal of Personality*, 77(6), 1777-1804.  
<https://doi.org/10.1111/j.1467-6494.2009.00600.x>

Rutter, M. (1979). Protective factors in children's responses to stress and disadvantage. In M. W. Kent & J. E. Rolf (Eds.), *Primary prevention of psychopathology, Volume 3: Social competence in children* (pp. 49–74). University Press of New England.

Treichler, E., Glorioso, D., Lee, E., Wu, T., Tu, X., Daly, R., . . . Jeste, D. (2020). A pragmatic trial of a group intervention in senior housing communities to increase resilience. *International Psychogeriatrics*, 32(2), 173-182.  
[doi:10.1017/S1041610219002096](https://doi.org/10.1017/S1041610219002096)

Ungar M. (2019). Designing resilience research: Using multiple methods to investigate risk exposure, promotive and protective processes, and contextually relevant outcomes for children and youth. *Child Abuse & Neglect*, 96, 104098.  
<https://doi.org/10.1016/j.chiabu.2019.104098>

Werner, E. E., & Smith, R. S. (1989). *Vulnerable but invincible: A longitudinal study of resilient children and youth*. Adams, Bannister & Cox.

Wister, A. V., Coatta, K. L., Schuurman, N., Lear, S. A., Rosin, M. and MacKey, D. (2016). A lifecourse model of multi-morbidity resilience: Theoretical and research developments.



*International Journal of Aging & Human Development*, 82(4), 290–313.  
<https://doi.org/10.1177/0091415016641686>

Wister, A., Cosco, T., Mitchell, B., & Fyffe, I. (2020). Health behaviors and multimorbidity resilience among older adults using the Canadian Longitudinal Study on Aging. *International Psychogeriatrics*, 32(1), 119-133. doi:10.1017/S1041610219000486

Wister, A., Klasa, K., & Linkov, I. (2022). A Unified Model of Resilience and Aging: Applications to COVID-19. *Frontiers in Public Health*, 10, 865459.  
<https://doi.org/10.3389/fpubh.2022.865459>

Yu, W., Zhu, F., Foo, M. D., & Wiklund, J. (2022). What does not kill you makes you stronger: Entrepreneurs' childhood adversity, resilience, and career success. *Journal of Business Research*, 151, 40-55. <https://doi.org/10.1016/j.jbusres.2022.06.035>