

CONTENTS.

Original Articles, p. 267.—Nasal Obstruction in its relation to the Administration of Anæsthetics, by Dr. J. F. W. SILK. On the Treatment of so-called Phthisis of the Larynx (continued from p. 238), by Dr. JOHN SEDZIAK (of Warsaw).

Instruments, Therapeutics, &c., p. 276.—VOLTOLINI: Illumination. GÄRTNER: A Pocket Lamp. ROTH: Electric Lighting. DIONISIO: Posterior Rhinoscopy. WÖLFLE: Cocaine Intoxication. SANDRAS: Alterations in the Voice produced by Cocaine. MAXWELL: Ammonium Chloride for Inhalation. CASH: Carbolic Acid, Antipyrin and Antifebrin. FERNET: Camphorated Naphthol in Tubercular Ulcers. DUBOVSQUET-LABORDERIE: Antipyrin in Whooping Cough. OLLIVIER: Treatment of Whooping Cough. HEYMANN: Phenacetin in Whooping Cough. SONNENBERGER: Specific Treatment of Whooping Cough. PERCEVAL: Antipyrin in Laryngismus Stridulus. ADAMS: Duration of Infectiveness of Scarlet Fever. PASTEUR: Pulmonary Gangrene. GAREL: Medical Electricity. THOMPSON: Voyage to the Cape. CHARAZAC: Sulphur Waters in Tuberculosis. TAYLOR: Ethylate of Soda in Lupus. HANSEN: Ice in Lupus.

Diphtheria, p. 280.—ROSS: Larynx and Trachea from a Case. IRVING: Epidemic Diphtheria. "BRITISH MEDICAL JOURNAL," *ibid.* WHITE: Croup. HALL, CALLIAS, LABBÉ, CARRIÈRE: Treatment. PALMER: Intubation in Diphtheria. LANGLOIS: Inhalations of Oxygen. GRELLLET: Longevity of Klebs' Bacillus.

Nose and Naso-Pharynx, p. 283.—MACDONALD: Mechanism of Nose. KERMET: Opening of Maxillary Sinus. TROUSSEAU: Nasal Ocular Reflexes. TROUSSEAU: Ozena and Corneal Ulcers. NATIER: Perforation of Septum, &c. ILLINGWORTH, RUAULT: Hay Fever. BADEN: Rhinolith. JUFFINGER: Tubercular Tumour of Nose. LABIT: Purulent Catarrh of Frontal Sinuses. ARONSOHN: Asthma Artificially Produced. SCHMIEGELOW: Asthma and Nasal Diseases. WOLKOWITSCH: Rhinoscleroma. RUAULT: Treatment of Atrophic Rhinitis. SIEBENMANN: Mycosis of Naso-Pharynx. GELLÉ: Luschka's Bursa. MICHEL: Affections of Voice and Naso-Pharyngeal Conditions. HOVELL, URUNUELA, LUC: Adenoid Vegetations.

Mouth, Tongue, Pharynx, &c., p. 288.—SUCKLING, JONES: Cancrum Oris. HUTCHINSON: Aptyalism. FOX: Epithelioma of Gum. LOCKWOOD: Epithelioma of Tongue. SUCKLING: Paralysis of Fifth, Sixth, and Third Cranial Nerves. SIMON: Neuropathic Tongue. GRIFFITH: Frederick's Ataxia. TERENCECZY: Tubercular Ulcer and Psoriasis of Tongue. GREEN: Dermoid Cyst of Tongue. SHEPHERD: Anomaly of Lingual Artery. REID: Smoker's Palate. WHETTLE: Cleft Palate. JONES, KRONACHER: Tumours of Hard Palate. SUCKLING: Peripheral Neuritis. SPICER: Hysterical Affections of Throat. SCHWARTZ: Treatment of Infectious Angina. RUAULT: Tonsillitis and Infectious Angina. NATIER: Chronic Abscess of Tonsil. SEACLIFF: Night Terrors Cured by Removal of Tonsils. MCPHEDRAN: Necrotic Tonsillitis. VERDÉS: Tonsillar Deafness. PLIQUE: Malignant Tumours of Tonsil. MASSE: Supra-Hyoid Dermoid Tumour. MACKENZIE: Ulceration of Oesophagus. BRAY: Oesophageal Obstruction. HOFFMANN: Foreign Body in Oesophagus. KLEMPERER: Cancer of Oesophagus.

Larynx, Trachea, &c., p. 293.—HOWARD, FOULIS, SMYTH: Raising the Epiglottis. STOKES: Hypertrophy of Epiglottis. DUKE: Position of Epiglottis in Inversion. AUDUBERT: Different Aspects of Epiglottis. ZIEGLER: Primary Erysipelas. LUC: Treatment of Laryngeal Tuberculosis. GOUQUENHEIM and TISSIER, CARTAZ: Tubercular Tumours. GRÜNWARD: Tubercular Abscess of Cricoid. TISSIER: Tertiary Syphilis. GNODY: New Experimental Method. CASADESUS: Hæmorrhagic Laryngitis. ZALESKI: Aluminium Canulas in Tracheotomy. THOMSON: Tracheotomy for Foreign Body. CARRIÈRE: Performance of Tracheotomy. LAFLEUR, MCPHEDRAN: Aneurism of Aorta. FOWLER: Aortic Stenosis with Tracheal Murmurs. ONODY: Argyria and Cancer. BEECHAM, FUNDARENA: Laryngeal Growths. SIMON: Spasms of Glottis. JUFFINGER, ORTEGA: Foreign Bodies in Air Passages. BALL: Hysterical Affections of Vocal Cords. SCHEINMANN, MILSOM: Hysterical Aphonia treated by Suggestion. PEYRISSAC: Hysterical Aphonia in a Child. BRESGEN: Hoarseness. PILTAN: Recording Movements of Vocalisation. BENNETT: Rupture of Trachea. MURRAY: Fracture of Larynx. SMITH: Perichondritis. BECKER: Laryngo-fissure.

Thyroid Gland, Neck, &c., p. 303.—ROBINSON: Cystic Bronchocele. STOKER, ROBSON: Goitre. DUNCAN: Electrolysis in Goitres. BROWNE: Partial Removal of Goitre. POLOSSON: Constricting Goitre. POLHERAT: Intra-thoracic Goitre. GUYOT, GÉRIN, PORTER: Exophthalmic Goitre. HARSANT: Cancer of Thyroid. MORTIMER: Early Myxœdema. SHEPHERD: Inferior Thyroid Arteries. SHEPHERD: Large Thyroidea Ima. DILL: Accessory Thyroid. SPANTON: Tumours of Neck. MEYER: Blood Cysts of Neck. ANNANDALE: Pathology and Treatment of Strumous Glands. SUTHERLAND: Ligature of Common Carotid. LLOYD: Congenital Gumma. CARPENTER: Wry Neck. SAUNDBY: Acromegaly. BOND: Surgery of Parotid. MORGAN: Frontal Meningocele. BARKER: Inoculation of Syphilis by Tattooing.

Reports of Societies, p. 310.—The British Laryngological and Rhinological Association.

New Preparations, p. 310.—Kronenquelle Water. Jeyes' Fluid. Vinolia Cream and Soap. Pure Hordeum Malt Extract.

PEPTONISED MILK IS SWEET AND PALATABLE

When prepared as directed with

ZYMINE (EXTRACTUM PANCREATIS) FAIRCHILD.

ITS PREPARATION

COSTS LESS THAN A PENNY A PINT.

ZYMINE (EXTRACTUM PANCREATIS), FAIRCHILD is a *dry powder*, containing in the most active and concentrated form all the Enzymes of the Pancreas, viz.: TRYPSIN, AMYLOPSIN, STEAP SIN, and the RENNET FERMENT.

IT WILL DIGEST ALL KINDS OF FOOD: 5 grains, with a little soda, will sufficiently peptonise a pint of milk in a few minutes; 30 grains, with a little soda, will peptonise 4 ozs. of beef, producing a concentrated, nutritious and delicious beef-tea.

ZYMINE PEPTONISING POWDER (FAIRCHILD) in glass tubes, is the most convenient form for preparing peptonised milk, gruels, jellies, custards, blanc-manges, etc., etc.

THE INSTRUCTIONS given in one of our direction slips enable even *inexperienced domestics* to quickly prepare any peptonised food.

FOR NUTRITIVE ENEMATA, any food can be thoroughly pre-digested with Zymine (Extractum Pancreatis); it is then readily absorbed by the rectum.

THE TABLOIDS OF ZYMINE (EXTRACTUM PANCREATIS) afford the most convenient form for direct internal administration. They are pleasing, and can easily be carried about in the pocket. Now extensively used as a food aid in intestinal derangements.

While the Peptonising Powders are the most *elegant* and *convenient* form for peptonising milk, they are somewhat more expensive than the simple Zymine; where cheapness is of first importance the latter may therefore be used.

Zymine (Extractum Pancreatis) supplied in $\frac{1}{2}$ oz. and 1 oz. bottles, at 1s. 8d. and 5s. each.

Zymine Peptonising Powders supplied in boxes of twelve tubes. One tube peptonises a pint of milk, 1s. 6d. per box.

Zymine Tabloids, 3 grs. each, supplied in bottles of 25 and 100, at 1s. 6d. and 4s. each.

BURROUGHS, WELLCOME & CO., Manufacturing Chemists, Snow Hill Buildings, E.C.,

Will be happy to supply trial Specimens to Medical Men free on request.