



Turnpike in infinite dimension

Paolo Leonetti and Michele Caprio

Abstract. Let Φ be a correspondence from a normed vector space X into itself, let $u : X \rightarrow \mathbf{R}$ be a function, and let \mathcal{J} be an ideal on \mathbf{N} . In addition, assume that the restriction of u on the fixed points of Φ has a unique maximizer η^* . Then, we consider feasible paths (x_0, x_1, \dots) with values in X such that $x_{n+1} \in \Phi(x_n)$, for all $n \geq 0$. Under certain additional conditions, we prove the following turnpike result: every feasible path (x_0, x_1, \dots) which maximizes the smallest \mathcal{J} -cluster point of the sequence $(u(x_0), u(x_1), \dots)$ is necessarily \mathcal{J} -convergent to η^* .

We provide examples that, on the one hand, justify the hypotheses of our result and, on the other hand, prove that we are including new cases which were previously not considered in the related literature.

1 Introduction

Let X be a normed real vector space, and fix a correspondence Φ from X into itself and a functional $u : X \rightarrow \mathbf{R}$ which may be interpreted as a utility function. Then, a sequence $\mathbf{x} = (x_0, x_1, \dots)$ with values in X is said to be feasible if $x_{n+1} \in \Phi(x_n)$, for all $n \geq 0$. Note that this sequence is simply the orbit of the starting point x_0 if Φ is singleton-valued. Fix also an ideal \mathcal{J} on the nonnegative integers \mathbf{N} , which will represent the family of “small” sets (see Section 1.1 for definitions). Assuming that \mathbf{x} belongs to a given constraint set \mathcal{C} of feasible sequences, we say that \mathbf{x} is \mathcal{J} -optimal if it maximizes the smallest \mathcal{J} -cluster point of the real sequence $(u(x_0), u(x_1), \dots)$; here, an \mathcal{J} -cluster point is, informally, an accumulation point which is not small with respect to \mathcal{J} .

Our aim is to study the asymptotic stability of \mathcal{J} -optimal paths, which is often referred to as *turnpike property* (see, e.g., [26, 27] for a textbook exposition). Roughly, this property states that an \mathcal{J} -optimal path spends “most” of the time within a small neighborhood of some optimal stationary point, which is an identified fixed point of Φ . Here, following the same lines of [5, 20, 25], the adjective “most” is intended with respect to the ideal \mathcal{J} . In particular, \mathcal{J} -optimal paths are potentially not convergent to the optimal stationary point. As remarked in [26], the turnpike property has the following interpretation: if one is looking for an optimal way to reach A from B by car, then he should enter onto a turnpike, spend most of the time there, and finally leave

Received by the editors March 2, 2021; revised May 21, 2021; accepted May 23, 2021.

Published online on Cambridge Core June 3, 2021.

P.L. is grateful to PRIN 2017 (grant 2017CY2NCA) for financial support. M.C. would like to acknowledge funding from NSF CCF-1934964.

AMS subject classification: 37C70, 37C99, 40A35, 40A05.

Keywords: Turnpike, ideal and statistical convergence, ideal cluster point, optimal stationary point, fixed point of correspondences.

the turnpike to reach the claimed point. There is an extensive literature which studies this phenomenon (see, e.g., [3, 4, 12, 18, 21, 23]).

Our main result (Theorem 2.1) generalizes the main ones obtained in [5, 20]. We discuss later how our assumptions are related to the ones in these articles. In addition, we show with some novel examples that:

- (i) The turnpike property provided in Theorem 2.1 does not hold without any restriction on the ideal \mathcal{J} (see Example 2.3);
- (ii) An \mathcal{J} -optimal path may not converge, in the classical sense, to the optimal stationary point (see Example 2.5);
- (iii) The turnpike property holds also in infinite dimension (see Example 5.1).

Our main result and its consequences follow in Section 2.

1.1 Preparation

An ideal $\mathcal{J} \subseteq \mathcal{P}(\mathbf{N})$ is a family closed under finite union and subsets. It is also assumed that \mathcal{J} contains the family of finite sets Fin and it is different from $\mathcal{P}(\mathbf{N})$. Let also $\mathcal{J}^* := \{S \subseteq \mathbf{N} : S^c \in \mathcal{J}\}$ be its dual filter and $\mathcal{J}^+ := \{S \subseteq \mathbf{N} : S \notin \mathcal{J}\}$ be the collection of \mathcal{J} -positive sets. We denote by \mathcal{Z} the ideal of asymptotic density zero sets, i.e.,

$$\mathcal{Z} = \{A \subseteq \mathbf{N} : |\{a \in A : a \leq n\}| = o(n) \text{ as } n \rightarrow \infty\}.$$

An ideal \mathcal{J} on \mathbf{N} is said to be *translation invariant* if $(A + k) \cap \mathbf{N} \in \mathcal{J}$ for all $A \in \mathcal{J}$ and all (possibly negative) integers $k \in \mathbf{Z}$. Note that the ideal \mathcal{Z} is translation invariant. Classes of translation invariant ideals have been widely studied (see, e.g., [6, 9] and the ideals generated by the upper densities considered in [17]). However, there exist ideals which are not translation invariant: for instance, all the maximal ideals (indeed, exactly one between the even and the odd integers belongs to a maximal ideal) and much simpler ones as the family of all sets $A \subseteq \mathbf{N}$ containing finitely many even integers.

Let $\mathbf{x} = (x_n)$ be a sequence taking values in a topological vector space S . Then, we say that \mathbf{x} is \mathcal{J} -convergent to $\eta \in S$, shortened as $\mathcal{J}\text{-}\lim \mathbf{x} = \eta$, if $\{n \in \mathbf{N} : x_n \in U\} \in \mathcal{J}^*$ for all open neighborhoods U of η . Moreover, we say that $\eta \in S$ is an \mathcal{J} -cluster point of \mathbf{x} if $\{n \in \mathbf{N} : x_n \in U\} \in \mathcal{J}^+$ for all open neighborhoods U of η . The set of \mathcal{J} -cluster points of \mathbf{x} is denoted by $\Gamma_{\mathbf{x}}(\mathcal{J})$. Usually, \mathcal{Z} -convergence and \mathcal{Z} -cluster points are referred to as *statistical convergence* and *statistical cluster points*, respectively (see, e.g., [10, 24]). Note that Fin -convergence coincides with the ordinary convergence and that $\Gamma_{\mathbf{x}}(\text{Fin})$ is the set of ordinary accumulation points of \mathbf{x} . It is worth noting that \mathcal{J} -cluster points have been studied much before under a different name. Indeed, as it follows by [16, Theorem 4.2] and [14, Lemma 2.2], they correspond to classical “cluster points” of a filter (depending on \mathbf{x}) on the underlying space (cf. [2, Definition 2, p. 69]).

Finally, following [11], for each real sequence \mathbf{x} such that $\{n \in \mathbf{N} : |x_n| \geq M\} \in \mathcal{J}$, for some $M \in \mathbf{R}$, we define its \mathcal{J} -limit inferior as

$$\mathcal{J}\text{-}\lim \inf \mathbf{x} := \inf\{r \in \mathbf{R} : \{n \in \mathbf{N} : x_n > r\} \in \mathcal{J}^+\}.$$

Simmetrically, we let $\mathcal{J}\text{-}\lim \sup \mathbf{x} := -\mathcal{J}\text{-}\lim \inf(-\mathbf{x})$ be the \mathcal{J} -limit superior. Again, it is easy to see that if $\mathcal{J} = \text{Fin}$, then they coincide with the ordinary limit inferior and

limit superior of \mathbf{x} , respectively. It is remarkable that they can be rewritten also as the smallest and the biggest \mathcal{J} -cluster points of \mathbf{x} , respectively (cf. Corollary 3.3 below).

Given sets A, B , we say that $\alpha : A \rightrightarrows B$ is a correspondence if $\alpha(x)$ is a (possibly empty) subset of B for each $x \in A$. Moreover, we denote the set of its fixed points by

$$\text{Fix}(\alpha) := \{x \in A : x \in \alpha(x)\}.$$

We recall that, if A and B are endowed with some topologies, then the correspondence α is upper hemicontinuous at $x \in A$ if for each open $U \supseteq \alpha(x)$, there exists an open neighborhood V of x such that $z \in V$ implies $\alpha(z) \subseteq U$. Moreover, α is lower hemicontinuous at $x \in A$ if for every open $U \subseteq B$ with $\alpha(x) \cap U \neq \emptyset$, there exists an open neighborhood V of x such that $z \in V$ implies $\alpha(z) \cap U \neq \emptyset$. Finally, the correspondence α is said to be *continuous* if it is both upper and lower hemicontinuous at each point $x \in A$ (see [1, Definition 17.2]).

Given a function $h : A \rightarrow B$ and a sequence $\mathbf{x} = (x_0, x_1, \dots)$ with values in A , we write $h(\mathbf{x})$ for the sequence $(h(x_0), h(x_1), \dots)$. Finally, if $B = \mathbf{R}$, we say that $x_0 \in A$ is a *maximizer* of h if $h(x) \leq h(x_0)$, for all $x \in A$ (and similarly for minimizers).

2 Main result

Let X be a real normed vector space and denote by \mathcal{K} the collection of its nonempty compact subsets. In addition, let \mathcal{J} be an ideal on the nonnegative integers \mathbf{N} , and fix a correspondence $\Phi : X \rightrightarrows X$ and a function $u : X \rightarrow \mathbf{R}$. In this setting, the function u will take the role of a utility function which induces a total preorder on X .

Let \mathcal{K} be the family of sequences $\mathbf{x} = (x_0, x_1, \dots)$ taking values in X which are \mathcal{J} -contained in a compact, that is, such that $\{n \in \mathbf{N} : x_n \notin K\} \in \mathcal{J}$, for some $K \in \mathcal{K}$. Moreover, we let \mathcal{F} be the collection of *feasible paths* \mathbf{x} which satisfy $x_{n+1} \in \Phi(x_n)$, for all n , that is,

$$\mathcal{F} = \{\mathbf{x} \in X^{\mathbf{N}} : \forall n \in \mathbf{N}, x_{n+1} \in \Phi(x_n)\}.$$

It is easy to see that, if u is continuous, then $\mathcal{J}\text{-lim inf } u(\mathbf{x})$ and $\mathcal{J}\text{-lim sup } u(\mathbf{x})$ are well defined for each sequence $\mathbf{x} \in \mathcal{F}_K$, where

$$\mathcal{F}_K := \mathcal{F} \cap \mathcal{K}$$

(cf. Section 3). Fix also a nonempty subset $\mathcal{C} \subseteq \mathcal{F}_K$, which will take the role of the collection of constraints. Note that the primitive elements of this system are represented by the tuple $\langle X, \Phi, u, \mathcal{J}, \mathcal{C} \rangle$. Finally, we say that a sequence $\mathbf{x} \in \mathcal{C}$ is *\mathcal{J} -optimal* if

$$(2.1) \quad \forall \mathbf{y} \in \mathcal{C}, \quad \mathcal{J}\text{-lim inf } u(\mathbf{x}) \geq \mathcal{J}\text{-lim inf } u(\mathbf{y}).$$

In other words, an \mathcal{J} -optimal path \mathbf{x} is a maxmin solution in a precise sense: it maximizes the minimal \mathcal{J} -cluster point of the sequence $(u(y_0), u(y_1), \dots)$ among all feasible paths \mathbf{y} in the constraint set \mathcal{C} (cf. Corollary 3.3 below).

The aim of this work, in the same spirit of [5, 20, 21, 25], is to find sufficient conditions on the system $\langle X, \Phi, u, \mathcal{J}, \mathcal{C} \rangle$ such that every \mathcal{J} -optimal path is necessarily \mathcal{J} -convergent to some identified fixed point of Φ (in this setting, a fixed point of Φ is

usually called *stationary point*). We are going to show that certain feasible paths satisfy this property whenever the following conditions on $\langle X, \Phi, u, \mathcal{J}, \mathcal{C} \rangle$ hold:

- (A1) Φ is continuous and takes values in \mathcal{K} ;
- (A2) u is continuous;
- (A3) \mathcal{J} is translation invariant;
- (A4) There exists a unique $\eta^* \in \text{Fix}(\Phi)$ which maximizes the restriction of u on $\text{Fix}(\Phi)$;
- (A5) There exists a continuous linear functional $T : X \rightarrow \mathbf{R}$ such that

$$(2.2) \quad \forall x \in F, \forall y \in \Phi(x), \quad Tx \leq Ty \implies x = y = \eta^*,$$

where $F := \{x \in X : u(x) \geq u(\eta^*)\}$;

- (A6) $\sup_{y \in \mathcal{C}} \mathcal{J}\text{-lim inf } u(y) \geq u(\eta^*)$.

We remark that condition (A1) is equivalent to the fact that the function $X \rightarrow \mathcal{K}$ defined by $x \mapsto \Phi(x)$ is continuous with respect to the Hausdorff metric (see [1, Theorem 17.15]).

Note that the separation property given in condition (A5) has been already used in [5, 20] for the case $X = \mathbf{R}^n$, replacing (2.2) with the weaker variant

$$(2.3) \quad \forall x \in F, \forall y \in \Phi(x), \quad Tx \leq Ty \implies x = \eta^*.$$

However, a careful analysis of their proofs reveals that, in fact, they were both implicitly using (2.2). Indeed, as pointed out by Piotr Szuca in a private communication [22], condition (2.3) is, in fact, *not* sufficient for their purposes. On this direction, see also Remark 4.1 below. Condition (2.3) appeared also in [19] in the study of turnpike theorems for integral functionals in a continuous time setting. A somehow related condition can be found in [21, Lemma 4.3].

Our main result follows.

Theorem 2.1 *Let $\langle X, \Phi, u, \mathcal{J}, \mathcal{C} \rangle$ be a system which satisfies conditions (A1)–(A6). Let also $\mathbf{x} \in \mathcal{C}$ be an \mathcal{J} -optimal path. Then, $\mathcal{J}\text{-lim } \mathbf{x} = \eta^*$.*

It is worth to remark that, differently from most of the literature on turnpike theorems, we do not assume neither the concavity of the utility function u nor the convexity of the images $\Phi(x)$, for each $x \in X$. Moreover, a sufficient condition to imply condition (A6) is the existence of a sequence $\mathbf{y} \in \mathcal{C}$ which is \mathcal{J} -convergent to η^* , which gives us the following corollary.

Corollary 2.2 *Let $\langle X, \Phi, u, \mathcal{J}, \mathcal{C} \rangle$ be a system which satisfies conditions (A1)–(A5) and suppose that there exists $\mathbf{y} \in \mathcal{C}$ such that $\mathcal{J}\text{-lim } \mathbf{y} = \eta^*$. Let also $\mathbf{x} \in \mathcal{C}$ be an \mathcal{J} -optimal path. Then, $\mathcal{J}\text{-lim } \mathbf{x} = \eta^*$.*

Corollary 2.2 generalizes the main results obtained in [5, 20]. Indeed, in [20], Mamedov and Pehlivan assumed, in addition, that: X is the finite dimensional vector space \mathbf{R}^k , F is compact, there exists a compact set containing (the image of) each feasible sequence in \mathcal{F}_K , the set of constraints \mathcal{C} depends on the sequence \mathbf{x} , so that \mathcal{C} is of the type $\{\mathbf{z} \in \mathcal{F}_K : x_0 = z_0\}$, and there exists $\mathbf{y} \in \mathcal{C}$ which is convergent to η^* (in

the place of the weaker assumption of \mathcal{J} -convergence). The same hypotheses have been also used by Das et al. in [5], where the authors considered certain correspondences $\Phi : \mathbf{R}^k \rightrightarrows \mathbf{R}^k$ such that $\Phi(x) = \{h(x, y) : y \in U\}$, for all $x \in \mathbf{R}^k$, where $h : \mathbf{R}^k \rightarrow \mathbf{R}^m$ is a continuous function and $U \subseteq \mathbf{R}^m$ is a fixed nonempty compact set (it is routine to show that all correspondences Φ of this type are continuous). Finally, Mamedov and Pehlivan [20] focused on the case $\mathcal{J} = \mathcal{Z}$. Hence, Corollary 2.2 proves that all these assumptions are not really needed.

In the next example, we show that Corollary 2.2 (and, hence, also Theorem 2.1) cannot be extended to all the ideals \mathcal{J} .

Example 2.3 Let \mathcal{J} be an ideal on \mathbf{N} such that $2\mathbf{N} \subseteq \mathcal{J}$. Note that such ideals exist, e.g., the family of subsets of \mathbf{N} containing finitely many odd integers, or the maximal ideals extending $2\mathbf{N}$. Now, let $X = \mathbf{R}$, and define $\Phi(x) = \{-x, x/2\}$ and $u(x) = x^3$, for each $x \in \mathbf{R}$. Moreover, set $\mathcal{C} := \{y \in \mathcal{F}_{\mathbf{K}} : y_0 = 1\}$. Then, the continuous correspondence Φ has a unique fixed point, i.e., $\text{Fix}(\Phi) = \{0\}$. It is easily seen that the system $\langle X, \Phi, u, \mathcal{J}, \mathcal{C} \rangle$ satisfies conditions (A1)–(A4). Moreover, also condition (A5) holds: indeed, notice that $F = \{x \in \mathbf{R} : u(x) \geq u(0)\} = [0, \infty)$. Then, setting $T(r) = r$, for all $r \in \mathbf{R}$, we obtain that $Ty < Tx$ for all $(x, y) \neq (0, 0)$ with $x \in F$ and $y \in \Phi(x)$, i.e., for all $x > 0$ and $y \in \{-x, x/2\}$.

At this point, let $\mathbf{x} = (x_0, x_1, \dots) \in \mathcal{C}$ be the sequence defined by $x_n = (-1)^n$, for all $n \in \mathbf{N}$. Then, $\mathbf{x} \in \mathcal{C}$ and $u(\mathbf{x}) = \mathbf{x}$, so that $\mathcal{J}\text{-lim } u(\mathbf{x}) = 1$. Because $|u(y_n)| \leq 1$, for all $y \in \mathcal{C}$ and $n \in \mathbf{N}$, it follows that \mathbf{x} is \mathcal{J} -optimal. In addition, the sequence \mathbf{y} defined by $y_n = 2^{-n}$, for all $n \in \mathbf{N}$, belongs to \mathcal{C} , and it is convergent (in the classical sense) to 0. Hence, all hypotheses of Corollary 2.2 hold. However, the sequence \mathbf{x} is clearly not \mathcal{J} -convergent to 0.

Note that the same construction given in Example 2.3 does not contradict Corollary 2.2 in the case that \mathcal{J} is a translation invariant ideal. Indeed, in such case, $2\mathbf{N} \in \mathcal{J}$ if and only if $2\mathbf{N} + 1 \in \mathcal{J}$. However, because their union is \mathbf{N} , then they are both \mathcal{J} -positive sets, so that $\Gamma_{u(\mathbf{x})}(\mathcal{J}) = \{1, -1\}$ and $\mathcal{J}\text{-lim inf } u(\mathbf{x}) = -1$. Therefore, \mathbf{x} would be not \mathcal{J} -optimal. Moreover, this provides an example of a system $\langle X, \Phi, u, \mathcal{J}, \mathcal{C} \rangle$ such that u is neither concave nor convex, Φ is not convex-valued, and F is not compact.

Suppose now that Φ is singleton-valued, that is, there exists a function $\phi : X \rightarrow X$ such that $\Phi(x) = \{\phi(x)\}$, for all $x \in X$. Note that the continuity of Φ in condition (A1) is equivalent to the continuity of ϕ (see [1, Lemma 17.6]). Here, let us identify Φ with ϕ . Notice also that a feasible sequence $\mathbf{x} \in \mathcal{F}$ is simply an orbit $(x_0, \phi(x_0), \phi^2(x_0), \dots)$. Hence, the constraint set \mathcal{C} can be identified with the set of starting values $C := \{x \in X : \exists \mathbf{x} \in \mathcal{C}, x_0 = x\}$. To sum up, the system can be identified with the tuple $\langle X, \phi, u, \mathcal{J}, C \rangle$, and a sequence \mathbf{x} with $x_0 \in C$ is \mathcal{J} -optimal provided that

$$\forall y \in C, \quad \mathcal{J}\text{-lim inf}_n u(\phi^n(x_0)) \geq \mathcal{J}\text{-lim inf}_n u(\phi^n(y)),$$

where $\phi^0(x) := x$, for all $x \in X$. With these premises, we have the following corollary.

Corollary 2.4 Let $\langle X, \phi, u, \mathcal{J}, C \rangle$ be a system which satisfies conditions (A1)–(A5), and suppose that there exists $y_0 \in C$ such that $\liminf_n u(\phi^n(y_0)) \geq u(\eta^*)$. Fix also $x_0 \in X$ such that the orbit $(\phi^n(x_0))$ is \mathcal{J} -optimal. Then, $\mathcal{J}\text{-lim}_n \phi^n(x_0) = \eta^*$.

At this point, one may wonder if the standing assumptions of Theorem 2.1 together with the \mathcal{J} -optimality of the sequence \mathbf{x} imply the stronger conclusion that $\lim \mathbf{x} = \eta^*$, so that, in a sense, it would be not necessary to speak about ideals. In the next example, we show that this is not the case. Indeed, there exists a system $\langle X, \Phi, u, \mathcal{J}, \mathcal{C} \rangle$ which satisfies conditions (A1)–(A6) and an \mathcal{J} -optimal sequence $\mathbf{x} \in \mathcal{C}$ which is not convergent in the ordinary sense (however, thanks to Theorem 2.1, it is \mathcal{J} -convergent to η^*).

Example 2.5 Set $X = \mathbf{R}$, $\mathcal{J} = \mathcal{Z}$, $\mathcal{C} = \mathcal{F}_{\mathbf{K}}$, $\Phi(x) := [-2x, -\frac{x}{2}]$, and $u(x) = x$, for all $x \in \mathbf{R}$. It is not difficult to see that conditions (A1)–(A6) hold, and $\text{Fix}(\Phi) = \{0\}$ (cf. Example 2.3).

At this point, let $\mathbf{x} = (x_0, x_1, \dots)$ be the sequence for which for all n $x_n = (-1)^n z_n$, where $\mathbf{z} = (z_0, z_1, \dots)$ is defined as it follows:

$$\begin{aligned} & \overbrace{(1, 1/2, 1, 1/2, 1/4, 1/4, 1/2, \dots)}^{B_1}, \quad \overbrace{(1, 1/2, 1, 1/2, 1/4, 1/4, 1/2, \dots)}^{B_2}, \dots, \\ & \overbrace{(1, 1/2, 1/4, \dots, 1/2^{k-1}, 1/2^k, 1/2^k, \dots, 1/2^k, 1/2^{k-1}, 1/2^{k-2}, \dots, 1/2, \dots)}^{B_k}. \end{aligned}$$

$k!$ times

Here, for each $k \geq 1$, the block B_k has $2k - 1 + k!$ terms and its middle part is made by $k!$ consecutive terms equal to $1/2^k$.

Let us show that \mathbf{x} is \mathcal{Z} -optimal: first of all, using the fact that

$$\sum_{k \leq n-1} |B_k| \leq \sum_{k \leq n-1} 3 \cdot (k-1)! = o(n!) \text{ as } n \rightarrow \infty,$$

thus \mathcal{Z} - $\lim \mathbf{x} = 0$ (cf. also [17, Lemma 1]). In particular, \mathcal{Z} - $\lim \inf \mathbf{x} = \mathcal{Z}$ - $\lim \inf u(\mathbf{x}) = 0$. Let us suppose, for the sake of contradiction, that there exists a sequence $\mathbf{y} \in \mathcal{F}_{\mathbf{K}}$ such that $\kappa := \mathcal{J}$ - $\lim \inf \mathbf{y} > 0$. Hence, κ is a statistical cluster point of \mathbf{y} (see Corollary 3.3 below). It follows that

$$A := \{n \in \mathbf{N} : y_n > \kappa/2\} \in \mathcal{J}^+.$$

However, by construction, we have that $y_n y_{n+1} < 0$ whenever $y_n \neq 0$. Therefore, $y_{n+1} < -\kappa/4$, for all $n \in A$. Considering that \mathcal{Z} is a translation invariant ideal, it follows that \mathbf{y} is a bounded sequence such that $\{n \in \mathbf{N} : y_n < -\kappa/4\} \supseteq A + 1 \in \mathcal{J}^+$. We conclude by Lemma 3.1(ii) below that the sequence \mathbf{y} has a negative statistical cluster point, contradicting the standing hypothesis that \mathcal{J} - $\lim \inf \mathbf{y} > 0$.

Hence, \mathbf{x} is \mathcal{Z} -optimal. However, because the length of block B_k is odd for each $k \geq 2$, it follows that $\lim \inf \mathbf{x} = -1$ and $\lim \sup \mathbf{x} = 1$.

With these premises, we give below a practical application of our main result in the context of (correspondences generated by) iterated function systems, a basic tool in fractal geometry (see, e.g., [8]). Additional examples can be found also in [22].

Example 2.6 Set $X = \mathbf{R}$, let \mathcal{J} be a translation invariant ideal on \mathbf{N} , and let $u : \mathbf{R} \rightarrow \mathbf{R}$ be a strictly increasing continuous function. In addition, let $\{\phi_1, \dots, \phi_k\}$ be an

iterated function system on \mathbf{R} , that is, a finite number of contractions on \mathbf{R} , and define the correspondence $\Phi : \mathbf{R} \rightrightarrows \mathbf{R}$ by

$$\forall x \in \mathbf{R}, \quad \Phi(x) := \{\phi_1(x), \dots, \phi_k(x)\}.$$

Accordingly, let \mathcal{C} be an arbitrary subset of bounded feasible sequences such that

$$(2.4) \quad \forall i = 1, \dots, k, \exists x \in \mathbf{R}, \quad (x, \phi_i(x), \phi_i^2(x), \dots) \in \mathcal{C}.$$

(It is remarkable that there exists a unique nonempty compact set $S \subseteq \mathbf{R}$, called *attractor*, such that $\lim_n H^n(\{x_0\}) = S$, for all $x_0 \in \mathbf{R}$, in the Hausdorff metric, where H stands for Hutchinson operator defined by $H(A) := \bigcup_{x \in A} \Phi(x)$, for all $A \subseteq \mathbf{R}$; see [13].)

For each $i = 1, \dots, k$, let η_i be the fixed point of ϕ_i , hence the restriction of u on $\text{Fix}(\Phi) = \{\eta_1, \dots, \eta_k\}$ is maximized at the unique point $\eta^* = \max\{\eta_1, \dots, \eta_k\}$. In particular, conditions (A1)–(A4) hold. At this point, note that $F = [\eta^*, \infty)$ and

$$\forall i = 1, \dots, k, \forall \eta > \eta_i, \quad \phi_i(\eta) - \eta_i \leq |\phi_i(\eta) - \phi_i(\eta_i)| < \eta - \eta_i,$$

hence $\phi_i(\eta) < \eta$ whenever $\eta > \eta_i$. In particular, $\max \Phi(\eta^*) = \eta^*$ and $\max \Phi(\eta) < \eta$, for all $\eta > \eta^*$. Therefore, condition (A5) holds letting T be the identity map. Finally, let j be an index such that $\eta_j = \eta^*$. Then, it follows by (2.4) and the Banach contraction theorem that there exists $x \in \mathbf{R}$ such that $(x, \phi_j(x), \phi_j^2(x), \dots) \in \mathcal{C}$ and $\lim_n \phi_j^n(x) = \eta^*$.

We conclude by Corollary 2.2 that, if a sequence \mathbf{x} in the constraint set \mathcal{C} is \mathcal{J} -optimal, then $\mathcal{J}\text{-lim } \mathbf{x} = \eta^*$. In particular, in the special case $\mathcal{C} = \mathcal{F}_K$, $u(x) = x$, and $\mathcal{J} = \text{Fin}$, we obtain that: if a bounded feasible sequence \mathbf{x} maximizes its smallest accumulation point, then it is convergent to the maximal fixed point η^* (this could be obtained, of course, also by a direct method.)

As a last motivation for the assumptions given in Theorem 2.1, we provide in Section 5 an example where our main result holds in an infinite dimensional vector space X (we postpone it because of its length). The proofs of our results are given in Section 4.

3 Preliminaries on \mathcal{J} -cluster points

We collect in the next lemma the basic properties of \mathcal{J} -cluster points and \mathcal{J} -convergence. These properties hold in greater generality, which we do not require here.

Lemma 3.1 *Let \mathbf{x} be a sequence taking values in a metric space S and fix an ideal \mathcal{J} . Then:*

- (i) $\Gamma_{\mathbf{x}}(\mathcal{J})$ is closed;
- (ii) $\Gamma_{\mathbf{x}}(\mathcal{J}) \cap K \neq \emptyset$, provided that there exists a compact $K \subseteq S$ such that $\{n \in \mathbf{N} : x_n \in K\} \in \mathcal{J}^+$;
- (iii) $\mathcal{J}\text{-lim } \mathbf{x} = \eta$ implies $\Gamma_{\mathbf{x}}(\mathcal{J}) = \{\eta\}$;

- (iv) \mathcal{J} -lim $\mathbf{x} = \eta$ if and only if $\Gamma_{\mathbf{x}}(\mathcal{J}) = \{\eta\}$, provided that there exists a compact $K \subseteq S$ such that $\{n \in \mathbf{N} : x_n \in K\} \in \mathcal{J}^*$;
- (v) $\Gamma_{\mathbf{x}}(\mathcal{J})$ is the smallest closed set C such that $\{n \in \mathbf{N} : x_n \in U\} \in \mathcal{J}^*$ for all open sets $U \supseteq C$, provided that there exists a compact $K \subseteq S$ such that $\{n \in \mathbf{N} : x_n \in K\} \in \mathcal{J}^*$.

Proof See [16, Lemma 3.1, Corollaries 3.2 and 3.4, and Theorem 4.3]. ■

To the best of our knowledge, the following result is the first one of this type, even if some consequences were known (cf. Corollary 3.3 below). Informally, it states that, for each sequence contained in a compact, the set of \mathcal{J} -cluster points of its continuous image coincides with the continuous image of its \mathcal{J} -cluster points.

Proposition 3.2 *Let S, S' be metric spaces, and let \mathcal{J} be an ideal. Fix also a continuous function $h : S \rightarrow S'$, and let \mathbf{x} be a sequence with values in S such that $\{n \in \mathbf{N} : x_n \in K\} \in \mathcal{J}^*$ for some compact $K \subseteq S$. Then, $h(\Gamma_{\mathbf{x}}(\mathcal{J})) = \Gamma_{h(\mathbf{x})}(\mathcal{J})$.*

Proof First, suppose that $\eta \in \Gamma_{\mathbf{x}}(\mathcal{J})$. Then, it follows by the continuity of h that $h(\eta) \in \Gamma_{h(\mathbf{x})}(\mathcal{J})$: indeed, for each open neighborhood U of $h(\eta)$, there exists an open neighborhood V of η such that $\{n \in \mathbf{N} : x_n \in V\} \subseteq \{n \in \mathbf{N} : h(x_n) \in U\}$. Hence, $h(\Gamma_{\mathbf{x}}(\mathcal{J})) \subseteq \Gamma_{h(\mathbf{x})}(\mathcal{J})$.

Conversely, suppose that $v \in \Gamma_{h(\mathbf{x})}(\mathcal{J})$. Note that $F := h^{-1}(\{v\})$ is closed and that $\{n \in \mathbf{N} : h(x_n) \in H\} \in \mathcal{J}^*$, where $H := h(K)$ is compact. Because v belongs to H , then $K_0 := F \cap K$ is a nonempty compact set. We claim that there exists $\eta \in K_0$ which is also an \mathcal{J} -cluster point of \mathbf{x} . To show this, for each $r > 0$, let V_r be the closed ball with center v and radius r . Moreover, for each $x \in F$, let $U_{x,r}$ be the open ball with center x and radius r .

Because h is continuous and K is compact, then $G_r := K \cap h^{-1}(V_r)$ is compact and contains K_0 for each $r > 0$. Let h_0 be the restriction of h to the compact set K , so that h_0 is uniformly continuous. It follows that h_0 admits a modulus of continuity, i.e., there exists a function $\omega : [0, \infty] \rightarrow [0, \infty]$ such that

$$\lim_{r \rightarrow 0} \omega(r) = 0 \quad \text{and} \quad \forall a, b \in K, \quad d'(h_0(a), h_0(b)) \leq \omega(d(a, b)),$$

where d and d' represent the metric on S and S' , respectively. In particular, ω is finite in a (right) neighborhood of 0, let us say $[0, \varepsilon]$. Replacing, if necessary, each $\omega(r)$ with $\sup_{q \leq r} \omega(q)$, we can assume without loss of generality that ω is nondecreasing. Finally, let ω^{-1} be the generalized inverse of ω , i.e., $\omega^{-1}(r) := \inf\{q : \omega(q) > r\}$, for each $r > 0$. For each $r > 0$, we obtain that $r \leq \sup \omega(d(a, b))$, where the supremum is taken with respect to all $a \in F$ and $b \in U_{a, \omega^{-1}(r)}$. This implies that

$$(3.1) \quad \forall r > 0, \quad G_r \subseteq \bigcup_{x \in F} U_{x, 2\omega^{-1}(r)}.$$

However, because G_r is compact, there exists $x_{r,1}, \dots, x_{r,m_r} \in F$ such that G_r is contained into $\bigcup_{i \leq m_r} U_{x_{r,i}, 2\omega^{-1}(r)}$. To conclude, for each $r > 0$, we have that $A_r := \{n \in \mathbf{N} : h(x_n) \in V_r\} \in \mathcal{J}^*$, hence it follows by (3.1) that

$$A_r \setminus I = \{n \in \mathbf{N} : x_n \in G_r\} \subseteq \bigcup_{i \leq m_r} \{n \in \mathbf{N} : x_n \in U_{x_{r,i}, 2\omega^{-1}(r)}\},$$

where $I := \{n \in \mathbf{N} : x_n \notin K\} \in \mathcal{J}$. Because $A_r \setminus I \in \mathcal{J}^+$ and \mathcal{J} is closed under finite unions, it follows that, for each $t \in \mathbf{N}$, there exists $k(t) \in \{1, \dots, m_{1/t}\}$ such that

$$B_t := \{n \in \mathbf{N} : x_n \in U_{x_{1/t, k(t)}, 2\omega^{-1}(1/t)}\} \in \mathcal{J}^+.$$

Because $(x_{1/t, k(t)} : t \in \mathbf{N})$ is a sequence in the compact set K_0 , there exists a convergent subsequence with limit, let us say, $\eta \in K_0$. Considering that $\lim_t 2\omega^{-1}(1/t) = 0$ and that, for every $r > 0$, the set $C_r := \{n \in \mathbf{N} : x_n \in U_{\eta, r}\}$ contains B_t for every t sufficiently large in the latter subsequence, we obtain that $C_r \in \mathcal{J}^+$, for all $r > 0$. In other words, $\eta \in \Gamma_x(\mathcal{J})$ and $h(\eta) = v$. This shows that $\Gamma_{h(x)}(\mathcal{J}) \subseteq h(\Gamma_x(\mathcal{J}))$, concluding the proof. ■

Corollary 3.3 *Let S be a metric space and \mathcal{J} be an ideal. In addition, fix a continuous function $h : S \rightarrow \mathbf{R}$ and a sequence \mathbf{x} in S such that $\{n \in \mathbf{N} : x_n \in K\} \in \mathcal{J}^*$ for some compact $K \subseteq S$. Then,*

$$(3.2) \quad \mathcal{J}\text{-lim inf } h(\mathbf{x}) = \min \Gamma_{h(x)}(\mathcal{J}) = \min_{\eta \in \Gamma_x(\mathcal{J})} h(\eta),$$

and, symmetrically,

$$(3.3) \quad \mathcal{J}\text{-lim sup } h(\mathbf{x}) = \max \Gamma_{h(x)}(\mathcal{J}) = \max_{\eta \in \Gamma_x(\mathcal{J})} h(\eta).$$

Proof The first equality of (3.2) is a consequence of [15, Corollary 2.3] (cf. also [11, Theorem 1'] for the case $\mathcal{J} = \mathcal{Z}$). The second equality of (3.2) follows directly by Proposition 3.2. The proof for the case $S = \mathbf{R}^n$ and $\mathcal{J} = \mathcal{Z}$ can be found also in [25, Lemma 3.1].

The proof of (3.3) is analogous. ■

4 Proofs

Proof of Theorem 2.1 Suppose that \mathbf{x} is an \mathcal{J} -optimal path. Because $\mathbf{x} \in \mathcal{C} \subseteq \mathcal{H}$, there exists a compact set $K \subseteq X$ such that $\{n \in \mathbf{N} : x_n \notin K\} \in \mathcal{J}$. It follows by (2.1), Corollary 3.3, and conditions (A2) and (A6) that

$$\min_{\eta \in \Gamma_x(\mathcal{J})} u(\eta) = \mathcal{J}\text{-lim inf } u(\mathbf{x}) \geq \sup_{y \in \mathcal{C}} \mathcal{J}\text{-lim inf } u(y) \geq u(\eta^*),$$

hence $\Gamma_x(\mathcal{J}) \subseteq K \cap F$, where we recall that F is the closed set $\{x \in X : u(x) \geq u(\eta^*)\}$. Because K is compact and $\Gamma_x(\mathcal{J})$ is closed by Lemma 3.1(i), we obtain that $\Gamma_x(\mathcal{J})$ is compact. In addition, it is nonempty by Lemma 3.1(ii), therefore $\Gamma_x(\mathcal{J}) \in \mathcal{K}$.

Suppose that $F = \{\eta^*\}$. Because $\Gamma_x(\mathcal{J})$ is a nonempty subset of F , it follows that $\Gamma_x(\mathcal{J}) = \{\eta^*\}$, hence $\mathcal{J}\text{-lim } \mathbf{x} = \eta^*$ by Lemma 3.1(iv).

Let us suppose hereafter that $|F| \geq 2$, so that the linear operator T in (A5) is nonzero. Replacing T with $T/\|T\|$, we can assume without loss of generality that $\|T\| = 1$.

Claim 1 *The map $\text{Gr}(\Phi) \rightarrow \mathbf{R} : (x, y) \mapsto T(x - y)$ is continuous.*

Proof The claimed map can be rewritten as the restriction on $\text{Gr}(\Phi)$ of the composition $T \circ g$, where g is the continuous function $g : X^2 \rightarrow X : (x, y) \mapsto x - y$. ■

Because T is continuous and $\Phi(x) \in \mathcal{K}$, for each $x \in X$, the function

$$\widehat{T} : X \rightarrow \mathbf{R} : x \mapsto \max_{y \in \Phi(x)} T(y - x)$$

is well defined. Because the maximum is reached, we have $\widehat{T}x < 0$, for all $x \in F \setminus \{\eta^*\}$, by (A5).

Claim 2 \widehat{T} is continuous and $\widehat{T}\eta^* = 0$.

Proof Because Φ is a continuous correspondence by (A1) and the map defined in Claim 1 is continuous, it follows by Berge’s maximum theorem [1, Theorem 17.31] that \widehat{T} is continuous.

For the second part, we obtain by (A5) that $T(y - \eta^*) < 0$, for all $y \in \Phi(\eta^*)$, with $y \neq \eta^*$. Because $\eta^* \in \text{Fix}(\Phi)$, we conclude that $\widehat{T}\eta^* = T(\eta^* - \eta^*) = 0$. ■

Because T is continuous and $x \in \mathcal{C} \subseteq \mathcal{X}$, it follows that there exists $M \in \mathbf{R}$ such that $\{n \in \mathbf{N} : |Tx_n| \geq M\} \in \mathcal{J}$. Hence, the \mathcal{J} -limit inferior and the \mathcal{J} -limit superior of the real sequence (Tx_n) are well defined.

Claim 3 Fix $\eta \in \Gamma_x(\mathcal{J})$ such that $\mathcal{J}\text{-lim inf}_n Tx_n = T\eta$. Then, $\eta = \eta^*$.

Proof Assume that there exists $\eta_0 \in \Gamma_x(\mathcal{J})$ different from η^* such that $\mathcal{J}\text{-lim inf}_n Tx_n = T\eta_0$. Because $\Gamma_x(\mathcal{J}) \subseteq F$ and $\eta_0 \neq \eta^*$, then $\widehat{T}\eta_0 < 0$. Because \widehat{T} is continuous by Claim 2, there exist $\varepsilon, \delta > 0$ such that $\widehat{T}x < -\varepsilon$ whenever $\|x - \eta_0\| < \delta$. Moreover, it can be assumed without loss of generality that $\delta < \varepsilon/2$. At this point, fix $x, y \in X$ such that $\|x - \eta_0\| < \delta$ and $y \in \Phi(x)$, and let π_y be a minimizer of $\|\pi - y\|$ with $\pi \in \Gamma_x(\mathcal{J})$. Because $\widehat{T}x < -\varepsilon$, we get

$$Ty < Tx - \varepsilon = T\eta_0 + T(x - \eta_0) - \varepsilon \leq T\eta_0 + \|T\|\|x - \eta_0\| - \varepsilon < T\eta_0 - \varepsilon/2.$$

At the same time, we have

$$Ty = T\pi_y + T(y - \pi_y) \geq T\eta_0 - \|y - \pi_y\|,$$

which implies that $\|y - \pi_y\| > \varepsilon/2$.

To sum up, if $\|x - \eta_0\| < \delta$, then $\|y - \pi_y\| > \varepsilon/2$, for all $y \in \Phi(x)$. Because η_0 is an \mathcal{J} -cluster point of x , we have $A := \{n \in \mathbf{N} : \|x_n - \eta_0\| < \delta\} \in \mathcal{J}^+$. Thus, because \mathcal{J} is translation invariant by (A3), then also $A + 1 \in \mathcal{J}^+$. However, considering that $x_{n+1} \in \Phi(x_n)$, for all $n \in A$, we obtain by the preceding part that $\|x_{n+1} - \pi_{x_{n+1}}\| > \varepsilon/2$. To sum up, the open set $U := \{z \in X : \exists \eta \in \Gamma_x(\mathcal{J}), \|z - \eta_0\| < \varepsilon/2\}$ contains $\Gamma_x(\mathcal{J})$ and it has empty intersection with the \mathcal{J} -positive set $A + 1$. This contradicts Lemma 3.1(v). ■

Claim 4 Fix $\eta \in \Gamma_x(\mathcal{J})$ such that $\mathcal{J}\text{-lim sup}_n Tx_n = T\eta$. Then, $T\eta = T\eta^*$.

Proof Assume that there exists $\eta_0 \in \Gamma_x(\mathcal{J})$ such that $\mathcal{J}\text{-lim sup}_n Tx_n = T\eta$ and $T\eta \neq T\eta^*$. Because η^* is a minimizer of $T\eta$ with $\eta \in \Gamma_x(\mathcal{J})$ by Claim 3, then $\kappa := T(\eta_0 - \eta^*) > 0$.

Because \widehat{T} is continuous and $\widehat{T}\eta^* = 0$ by Claim 2, there exist $\varepsilon, \delta \in (0, \kappa/4)$ such that $\widehat{T}x < \varepsilon$ whenever $\|x - \eta^*\| < \delta$. Therefore, for each $y \in \Phi(x)$ such that $\|x - \eta^*\| < \delta$, we obtain

$$Ty \leq Tx + \varepsilon = T\eta^* + T(x - \eta^*) + \varepsilon < T\eta^* + \delta + \varepsilon < T\eta^* + \kappa/2,$$

and, at the same time,

$$Ty = T\eta_0 + T(y - \eta_0) \geq T\eta^* + \kappa - \|y - \eta_0\|.$$

Therefore, $\|y - \eta_0\| > \kappa/2$ whenever $\|x - \eta^*\| < \delta$ and $y \in \Phi(x)$. It follows that, if $x \in X$ is chosen such that $\|x - \eta\| < \delta/2$ and $\|\eta - \eta^*\| < \delta/2$, for some $\eta \in \Gamma_x(\mathcal{J})$, then $\|x - \eta^*\| \leq \|x - \eta\| + \|\eta - \eta^*\| < \delta$ and hence $\|y - \eta_0\| < \kappa$, for all $y \in \Phi(x)$.

At this point, note that the set $Q := \{\eta \in \Gamma_x(\mathcal{J}) : \|\eta - \eta^*\| \geq \delta/2\}$ is compact. Suppose that $Q \neq \emptyset$. By the continuity of \widehat{T} , we have $\max_{\eta \in Q} \widehat{T}\eta < 0$. It follows there exist $\lambda, \tau > 0$ such that $\widehat{T}x < -\lambda$ whenever $\|x - \eta\| < \tau$, for some $\eta \in Q$. In addition, it can be assumed without loss of generality that $\tau < \lambda/2$. Now, let us suppose that $x \in X$ is chosen such that $\|x - \eta\| < \tau$ for some fixed $\eta \in Q$. Fix $y \in \Phi(x)$; then,

$$T\eta_0 - \|y - \eta_0\| \leq Ty < Tx - \lambda \leq T\eta_x + \|x - \eta\| - \lambda \leq T\eta_0 - \lambda/2,$$

so that $\|y - \eta_0\| > \lambda/2$.

Set $\nu := \min\{\delta/2, \kappa/2, \tau, \lambda/2\} > 0$ and fix $x, y \in X$ with $y \in \Phi(x)$. To sum up the previous observations, we have that:

- (i) If there exists $\eta \in \Gamma_x(\mathcal{J})$ such that $\|x - \eta\| < \nu$ and $\|\eta - \eta^*\| < \delta/2$, then $\|y - \eta_0\| > \nu$;
- (ii) If there exists $\eta \in \Gamma_x(\mathcal{J})$ such that $\|x - \eta\| < \nu$ and $\|\eta - \eta^*\| \geq \delta/2$ (so that $Q \neq \emptyset$), then $\|y - \eta_0\| > \nu$.

Putting everything together, if $\|y - \eta_0\| \leq \nu$, then $\|x - \eta\| > \nu$, for all $\eta \in \Gamma_x(\mathcal{J})$.

We conclude as in the proof of Claim 3: because $A := \{n \in \mathbf{N} : \|x_{n+1} - \eta_0\| \leq \nu\} \in \mathcal{J}^+$ and \mathcal{J} is translation invariant by (A3), then $A - 1 \in \mathcal{J}^+$. However, $A - 1$ is a subset of $\{n \in \mathbf{N} : \forall \eta \in \Gamma_x(\mathcal{J}), \|x_n - \eta\| > \nu\}$, which belongs to \mathcal{J} thanks to Lemma 3.1(v). This contradiction concludes the proof. ■

To complete the proof, note that by Corollary 3.3 there exist nonempty compact sets $\Gamma_{\min}, \Gamma_{\max} \subseteq \Gamma_x(\mathcal{J})$ such that $\mathcal{J}\text{-lim inf}_n Tx_n = T\eta$, for all $\eta \in \Gamma_{\min}$, and $\mathcal{J}\text{-lim sup}_n Tx_n = T\eta$, for all $\eta \in \Gamma_{\max}$. Hence, Claims 3 and 4 imply that

$$\{\eta^*\} = \Gamma_{\min} \quad \text{and} \quad \eta^* \in \Gamma_{\min} \cap \Gamma_{\max}.$$

In other words, the function $\Gamma_x(\mathcal{J}) \rightarrow \mathbf{R}$ defined by $\eta \mapsto T\eta$ has a unique point of minimum which is also a maximizer. Therefore, $\Gamma_x(\mathcal{J}) = \{\eta^*\}$, which is equivalent to $\mathcal{J}\text{-lim } x = \eta^*$ by Lemma 3.1(iv). ■

Remark 4.1 As it is evident from the proof of Theorem 2.1, the full strength of condition (A5) has not been used. Indeed, we needed it only in Claim 3 to show that

$\widehat{T}\eta_0 < 0$ for some $\eta_0 \in \Gamma_x(\mathcal{J})$ and in Claim 4 to show that $\max_{\eta \in Q} \widehat{T}\eta < 0$ for a suitable subset $Q \subseteq \Gamma_x(\mathcal{J})$. Therefore, it is enough to replace (2.2) with the weaker condition

$$\forall x \in \Gamma_x(\mathcal{J}), \forall y \in \Phi(x), \quad Tx \leq Ty \implies x = y = \eta^*.$$

However, this condition, differently from (A5), depends on a given sequence \mathbf{x} .

Proof of Corollary 2.2 Thanks to Theorem 2.1, it is sufficient to show that the existence of a sequence $\mathbf{y} \in \mathcal{C}$ which is \mathcal{J} -convergent to η^* implies condition (A6). To this aim, observe that $\Gamma_{\mathbf{y}}(\mathcal{J}) = \{\eta^*\}$ by Lemma 3.1(iii). It follows by Corollary 3.3 that

$$\mathcal{J}\text{-}\liminf u(\mathbf{y}) = \min_{\eta \in \Gamma_{\mathbf{y}}(\mathcal{J})} u(\eta) = u(\eta^*),$$

concluding the proof. ■

Proof of Corollary 2.4 It is enough to note that every \mathcal{J} -cluster point is an ordinary accumulation point, so that by Corollary 3.3 we obtain

$$\begin{aligned} \sup_{\mathbf{y} \in \mathcal{C}} \mathcal{J}\text{-}\liminf_n u(\phi^n(\mathbf{y})) &\geq \mathcal{J}\text{-}\liminf_n u(\phi^n(\mathbf{y}_0)) \\ &= \min \Gamma_{(u(\phi^n(\mathbf{y}_0)))}(\mathcal{J}) \geq \liminf_n u(\phi^n(\mathbf{y}_0)) \geq u(\eta^*). \end{aligned}$$

Hence, condition (A6) holds, and the conclusion follows by Theorem 2.1. ■

5 An infinite dimensional example

As promised, we provide a practical example where Theorem 2.1 holds in infinite dimension.

Example 5.1 Let X be the Hilbert space ℓ_2 of square summable real sequences, i.e., sequences $\mathbf{x} = (x_0, x_1, \dots)$ such that

$$\|\mathbf{x}\| := \sqrt{\sum_{i \geq 0} x_i^2} < \infty.$$

Fix a sequence $\mathbf{x}^* \in \ell_2$, and define $\mathcal{C} = \{(\mathbf{x}^{(n)}) \in \mathcal{F}_K : \mathbf{x}^{(0)} = \mathbf{x}^*\}$. Let also \mathcal{J} be an arbitrary translation invariant ideal, and, for each $\mathbf{x} \in \ell_2$, set $u(\mathbf{x}) = x_0$ and

$$(5.1) \quad \Phi(\mathbf{x}) = \left\{ \left(-\sum_{i \geq 1} x_i^2, y_1, y_2, \dots \right) : 2x_i \leq y_i \leq x_i + 1/i \text{ for all } i \geq 1 \right\} \cup \left\{ \frac{1}{2}\mathbf{x} \right\},$$

where $\frac{1}{2}\mathbf{x} = (x_0/2, x_1/2, \dots)$. First of all, let us show that Φ is well defined. To do this, fix $\mathbf{x} \in \ell_2$ and let us prove that $\Phi(\mathbf{x}) \subseteq \ell_2$. If $\mathbf{y} \in \Phi(\mathbf{x})$ is equal to $\frac{1}{2}\mathbf{x}$, then it clearly belongs to ℓ_2 . Otherwise, because ℓ_2 is a vector space, it is sufficient to show that $\mathbf{z} = \mathbf{y} - \mathbf{x} = (-x_0 - \sum_{i \geq 1} x_i^2, z_1, z_2, \dots) \in \ell_2$, where $x_i \leq z_i \leq 1/i$, for all $i \geq 1$. Therefore,

$$\begin{aligned} \sum_{i \geq 0} z_i^2 &\leq z_0^2 + \sum_{i \geq 1} \left(|x_i| + \frac{1}{i} \right)^2 \leq z_0^2 + \sum_{i \geq 0} x_i^2 + \sum_{i \geq 1} \frac{1}{i^2} + 2 \sum_{i \geq 1} \frac{|x_i|}{i} \\ &\leq z_0^2 + \sum_{i \geq 0} x_i^2 + \sum_{i \geq 1} \frac{1}{i^2} + 2\sqrt{\sum_{i \geq 1} x_i^2 \cdot \sum_{i \geq 1} \frac{1}{i^2}} < \infty, \end{aligned}$$

where the last \leq follows by the Cauchy–Schwarz inequality. In addition, $\Phi(\mathbf{x})$ is compact. To this aim, because $\{\frac{1}{2}\mathbf{x}\}$ is compact, it is sufficient to show that a translation of the first set in the definition (5.1) of $\Phi(\mathbf{x})$ is compact. Let us define

$$(5.2) \quad \varphi(\mathbf{x}) := \left\{(-x_0 - \sum_{i \geq 1} x_i^2, z_1, z_2, \dots) : x_i \leq z_i \leq 1/i \text{ for all } i \geq 1\right\}.$$

Let \mathbf{a} be the sequence defined by $a_0 := -x_0 - \sum_{i \geq 1} x_i^2$ and $a_i := |x_i| + \frac{1}{i}$. Note that $\mathbf{a} \in \ell_2$ and that $\varphi(\mathbf{x})$ is a closed subset of $\{\mathbf{z} \in \ell_2 : |z_i| \leq a_i \text{ for all } i \geq 0\}$. However, the latter set is compact thanks to [7, p. 453], hence $\varphi(\mathbf{x})$ is compact too. To sum up, $\Phi(\mathbf{x})$ is compact subset of ℓ_2 which is nonempty (because it contains $\frac{1}{2}\mathbf{x}$).

At this point, let us show that Φ is continuous. Reasoning as above, it is sufficient to show that the correspondence φ defined in (5.2) is continuous at \mathbf{x} . Assume that $\varphi(\mathbf{x}) \neq \emptyset$, i.e., $x_i \leq 1/i$, for all $i \geq 1$, otherwise the claim is trivial.

First, let us show that φ is upper hemicontinuous. Fix $\varepsilon > 0$ and define the open set $U_\varepsilon := \{\mathbf{z} \in \ell_2 : \exists \mathbf{y} \in \varphi(\mathbf{x}), \|\mathbf{z} - \mathbf{y}\| < \varepsilon\}$. We need to find a constant $\delta > 0$ such that, for each $\mathbf{x}' \in \ell_2$, if $\|\mathbf{x} - \mathbf{x}'\| < \delta$, then $\varphi(\mathbf{x}') \subseteq U_\varepsilon$. Hence, fix also $\mathbf{x}' \in \ell_2$ such that $\|\mathbf{x} - \mathbf{x}'\| < \delta$ for a suitable $\delta > 0$ that will be chosen later and pick $\mathbf{y}' \in \varphi(\mathbf{x}')$. In particular, $|x_i - x'_i| < \delta$, for all $i \geq 0$. Similarly, we can assume without loss of generality that $x'_i \leq 1/i$, for all $i \geq 1$, otherwise $\varphi(\mathbf{x}') = \emptyset$. Then,

$$(5.3) \quad \begin{aligned} \forall \mathbf{y} \in \varphi(\mathbf{x}), \quad |y_0 - y'_0| &\leq |x_0 - x'_0| + |\sum_{i \geq 1} x_i^2 - \sum_{i \geq 1} (x'_i)^2| \\ &\leq |x_0 - x'_0| + |x_0^2 - (x'_0)^2| + \|\mathbf{x}\|^2 - \|\mathbf{x}'\|^2 \\ &\leq \delta + \delta|x_0 + x'_0| + \delta(\|\mathbf{x}\| + \|\mathbf{x}'\|) \\ &\leq \delta(1 + 2(\|\mathbf{x}\| + \|\mathbf{x}'\|)) \\ &\leq \delta(1 + 4\|\mathbf{x}\| + 2\delta). \end{aligned}$$

Now, recall that, for each integer $i \geq 1$, we have $x'_i \leq y'_i \leq 1/i$ and $x_i \leq y_i \leq 1/i$. In particular, there exists $\mathbf{y} \in \varphi(\mathbf{x})$ such that $|y_i - y'_i| \leq |x_i - x'_i|$, for all $i \geq 1$. It follows that

$$(5.4) \quad \sum_{i \geq 1} (y_i - y'_i)^2 \leq \sum_{i \geq 1} (x_i - x'_i)^2 \leq \|\mathbf{x} - \mathbf{x}'\|^2 \leq \delta^2.$$

Putting together the above estimates, we obtain that, for each given $\mathbf{y}' \in \varphi(\mathbf{x}')$, there exists $\mathbf{y} \in \varphi(\mathbf{x})$ such that

$$(5.5) \quad \|\mathbf{y} - \mathbf{y}'\| = \sqrt{|y_0 - y'_0|^2 + \sum_{i \geq 1} (y_i - y'_i)^2} \leq \delta \sqrt{(1 + 4\|\mathbf{x}\| + 2\delta)^2 + 1} < \varepsilon,$$

where the last inequality holds if δ is sufficiently small.

Second, let us show that φ is lower hemicontinuous. To this aim, fix an arbitrary $\mathbf{y} \in \varphi(\mathbf{x})$ and some $\varepsilon > 0$. We claim that there exists $\delta > 0$ such that if $\|\mathbf{x} - \mathbf{x}'\| < \delta$, then $\|\mathbf{y} - \mathbf{y}'\| < \varepsilon$, for some $\mathbf{y}' \in \varphi(\mathbf{x}')$. Note that estimates (5.3) and (5.4) hold symmetrically also in this case, with the conclusion that we have exactly the same upper bound computed in (5.5) for $\|\mathbf{y} - \mathbf{y}'\|$.

Therefore, Φ is a nonempty compact-valued continuous correspondence on ℓ_2 , \mathbb{J} is translation invariant, and u is clearly continuous. Hence, conditions (A1)–(A3) hold.

The set of fixed points of Φ is neither convex nor compact, and it is equal to

$$\text{Fix}(\Phi) = \{\mathbf{x} \in \ell_2 : x_0 = -\sum_{i \geq 1} x_i^2 \text{ and } x_j \leq 0 \text{ for all } j \geq 1\}.$$

It follows that the restriction of u on $\text{Fix}(\Phi)$ has a unique maximizer, which is the zero sequence $\mathbf{0}$ of ℓ_2 , hence condition (A4) holds. Moreover, we have that

$$F = \{\mathbf{x} \in \mathcal{C} : u(\mathbf{x}) \geq u(\mathbf{0})\} = \{\mathbf{x} \in \mathcal{C} : x_0 \geq 0\}.$$

Fix sequences $\mathbf{x} \in F$ and $\mathbf{y} \in \Phi(\mathbf{x})$ such that $(\mathbf{x}, \mathbf{y}) \neq (\mathbf{0}, \mathbf{0})$. Note that $\Phi(\mathbf{0}) = \{\mathbf{0}\}$, hence $\mathbf{x} \neq \mathbf{0}$ (in particular, if $x_0 = 0$, then $x_i \neq 0$, for some $i \geq 1$). Thus, setting $T = u$ (which is a continuous linear functional on ℓ_2), we obtain that $T\mathbf{y} = -\sum_{i \geq 1} x_i^2 < x_0 = T\mathbf{x}$, i.e., condition (A5) holds. Finally, note that the sequence $(\mathbf{x}^*, \frac{1}{2}\mathbf{x}^*, \frac{1}{4}\mathbf{x}^*, \dots)$ is convergent to $\mathbf{0}$ and belongs to \mathcal{C} (indeed, it starts at \mathbf{x}^* , it is feasible, and its image is contained in the compact set $\{k\mathbf{x}^* : k \in [0, 1]\}$).

We conclude by Corollary 2.2 that each \mathcal{J} -optimal sequence $(\mathbf{x}^{(n)}) \in \mathcal{C}$ in the system $(\ell_2, \Phi, u, \mathcal{J}, \mathcal{C})$ is necessarily \mathcal{J} -convergent to $\mathbf{0}$.

Acknowledgment We would like to express our sincere gratitude for Marek Balcerzak (Lodz University of Technology, P.L.) and Piotr Szuca (University of Gdańsk, P.L.) for useful discussions and for letting us know about the manuscript [22].

References

- [1] C. D. Aliprantis and K. C. Border, *Infinite dimensional analysis*. 3rd ed., Springer, Berlin, 2006, A hitchhiker's guide.
- [2] N. Bourbaki, *General topology*. Chapters 1–4, Elements of Mathematics (Berlin), Springer, Berlin, 1998, Translated from the French, reprint of the 1989 English translation.
- [3] T. Breiten and L. Pfeiffer, *On the turnpike property and the receding-horizon method for linear-quadratic optimal control problems*. SIAM J. Control Optim. 58(2020), no. 2, 1077–1102.
- [4] T. Damm, L. Grüne, M. Stieler, and K. Worthmann, *An exponential turnpike theorem for dissipative discrete time optimal control problems*. SIAM J. Control Optim. 52(2014), no. 3, 1935–1957.
- [5] P. Das, S. Dutta, S. A. Mohiuddine, and A. Alotaibi, *A-statistical cluster points in finite dimensional spaces and application to turnpike theorem*. Abstr. Appl. Anal. 2014(2014), Article ID 354846, 7 pages.
- [6] M. Di Nasso and R. Jin, *Abstract densities and ideals of sets*. Acta Arith. 185(2018), no. 4, 301–313.
- [7] N. Dunford and J. T. Schwartz, *Linear operators. Part I: general theory*. Wiley Classics Library, John Wiley & Sons, Inc., New York, 1988, with the assistance of William G. Bade and Robert G. Bartle, reprint of the 1958 original, a Wiley-Interscience Publication.
- [8] K. Falconer, *Fractal geometry*. 3rd ed., Mathematical Foundations and Applications, John Wiley & Sons, Ltd., Chichester, 2014.
- [9] R. Filipów and J. Tryba, *Densities for sets of natural numbers vanishing on a given family*. J. Number Theory 211(2020), 371–382.
- [10] J. A. Fridy, *Statistical limit points*. Proc. Amer. Math. Soc. 118(1993), no. 4, 1187–1192.
- [11] J. A. Fridy and C. Orhan, *Statistical limit superior and limit inferior*. Proc. Amer. Math. Soc. 125(1997), no. 12, 3625–3631.
- [12] M. Gugat and F. M. Hante, *On the turnpike phenomenon for optimal boundary control problems with hyperbolic systems*. SIAM J. Control Optim. 57(2019), no. 1, 264–289.
- [13] J. E. Hutchinson, *Fractals and self-similarity*. Indiana Univ. Math. J. 30(1981), no. 5, 713–747.
- [14] V. Kadets and D. Seliutin, *On relation between the ideal core and ideal cluster points*. J. Math. Anal. Appl. 492(2020), no. 1, Article ID 124430, 7 pages.
- [15] P. Leonetti, *Characterizations of the ideal core*. J. Math. Anal. Appl. 477(2019), no. 2, 1063–1071.
- [16] P. Leonetti and F. Maccheroni, *Characterizations of ideal cluster points*. Analysis (Berlin) 39(2019), no. 1, 19–26.

- [17] P. Leonetti and S. Tringali, *On the notions of upper and lower density*. Proc. Edinb. Math. Soc. (2) **63**(2020), no. 1, 139–167.
- [18] M. E. Lewis and A. Paul, *Uniform turnpike theorems for finite Markov decision processes*. Math. Oper. Res. **44**(2019), no. 4, 1145–1160.
- [19] M. A. Mamedov, *Turnpike theorems in continuous systems with integral functionals*. Dokl. Akad. Nauk. **323**(1992), no. 5, 830–833.
- [20] M. A. Mamedov and S. Pehlivan, *Statistical cluster points and turnpike theorem in nonconvex problems*. J. Math. Anal. Appl. **256**(2001), no. 2, 686–693.
- [21] M. Mammadov and R. J. Evans, *Turnpike theorem for terminal functionals in infinite horizon optimal control problems*. J. Math. Anal. Appl. **428**(2015), no. 2, 1147–1160.
- [22] M. Mammadov and P. Szuca, *Invariant ideals and its applications to the turnpike theory*. Manuscript.
- [23] M. Marena and L. Montrucchio, *Neighborhood turnpike theorem for continuous-time optimization models*. J. Optim. Theory Appl. **101**(1999), no. 3, 651–676.
- [24] H. I. Miller, *A measure theoretical subsequence characterization of statistical convergence*. Trans. Amer. Math. Soc. **347**(1995), no. 5, 1811–1819.
- [25] S. Pehlivan and M. A. Mamedov, *Statistical cluster points and turnpike*. Optimization. **48**(2000), no. 1, 93–106.
- [26] A. J. Zaslavski, *Turnpike properties in the calculus of variations and optimal control*. Nonconvex Optimization and its Applications, 80, Springer, New York, 2006.
- [27] A. J. Zaslavski, *Turnpike conditions in infinite dimensional optimal control*. Springer Optimization and its Applications, 148, Springer, Cham, 2019.

Department of Statistics, Università Luigi Bocconi, via Roentgen 1, 20136 Milan, Italy
e-mail: leonetti.paolo@gmail.com

Department of Statistical Science, Duke University, 415 Chapel Drive, Durham, NC 27708-0251, USA
e-mail: michele.caprio@duke.edu