

Beneficial effects of probiotic bacteria isolated from breast milk

Federico Lara-Villoslada¹, Mónica Olivares¹, Saleta Sierra¹, Juan Miguel Rodríguez², Julio Boza¹ and Jordi Xaus^{1*}

¹Nutrition and Health Department, Puleva Biotech. Cno, De Purchil no 66, 18004, Granada (Spain)

²Nutrition and Bromatology Department, Universidad Complutense, Ciudad Universitaria, 28040 Madrid (Spain)

Breast milk is the best food for the neonate because it provides a unique combination of proteins, carbohydrates, lipids, minerals and vitamins that ensures the correct growth and development of the infant. In addition, it also contains bioactive compounds responsible for a wide range of beneficial effects such as the promotion of immune system maturation and the protection against infections. Among these bioactive agents, probiotic bacteria have been recently isolated from human milk. The present work reviews the beneficial effects of these bacteria both in animal models and in clinical trials. The promotion of immune system maturation and defence against infections as well as the anti-inflammatory properties are among the main healthy effects of these bacteria. The isolation of probiotic bacteria with beneficial effects for the host provides scientific support for the supplementation of infant formula with these bacteria, in order to advance the pursuit of the main goal of formula: to mimic breast milk and its functional effects as closely as possible.

Probiotic: Human milk: Infant nutrition: Immune system

Human milk is a complex species-specific biological fluid adapted to perfectly satisfy the nutritional and immunological needs of the neonate. It has been demonstrated that breast milk confers protection against different infectious diseases since the incidence of these disorders is lower in breast-fed than in formula-fed infants^{1,2}. It has been suggested that this anti-infective effect is due to several bioactive compounds present in colostrum and/or in mature milk. These include immunoglobulins, immune cells, antimicrobial acids, polyamines, oligosaccharides, lysosyme, glycoproteins such as lactoferrin and bioactive peptides, which, acting individually or synergistically, could inhibit pathogenic microorganisms^{3,4}.

Recent studies have demonstrated that human milk, far from being a sterile fluid, constitutes an excellent and continuous source of commensal bacteria for the infant gut^{5,6}. These bacteria could also play an important role in the reduction of incidence and severity of infectious diseases in breastfed children. This hypothesis is supported by relatively old studies reporting the loss of antimicrobial activity in pasteurised human milk⁷.

Among the bacteria found in human milk, those belonging to the species *Staphilococcus*, *Lactococcus*, *Enterococcus* and *Lactobacillus* are the most frequent^{5,6} (Table 1). There is increasing interest in some of these breast milk lactobacilli, such as *L. gasseri*, *L. salivarius*, *L. rhamnosus*, *L. plantarum* and *L. fermentum*, because they are considered as potentially probiotic species (Table 1).

Breastfeeding and protection against diseases

In developing countries, one of the main causes of death in the paediatric age group is the infectious disease, specially

gastroenterocolitis and respiratory infections. Newborns who have not been breastfed show a 17-fold higher risk of being hospitalised due to pneumonia than those who exclusively received human milk⁸. Similarly, the risk of death due to diarrhoea increases 14.2-fold in weaned infants⁹. Breastfeeding has also been related to a lower incidence of acute otitis media¹⁰, urinary tract infection¹¹ and meningitis caused by *Haemophilus influenzae*¹².

Besides its anti-infective properties, it has been demonstrated that human milk modulates the immune system of the newborn¹³. Although an anti-inflammatory activity has not yet been demonstrated *in vivo*, several epidemiologic studies suggest that breast-fed children are protected against infections without the observation of evident lesion of the intestinal or respiratory mucosa due to an inflammatory response¹⁴. This is probably the result of an anti-inflammatory system better regulated by bioactive components of human milk.

The immunomodulatory action of breast milk could also explain the better antibody production response in breast-fed compared to formula-fed infants after vaccination against poliomyelitis, tetanus and diphtheria¹⁵.

Neonates who received breast milk also have a more favourable intestinal microbiota than those fed infant formula¹⁶, which is probably due to presence of lactic acid bacteria in human milk, besides other bifidogenic compounds such as oligosaccharides¹⁷. It has been suggested that these differences in intestinal microbiota could be responsible for some of the beneficial effects seen in breast-fed infants. It has been known for several decades that lactobacilli and bifidobacteria inhibit the growth of pathogen microorganisms such as *Staphylococcus aureus*, *Salmonella typhimurium*,

*Corresponding author: Jordi Xaus Pey, fax + 34958240160, email jxaus@pulevabiotech.es

Table 1. Bacterial species generally isolated from the breast milk of healthy women

Bacterial group	Main species
<i>Staphylococcus spp.</i>	<i>S. epidermidis</i> <i>S. hominis</i> <i>S. capitis</i> <i>S. aureus</i>
<i>Streptococcus spp.</i>	<i>S. salivarius</i> <i>S. mitis</i> <i>S. parasanguis</i> <i>S. peores</i>
<i>Lactobacillus spp.</i>	<i>L. gasseri</i> <i>L. rhamnosus</i> <i>L. acidophilus</i> <i>L. plantarum</i> <i>L. fermentum</i> <i>L. salivarius</i> <i>L. reuteri</i>
<i>Enterococcus sp.</i>	<i>E. faecium</i> <i>E. faecalis</i>

Modified from Martin *et al.*³².

Yersinia enterocolitica and *Clostridium perfringens*¹⁸. These bacteria competitively colonise the intestine of the child, thus preventing the adhesion of pathogens. Moreover, a competition for nutrients is established and this is another mechanism that inhibits the growth of pathogenic microorganisms¹⁹.

Intestinal colonisation by commensal bacteria also plays a key role in the maintenance of immune system homeostasis. These bacteria stimulate T_{H1} responses and compensate the trend towards T_{H2} responses characteristic of the neonatal immune system. It has been reported that the administration of specific probiotics to newborns reduces the incidence of atopic manifestations²⁰ and also of inflammatory processes where a T_{H2} response is involved, such as necrotizing enterocolitis²¹.

Probiotics isolated from breast milk

The description of the presence of bacteria in human milk dates back 30 years, but then it was assumed to be a contamination occurring during sample extraction²². At the beginning of the 21st century, two European groups independently demonstrated the presence of lactic acid bacteria in human milk and their probiotic potential^{5,6}. Thus Heikkilä *et al.* reported that these human milk bacteria protect both mother and newborn from infections caused by *Staphylococcus aureus*⁵. In a similar way and in a series of reports, Martín *et al.*⁶ described the isolation of lactic acid bacteria from human milk, namely, *L. gasseri* CECT5714²³, *L. salivarius* CECT5713²⁴ and *L. fermentum* CECT5716²³. Besides these strains, there are other commercial strains related to human milk. One of them is *L. reuteri* ATCC55730, which is claimed to be derived from human milk but its origin has not been published yet. *L. rhamnosus* LGG, although originally isolated from intestinal sources²⁵, has also been found in human milk by the Finnish group⁵. Nevertheless, this work focuses on the beneficial effects of *L. gasseri* CECT5714, *L. salivarius* CECT5713 and *L. fermentum* CECT5716, which have been summarised in Table 2.

Anti-microbial effects

Protection against viral or bacterial infections is one of the most frequent claims made for probiotic consumption. Different mechanisms have been suggested to explain this anti-microbial activity (Fig. 1). *In vitro* studies demonstrate that certain probiotic strains produce anti-microbial compounds, such as organic acids, H₂O₂ and/or bacteriocins²⁶, that have been reported to inhibit the growth of *E. coli*, *Salmonella spp.* and *Listeria monocytogenes*²⁷. No bacteriocin-producing *lactobacillus* has been found in human milk, although a high production of H₂O₂ has been reported²³.

Table 2. Beneficial effects of some breast milk-isolated probiotic strains

Strain	Beneficial effect	Reference
<i>L. salivarius</i> CECT5713	Intestinal colonisation	Martin <i>et al.</i> , 2006 ²⁴
	Production of antimicrobial compounds	Martin <i>et al.</i> , 2006 ²⁴
	No D-lactic production	Martin <i>et al.</i> , 2006 ²⁴
	Anti-microbial effect	Olivares <i>et al.</i> , 2006 ²⁷
	Immunomodulatory effect	Díaz-Ropero <i>et al.</i> , 2006 ³⁰
	Anti-inflammatory effect	Peran <i>et al.</i> , 2005 ³⁷
	Intestinal colonisation	Martin <i>et al.</i> , 2005 ²³
<i>L. gasseri</i> CECT5714	Improved gastrointestinal function	Olivares <i>et al.</i> , 2006 ³⁸ Lara-Villoslada <i>et al.</i> , 2007 ³⁹
	Production of antimicrobial compounds	Olivares <i>et al.</i> , 2006 ³⁸
	Anti-microbial effect	Martin <i>et al.</i> , 2005 ²³
	Immunomodulatory effect	Olivares <i>et al.</i> , 2006 ²⁷ Olivares <i>et al.</i> , 2006 ³⁵
	Anti-allergic effect	Lara-Villoslada <i>et al.</i> , 2007 ³⁹ Olivares <i>et al.</i> , 2005 ³³
	Intestinal colonisation	Martin <i>et al.</i> , 2005 ²³ Olivares <i>et al.</i> , 2006 ³⁶
	Production of antimicrobial compounds	Martin <i>et al.</i> , 2005 ²³
<i>L. fermentum</i> CECT5716	Anti-microbial effect	Olivares <i>et al.</i> , 2006 ²⁷
	Immunomodulatory effect	Díaz-Ropero <i>et al.</i> , 2006 ³⁰
	Enhanced effects of vaccination	Olivares <i>et al.</i> , 2007 ³⁶
	Anti-inflammatory effect	Díaz-ropero <i>et al.</i> , 2006 ³⁰ Peran <i>et al.</i> , 2005 ³⁷
		Peran <i>et al.</i> , 2007 ³⁴




Fig. 1. Intestinal anti-infective mechanisms of probiotic bacteria.

In addition, those strains belonging to the species *L. reuteri* produce reuterin, another antimicrobial compound²⁸. It has also been shown that some bacteria present in human milk improve the intestinal barrier function by increasing mucine production and reducing intestinal permeability²⁷. However, competition with entero-toxigenic bacteria for nutrients and for epithelial intestinal cell receptor binding sites is probably the main anti-infective mechanism of probiotic bacteria^{26,27} (Fig. 1).

The human milk-isolated probiotics *L. gasseri* CECT5714, *L. salivarius* CECT5713 and *L. fermentum* CECT5714 have been reported to inhibit the adhesion of *Salmonella choleraesuis* to mucins and to increase the survival of mice infected with this pathogen²⁷. It was demonstrated that the protective effect of *L. salivarius* CECT5713 is significantly higher than the effect of a reuterin producing strain²⁹. This is probably due to the combination of the immunomodulatory role and the competitive activity reported for *L. salivarius* CECT5713³⁰.

Different clinical trials have demonstrated that, when breastfeeding is not possible, infant formula supplemented with probiotics protect children from infectious diseases. To our knowledge, most of the studies have involved supplementation with *L. rhamnosus* LGG, which have demonstrated a reduction in the incidence of rotavirus infection and in the duration of diarrhoea³¹. Currently, clinical studies are in progress to evaluate the tolerance and effectiveness of other breast milk strains, such as *L. reuteri* ATCC55730 and *L. salivarius* CECT5713.

Immunomodulatory properties

Intestinal colonisation is often the result of the first contact of the newborn with microorganisms, which is crucial for the development of the immune system of the neonate. It has been reported that differences in the composition of intestinal microbiota influence the incidence of certain pathologies with an important immunological component, such as allergic or inflammatory processes³². The anti-allergic effect of probiotics could be explained on the basis of the Hygiene Hypothesis

and the T_H1/T_H2 balance. Probiotics induce a T_H1 response, and thus down-regulate the production of T_H2 cytokines, responsible for the allergic response.

In contrast, the anti-inflammatory effect of probiotics is more difficult to explain. *In vitro* studies have demonstrated that the immunomodulatory effects of probiotics depend on the cell environment. Thus, in the absence of additional stimulus, the breast milk probiotics *L. salivarius* CECT5713 and *L. fermentum* CECT5716 enhance the production of T_H1 cytokines such as IL-2 and IL-12 and the inflammatory mediator TNF- α ³⁰. However, when cells are incubated in the presence of lipopolysaccharide, together with the probiotics, a reduction of T_H1 cytokines is observed³⁰. This regulatory mechanism is probably based on the production of IL-10, an immunosuppressive cytokine, which has been reported to be increased by these probiotic strains³⁰.

The immunomodulatory effects of probiotics have also been reported in animal models of pathologies where the immune system is involved. Different probiotic strains isolated from human milk have been reported to enhance the immune defence of mice, increasing both natural and acquired immune responses³⁰. This immune-stimulating activity could be also involved in the anti-infective role previously mentioned for these bacteria in an animal model of *Salmonella* infection²⁷. In addition, the breast milk probiotic *L. gasseri* CECT5714 in combination with *L. coryniformis* CECT5711 reduces the incidence and severity of the allergic response in an animal model of cow's milk protein allergy³³. In a recent report, *L. fermentum* CECT5716 showed a beneficial effect in an animal model of intestinal inflammation, reducing the inflammatory response and the intestinal damage³⁴.

Probiotics have also been reported to modulate the immune response of healthy humans, as shown by a recent study which reports an increase in phagocytic activity, in the number of natural killer cells and in the plasma concentration of IgA in healthy humans consuming human milk-isolated probiotics daily for 3 months³⁵. A more recent report demonstrates that the consumption of *L. fermentum* CECT5716 enhances the response to

influenza vaccination in healthy volunteers aged 26-40 and reduces the incidence of influenza-like illness³⁶.

In addition, the beneficial effect of probiotics in allergic processes has been widely reported. In this sense, the consumption of probiotics present in human milk, especially *L. rhamnosus* LGG, has been shown to reduce the incidence and severity of atopic dermatitis in children²⁰. Although less is known about other allergic disorders, there is data to support a positive effect of *L. gasseri* CECT5714 in adults with respiratory allergy³³.

Gastrointestinal benefits

There is increasing interest in the manipulation of intestinal microbiota with the aim of improving gastrointestinal function and nutrient absorption. Different reports demonstrate that human milk probiotics colonise the intestine and increase faecal lactobacilli counts thus modifying intestinal microbiota both in rodents³⁷ and humans³⁸. In addition, molecular analysis show that these bacteria are metabolically active in the human gut, increasing the production of functional metabolites such as butyrate³⁸, which is the main energy source for colonocytes and plays a key role in the modulation of intestinal function. In the previously mentioned clinical trial³⁸, an increase in faecal moisture, and in stool frequency and volume was observed which could be related to the increase in the faecal concentration of butyric acid.

Similarly, the administration of *L. gasseri* CECT5714 also caused an increase in faecal lactobacilli counts in a clinical trial in children aged 3-12³⁹. In the same study the cytotoxicity of the faecal water of children who received the probiotic has been shown to be lower than that of the control children³⁹. Finally, in another clinical trial the supplementation of infant formulas with *L. rhamnosus* LGG has been demonstrated to improve neonate growth pattern, which could suggest an increased bioavailability of nutrients in these infants⁴⁰.

Conclusions

Breastfeeding is the main determinant of the intestinal colonisation of the neonate, which, apart from other components, is due to the recently discovered presence of probiotic bacteria in human milk. In addition to gastrointestinal benefits, modulation of microbiota by probiotic bacteria has been shown to regulate the immune function and to enhance defence against intestinal pathogens. Thus, the addition of breast milk probiotics to infant formulas could be a new alternative to mimic some of the functional effects of human milk in children who are not breastfed.

Conflict of interest statement

All the authors except JMR are employees at Puleva Biotech SA. All the studies presented have been funded by Puleva Biotech's own funds. This review has mainly been written by JX with collaboration of all the other authors.

References

1. Lopez-Alarcon M, Villalpando S & Fajardo A (1997) Breastfeeding lowers the frequency and duration of acute respiratory infection and diarrhea in infants under six months of age. *J Nutr* **127**, 436-443.

2. Wright AL, Bauer M, Naylor A, Sutcliffe E & Clark L (1998) Increasing breastfeeding rates to reduce infant illness at the community level. *Pediatrics* **101**, 837-844.
3. Saavedra JM (2002) Probiotic agents: clinical applications in infants and children. In *Infant formula: closer to reference*, pp. 15-27 [NCR Rähä and FF Rubaltelli, editors]. Philadelphia: Lippincott Williams & Wilkins.
4. Isaacs CE (2005) Human milk inactivates pathogen individually, additively and synergistically. *J Nutr* **51**, 1286-1288.
5. Hekkilä MP & Saris PEJ (2003) Inhibition of *Staphylococcus aureus* by the commensal bacteria of human milk. *J Appl Microbiol* **95**, 471-478.
6. Martin R, Langa S, Reviriego C, Jimenez E, Marin ML, Xaus J, Fernandez L & Rodriguez JM (2003) Human milk is a source of lactic acid bacteria for the infant gut. *J Pediatr* **143**, 754-758.
7. Ford JE, Law BA, Marshall VME & Reiter B (1977) Influence of the heat treatment of human milk on some of its protective constituents. *J Pediatr* **91**, 29-35.
8. Cesar JA, Victoria CG, Barros FC, *et al.* (1999) Impact of breastfeeding on admission for pneumonia in postneonatal period in Brazil: nested case-control study. *BMJ* **318**, 1316-1320.
9. Victoria CG, Smiyh PG, Vaughan JP, *et al.* (1987) Evidence for protection by breast-feeding against infant deaths from infectious diseases in Brazil. *Lancet II* 319-322.
10. Duncan B, Ey J, Holberg CJ, Wright AL, Martinez FD & Taussing LM (1994) Exclusive breast-feeding for at least 4 months protects against otitis media. *Pediatrics* **91**, 867-871.
11. Pisacane A, Graziano L, Mazzarella G, Scarpellino B & Zona G (1992) Breast-feeding and urinary tract infections. *J Pediatr* **121**, 331-332.
12. Silfverdal AS & Olcen P (1999) Protective effect of breastfeeding: an ecologic study of *Haemophilus influenzae* meningitis and breast-feeding in a Swedish population. *Int J Epidemiol* **28**, 152-156.
13. Grazioso CF, Werner AL, Alling DW, *et al.* (1997) Antiinflammatory effects of human milk on chemically induced colitis in rats. *Pediatr Res* **42**, 639-643.
14. Garofalo RP & Goldman AP (1999) Expression of functional immunomodulatory and antiinflammatory factors in human milk. *Clin Perinatol* **26**, 361-367.
15. Hahn-Zoric M, Fulconis F, Minoli L, *et al.* (1990) Antibody response to parenteral and oral vaccines are impaired by conventional and low protein formulas as compared to breast-feeding. *Acta Paediatr Scand* **79**, 1137-1142.
16. Harmsen HJM, Wildeboer-Veloo ACM, Raangs GC, *et al.* (2000) Analysis of intestinal flora development in breast-fed infants by using molecular identification and detection methods. *J Pediatr Gastroenterol Nutr* **30**, 61-67.
17. Kunz C & Rudloff S (1993) Biological functions of oligosaccharides in human milk. *Acta Paediatr* **82**, 903-912.
18. Gilliland SE & Speck ML (1977) Antagonistic action of *Lactobacillus acidophilus* toward intestinal and food borne pathogens in associative cultures. *J Food Prot* **40**, 820-823.
19. Conway PL (1996) Selection criteria for probiotic microorganisms. *Asia pacific J Clin Nutr* **5**, 10-14.
20. Kalliomaki M, Salminen S, Arvilommi H, Kero P, Koskinen P & Isolauri E (2001) Probiotics in primary prevention of atopic disease: a randomized placebo-controlled trial. *Lancet* **357**, 1076-1079.
21. Dani C, Biandaioli R & Firmito FR (2002) Potential role of probiotics in the prevention of necrotizing enterocolitis. In *Infant formula: Closer to the reference*. Philadelphia: Lippincott Williams & Wilkins.
22. Gavin A & Ostovr K Microbiological characterization of human milk (1977) *J Food Prot* **40**, 614-616.
23. Martin R, Olivares M, Marin ML, Fernandez L, Xaus J & Rodriguez JM (2005) Probiotic potential of 3 lactobacilli strains isolated from breast milk. *Hum Lact* **21**, 8-17.

24. Martin R, Jimenez E, Olivares M, Marin ML, Fernandez L, Xaus J & Rodriguez JM (2006) *Lactobacillus salivarius* CECT5713, a potential probiotic strain isolated from infant feces and breast milk of a mother-child pair. *Int J Microbiol* **112**, 35–43.
25. Conway PL, Gorbach SL & Goldin BR (1987) Survival of lactic acid bacteria in the human stomach and adhesion to intestinal cells. *J Dairy Sci* **70**, 1–12.
26. Fons M, Gomez A & Karjalainen T (2000) Mechanisms of colonization and colonization resistance of the digestive tract. *Microb Ecol Health Dis* **2**, 240–246.
27. Olivares M, Diaz-Ropero MP, Martin R, Rodriguez JM & Xaus J (2006) Antimicrobial potential of four lactobacillus strains isolated from breast milk. *J Appl Microbiol* **101**, 72–79.
28. Talarico TL & Dobrogosz WJ (1989) Chemical characterization of an antimicrobial substance produced by *Lactobacillus reuteri*. *Antimicrob Agents Chemother* **33**, 674–679.
29. Martin R, Olivares m, Marin ML, Xaus J, Fernandez L & Rodriguez JM (2005) Characterization of a reuterin-producing *Lactobacillus coryniformis* strain isolated from a goat's milk cheese. *Int J Food Microbiol* **104**, 267–277.
30. Diaz-Ropero MP, Martin R, Sierra S, Lara-Villoslada F, Rodriguez JM, Xaus J & Olivares M (2006) Two *Lactobacillus* strains, isolated from breast milk, differently modulate the immune response. *J Appl Microbiol* **102**, 337–343.
31. Senok AC, Ismaeel AY & Botta GA (2005) Probiotics: facts and myths. *Clin Microb Infect* **11**, 958–966.
32. Björkstén B, Naaber P, Sepp E & Mikelsaar M (1999) The gut microflora in allergic Estonian and swedish 2-year-old children. *Clin Exp Allergy* **29**, 342–346.
33. Olivares M, Díaz-Ropero MP, Lara-Villoslada F, Rodriguez JM & Xaus J (2005) Effectiveness of probiotics in allergy: child's game or adult affair? *Nutrafoods* **4**, 59–64.
34. Peran L, Sierra S, Comalada M, *et al.* (2007) A comparative study of the preventative effects exerted by two probiotics, *Lactobacillus reuteri* and *Lactobacillus fermentum*, in the trinitrobenzenesulfonic acid model of rat colitis. *Br J Nutr* **97**, 96–103.
35. Olivares M, Diaz-Ropero MP, Gomez N, Lara-Villoslada F, Sierra S, Maldonado JM, Martin R, Rodriguez JM & Xaus J (2006) The consumption of two new probiotic strains, *Lactobacillus gasseri* CECT5714 and *Lactobacillus coryniformis* CECT5711, boost the immune system of healthy adults. *Int Microbiol* **9**, 47–52.
36. Olivares M, Díaz-Ropero MP, Sierra S, Lara-Villoslada F, Fonolla J, Navas M, Rodriguez JM & Xaus J (2007) Oral intake of *Lactobacillus fermentum* CECT5716 enhances the effect of influenza vaccination. *Nutrition* **23**, 254–260.
37. Peran L, Camuesco D, Comalada M, Nieto A, Concha A, Diaz-Ropero MP, Olivares M, Xaus J, Zarzuelo A & Galvez J (2005) Preventative effects of a probiotic, *Lactobacillus salivarius ssp salivarius*, in the TNBS model of rat colitis. *World J Gastroenterol* **11**, 5185–5192.
38. Olivares M, Díaz-Ropero MP, Gomez N, Lara-Villoslada F, Sierra S, Maldonado JA, Martin R, Lopez-Huertas E, Rodriguez JM & Xaus J (2006) Oral administration of two probiotic strains, *Lactobacillus gasseri* CECT5714 and *Lactobacillus coryniformis* CECT5711, enhances the intestinal function of healthy adults. *Int J Microbiol* **107**, 104–111.
39. Lara-Villoslada F, Sierra S, Boza J, Xaus J & Olivares M (2007) Beneficial effects of consumption of a dairy product containing two probiotic strains, *Lactobacillus coryniformis* CECT5711 and *Lactobacillus gasseri* CECT5714n in healthy children. *Nutr Hosp* **4**, **22**, 496–502.
40. Vendt N, Grünberg H, Tuure T, Malminiemi O, Wuolijoki E, Tillmann V, Sepp E & Korpela R (2006) Growth during the first 6 months of life in infants using formula enriched with *Lactobacillus rhamnosus* GG: double-blind, randomized trial. *J Hum Nutr Dietet* **19**, 51–58.