

**Conclusions.** The use of surrogate outcomes is common in NICE HST evaluations and the challenges in supporting surrogate relationships with more than biological plausibility are recognized. However, our review indicates that, the committee considers more than just biological plausibility and will take into account other related factors.

## OP128 Improving Literature Searching For Evidence On Health Apps: The National Institute For Health And Care Excellence (NICE) MEDLINE And Embase (Ovid) Health Apps Search Filters

Lynda Ayiku ([Lynda.ayiku@nice.org.uk](mailto:Lynda.ayiku@nice.org.uk)) and Sarah Glover

**Introduction.** Literature searching for evidence on apps in bibliographic databases is challenging because they are often described with inconsistent terminology. Information Specialists from the United Kingdom's National Institute for Health and Care Excellence (NICE) have developed validated search filters for retrieving evidence about apps from MEDLINE and Embase (Ovid) reliably.

**Methods.** Medical informatics journals were hand-searched to create a 'gold standard' set of app references. The gold standard set was divided into two sets. The development set provided the search terms for the filters. The filters were validated by calculating their recall against the validation set. Target recall was >90%.

A case study was then conducted to compare the number-needed-to-read (NNR) of the filters with previous non-validated MEDLINE and Embase app search strategies used for the 'MIB214 myCOPD app' NICE topic. NNR is the number of references screened to find each relevant reference.

**Results.** The MEDLINE and Embase filters achieved 98.6 percent and 98.5 percent recall against the validation set, respectively. In the case study they achieved 100 percent recall, reducing NNR from 348 to 147 in MEDLINE and from 456 to 271 in Embase.

**Conclusions.** The novel NICE health apps search filters retrieve evidence on apps from MEDLINE and Embase effectively and more efficiently than previous non-validated search strategies used at NICE.

## OP129 The Use Of A Text-Mining Screening Tool For Systematic Review Of Treatments For Relapsed/Refractory Diffuse Large B-Cell Lymphoma

Niamh Carey ([nicarey@tcd.ie](mailto:nicarey@tcd.ie)), Marie Harte and Laura McCullagh

**Introduction.** Human screening of title and abstracts in a systematic literature review (SLR) is labor intensive and time-consuming. In many instances, thousands of citations may be retrieved; the vast majority excluded upon screening. Text-mining semi-

automates and accelerates screening by identifying patterns in relevant and irrelevant citations, as labelled by the screener. One such text-mining tool, Abstrackr, uses an algorithm within an active-learning framework to predict the likelihood of citations being relevant. The objective of this study was to assess the performance of Abstrackr for title and abstract screening in an SLR of treatments for relapsed/refractory diffuse large B-cell lymphoma.

**Methods.** Citations identified from searches of electronic databases were imported to Abstrackr. An investigator-selected database of terms indicating relevance of title and abstract to the research question were uploaded. These terms were partly informed by the SLR inclusion/exclusion criteria. Citations deemed most relevant by Abstrackr were screened first (screening prioritization). Screening was carried out until a maximum prediction score of 0.4 or less, based on previous experience in the literature, was reached. Remaining citations were deemed unlikely to be relevant and did not undergo screening (screening truncation). Separately, a single-human screener screened all citations using Covidence.

**Results.** A total of 7,723 citations and 154 initial terms were uploaded to Abstrackr. Of these citations, 2,572 (33 percent) were screened before a prediction score of 0.39 was reached. Compared to single-human screening (conducted on all citations), the workload saving associated with Abstrackr was 5 days. A total of 451 (6 percent) citations proceeded to full-text screening; ten (0.1 percent) were included in the final evidence base. No citations predicted to be irrelevant by Abstrackr were included in the final evidence base.

**Conclusions.** Text-mining tools such as Abstrackr have the potential to reduce workload associated with title and abstract screening, without missing relevant citations.

## OP130 Economic Evaluation Of High-Cost Drugs For Relapsing-Remitting Multiple Sclerosis In Thailand

Sarayuth Khuntha, Nuttakarn Budtarad ([nuttakarn.b@hitap.net](mailto:nuttakarn.b@hitap.net)), Pritaporn Kingkaew, Phorntida Hadnorntun and Pattara Leelahavarong

**Introduction.** Drugs for relapsing-remitting multiple sclerosis (RRMS) are costly and not included in the National List of Essential Medicines of Thailand yet. This study aims to conduct an economic evaluation of high-cost drugs for RRMS.

**Methods.** The Markov model was used to estimate lifetime costs and quality-adjusted life years (QALYs) gained. The treatment options include Interferon beta-1a (IFN) and Teriflunomide (TERI) (first-line), Fingolimod (FIN) and Natalizumab (NATA) (second-line), and Alemtuzumab (ALEM) (third-line) compared with usual care. The effectiveness of drugs was retrieved by network meta-analysis. The probability of health state transition was obtained from primary data. Treatment-related costs were derived from the national database. Other costs and utilities were obtained from the study in Thai RRMS patients.

**Results.** The lowest lifetime costs option was usual care (THB2 million) (USD65,808), while the highest QALY gained option

was IFN-NATA-ALEM (8.6 QALY gained). All treatment options were not cost-effective compared with usual care at the threshold of THB160,000 (USD5,300) per QALY gained. However, the option of IFN-NATA-ALEM yielded the lowest incremental cost-effectiveness ratio (ICER), which was THB4.4 million (USD144,778) per QALY gained.

**Conclusions.** High-cost drugs were not cost-effective; nonetheless, the IFN-NATA-ALEM option could increase QALY gained with the lowest additional budget. The government should negotiate the price of IFN to decrease by eighty percent.

### OP135 Impact Of Updated Trial Data On The Cost-effectiveness Of Health Technologies: A Case Study On Percutaneous Mitral Repair

Xavier Armoiry ([xavier.armoiry@univ-lyon1.fr](mailto:xavier.armoiry@univ-lyon1.fr)), Peter Auguste, Jean-François Obadia, Daniel Grinberg and Martin Connock

**Introduction.** Extrapolation methods are commonly used to model the cost-effectiveness of health technologies beyond observed data. Reassessing cost-effectiveness estimates using updated clinical trial data has the potential to reduce uncertainty and optimize decision-making. We present a case study based on percutaneous repair (PR) with the Mitraclip system, a technology to treat severe secondary mitral regurgitation (MR). For the study purpose, we considered the COAPT trial that evaluated the effectiveness of adding PR to medical treatment versus medical treatment alone.

**Methods.** We developed a time-varying Markov model to assess the cost-effectiveness of PR. Clinical inputs were based on reconstructed individual patient data from the COAPT trial results reported at 2 years, and at 3 years.

We developed parametric modeling for overall survival (OS) and heart failure hospitalizations (HFH) to obtain clinically plausible extrapolations beyond observed data. We adopted the French perspective and used a 30-year time horizon. We expressed incremental cost-effectiveness ratios (ICERs) as cost per quality-adjusted life year (QALY).

**Results.** Based on 2 year-data, preferred parametric models for OS and HFH were exponential and log-logistic respectively, yielding an ICER of EUR21,918/QALY and >0.5 probability of PR being cost-effective (EUR50,000/QALY threshold).

Updated analyses at 3 years showed a change of OS trajectory for PR that justified the use of piecewise modelling, yielding an updated ICER that went up to EUR77,904/QALY (base-case), and to a minimum of EUR58,175/QALY (scenario analysis). Using data at 3 years, PR had <0.5 probability of being cost-effective.

**Conclusions.** In this case study, the availability of updated survival analyses of the main trial is likely to have some impact on decision-making and/or pricing discussion as part of health-technology assessment (HTA). We aim to provide further updated analyses as 4 years results of the COAPT study become available.

More broadly, original technology appraisals are frequently undertaken when mid/long-term follow-up trial data may be

lacking. Our example suggests the need for continuous HTA review as new clinical data are released.

### OP164 Extracorporeal Cytokine Adsorption Therapy: An Update Systematic Review Of Clinical Efficacy And Safety For Two Indications

Gregor Goetz ([gregor.goetz@aihta.at](mailto:gregor.goetz@aihta.at)), Katharina Hawlik and Claudia Wild

**Introduction.** The idea of using extracorporeal cytokine adsorption therapy (ECAT) is to remove cytokines from the blood in order to restore a balanced immune response. Yet, it is unclear as to whether the use of ECAT improves patient-relevant outcomes. Hence, the aim of this article is to synthesize the currently available evidence with regard to a potential clinical benefit of ECAT used in cardiac surgery or sepsis.

**Methods.** We conducted an updated systematic review summarizing the body of evidence with regard to a potential clinical benefit of ECAT. The study followed the PRISMA statement and the European Network for Health Technology Assessment (EUnetHTA) guidelines. The quality of the individual studies and the strength of the available evidence was assessed using the Cochrane risk of bias tool (v.1) and the GRADE approach respectively. Mortality, organ function, length of stay in the intensive care unit and length of hospitalization, as well as adverse events, were defined as critical outcomes.

**Results.** For the preventive treatment of ECAT in patients undergoing cardiac surgery, we found very low-quality inconclusive evidence for mortality (5 randomized controlled trials (RCTs), n = 163), length of stay in the intensive care unit (5 RCTs, n = 163), and length of hospitalization (3 RCTs, n = 101). In addition, very low-quality inconclusive evidence was found for (serious) adverse events (4 RCTs, n = 148). For the therapeutic treatment of ECAT in patients with sepsis/ septic shock, we found very low-quality inconclusive evidence for mortality up to 60-day follow-up (2 RCTs, n = 117), organ function (2 RCTs, n = 117) and length of stay in the intensive care unit (1 study, n = 20). Similarly, very low-quality inconclusive evidence was found for (serious) adverse events (2 RCTs, n = 117). There are currently eighteen ongoing RCTs on the use of ECAT.

**Conclusions.** There is a lack of reliable data on the clinical benefit of using ECAT as an add-on treatment preventively in cardiac surgery and therapeutically in patients with sepsis or septic shock. While theoretical advantages are anticipated, the current available evidence is inconclusive and was not able to establish the efficacy and safety of ECAT in combination with standard care in the investigated indications. In light of the available RCTs, we strongly recommend the consideration of studies with patient-relevant endpoints and adequate statistical power, instead of investing further research funds on small studies that may not shed more light onto the potential clinical benefit of ECAT. The results of ongoing RCTs are awaited to guide the decision on whether further research funds should be invested in ECAT research or to conclude that the intervention may not show clinical benefits for patients.