

Original Research


Cite this article: Maimaitiming M, Xie J, Zheng Z, Zhu Y. Effect of the announcement of human-to-human transmission on telemedicine services in China during COVID-19. *Disaster Med Public Health Prep.* 17(e311), 1–7. doi: <https://doi.org/10.1017/dmp.2022.278>.

Keywords: COVID-19; teleconsultation; telehealth; telemedicine

Abbreviations: COVID-19, coronavirus disease; DID, difference-in-differences; TCM, traditional Chinese medicine

Corresponding author: Yongjian Zhu, Email: ustczyj@mail.ustc.edu.cn.

Effect of the Announcement of Human-to-Human Transmission on Telemedicine Services in China During COVID-19

Mairehaba Maimaitiming MSc¹, Jingui Xie PhD², Zhichao Zheng PhD³ and Yongjian Zhu PhD¹ 

¹School of Management, University of Science and Technology of China, Hefei, China; ²School of Management, Technical University of Munich, Heilbronn, Germany and ³Lee Kong Chian School of Business, Singapore Management University, Singapore

Abstract

Objectives: Telemedicine enables patients to communicate with physicians effectively, especially during the coronavirus disease (COVID-19) pandemic. However, few studies have explored the use of online health care platforms for a comprehensive range of specialties during the COVID-19 pandemic. This study aimed to investigate how telemedicine services were affected by the announcement of human-to-human transmission in China.

Methods: Telemedicine data from haodf.com in China were collected. A difference-in-differences analysis compared the number of telemedicine use and the number of active online physicians for different specialties in 2020 with the numbers in 2019, before and after the announcement of human-to-human transmission.

Results: Data from 2 473 734 telemedicine use during the same calendar time in 2020 and 2019 were collected. Telemedicine use in 2020 increased by 349.9% after the announcement of human-to-human transmission in China, and the number of active online physicians increased by 23.2%. The difference-in-differences analysis indicated that the announcement had statistically significant positive effects on the numbers of telemedicine use for almost all specialties, except cosmetic dermatology, pathology, occupational diseases, sports medicine, burn, medical imaging, and interventional medicine.

Conclusion: Telemedicine services increased significantly after the announcement of human-to-human transmission of COVID-19. Online activities of most specialties increased, except where providers had to conduct in-person testing and provide bedside therapies.

The novel coronavirus disease (COVID-19) was first identified in Wuhan, China, in 2019. On January 20, 2020, the National Health Commission of China confirmed human-to-human transmission of COVID-19 and infections among medical staff.¹ Before the announcement, limited attention was given to COVID-19;² however, concerns and panic arose significantly among the public after the announcement.³ Masks and hand hygiene were encouraged in response to the pandemic.⁴ Furthermore, many countries have implemented travel restrictions and asked citizens to stay at home because avoiding contact with others seemed to be the most effective way to control the rapid transmission of the virus.^{5,6}

Under these circumstances, traditional offline health care services faced significant challenges. Face-to-face service was difficult to provide because of government-enforced policies (such as lockdown) and pandemic-related individual motivations such as fear of infection.^{7,8} Conversely, medical care is in higher demand because people tend to care more about their physical health and mental health during community-wide disasters.^{9,10} To cope with this crisis, the Chinese Government issued several mandates to encourage Internet hospitals to provide timely medical services for the public during the pandemic.^{11–13}

Several telemedicine services have been established to address health needs in China during the COVID-19 pandemic, including COVID-19 diagnoses,^{11,12} psychological counseling,¹⁴ online prescription services,¹⁵ and general teleconsultation (such as that provided by Hao Dai Fu: haodf.com, Ding Xiang Yi Sheng: dxy.com, and Chun Yu Yi Sheng: chunyuisheng.com). In-person care provided by some medical institutions in the United States also transitioned to telehealthcare (through video or telephone) due to the COVID-19 pandemic.^{16–18} However, few studies assessed the effect of COVID-19 on online health services.

This study aimed to investigate how telemedicine service was affected by the onset of the COVID-19 pandemic in China, which is recognized as the announcement of human-to-human transmission on January 20, 2020. Using the difference-in-differences (DID) estimation strategy and data from haodf.com, an online health platform, this study investigated the effect of the announcement on telemedicine services from two perspectives, demand and supply. For

demand, this study determined the change in the number of telemedicine services used by various patients. For supply, this study analyzed how the number of active online physicians was affected by the announcement. This study also evaluated the heterogeneous effect of the announcement on different specialties.

Methods

Data

This study focused on one of the most popular online health platforms in China, haodf.com. Since 2006, haodf.com has provided professional medical services online for more than 58 million patients. According to the official website, haodf.com had approximately 610 000 physicians from 9917 hospitals at the end of 2019. Among these, approximately 230 000 physicians registered on the website to provide timely online medical services. Telemedicine service was chosen as the context of this research due to its impact on health care.

The platform provides various medical services, which primarily include telemedicine (online picture and text consultations), telephone consultations, video consultations, family doctor consultations, and private doctor consultations. Patients can leave comments after the service. After selecting a telemedicine service, patients can browse lists of physicians according to disease conditions or preferred hospitals. They can then decide which physician to consult with based on detailed information on the physician's personal website, including the physician's clinical titles, specialties, patient votes and reviews, and the number of previous online consultations. Patients can only begin treatment after registration, and their personal information, such as name, gender, address, and phone number, is not revealed to other users.

This study developed a Python crawling program to download public telemedicine data from different specialties. For each consultation, this study extracted physician-related information (name, hospital, specialty area) and date of consultation. The consultation data set was extracted for the period from January 1, 2020, to February 16, 2020, which covers 24 days before and 22 days after the Chinese New Year on January 25, 2020. In addition, to avoid the influence of the Chinese New Year, data from all online consultations for the period from January 12, 2019, to February 27, 2019, matched on the lunar calendar, were collected.

This study computed the daily number of telemedicine use in all specialties and the daily number of telemedicine use from individual specialties. This study then calculated the number of physicians consulted each day and defined it as the daily number of active online physicians.

Empirical Model

To examine the causal impact of the announcement of human-to-human transmission on telemedicine services in China, this study applied a DID method using 2019 data as the control group and 2020 data as the treatment group, which is a common method used in COVID-19 studies.^{5,19} Specifically, this study compared the difference in the number of telemedicine use or active online physicians before and after the announcement in 2020 with the difference in the corresponding period in 2019 (Supplement Figure 1). The model is specified as follows:

Table 1. Definitions of variables

Variables	Definitions
$\ln(N_{i,t})$	The number of telemedicine use or active online physicians on day t year i (logarithm)
$Treat_t$	Binary, 1 if the year is 2020, 0 if the year is 2019
$After_t$	Binary, 1 if the day is after known human transmission in 2020 or the corresponding date in 2019, and 0 otherwise
$Date_t$	Date fixed effect
i	Year
t	Day

$$\ln(N_{i,t}) = \alpha + \beta * Treat_t * After_t + Treat_t + Date_t + \varepsilon_{i,t} \quad (1)$$

Table 1 illustrates the definition of each variable in model (1). Here, i denotes the year and t denotes the day. The dependent variable $\ln(N_{i,t})$ represents the logarithmic number of telemedicine use or active online physicians for the year i on day t . The dummy variable $Treat_t$ equals 1 if the year is 2020 and 0 if the year is 2019, which suggests that the control data are the year 2019. The dummy variable $After_t$ equals 1 for the period from January 20, 2020, to February 16, 2020, and the same lunar calendar period from January 31, 2019, to February 27, 2019, and 0 for the period from January 1, 2020, to January 19, 2020, and the same lunar calendar period from January 12, 2019, to January 30, 2019. $Date_t$ is the date fixed effect, which includes a set of time dummies to control for shocks that are common to treatment and control in a given day, such as the Chinese New Year and the Lantern Festival. The $\varepsilon_{i,t}$ is the error term. Our coefficient of interest is β , which can be used to estimate the difference in the number of telemedicine use or active online physicians between the treatment year 2020 and the control year 2019 before and after the announcement of human-to-human transmission. Then analysis was repeated using equation (1) for each specialty separately.

Test for Parallel Trends

The assumption for the DID estimation is that before the announcement of human-to-human transmission, the telemedicine use and active online physicians in 2019 and 2020 would show parallel trends, indicating the difference between the treatment year and the control year is constant over time. This study tested this assumption by using the same method in He et al.'s study,²⁰ which also helped capture the changes in the effect over time:

$$\ln(N_{i,t}) = \alpha + \sum_k \beta^k * Treat_t * After_{t,k} + Treat_t + Date_t + \varepsilon_{i,t} \quad (2)$$

where $After_{t,k}$ are a set of dummy variables and $k \in \{-3, -2, -1, 0, 1, 2, 3\}$ indicates the relative k th week of the announcement of human-to-human transmission. The benchmark period is 1 week before the announcement so that the coefficient β^k can be used to measure the difference in the difference in telemedicine use or active online physicians in 2019 and 2020 in period k and the difference of 1 week before the announcement. If the preannouncement trends are parallel, β^k would be statistically insignificant when $k < -1$.




Figure 1. The daily number of telemedicine use during the same lunar calendar periods in 2019 and 2020 from 24 days before the Chinese New Year to 22 days after the Chinese New Year on January 25, 2020. The study periods were from January 1, 2020, to February 16, 2020, and from January 12, 2019, to February 27, 2019 (matched by the lunar calendar). The x-axis shows the number of days before the Chinese New Year and after the Chinese New Year, based on the 2020 solar calendar.




Figure 2. The number of telemedicine use in each specialty during the study periods in 2019 and 2020. The study periods were from January 1, 2020, to February 16, 2020, and from January 12, 2019, to February 27, 2019 (matched by the lunar calendar).

Results

After removing the duplicated and unqualified data (consultation sets visible to the physician only), this study obtained data on 2 473 734 telemedicine use. Figure 1 shows the daily number of telemedicine use during the same lunar calendar periods in 2019 and 2020. Internal medicine has the highest consultation percentage at 43% among all specialties in 2020 (Figure 2). Compared with the same period in 2019, telemedicine use in 2020 had increased in 12 specialties, including traditional Chinese medicine (TCM); integrated TCM and Western medicine; infectious diseases; pediatrics; internal medicine; psychiatry; obstetrics and gynecology; rehabilitation medicine; dermatology and sexually

transmitted diseases; head and neck otorhinolaryngology; and nutriology.

Figure 3 plots the daily number of active online physicians during the study periods. Although the daily number of active online physicians was generally fewer in 2020, it increased at a faster rate after the announcement of human-to-human transmission. In line with the number of telemedicine use in 2020, physicians practicing internal medicine accounted for the largest proportion of total active online physicians (Figure 4).

Table 2 shows the DID estimates for the number of telemedicine use and the number of active online physicians, using the model (1). The coefficients are significantly positive, which show




Figure 3. The daily number of active online physicians during the study periods in 2019 and 2020, from 24 days before to 22 days after the Chinese New Year on January 25, 2020. The study periods were from January 1, 2020, to February 16, 2020, and from January 12, 2019, to February 27, 2019 (matched by the lunar calendar). The x-axis shows the number of days before the Chinese New Year and after the Chinese New Year, based on the 2020 solar calendar.




Figure 4. The number of active online physicians in each specialty during the study periods in 2019 and 2020. The study periods were from January 1, 2020, to February 16, 2020, and from January 12, 2019, to February 27, 2019 (matched by the lunar calendar).

that the announcement of human-to-human transmission significantly increased telemedicine use by 349.9% ($= \exp(1.504) - 1$) relative to the same period in 2019, and increased the number of active online physicians by 23.2% ($= \exp(0.209) - 1$) relative to the same period in 2019.

Table 3 reports the coefficients of *Treat_t* * *After_t* in each specialty for the number of telemedicine use and the number of active online physicians. This study found that the announcement of human-to-human transmission had different impacts across specialties for both outcomes. For the number of telemedicine, the effect was positive in internal medicine; TCM; integrated TCM and Western medicine; pediatrics; head and neck otorhinolaryngology;

nutriology; dermatology and sexually transmitted diseases; obstetrics and gynecology; rehabilitation medicine; infectious diseases; stomatology; ophthalmology; psychiatry; andrology; orthopedic surgery; surgery; anesthesiology; tuberculosis; reproductive medicine; oncology; and others. The number of teleconsultations for cosmetic dermatology was negatively impacted by the announcement of human-to-human transmission, and the consultations from the rest of the specialties did not differ between 2020 and 2019.

Similarly, the announcement increased the number of active online physicians from internal medicine, TCM, integrated TCM and Western medicine, pediatrics, head and neck otorhinolaryngology, nutriology, dermatology and sexually transmitted

Table 2. Impact of the announcement of human-to-human transmission on the number of telemedicine use and active online physicians

Variables	Dependent variable	
	Ln (number of telemedicine)	Ln (number of active online physicians)
Treat*After	1.504*** (0.149)	0.209*** (0.050)
Treat	Yes	Yes
Date fixed effects	Yes	Yes
Observations	94	94
R ²	0.8523	0.9333

Note: The dependent variables are the logarithm number of telemedicine use (left column) and the logarithm number of active online physicians (right column).

Robust standard errors are shown in parentheses.

*P < 0.1; **P < 0.05; ***P < 0.01.

diseases, obstetrics and gynecology, psychiatry, tuberculosis, reproductive medicine, and others. However, in specialties for which telemedicine use increased significantly, such as rehabilitation medicine and stomatology, the number of active online physicians did not differ and even decreased in infectious diseases. The number of active online physicians also decreased in medical imaging, burn, sports medicine, occupational diseases, and cosmetic dermatology.

The results of the parallel trends tests showed that no preannouncement differences occurred in the telemedicine trend and active online physicians trend between 2019 and 2020 (Supplement Figure 2). The estimated coefficients β^k were all statistically insignificant when $k < -1$, which supported the parallel trends assumption. The positive effect on telemedicine use was significant during the week of the announcement of human-to-human transmission and became stronger over time during the study period after the announcement (k_0). However, for active online physicians, the positive effect only became significant 2 weeks after the announcement, indicating that the number of active online physicians increased 2 weeks later (k_2). This may be due to the increased offline pandemic-related workload experienced by physicians at the initial stage of the announcement.²¹

Discussion

This study quantified the influence of the announcement of human-to-human transmission on an online health care platform in China. This study found that, after the announcement on January 20, 2020, there was a significant increase in the use of telemedicine services on haodf.com compared with the same period in 2019. Specifically, the top 5 teleconsultation specialties were internal medicine, TCM, integrated TCM and Western medicine, pediatrics, and head and neck otorhinolaryngology. In addition, this study observed that the announcement also led to an increase in active physicians on this online health care platform.

For the impact on telemedicine use, there are several possible mechanisms. First, before the announcement of human-to-human transmission on January 20, 2020, the coronavirus was believed to be controllable, and there were no effective measures taken to prevent the spread of the virus.⁵ However, the increased risk of this novel virus after the announcement gained the attention of the Chinese Government and the public. The lockdown policy was imposed on Wuhan, China, on January 23, 2020, and travel

Table 3. Impact of the announcement of human-to-human transmission on the number of telemedicine use and the number of active online physicians in each specialty

Specialty	β for the number of telemedicine use	β for the number of active online physicians
Internal medicine	2.318***	0.452***
TCM	2.083***	0.394***
Integrated TCM and Western medicine	1.941***	0.341***
Pediatrics	1.828***	0.431***
Head and neck otorhinolaryngology	1.391***	0.249***
Nutrition	1.168***	0.587**
Dermatology and sexually transmitted diseases	1.070***	0.346***
Obstetrics and gynecology	0.958***	0.262***
Rehabilitation medicine	0.838***	-0.072
Stomatology	0.734***	-0.112
Ophthalmology	0.715***	0.068
Psychiatry	0.682***	0.124**
Infectious diseases	0.669***	-0.146**
Andrology	0.646***	-0.001
Orthopedic surgery	0.631***	-0.055
Surgery	0.519***	-0.032
Anesthesiology	0.426**	-0.101
Tuberculosis	0.391**	0.255**
Reproductive medicine	0.305**	0.249***
Oncology	0.290**	0.082
Interventional medicine	0.208	-0.062
Medical imaging	0.208	-0.184*
Burn	-0.144	-0.246**
Sports medicine	-0.233	-0.246**
Occupational diseases	-0.326	-0.561***
Pathology	-0.331	-0.218
Cosmetic dermatology	-0.515***	-0.264***
Others	2.777***	0.857***

Note: Estimated coefficients β are indicated for interaction term *Treat_t * After_t* in equation (1) for each specialty.

*P < 0.1; **P < 0.05; ***P < 0.01.

restrictions were subsequently imposed in other cities.²² Social gathering was also discouraged.²³ Although such policies effectively curbed the spread of the virus, they also inhibited in-person doctor visits. Thus, telemedicine services have become more essential than previously because they can provide timely services regardless of logistical restrictions.

Second, due to the spread of the novel coronavirus, the demand for medical services increased. Reduced availability of offline medical resources could only be used by limited numbers of patients with symptoms related to COVID-19.²⁴ Patients with common diseases found it hard to get treatment offline. In addition, the novelty of the virus and the overwhelming news coverage of COVID-19 caused public panic.²⁵ People were also worried about being infected by patients with undetected COVID-19 due to the lack of testing equipment.²⁶ Online consultation became the most efficient approach to meeting patients' medical needs under these circumstances.

This study also showed that several specialties were most likely to be consulted after the announcement of human-to-human transmission. The primary symptoms of COVID-19 are fever, fatigue, and cough, sometimes accompanied by a runny nose, nasal congestion, and other upper respiratory tract symptoms.^{4,27} Patients with any of these symptoms were anxious to determine whether they had been infected. Because internal medicine included respiratory medicine on haodf.com, consultations in this specialty accounted for the largest number of consultations among all specialties. Traditional Chinese medicine is widely used in China in treating viral and bacterial infections.^{28,29} In severe acute respiratory syndrome and the Middle East respiratory syndrome coronavirus outbreaks, TCM was used as a supplementary treatment resulting in significant patient improvement.³⁰ Several meta-analyses showed that TCM integrated with Western medicine had better effects against COVID-19, such as reducing clinical deterioration compared with using Western medicine alone in clinical cure.^{31,32} Because no specific anti-virus drugs or vaccines were available at the onset of the pandemic, TCM was used to ameliorate COVID-19-related symptoms.²⁸ Traditional Chinese medicine was also included in the diagnosis and treatment plan for COVID-19 issued on January 22, 2020.^{27,33} Thus, many people chose to consult physicians of TCM for treatment.

The increase in the number of active physicians attributed to the announcement shows that physicians from haodf.com actively responded to the unprecedented medical demands during the pandemic. The novelty of the virus and the scarcity of medical resources during the initial outbreak caught physicians unprepared, and, thus, the number of active physicians did not increase immediately. With the growing knowledge about disease control and the improvement in medical resource management,³⁴ physicians could better allocate their time to cope with the high volume of consultations.

There were studies reporting the changes in the use of telemedicine during the initial phase of the COVID-19 pandemic. Internet consultations increased in 2 hospitals in China after promoting telehealth platforms in response to the pandemic.^{11,24} Similarly, post-COVID-19 video/telephone visits increased significantly after the implementation of video/telephone consultation services in the United States in March 2020.^{16,17,35,36} This study also indicated a significant increase in telemedicine use during COVID-19. This study adds to this body of literature by quantifying the relative change in telemedicine services due to COVID-19 from the perspectives of both patients and physicians. Compared with the previous studies that mainly evaluate telemedicine use in 1 hospital/medical institution, this studied platform includes hospitals nationwide and 28 medical specialties. In addition to considering the pre-COVID-19 period in 2020, which was included in other studies, this study used telemedicine services provided in 2019 as the control year and applied the DID method to compare the number of telemedicine use and the number of active online physicians. Moreover, this study evaluated the telemedicine service during the pandemic by specialty.

Our findings have several practical implications. First, the increased telemedicine use indicated that telemedicine service had become an important approach to satisfying people's medical needs during COVID-19. The government should further encourage the use of telemedicine to cope with the COVID-19 situation and other public health care emergencies. Second, telemedicine use increased significantly during the week of the announcement of human-to-human transmission, whereas the number of active online physicians increased only 2 weeks after the announcement.

This may suggest the lack of online physicians and lower service quality. The platform should design an effective mechanism to attract physicians to provide online services as early as possible. Third, the heterogeneous effect indicated that several specialties were most likely to be affected during the pandemic, so the platform should pay more attention to encouraging the participation of online physicians from these specialties.

Limitations

Several limitations should be acknowledged. First, the telemedicine data were collected from a single website. Although it is one of the most popular online health platforms in China, the website can only serve a limited number of patients. Second, the online consultation process was anonymous, and, thus, this study could not collect detailed information about patient gender, age, and address for additional analysis. Third, this study did not analyze the unstructured data of patient-to-physician interactions, such as the consultation text.

Conclusions

In conclusion, this study has shown that the use of telemedicine and the number of active online physicians increased by 349.9% and 23.2%, respectively, after the announcement of human-to-human transmission of COVID-19. Specifically, telemedicine usage increased immediately after the announcement, whereas active online physicians only increased significantly 2 weeks after the announcement. Online activities of most specialties (including internal medicine, TCM, integrated TCM and Western medicine, pediatrics, etc) increased, except where providers must conduct in-person testing and provide bedside therapies (eg, cosmetic dermatology, pathology, occupational diseases, sports medicine). Therefore, telemedicine could be leveraged to meet the patient demand for most health issues during the pandemic, and telemedicine services should be well supported by institutions and governments when facing a public health crisis.

Supplementary material. To view supplementary material for this article, please visit <https://doi.org/10.1017/dmp.2022.278>.

Author contributions. All authors contributed to the study conception and design. Mairehaba Maimaitiming: conceptualization, data curation, formal analysis, investigation, methodology, software, visualization, and writing – original draft; Yongjian Zhu: conceptualization, validation, and writing – original draft; Jingui Xie: funding acquisition, supervision, writing – reviewing and editing; Zhichao Zheng: funding acquisition, supervision, writing – reviewing and editing.

Consent to participate. The website indicates that the telemedicine contents are available online after anonymization. Patients can set the consultation content only visible to the physician if they do not want to display the information on the platform.

Funding statement. This work was supported by the National Natural Science Foundation of China (grant numbers 71921001, 71771202) and by the Ministry of Education, Singapore, under its Academic Research Fund (AcRF) Tier 2 (grant number MOE2019-T2-1-185). The funder has no involvement in conducting or preparing the article.

Conflict(s) of interest. None.

Ethical standards. The telemedicine data from the website haodf.com are displayed after anonymization. The website indicates that patients' personal information is collected and used in accordance with the network security law of the

People's Republic of China, Information Security Technology – Personal Information Security Specification (GB/T 35273-2017), and other relevant laws, regulations, and technical specifications.

References

1. **The National Health Commission of the People's Republic of China and Relevant Departments Jointly Responded to the Coronavirus.** National Health Commission of the People's Republic of China. Published January 21, 2020. Accessed September 3, 2021, <http://www.nhc.gov.cn/xcs/fkdt/202001/d9570f3a526141133ae0093df51509684.shtml>
2. **Zhu Y, Fu KW, Grepin KA, et al.** Limited early warnings and public attention to coronavirus disease 2019 in China, January–February, 2020: a longitudinal cohort of randomly sampled Weibo users. *Disaster Med Public Health Prep.* 2020;14(5):e24–e27. doi: [10.1017/dmp.2020.68](https://doi.org/10.1017/dmp.2020.68)
3. **Wu Y, Zhang C, Liu H, et al.** Perinatal depressive and anxiety symptoms of pregnant women during the coronavirus disease 2019 outbreak in China. *Am J Obstet Gynecol.* 2020;223(2):240–e1. doi: [10.1016/j.ajog.2020.05.009](https://doi.org/10.1016/j.ajog.2020.05.009)
4. **Pascarella G, Strumia A, Piliago C, et al.** COVID-19 diagnosis and management: a comprehensive review. *J Intern Med.* 2020;288(2):192–206. doi: [10.1111/joim.13091](https://doi.org/10.1111/joim.13091)
5. **Fang H, Wang L, Yang Y.** Human mobility restrictions and the spread of the novel coronavirus (2019-nCoV) in China. *J Public Econ.* 2020;191:104272. doi: [10.1016/j.jpubeco.2020.104272](https://doi.org/10.1016/j.jpubeco.2020.104272)
6. **Aliakbar Kabiri AD, Zhou W, Sun Q, Zhang L.** How different age groups responded to the COVID-19 pandemic in terms of mobility behaviors: a case study of the United States. 2020;arXiv preprint arXiv:2007.10436.
7. **Chudasama YV, Gillies CL, Zaccardi F, et al.** Impact of COVID-19 on routine care for chronic diseases: a global survey of views from healthcare professionals. *Diabetes Metab Syndr.* 2020;14(5):965–967. doi: [10.1016/j.dsx.2020.06.042](https://doi.org/10.1016/j.dsx.2020.06.042)
8. **Guarino M, Cossiga V, Fiorentino A, et al.** Use of telemedicine for chronic liver disease at a single care center during the COVID-19 pandemic: prospective observational study. *J Med Internet Res.* 2020;22(9):e20874. doi: [10.2196/20874](https://doi.org/10.2196/20874)
9. **Zhong B, Huang Y, Liu Q.** Mental health toll from the coronavirus: social media usage reveals Wuhan residents' depression and secondary trauma in the COVID-19 outbreak. *Comput Human Behav.* 2021;114:106524. doi: [10.1016/j.chb.2020.106524](https://doi.org/10.1016/j.chb.2020.106524)
10. **Niu Z, Wang T, Hu P, et al.** Chinese public's engagement in preventive and intervening health behaviors during the early breakout of COVID-19: cross-sectional study. *J Med Internet Res.* 2020;22(8):e19995. doi: [10.2196/19995](https://doi.org/10.2196/19995)
11. **Li L, Liu G, Xu W, et al.** Effects of Internet hospital consultations on psychological burdens and disease knowledge during the early outbreak of COVID-19 in China: cross-sectional survey study. *J Med Internet Res.* 2020;22(8):e19551. doi: [10.2196/19551](https://doi.org/10.2196/19551)
12. **Gong K, Xu Z, Cai Z, et al.** Internet hospitals help prevent and control the epidemic of COVID-19 in China: multicenter user profiling study. *J Med Internet Res.* 2020;22(4):e18908. doi: [10.2196/18908](https://doi.org/10.2196/18908)
13. **Notice of Providing Internet Diagnosis and Consulting Services During the COVID-19 from the National Health Commission of the People's Republic of China.** National Health Commission of the People's Republic of China. Published February 7, 2020. Accessed September 3, 2021. <http://www.nhc.gov.cn/yzygj/s7653p/202002/ec5e345814e744398c2adef17b657fb8.shtml>
14. **Liu S, Yang L, Zhang C, et al.** Online mental health services in China during the COVID-19 outbreak. *Lancet Psychiatry.* 2020;7(4):e17–e18. doi: [10.1016/S2215-0366\(20\)30077-8](https://doi.org/10.1016/S2215-0366(20)30077-8)
15. **Ding L, She Q, Chen F, et al.** The Internet hospital plus drug delivery platform for health management during the COVID-19 pandemic: observational study. *J Med Internet Res.* 2020;22(8):e19678. doi: [10.2196/19678](https://doi.org/10.2196/19678)
16. **Sinha S, Kern LM, Gingras LF, et al.** Implementation of video visits during COVID-19: lessons learned from a primary care practice in New York City. *Front Public Health.* 2020;8:514. doi: [10.3389/fpubh.2020.00514](https://doi.org/10.3389/fpubh.2020.00514)
17. **Baum A, Kaboli PJ, Schwartz MD.** Reduced in-person and increased telehealth outpatient visits during the COVID-19 pandemic. *Ann Intern Med.* 2021;174(1):129–131. doi: [10.7326/M20-3026](https://doi.org/10.7326/M20-3026)
18. **Mann DM, Chen J, Chunara R, et al.** COVID-19 transforms health care through telemedicine: evidence from the field. *J Am Med Inform Assoc.* 2020;27(7):1132–1135. doi: [10.1093/jamia/ocaa072](https://doi.org/10.1093/jamia/ocaa072)
19. **Tanaka T, Okamoto S.** Increase in suicide following an initial decline during the COVID-19 pandemic in Japan. *Nat Hum Behav.* 2021;5(2):229–238. doi: [10.1038/s41562-020-01042-z](https://doi.org/10.1038/s41562-020-01042-z)
20. **He G, Pan Y, Tanaka T.** The short-term impacts of COVID-19 lockdown on urban air pollution in China. *Nat Sustain.* 2020;3(12):1005–1011. doi: [10.1038/s41893-020-0581-y](https://doi.org/10.1038/s41893-020-0581-y)
21. **Wang L, Yan L, Zhou T, et al.** Understanding physicians' online-offline behavior dynamics: an empirical study. *Inform Syst Res.* 2020;31(2):537–555. doi: [10.1287/isre.2019.0901](https://doi.org/10.1287/isre.2019.0901)
22. **Kraemer MUG, Yang CH, Gutierrez B, et al.** The effect of human mobility and control measures on the COVID-19 epidemic in China. *Science.* 2020;368(6490):493–497. doi: [10.1126/science.abb4218](https://doi.org/10.1126/science.abb4218)
23. **Chen S, Yang J, Yang W, et al.** COVID-19 control in China during mass population movements at New Year. *Lancet.* 2020;395(10226):764–766. doi: [10.1016/s0140-6736\(20\)30421-9](https://doi.org/10.1016/s0140-6736(20)30421-9)
24. **Li P, Liu X, Mason E, et al.** How telemedicine integrated into China's anti-COVID-19 strategies: case from a national referral center. *BMJ Health Care Inform.* 2020;27(3):e100164. doi: [10.1136/bmjhci-2020-100164](https://doi.org/10.1136/bmjhci-2020-100164)
25. **Laato S, Islam AKMN, Islam MN, Whelan E.** What drives unverified information sharing and cyberchondria during the COVID-19 pandemic? *Eur J Inform Syst.* 2020;29(3):288–305. doi: [10.1080/0960085x.2020.1770632](https://doi.org/10.1080/0960085x.2020.1770632)
26. **Shangguan Z, Wang MY, Sun W.** What caused the outbreak of COVID-19 in China: from the perspective of crisis management. *Int J Environ Res Public Health.* 2020;17(9):3279. doi: [10.3390/ijerph17093279](https://doi.org/10.3390/ijerph17093279)
27. **Notification on the Diagnosis and Treatment Plan for the Coronavirus Infection (the Third Trial Version).** National Health Commission of the People's Republic of China. Published January 23, 2020. Accessed September 3, 2021. <http://www.nhc.gov.cn/yzygj/s7653p/202001/f492c9153ea9437bb587ce2ffcbee1fa.shtml>
28. **Yang Y, Islam MS, Wang J, et al.** Traditional Chinese medicine in the treatment of patients infected with 2019-new coronavirus (SARS-CoV-2): a review and perspective. *Int J Biol Sci.* 2020;16(10):1708–1717. doi: [10.7150/ijbs.45538](https://doi.org/10.7150/ijbs.45538)
29. **Zhao Z, Li Y, Zhou L, et al.** Prevention and treatment of COVID-19 using Traditional Chinese medicine: a review. *Phytomedicine.* 2021;85:153308. doi: [10.1016/j.phymed.2020.153308](https://doi.org/10.1016/j.phymed.2020.153308)
30. **Ren JL, Zhang AH, Wang XJ.** Traditional Chinese medicine for COVID-19 treatment. *Pharmacol Res.* 2020;155:104743. doi: [10.1016/j.phrs.2020.104743](https://doi.org/10.1016/j.phrs.2020.104743)
31. **Liu M, Gao Y, Yuan Y, et al.** Efficacy and safety of integrated traditional Chinese and Western medicine for corona virus disease 2019 (COVID-19): a systematic review and meta-analysis. *Pharmacol Res.* 2020;158:104896. doi: [10.1016/j.phrs.2020.104896](https://doi.org/10.1016/j.phrs.2020.104896)
32. **Wang H, Xu B, Zhang Y, et al.** Efficacy and safety of traditional Chinese medicine in coronavirus disease 2019 (COVID-19): a systematic review and meta-analysis. *Front Pharmacol.* 2021;12:609213. doi: [10.3389/fphar.2021.609213](https://doi.org/10.3389/fphar.2021.609213)
33. **Zhang L, Yu J, Zhou Y, et al.** Becoming a faithful defender: traditional Chinese medicine against coronavirus disease 2019 (COVID-19). *Am J Chin Med.* 2020;48(4):763–777. doi: [10.1142/S0192415X2050038X](https://doi.org/10.1142/S0192415X2050038X)
34. **Pan A, Liu L, Wang C, et al.** Association of public health interventions with the epidemiology of the COVID-19 outbreak in Wuhan, China. *JAMA.* 2020;323(19):1915–1923. doi: [10.1001/jama.2020.6130](https://doi.org/10.1001/jama.2020.6130)
35. **Loneragan PE, Washington Iii SL, Branagan L, et al.** Rapid utilization of telehealth in a comprehensive cancer center as a response to COVID-19: cross-sectional analysis. *J Med Internet Res.* 2020;22(7):e19322. doi: [10.2196/19322](https://doi.org/10.2196/19322)
36. **Gilson SF, Umscheid CA, Laiteerapong N, et al.** Growth of ambulatory virtual visits and differential use by patient sociodemographics at one urban academic medical center during the COVID-19 pandemic: retrospective analysis. *JMIR Med Inform.* 2020;8(12):e24544. doi: [10.2196/24544](https://doi.org/10.2196/24544)