

NR7. Liaison psychiatry

Chairmen: A Farmer, R Mayou

PSYCHIATRIC MORBIDITY IN THE BONE MARROW TRANSPLANTATION SETTING

J. Blanch, J.M. Prieto, J. Atala, C. Gastó, E. Cirera. *Department of Consultation-Liaison Psychiatry, University of Barcelona, Villarroel 170, 08036, Barcelona, Spain*

Objective: This prospective in-patient study was conducted to determine the nature, extent, and timing of psychiatric morbidity in a group of patients hospitalized for bone marrow transplantation.

Method: A consecutive series of 103 patients admitted at the Hospital Clinic in Barcelona were evaluated on admission and weekly until discharge. We used DSM-IV criteria for diagnosis as well as the Hospital Anxiety Depression Scale to monitor affective symptomatology. Endicott's revised criteria (substitution of somatic symptoms by non-somatic alternatives) were used to diagnose major depression. **Results:** Forty-nine (48%) patients received at least one DSM-IV diagnosis. We found the following prevalence rates: adjustment disorders 29%, major depression 10%, other anxiety and mood disorders 3%, medication-induced disorders 16%, delirium 6%, and personality disorders 3%. When only considering anxiety and mood disorders, we diagnosed 28 (27%) patients on admission and 15 (15%) during hospitalization; 28% of these affective disorders persisted until discharge whereas 37% were considered to last \leq than 2 weeks. Psychotropic medication was prescribed in a high percentage of patients: antihistamines 69%, benzodiazepines 64%, methadone 46%, neuroleptics 25%, and antidepressants 12%. **Conclusion:** High prevalence of psychiatric disturbances (mainly affective disorders) is observed in this population. Consultation-liaison psychiatry plays an important role in the management of these patients.

THE CARDIFF CHRONIC FATIGUE SYNDROME (CFS)

Anne Farmer, Irene Jones, Andy Smith, Lesek Borysiewicz, Meirion Llewellyn. *University of Wales College of Medicine, Department of Psychological Medicine Heath Park, Cardiff, South Glamorgan, CF4 4XN, UK*

The Cardiff Chronic Fatigue Syndrome (CFS) study consists of medical, psychological and psychiatric research into this complex disorder. Initial assessments comparing the first cohort of 100 CFS subjects and 50 healthy controls will be presented.

Associated psychiatric morbidity as defined by ICD-10 and DSM-III-R operational criteria in the Cardiff group will be compared with other studies and the problems of using self rating screening questionnaires for case finding will be highlighted.

Lastly the preliminary psychometric test results will be discussed which show increased fatigability for repeated motor tests in CFS subjects with associated severe sleep disorder compared to those with only mild sleep disturbance and normal controls.

EMOTIONAL ADAPTATION OF CHILDREN IN THE LIFE ISLAND AFTER BONE MARROW TRANSPLANTATION

M. Günter¹, M. Karle¹, A. Werning¹, T. Klingebiel². ¹ *Dept. of Child and Adolescent Psychiatry and Psychotherapy, Osianderstr. 14, D-72076 Tübingen, Germany;* ² *Pediatric Hospital; University of Tübingen, Osianderstr. 14, D-72076 Tübingen, Germany*

Stress reactions and psychic adaptation of 8 to 12-year-old children

in the life island after bone marrow transplantation (BMT) were examined by free diagnostic interviews and by a large test battery. The study's design and initial results from the intense examination are given. As of yet, a mixture of projective tests (Rorschach, Thematic Apperception Test, Draw-a-person-test), questionnaires, two tests recording the body image and an intelligence test (HAWIK-R) proved to be very instructive. The questionnaires show clearly the strong tendencies to adapt to the situation and to normalise behaviour under isolation conditions. In contrast, the psychoanalytic interview and the projective tests prove a very differentiated emotional adjustment: Before transplantation, coming to terms with the life-threatening situation must be deferred. After transplantation one can detect a very intensive emotional examination of the illness, the future prospects, the isolation and the loneliness, the somatic problems and fantasies of guilt and punishment in addition to a superficial adaptation in the sense of a protective denial. Experiencing loving care seems to be most important for emotional stabilisation in this situation. Very impressive documents of these psychic processes are the patients pictures.

EMOTIONAL DISTRESS AND OCCUPATIONAL BURNOUT IN NURSES AND PHYSICIANS DEALING WITH H.I.V. PATIENTS: A COMPARISON WITH ONCOLOGY AND INTERNAL MEDICINE SERVICES

J.J. López-Castillo¹, M. Gurpegui², J.L. Ayuso-Mateos³, J.D. Luna⁴. ¹ *Dept. of Psychiatry, Ciudad de Baza Hospital, Granada, Spain;* ² *Dept. of Psychiatry, San Cecilio University Hospital, Granada, Spain;* ³ *Dept. of Psychiatry, Valdecilla University Hospital, Santander, Spain;* ⁴ *Dept. of Statistics, Universidad de Granada, Granada, Spain*

Objectives: To determine levels of emotional distress and professional burnout in staff working with HIV patients.

Methods: A cross-sectional anonymous survey was collected from 196 individuals in four different public hospitals. Participants were 55 subjects from three Infectious Disease Units, 30 from an Haemophilia Unit (both groups dealing with HIV patients), 41 from two Oncology Units, and 70 from three Internal Medicine Units. Psychological condition, occupational stress, work and social adjustment, and information about stressors, supports and coping methods of the health care workers were evaluated by means of the General Health Questionnaire (GHQ-28), the Maslach Burnout Inventory (MBI), the Modified Social Adjustment Scale (SAS), and other self-report questionnaires.

Results: High levels of psychological morbidity were found in 38.5% of the participants, with significant differences between groups (Haemophilia staff were less affected). Stress level was significantly higher in both Oncology and Internal Medicine staff, with significant differences among groups. On multivariate analysis, HIV groups were more affected than the other two groups. Training was the principal method of coping in all the groups.

Conclusion: Health care workers suffer from significant levels of emotional distress and occupational burnout. HIV staff were more affected, a fact probably due to distinct HIV risk group in Spain, where drug users are predominant among the HIV-affected population.

AFFECTIVE DISORDERS IN THE PARTNERS OF CANCER PATIENTS: PREVALENCE, COURSE & RISK FACTORS

Peter Haddad, Carolyn Pitceathly, Beverley Jones, Peter Maguire. *Cancer Research Campaign Psychological Medicine Group, Stanley House, Wilmslow Road, Withington, Manchester, M20 9BX, England*

A representative sample of newly diagnosed cancer patients was fol-

lowed for 26 months after cancer diagnosis. They and their partners were assessed using semistructured interviews conducted by trained interviewers. Period prevalence rates were calculated for the 26 months following cancer diagnosis for DSM-III-R major depressive disorder (MDD) and total affective disorders (i.e. DSM-III-R MDD, DSM-III generalised anxiety disorder and minor affective disorders). Potential risk factors were assessed.

The 26 month period prevalence of total affective disorder (TAD) in female partners (12/50, 24.0%) did not differ significantly to that in male cancer patients (13/50, 26.0%), McNemar, $p = 1.000$. The prevalence of TAD in male partners (8/118, 6.8%) was significantly less than in female cancer patients (27/118, 22.9%), McNemar, $p < 0.001$. In both male and female partners the majority of affective disorders were accounted for by MDD. The prevalence of MDD in female partners was particularly high (10/50, 20.0%).

Most disorders in partners commenced 14–26 months after cancer diagnosis, significantly later than for cancer patients (McNemar, $p < 0.001$). The median duration of disorder in partners was 12 weeks, though this was a conservative estimate as nearly half of disorders were ongoing at the end of the study. Most of the 20 partners with affective disorder regarded their spouses' cancer as the cause of their disorder but only 12 consulted a doctor. Affective disorder in a partner was significantly associated with affective disorder in the corresponding cancer patient. Multivariate analysis showed three independent contributions to partners' affective disorder i.e. lower social class, female sex and past psychiatric history.

This is the only existing study to have used a standardised interview and operational criteria to compare rates of psychiatric disorder in cancer patients and their partners. Partners, particularly females, are at high risk of affective disorders. This raises the question of whether relatives/carers of patients with other chronic illnesses also suffer from high rates of psychiatric morbidity. Nearly half of partners with affective disorder did not receive treatment emphasising the importance of education for the public and health care professionals. The identification of risk factors may facilitate primary and secondary prevention of these disorders.

A DOUBLE-BLIND PLACEBO CONTROLLED TREATMENT TRIAL OF FLUOXETINE AND GRADED EXERCISE FOR CHRONIC FATIGUE SYNDROME (CFS)

R. Morriss, A. Wearden, R. Mullis, P. Strickland, L. Appleby, I. Campbell, D. Pearson. *University of Manchester, Department of Psychiatry, Medicine, Manchester M20 8LR United Kingdom; Anaesthetics, Withington Hospital, Nell Lane, West Didsbury, Manchester M20 8LR United Kingdom*

Objectives: To test the efficacy and acceptability to patients of two widely available treatments for chronic fatigue syndrome (CFS), with or without DSM-III-R depressive disorders, fluoxetine (20 mg/day) and graded exercise.

Methods: Six month prospective placebo and time with therapist controlled randomised trial with allocation to one of four treatment cells: exercise and fluoxetine; exercise and placebo drug; appointments and fluoxetine; appointments and placebo drug. The drug treatment was double-blind; patients were blind to the exercise programme. Graded exercise was delivered by a physiotherapist. 136 patients meeting Oxford research criteria for CFS were recruited from a University department of medicine outpatient clinic.

Results: 90 (66%) patients completed the trial. Patients were more likely to drop out of exercise than non-exercise treatment ($p = 0.05$). There was a non-specific treatment effect on most measures. When patients complied fully with exercise, there was a significant effect of exercise on fatigue ($p = 0.04$), functional work capacity ($p < 0.001$) and health perception ($p = 0.03$). In patients who complied, fluoxetine had a significant effect on fatigue ($p = 0.02$) and depression (p

$= 0.03$). On intention to treat, 12 (18%) patients had fully recovered from their fatigue symptoms in the exercise groups at six months compared to four (6%) who did not receive exercise ($p = 0.004$). Significant improvement with exercise only occurred in patients with DSM-III-R depressive disorders who were prescribed fluoxetine.

Conclusions: CFS patients benefit from graded exercise and fluoxetine if they can tolerate these treatments. Depressed CFS patients do not benefit from graded exercise unless they also receive an antidepressant. Graded exercise and fluoxetine have clinically distinct effects in CFS patients.

PREVALENCE OF DEPRESSION DURING HOSPITALIZATION FOR BONE MARROW TRANSPLANTATION: EFFECTS OF DIAGNOSTIC CRITERIA

J.M. Prieto, J. Blanch, J. Atala, C. Gastó, E. Cirera. *Department of Consultation-Liaison Psychiatry, University of Barcelona, Villarroel 170, 08036, Barcelona, Spain*

Objective: This study used different approaches to case identification to examine prevalence rates for major (MD) and minor depression (md) in a group of patients with hematological malignancies.

Method: A consecutive series of 103 patients hospitalized for bone marrow transplantation at the Hospital Clinic in Barcelona were evaluated on admission and weekly until discharge. **Results:** Diagnosis according to DSM-IV (exclude somatic symptoms related to a physical condition), DSM-III (include all somatic symptoms), and Endicott's revised criteria (replace somatic symptoms with non-somatic ones) are shown in the following table:

Diagnosis	DSM-IV		DSM-III		Endicott Criteria	
	N	%	N	%	N	%
MD	3	3	26	25	10	10
md	31	30	16	15	25	24

Taking in account the existing literature, a low rate of major depression was obtained when using DSM-IV criteria. The higher rate obtained with DSM-III can be related to the prevalence of somatic symptoms. After receiving intensive treatment and during the following 4 weeks, 95–81% of patients reported an energy loss $\geq 30\%$, with 98–78% reporting a moderate decrease in appetite.

Conclusions: When somatic symptoms are prevalent, an approach that relies more in psychological features can increase the diagnostic accuracy for mood disorders.

PSYCHOSOCIAL ASSESSMENT OF TRANSPLANT PATIENTS. THE GREEK EXPERIENCE

M. Syngelakis, A. Vidalis, G. Kaprinis. *Psychiatric Department, Hippocrates General Peripheral Hospital, Konstantinoupoleos 49, TK 546 42 Thessaloniki, Greece*

Research on psychiatric aspects of transplantation has been undertaken as transplantation is linked with increased psychiatric morbidity and also due to evidence that psychosocial factors influence the clinical outcome and the quality of life. The biopsychosocial model seems to be a necessary approach to these patients.

Since October 1994 we have evaluated and supported transplant patients referred from the two major Transplant Units in Northern Greece.

Our sample consist of all heart- ($n = 25$), lung- ($n = 5$), liver- ($n = 22$) candidates and recipients as well as a number of renal transplant patients ($n = 60$).

Results are presented using an 10-item rating scale for psychosocial screening of transplant patients (Psychosocial Assessment of Candidates for Transplantation, Olbrisch 1989). Mental health