

borough Rehabilitation Unit, Oxford. This is comparable with the figure of 7% reported by Kaminer for his out-patients in Israel and by Pakes for his in the United States.

G. P. PULLEN  
N. R. BEST  
J. MAGUIRE

*Littlemore Hospital,  
Littlemore, Oxford OX4 4XN*

#### References

- JAKUBCZYK, A., SKOWRONSKA, J., SPASOWICZ, E. & SZELENBERGER, W. (1972) Delirious syndromes following ingestion of trihexyphenidyl as psychodysleptic. *Psychiatria Polska*, **6**, 249–53.
- KAMINER, Y., MUNITZ, H. & WIJSENBEK, H. (1982) Trihexyphenidyl (Artane) abuse: euphoriant and anxiolytic. *British Journal of Psychiatry*, **140**, 473–4.
- PAKES, G. E. (1978) Abuse of trihexyphenidyl (letter). *Journal of American Medical Association*, **240**, 2434.
- PULLEN, G. P., BEST, N. R. & MAGUIRE, J. (1984) Anticholinergic drug abuse: a common problem? *British Medical Journal*, **289**, 612–613.

#### ROCK AND ROLL DELUSIONS

DEAR SIR,

Further to the account by Glass and Campbell of two female patients with delusions involving rock performers (*Journal*, July 1984, **145**, 95–96). I report four patients who have had delusions and/or hallucinations concerning pop stars.

*Case 1:* A 22 year old single man with schizophrenia believed falsely that he was famous as a celebrity and that he had met Davie Bowie and John Lennon. He was upset by Lennon's death in December 1980 but continued to hear voices of famous people including Lennon talking to him. He died by committing suicide in 1983.

*Case 2:* This 25 year old single man whose mother suffered from schizophrenia presented with the belief that Hank Marvin owed him a large sum of money. This idea took seed some years earlier when he had been at a Hank Marvin concert and the star had given him a special look. He refused treatment after experiencing side effects from a depot neuroleptic. When last seen he intended to go to the next local Hank Marvin concert and he expected that Hank would make a sign to invite him for a drink.

*Case 3:* This 30 year old single man had a history of delinquency, multiple drug abuse and overdose in his teens. Schizophrenia was diagnosed when he was 21 and he has required continuous in-patient treatment since the age of 22. He has experienced voices coming from the television which say that he is David Cassidy's double. At times he has been convinced that he is David Cassidy, Elvis Presley or David Essex. He believed that John Lennon was trying to get him out of hospital and blamed hospital staff for killing Lennon and Presley.

*Case 4:* A 30 year old married girl has suffered from schizophrenia for five years. There is a past history of alcohol and multiple drug abuse. In 1980 she burnt her wrist with a

cigarette after hearing Linda McCartney's voice telling her to do it. She also heard Paul McCartney's voice telling her to wait for him. Recently she relapsed nine months after stopping treatment and she has heard several voices including those of Boy George and Sting commenting on her actions.

All these patients have had severe illnesses characterised by many other delusions and hallucinations in addition to their experiences concerning pop stars.

I thank Dr. John Le Gassicke for permission to report case 3.

A. J. ROBINSON

*St. George's Hospital,  
Morpeth, Northumberland NE61 2NU*

DEAR SIR,

The review of *Sexual Strands: Understanding and Treating Sexual Anomalies in Men*, by F. E. Kenyon, is misleading (*Journal*, May 1984, **144**, 563). The book discusses theories and their supporting empirical evidence on the causes and treatments for a wide range of sexually anomalous behaviours.

Dr. Kenyon's review suggests that he has not read the book carefully for its intended purpose or its content. I object to his statement that the book is "within the limitations of a committed behaviourist's standpoint", when most major psychological theories and treatments, from castration and Provera to psychoanalysis, have been discussed and evaluated.

He further states that my "ultra critical approach leaves little scope for treatment so that what is eventually advocated appears rather impersonal and mechanistic." This statement epitomises the inaccuracy of the review. My attempt has been to humanise our clinical practice. As I said in *Sexual Strands*: "One factor has aroused my anger and perplexed me. It is the selection of aversion therapy as a first and most frequent choice in treating sexual anomalies . . . aversion methods have no greater effect in changing behaviour than the less noxious positive behaviour therapies or psychotherapies. Can we continue to permit patients to suffer unnecessarily? . . . *A range of behavioural and psychotherapy methods is available and should be tried out.* Any growth experience or change can be painful, but if the patient feels he is the master of his destiny, it is a mutual concern of therapist and client to change and the patient has dignity and control." (p. 498).

R. LANGEVIN

*Clarke Institute of Psychiatry,  
250 College Street, Toronto M5T 1R8*

#### CORRECTION

The heading to the letter from Drs Addonizio and Susman (*Journal*, November 1984, **145**, 556–557) should read NEUROLEPTIC MALIGNANT SYNDROME AND HEAT STROKE.