

# MODULE OF ANNULUS

TOHRU AKAZA and TADASHI KURODA

1. Let  $C$  and  $C'$  be two simple closed curves in the complex  $z$ -plane which have no point in common and surround the origin. Denote by  $D$  the annulus bounded by  $C$  and  $C'$ . Consider a family  $\{\gamma\}$  of rectifiable curves  $\gamma$  in  $D$  and the family  $P$  of all non-negative lower semi-continuous functions  $\rho = \rho(z)$  in  $D$ . Put

$$L_\rho\{\gamma\} = \inf_{\gamma \in \{\gamma\}} \int_\gamma \rho |dz|.$$

Understanding  $\frac{0}{0} = \frac{\infty}{\infty} = 0$ , we call the quantity

$$\lambda\{\gamma\} = \sup_{\rho \in P} \frac{(L_\rho\{\gamma\})^2}{\iint_D \rho^2 d\sigma}$$

the extremal length of the family  $\{\gamma\}$ , where  $d\sigma$  denotes the area element. Let  $\{\gamma'\}$  be the family of all rectifiable curves  $\gamma'$  in  $D$  joining  $C$  with  $C'$  and let  $\{\gamma''\}$  be that of all rectifiable curves  $\gamma''$  in  $D$  separating  $C$  from  $C'$ . Then it is known that

$$(1) \quad \lambda\{\gamma'\} = \frac{1}{\lambda\{\gamma''\}}$$

and that the quantity

$$(2) \quad \mu = 2\pi\lambda\{\gamma'\}$$

is the module of  $D$ . In this note, we give some estimates of  $\mu$ .

2. Let  $D$  be an annulus stated in §1. We denote by  $l_\theta$  the intersection of the half straight line  $\arg z = \theta$  ( $0 \leq \theta \leq 2\pi$ ) with  $D$  and by  $l(\theta)$  the logarithmic length of  $l_\theta$ , that is,

$$l(\theta) = \int_{l_\theta} \frac{dr}{r}, \quad z = re^{i\theta}.$$

---

Received October 17, 1960.

The following was proved by Rengel [3].

**THEOREM.** *The module  $\mu$  of  $D$  satisfies the inequality*

$$(3) \quad \mu \leq \frac{1}{2\pi} \int_0^{2\pi} l(\theta) d\theta.$$

Now we shall prove the following which implies Rengel's theorem stated above.

**THEOREM 1.** *For the module  $\mu$  of  $D$ , the inequality*

$$(4) \quad \mu \leq \frac{2\pi}{\int_0^{2\pi} \frac{d\theta}{l(\theta)}}$$

holds.

*Proof.* Let  $\{\gamma'\}$  be the family of all rectifiable curves  $\gamma'$  joining  $C$  with  $C'$  in  $D$ . Then it is obvious that

$$L_\rho\{\gamma'\} \leq \int_{I_\theta} \rho dr$$

for any  $\rho \in P$  and for any  $\theta$  ( $0 \leq \theta \leq 2\pi$ ). By the Schwarz inequality, we have

$$\begin{aligned} (L_\rho\{\gamma'\})^2 &\leq \left( \int_{I_\theta} \rho dr \right)^2 \\ &\leq \int_{I_\theta} \frac{dr}{r} \int_{I_\theta} \rho^2 r dr = l(\theta) \int_{I_\theta} \rho^2 r dr, \end{aligned}$$

or

$$(L_\rho\{\gamma'\})^2 \frac{1}{l(\theta)} \leq \int_{I_\theta} \rho^2 r dr.$$

Integrating both sides with respect to  $\theta$ , we get

$$\frac{(L_\rho\{\gamma'\})^2}{\iint_D \rho^2 d\sigma} \leq \frac{1}{\int_0^{2\pi} \frac{d\theta}{l(\theta)}},$$

which gives

$$\lambda\{\gamma'\} \leq \frac{1}{\int_0^{2\pi} \frac{d\theta}{l(\theta)}}.$$

From (2), we obtain our theorem.

*Remark.* The Schwarz inequality yields

$$(5) \quad (2\pi)^2 = \left( \int_0^{2\pi} d\theta \right)^2 \leq \int_0^{2\pi} \frac{d\theta}{l(\theta)} \int_0^{2\pi} l(\theta) d\theta,$$

from which Rengel's theorem is obtained immediately by using Theorem 1. In (5), the equality holds if and only if  $l(\theta)$  is a constant. In this case, the curve  $C'$  is obtained as a set of points  $\alpha z (z \in C)$ , where  $\alpha$  is a positive constant, and Rengel's inequality (3) and ours (4) are identical.

3. Here we give an estimate, from below, of the module of an annulus of a special type.

Let  $C$  be a simple closed curve in the  $z$ -plane surrounding the origin and let  $(C)$  be a domain bounded by  $C$  and containing the origin. If, for any point  $z \in C$  and for any  $t (0 \leq t < 1)$ , the point  $tz$  lies in  $(C)$ , then we say that  $C$  is *strictly star-like* with respect to the origin.

Consider a curve  $C$  strictly star-like with respect to the origin. We assume that  $C$  consists of a finite number of arcs  $C^k : r = r_k(\theta), \theta_{k-1} \leq \theta \leq \theta_k (k = 1, 2, \dots, n)$ , where  $\theta_0 = 0, \theta_n = 2\pi$  and each  $r_k(\theta)$  has a continuous derivative  $r'_k(\theta)$  in  $\theta_{k-1} \leq \theta \leq \theta_k$ . The expression  $r = r(\theta) (0 \leq \theta \leq 2\pi)$  defined by putting  $r(\theta) = r_k(\theta)$  for  $\theta_{k-1} \leq \theta \leq \theta_k$  is a representation of  $C$  in the polar form. In such a case, we say that the curve  $C$  is *piecewise smooth*.

Denote by  $C'$  the curve defined by  $r = \alpha r(\theta)$ , where  $\alpha$  is a real constant such that  $0 < \alpha < 1$ .

We can prove

**THEOREM 2.** *Let  $C$  and  $C'$  be defined as above. Then the module  $\mu$  of the annulus  $D$  bounded by  $C$  and  $C'$  is estimated from below as follows:*

$$\mu \geq \frac{1}{K(C)} \log \frac{1}{\alpha},$$

where  $K(C) = \frac{1}{2\pi} \int_0^{2\pi} \left\{ 1 + \left( \frac{r'(\theta)}{r(\theta)} \right)^2 \right\} d\theta$  is a constant depending only on the curve  $C$ .

*Proof.* Let  $C_t (0 \leq t \leq 1)$  be a curve having a representation in the polar form

$$C_t : r = R(\theta, t) = r(\theta) \{ \alpha + (1 - \alpha)t \}.$$

The curves  $C_1$  and  $C_0$  are identical with  $C$  and  $C'$  respectively. Consider the family  $\{r''\}$  of all rectifiable curves  $r''$  separating  $C$  from  $C'$  in  $D$  and the family  $P$  in §1. It is easy to see that

$$\begin{aligned} ds &= \sqrt{(R(\theta, t))^2 + \left(\frac{\partial R(\theta, t)}{\partial \theta}\right)^2} \\ &= r(\theta) \{\alpha + (1 - \alpha)t\} \sqrt{1 + \left(\frac{r'(\theta)}{r(\theta)}\right)^2} d\theta \end{aligned}$$

is the line-element along  $C_t$  ( $0 < t < 1$ ) and that

$$d\sigma = (r(\theta))^2 (1 - \alpha) \{\alpha + (1 - \alpha)t\} d\theta dt$$

is the area-element. It is evident that

$$L_p\{r''\} \leq \int_{C_t} \rho ds$$

for  $0 < t < 1$ . Hence, by the Schwarz inequality, we have

$$\begin{aligned} (L_p\{r''\})^2 &\leq \left(\int_{C_t} \rho ds\right)^2 \\ &\leq \frac{\alpha + (1 - \alpha)t}{1 - \alpha} \int_0^{2\pi} \left\{1 + \left(\frac{r'(\theta)}{r(\theta)}\right)^2\right\} d\theta \int_0^{2\pi} \rho^2 (r(\theta))^2 (1 - \alpha) \{\alpha + (1 - \alpha)t\} d\theta. \end{aligned}$$

Therefore, using the same argument as in the proof of Theorem 1 and noting (1) and (2), we get

$$\frac{(L_p\{r''\})^2}{\iint_D \rho^2 d\sigma} \leq K_0(C) \frac{1}{\log \frac{1}{\alpha}},$$

where  $K_0(C) = \int_0^{2\pi} \left\{1 + \left(\frac{r'(\theta)}{r(\theta)}\right)^2\right\} d\theta$ . Putting  $K(C) = \frac{1}{2\pi} K_0(C)$ , we have our theorem.

**EXAMPLE.** Let  $\Pi_n$  be a regular polygon of center at the origin and with  $n$  sides of equal length and let  $\Pi'_n$  be another regular polygon obtained from  $\Pi_n$  by a transformation  $z = \alpha z'$  ( $z' \in \Pi_n$ ), where  $0 < \alpha < 1$ . If we denote by  $\mu$  the module of the annulus bounded by  $\Pi_n$  and  $\Pi'_n$ , then, using Theorems 1 and 2, we get

$$\frac{1}{K(\Pi_n)} \log \frac{1}{\alpha} \leq \mu \leq \log \frac{1}{\alpha},$$

where

$$K(\Pi_n) = \frac{n}{\pi} \tan \frac{\pi}{n}.$$

4. Applying Theorem 2, we prove the following

**THEOREM 3.** *Let  $\Delta$  be a domain in the  $z$ -plane whose boundary consists of the origin and of an enumerable number of sets  $E_k$  ( $k=1, 2, \dots$ ), where  $E_k$  lies on a simple closed curve  $C^k$  strictly star-like with respect to the origin and may consist of arcs and points. If there exists a simple closed curve  $C$  which is piecewise smooth in the sense stated in §3 and surrounds the origin and if, for each  $k$ , there exists a positive number  $\alpha_k$  such that the set of points  $\alpha_k z$  ( $z \in C$ ) is contained in  $C_k$  and such that  $\alpha_k > \alpha_{k+1}$ ,  $\lim_{k \rightarrow \infty} \alpha_k = 0$ , then the origin is a weak boundary component of  $\Delta$ .*

*Proof.* Let us denote by  $D_k$  the annulus bounded by  $C^k$  and  $C^{k+1}$ . Then  $D_k$  is contained in  $\Delta$ . Denoting by  $\mu_k$  the module of  $D_k$ , we see by Theorem 2 that there exists a constant  $K(C)$  depending only on  $C$  such that

$$\mu_k \geq \frac{1}{K(C)} \log \frac{\alpha_k}{\alpha_{k+1}}.$$

Hence we get

$$\sum_{k=1}^{\infty} \mu_k \geq \frac{1}{K(C)} \sum_{k=1}^{\infty} \log \frac{\alpha_k}{\alpha_{k+1}},$$

whose right hand side diverges. By Grötzsch's theorem [2] (Cf. Savage [4]), we have our assertion.

*Remark.* This theorem implies Theorem 1 in [1].

#### REFERENCES

- [1] T. Akaza: On the weakness of some boundary component, Nagoya Math. Journ., **17** (1960), 219–223.
- [2] H. Grötzsch: Eine Bemerkung zum Koebeschen Kreisnormierungsprinzip, Berichte Verh. Sächs. Akad. Wiss. Leipzig Math. Nat., Kl. **87** (1935), 319–324.
- [3] E. Rengel: Über einige Schlichttheoreme der konformer Abbildung, Schrif. d. Math. Sem. u. Inst. f. angew. Math., Univ. Berlin, **1** (1932–33), 141–162.
- [4] N. Savage: Weak boundary components of an open Riemann surface, Duke Math. Journ., **24** (1957), 79–95.

*Kanazawa University*

*and*

*Nagoya University*