

# Food insecurity, coping strategies and glucose control in low-income patients with diabetes

Victoria L Mayer<sup>1,2,\*</sup>, Kevin McDonough<sup>3</sup>, Hilary Seligman<sup>4</sup>, Nandita Mitra<sup>5,6</sup> and Judith A Long<sup>6,7,8</sup>

<sup>1</sup>Department of Population Health Science and Policy, Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Place, Box 1077, New York, NY 10029, USA: <sup>2</sup>Division of General Internal Medicine, Department of Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, USA: <sup>3</sup>School of Arts and Sciences, University of Pennsylvania, Philadelphia, PA, USA: <sup>4</sup>Division of General Internal Medicine, University of California San Francisco, San Francisco, CA, USA: <sup>5</sup>Department of Biostatistics and Epidemiology, University of Pennsylvania, Philadelphia, PA, USA: <sup>6</sup>Leonard Davis Institute of Health Economics, University of Pennsylvania, Philadelphia, PA, USA: <sup>7</sup>Division of General Internal Medicine, Department of Medicine, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA, USA: <sup>8</sup>Center for Health Equity Research and Promotion, Philadelphia Veterans Affairs Medical Center, Philadelphia, PA, USA

Submitted 27 February 2015: Final revision received 8 June 2015: Accepted 1 July 2015: First published online 2 September 2015

## Abstract

**Objective:** To examine the relationship between food insecurity and coping strategies (actions taken to manage economic stress) hypothesized to worsen glucose control in patients with diabetes.

**Design:** Using a cross-sectional telephone survey and clinical data, we compared food-insecure and food-secure individuals in their use of coping strategies. Using logistic regression models, we then examined the association between poor glucose control (glycated Hb, HbA1c  $\geq 8.0\%$ ), food insecurity and coping strategies.

**Setting:** An urban medical centre, between June and December 2013.

**Subjects:** Four hundred and seven adults likely to be low income (receiving Medicaid or uninsured and/or residing in a zip code with  $>30\%$  of the population below the federal poverty level) with type 2 diabetes.

**Results:** Of respondents, 40.5% were food insecure. A significantly higher percentage of the food-insecure group reported use of most examined coping strategies, including foregone medical care, participation in the Supplemental Nutrition Assistance Program (SNAP) and use of emergency food programmes. Food insecurity was associated with poor glucose control (OR = 2.23; 95% CI 1.22, 4.10); coping strategies that were more common among the food insecure were not associated with poor glucose control. Among the food insecure, receipt of SNAP was associated with lower risk of poor glucose control (OR = 0.27; 95% CI 0.09, 0.80).

**Conclusions:** While food insecurity was associated with poor glucose control, most examined coping strategies did not explain this relationship. However, receipt of SNAP among food-insecure individuals was associated with better diabetes control, suggesting that such programmes may play a role in improving health.

## Keywords

Diabetes  
Nutrition  
Vulnerable populations  
Socio-economic factors  
Disease management

Food insecurity is an important issue in the USA: 14.3% of households experienced food insecurity in 2013<sup>(1)</sup>. Food insecurity exists ‘whenever the availability of nutritionally adequate and safe foods or the ability to acquire acceptable food in socially acceptable ways is limited or uncertain’<sup>(2)</sup>. Food insecurity has been associated with poor overall health, poor mental health, obesity and chronic diseases<sup>(3–5)</sup>. Among those with diabetes, studies

have shown an association between food insecurity and higher risk of poor glucose control<sup>(6–11)</sup>.

Food-insecure households may employ coping strategies to manage economic stress that in turn may cause poor control of type 2 diabetes. Coping strategies as defined by the WHO refer to ‘remedial actions undertaken by people whose survival and livelihood are compromised or threatened’<sup>(12)</sup>. These include consumption of low-cost,

\*Corresponding author: Email victoria.mayer@mountsinai.org

energy-dense sugars, fats and grains<sup>(13,14)</sup>; decreased consumption of high-cost foods such as fruits and vegetables<sup>(15,16)</sup>; overconsumption in times of adequacy alternating with meal reduction and skipping in times of inadequacy<sup>(17)</sup>; use of food assistance and emergency food services (which may be associated with poorer-quality diets)<sup>(18–22)</sup>; and prioritizing food purchasing over other competing demands, including medications and medical care. Food insecurity in patients with diabetes has previously been associated with foregoing medications<sup>(8,23)</sup>. However, few studies have examined the role of coping strategies other than foregoing medications specifically among individuals with diabetes and their association with glucose control.

Greater understanding of the relationship between food insecurity and type 2 diabetes can potentially allow physicians and policy makers to direct interventions towards modifiable determinants of disease control. In order to further explore the mechanisms by which food insecurity impacts diabetes control, we performed a cross-sectional study of likely low-income patients with diabetes to examine the use of coping strategies and to determine if those strategies are associated with food insecurity and poor glycaemic control. Because of the important role of diet in glucose control, we focused primarily on food-related coping strategies. We hypothesized that greater use of coping strategies would be associated with worse glycaemic control.

## Methods

We used electronic health records to identify potential participants who were between 30 and 80 years old, had at least one diagnosis code for type 2 diabetes in the past year and had a laboratory result for glycated Hb (HbA1c) in the previous 7 d. Potential participants were seen at the University of Pennsylvania Health System in Philadelphia, PA, USA, an urban health system that includes primary care and specialty providers throughout the city. We selected patients likely to be low income by including only those insured by Medicaid or uninsured, and/or residing in a zip code where over 30% of the population is below the federal poverty level<sup>(24)</sup>. In order to ensure that our sample included only patients with type 2 diabetes, we excluded from our analyses all patients who were diagnosed before age 20 years, assuming those individuals were likely to have type 1 diabetes.

We mailed potential participants a letter about the study and within one month called and invited them to participate. We made up to six call attempts. Those who agreed to participate were read a verbal consent form. We excluded participants who were non-English speaking, as our instrument was not translated into other languages. After survey completion, we mailed participants a \$US 10 gift card in appreciation along with area resource

information from the Greater Philadelphia Coalition Against Hunger. We collected data between June and December 2013. The study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the University of Pennsylvania Institutional Review Board. Verbal consent was obtained from all participants and formally recorded.

### **Glycaemic control**

Our primary outcome was glycaemic control, measured by HbA1c. This is the American Diabetes Association's recommended metric by which to guide medical treatment and is strongly associated with important clinical outcomes<sup>(25)</sup>. We defined poor glycaemic control as HbA1c  $\geq 8.0\%$ , as this goal meets the American Diabetes Association's guidelines for the majority of patients<sup>(26)</sup>.

### **Survey instrument**

The survey contained seventy items (see online Supplementary Appendix A). We measured food insecurity using the US Department of Agriculture's eighteen-item Adult Food Security Survey – Core Module (FSS), a validated and commonly used tool in the USA<sup>(27)</sup>. Each respondent is classified as 'food secure' (0–2 affirmative responses) or 'food insecure' (>2 affirmative responses). The FSS asks about food budget, food supply and food quality. The FSS contains skip patterns and only individuals with children under 18 years of age in the household are asked eight of the questions. Only those who respond affirmatively to initial questions about food insecurity are asked questions about more severe manifestations of food insecurity.

Other variables included coping strategies proposed in the literature as probable mechanisms for the relationship between food insecurity and glucose control, including: cost-related medication non-adherence; foregone medical care; use of emergency food programmes; receipt of food assistance in the form of the Supplemental Nutrition Assistance Program (SNAP; formerly known as Food Stamps); fruit, vegetable and added sugar intake; and food management practices.

We used a five-question measure of cost-related medication non-adherence developed by Pierre-Jacques *et al.* and adapted by Ngo-Metzger *et al.*<sup>(28,29)</sup>. We asked one question from the National Health Interview Survey to assess delayed or foregone medical care<sup>(30)</sup>.

To assess use of emergency food programmes and food assistance programmes we employed questions from the Food Security Supplement of the Current Population Survey, conducted by the US Census Bureau and the Bureau of Labor Statistics<sup>(31)</sup>.

To assess diet we focused on fruit, vegetable and added sugar intake, rather than total dietary intake, for several reasons. Greater fruit and vegetable intake has been associated with lower HbA1c and is an important component of diet recommendations for patients with

diabetes<sup>(32,33)</sup>. Added sugars, on the other hand, contribute to overconsumption of energy, without providing nutritional benefit<sup>(34)</sup>. However, fruits and vegetables cost more than many other foods, including those with added sugars<sup>(35)</sup>. Given the limited survey time, we chose the Dietary Screener in the California Health Interview Survey from 2009, a ten-item instrument that measures intake over the previous month of fruits, vegetables and added sugars<sup>(36)</sup>. We used scoring algorithms developed by the National Center for Health Statistics to convert responses to estimates of daily intake and calculate variance-adjusted aggregate estimates. We calculated two variance-adjusted aggregate estimates for fruits and vegetables excluding beans (one including fried potatoes and the other excluding fried potatoes), and used each in separate regression models. Validation results from the National Cancer Institute indicate that this screener underestimates fruit and vegetable intake by 0 to 2/3 cup equivalents/d, while misestimates (over and under) of added sugar intake range from 0.7 to 1.6 teaspoons/d. However, the National Cancer Institute concludes that because misestimates are small, screener data can still be used to compare intake between different groups<sup>(36)</sup>.

After review of the literature, we did not identify any standard survey items addressing food management practices; that is, behavioural modifications of eating and shopping patterns that individuals may utilize in the face of hardship. Given the possible effects on health of such behaviours, we developed ten questions based on findings from qualitative studies<sup>(37–41)</sup>. The questions ask how often participants overeat in times of adequacy and to avoid hunger, purchase fresh foods in times when funds are adequate, and purchase processed foods in times of shortage. The questions also ask how often participants follow a food budget and plan meals. Finally, the questions ask how often participants eat with relatives, friends or neighbours in times of need, or eat food that is not fresh when necessary. We employed five categorical answer options from 'never true' to 'always true' (see online Supplementary Appendix B). We piloted these questions with 100 respondents from primary-care waiting rooms. Each question demonstrated adequate spread (no variable had more than 80% of answers in one category). Internal consistency was indicated by a Cronbach's  $\alpha$  of 0.73. Validity was indicated by correlation between more frequent use of the food management practices examined and food insecurity. All ten questions were included in the main survey instrument. Using the results from the main survey, we performed an exploratory factor analysis (using principal factor analysis with varimax rotation) and maximum likelihood methods as a sensitivity analysis.

We included questions on age, race, ethnicity, marital status, number of children in the household, education, employment and income. We also asked patients their age at diagnosis of diabetes and whether they take insulin. Finally, we abstracted height and weight from the medical record.

## Analyses

We performed comparisons of sociodemographic and clinical characteristics between those with well-controlled and poorly controlled diabetes, using  $\chi^2$  tests and *t* tests. We also evaluated the association of each coping strategy with glucose control and with food insecurity. We then examined the role of coping strategies as mediators of the relationship between food insecurity and glucose control. Next, we developed logistic regression models to examine the association of food insecurity with glucose control, after adjusting for sociodemographic factors, clinical covariates and coping strategies. We then examined potential interactions between food insecurity and coping strategies (by including a multiplicative term in the logistic regression model), hypothesizing that coping strategies may modify the relationship between food insecurity and glucose control. Finally, we performed a stratified analysis to further examine the association between coping strategies and glucose control within the food-insecure and food-secure groups. All analyses were performed using the statistical software package Stata version 12.

## Results

The team attempted to contact 1247 patients. Thirty-four per cent completed the survey ( $n$  427), 33% refused ( $n$  408) and 33% were not successfully contacted ( $n$  412). We excluded patients likely to have type 1 diabetes (diagnosed with diabetes before age 20 years;  $n$  7) and those missing data on key variables (including food security, any coping strategy or insulin status;  $n$  13). Our final analyses included 407 participants.

Participants were largely non-Hispanic Black (82.8%), female (73.7%) and low income (Table 1). The mean BMI was in the obese range (38.7 kg/m<sup>2</sup>). The majority of patients (59.2%) had HbA1c  $\geq$  8%. Those we were unable to contact were similar with regard to race/ethnicity (84% Black), BMI (mean 36.6 kg/m<sup>2</sup>) and glucose control (57.8% had HbA1c  $\geq$  8%). A smaller proportion of the group we were unable to contact were female (63.3%).

Of participants, 40.5% were food insecure. In comparison to food-secure participants, those who were food insecure were younger (53.5 *v.* 59.0 years,  $P < 0.001$ ), more likely to be below 100% of the federal poverty level (67.9% *v.* 52.5%,  $P < 0.001$ ), more likely to be disabled (70.3% *v.* 49.6%,  $P < 0.001$ ), more likely to be on Medicaid (50.9% *v.* 30.3%,  $P = 0.001$ ), more likely to be on insulin (50.9% *v.* 40.5%,  $P = 0.04$ ) and had a higher BMI (40.3 *v.* 37.7 kg/m<sup>2</sup>,  $P = 0.004$ ; Table 1). Those in the food-insecure group were more likely to have poorly controlled glucose (68.5% *v.* 52.9%,  $P = 0.002$ ). The mean and interquartile range (IQR) for HbA1c in the food-insecure group was 8.5% (IQR 7.3–9.6%) *v.* 8.2% (IQR 6.9–9.4%) in the food-secure group.

Food management practices items showed adequate spread (no variable had more than 80% of answers in one

**Table 1** Sample characteristics, overall and by food security status, among low-income patients with diabetes from an urban US medical centre, June–December 2013

Characteristic	Total sample (n 407)		Food insecure (n 165)		Food secure (n 242)		P value*
	Mean or n	SD or %	Mean or n	SD or %	Mean or n	SD or %	
Age (years), mean and SD	56.7	11.4	53.5	8.9	59.0	12.3	<0.001
Sex, n and %							
Female	300	73.7	128	77.6	172	71.1	0.144
Race/ethnicity, n and %							0.1
Non-Hispanic Black	337	82.8	130	78.8	207	85.5	
Non-Hispanic White	33	8.1	14	8.5	19	7.9	
All others	37	9.1	21	12.7	16	6.6	
Income, percentage of FPL, n and %							<0.001
<100 %	239	58.7	112	67.9	127	52.5	
100–200 %	89	21.9	38	23.0	51	21.1	
>200 %	58	14.3	6	3.6	52	21.5	
Missing	21	5.2	9	5.5	12	5.0	
Education level, n and %							0.06
Less than high school	73	17.9	27	16.4	46	19.0	
High school	137	33.7	48	29.1	89	36.8	
Some college or technical degree	153	37.6	75	45.5	78	32.2	
College degree or higher	44	10.8	15	9.1	29	12.0	
Employment status, n and %							<0.001
Employed	83	20.4	26	15.8	57	23.6	
Unemployed, in school, retired	88	21.6	23	13.9	65	26.9	
Disabled	236	58.0	116	70.3	120	49.6	
Number of occupants in household, mean and SD	2.3	1.7	2.5	1.6	2.2	1.8	0.09
Insurance type, n and %							0.001
Medicare	115	28.3	38	23.0	77	31.8	
Medicaid	157	38.6	84	50.9	73	30.2	
Medicare and Medicaid	57	14.0	20	12.1	37	15.3	
Private/VA/military insurance	76	18.7	22	13.3	54	22.3	
No insurance	2	0.5	1	0.6	1	0.4	
Use insulin, n and %	182	44.7	84	50.9	98	40.5	0.04
BMI (kg/m <sup>2</sup> ), mean and SD	38.7	9.2	40.3	9.9	37.7	8.5	0.004
Years with diabetes, mean and SD	11.1	9.7	10.3	9.7	11.6	9.7	0.19
HbA1c $\geq$ 8.0, n and %	241	59.2	113	68.5	128	52.9	0.002

FPL, federal poverty level; VA, Veterans Affairs; HbA1c, glycated Hb.

\*For comparison between food-insecure and food-secure groups.

category; see online Supplementary Appendix B). Principal factor analysis showed a one-factor solution with an eigenvalue  $>1$  (1.79), indicating one underlying construct accounting for 0.87 of overall variance. Only seven of the ten items demonstrated adequate factor loading ( $\geq 0.4$ ; unrotated and rotated). These seven items had a Cronbach's  $\alpha$  of 0.68. The other three items (meal planning, budgeting and buying food in bulk) were thus excluded from all analyses and we combined the remaining items into a Food Management Practices Scale, with a range from 7 to 35, with higher scores indicating more frequent use of these strategies. Confirmatory factor analysis supported our findings: all scale items had adequate loading onto one factor ( $\geq 0.4$ ).

Many of the coping strategies were employed significantly more frequently among those who were food insecure (Table 2), including cost-related medication non-adherence (60.6% of food insecure *v.* 37.5% of food secure,  $P < 0.001$ ), foregone medical care (35.2% *v.* 9.1%,  $P < 0.001$ ), use of emergency food programmes (53.3% *v.* 24.0%,  $P < 0.001$ ), receipt of SNAP (76.4% *v.* 59.1%,  $P < 0.001$ ) and frequency of food management practices (mean score 18.8 *v.* 13.1,  $P < 0.001$ ). Mean daily intakes of

fruits, vegetables and added sugar were not significantly different between the two groups.

Table 3 shows unadjusted comparisons between those with well-controlled glucose and with poorly controlled glucose, and odds ratios from the fully adjusted logistic regression for each coping strategy, along with the sociodemographic and clinical factors included in the model. In unadjusted analysis, those with poorly controlled glucose were more likely to be food insecure (46.9% of poorly controlled *v.* 31.3% of controlled group,  $P = 0.002$ ), younger in age (55.2 *v.* 58.9 years,  $P = 0.001$ ), disabled (62.7% *v.* 51.2%,  $P = 0.04$ ), insulin users (61.8% *v.* 19.9%,  $P < 0.001$ ), and to eat fewer fruits (0.5 *v.* 0.8 cup equivalents/d,  $P = 0.005$ ) and vegetables (0.4 *v.* 0.5 cup equivalents/d,  $P = 0.01$ ). Aside from fruit and vegetable intake, none of the coping strategies differed significantly by glucose control. As none of the coping strategies were associated with both food insecurity and glucose control, they did not meet the definition of mediator and we did not proceed through further steps in a mediation analysis<sup>(42)</sup>.

In the adjusted model, those who were food insecure were more likely to have poorly controlled glucose

**Table 2** Coping strategies and food insecurity among low-income patients with diabetes from an urban US medical centre, June–December 2013

Characteristic	Total sample ( <i>n</i> 407)		Food insecure ( <i>n</i> 165)		Food secure ( <i>n</i> 242)		<i>P</i> value*
	<i>n</i> or mean	%, IQR or SD	<i>n</i> or mean	%, IQR or SD	<i>n</i> or mean	%, IQR or SD	
Cost-related medication non-adherence, <i>n</i> and %	160	39.3	100	60.6	60	37.5	<0.001
Foregone medical care, <i>n</i> and %	80	19.7	58	35.2	22	9.1	<0.001
Used emergency food programmes, <i>n</i> and %	146	35.9	88	53.3	58	24.0	<0.001
Received SNAP, <i>n</i> and %	269	66.1	126	76.4	143	59.1	<0.001
Diet							
Fruit (cup equivalents/d), mean and IQR	0.6	0.1–0.8	0.6	0.1–0.7	0.7	0.1–0.9	0.709
Vegetables (cup equivalents/d), mean and IQR	0.4	0.2–0.5	0.4	0.2–0.5	0.4	0.2–0.5	0.486
Added sugar (teaspoons/d), mean and IQR	4.4	0.9–5.5	5.1	1.2–6.1	3.9	0.8–4.7	0.054
Score on Food Management Practices Scale (range 7–35), mean and SD	15.4	5.4	18.8	5.0	13.1	4.4	<0.001

IQR, interquartile range; SNAP, Supplemental Nutrition Assistance Program.

\*For comparison between food-insecure and food-secure groups.

**Table 3** Associations between food insecurity, coping strategies and glucose control among low-income patients with diabetes from an urban US medical centre, June–December 2013

Characteristic	Controlled glucose ( <i>n</i> 166)		Poorly controlled glucose ( <i>n</i> 241)		<i>P</i> value*	Adjusted OR†	95 % CI	<i>P</i> value‡
	<i>n</i> or mean	%, SD or IQR	<i>n</i> or mean	%, SD or IQR				
Food insecure, <i>n</i> and %	52	31.3	113	46.9	0.002	2.23	1.22, 4.10	0.01
Age (years), mean and SD	58.9	13.1	55.2	9.8	0.001	0.98	0.97, 1.01	0.16
Income, percentage of FPL, <i>n</i> and %					0.16			
<100 %	91	54.8	148	61.4		Ref.		
100–200 %	39	23.5	50	20.8		0.89	0.49, 1.62	0.70
>200 %	30	18.1	28	11.6		0.86	0.39, 1.91	0.72
Missing	6	3.6	15	6.2		1.30	0.41, 4.15	0.66
Employment status, <i>n</i> and %					0.04			
Employed, <i>n</i> and %	36	21.7	47	19.5		Ref.		
Unemployed, in school, retired	45	27.1	43	17.8		0.84	0.39, 1.82	0.66
Disabled	85	51.2	151	62.7		1.29	0.66, 2.54	0.46
Use insulin, <i>n</i> and %	33	19.9	149	61.8	<0.001	6.44	3.95, 10.50	<0.001
BMI (kg/m <sup>2</sup> ), mean and SD	38.5	9.7	38.9	8.8	0.68	0.99	0.97, 1.02	0.63
Cost-related medication non-adherence, <i>n</i> and %	66	39.8	94	39.0	0.88	0.66	0.39, 1.12	0.12
Foregone medical care, <i>n</i> and %	34	20.5	46	19.1	0.73	0.89	0.48, 1.67	0.72
Used emergency food programmes, <i>n</i> and %	59	35.5	87	36.1	0.91	0.76	0.45, 1.28	0.30
Received SNAP, <i>n</i> and %	105	63.3	164	68.1	0.32	0.77	0.43, 1.38	0.38
Diet								
Fruit (cup equivalents/d), mean and IQR	0.8	0.1–0.9	0.5	0.1–0.7	0.005	0.51§	0.32, 0.82	0.005
Vegetables (cup equivalents/d), mean and IQR	0.5	0.2–0.5	0.4	0.1–0.5	0.01			
Added sugar (teaspoons/d), mean and IQR	4.2	0.9–5.3	4.5	1.0–5.6	0.56	1.29§	0.77, 2.16	0.33
Score on combined Food Management Practices Scale (range 7–35), mean and SD	15.0	5.4	15.7	5.4	0.21	0.97	0.93, 1.03	0.32

IQR, interquartile range; FPL, federal poverty level; SNAP, Supplemental Nutrition Assistance Program; Ref., referent group;

\*For unadjusted comparison of well-controlled and poorly controlled groups.

†Odds ratio for poor glucose control, from fully adjusted logistic regression analysis, all covariates shown.

‡From fully adjusted logistic regression analysis.

§Using variance-adjusted aggregate estimates; fruit and vegetable intake combined (excluding beans and fried potatoes).

(OR = 2.23; 95 % CI 1.22, 4.10, *P* = 0.01). Use of insulin was also significantly associated with poor glucose control (OR = 6.44; 95 % CI 3.95, 10.50, *P* < 0.001), while a greater intake of fruits and vegetables was associated with lower risk of poor glucose control (OR = 0.51; 95 % CI 0.32, 0.82, *P* = 0.005). Models using a variance-adjusted aggregate estimate for fruits and vegetables excluding beans and excluding (shown in Table 3) or including (not shown) fried potatoes yielded similar results.

We found a statistically significant interaction between SNAP and food insecurity, indicating the association between food insecurity and glucose control is modified by receipt of SNAP (Table 4). Compared with food-insecure participants not receiving SNAP, food-insecure participants receiving SNAP had lower risk of poor glucose control (OR = 0.35; 95 % CI 0.13, 0.91, *P* = 0.03). Compared with those who were food secure and receiving SNAP, food-insecure participants receiving SNAP had a greater



risk of poor glucose control (OR = 1.68; 95 % CI 0.87, 3.25,  $P = 0.12$ ); however, this difference was not statistically significant at  $P < 0.05$ . In the adjusted stratified model (Table 5), among those who were food insecure, those who received SNAP were at lower risk of poor glucose control than those not receiving SNAP (OR = 0.27; 95 % CI 0.09, 0.80,  $P = 0.02$ ).

## Discussion

Among this population of largely low-income patients with diabetes, food insecurity was associated with greater risk of poor glucose control, after adjusting for sociodemographic and clinical characteristics. The use of coping strategies was more common among food-insecure participants, apart from intakes of fruits, vegetables and added sugars, which were similar between the two groups. However, intake of fruits and vegetables was the only coping strategy associated with glucose control in the full sample.

SNAP receipt was associated with lower risk of poor glucose control among those who were food insecure. It is possible that individuals who are food insecure but receiving SNAP have more funds to spend on food and therefore may have: (i) less stress; (ii) more ability to purchase healthier foods shown in previous studies to be more expensive (although not assessed in our study)<sup>(35)</sup>; or (iii) more ability to spend funds on non-food items with

the potential to improve diabetes control. Alternatively, our results may be attributable to non-random entry into SNAP. SNAP receipt may be a marker for characteristics associated with better disease control, such as skills in navigating complex public systems. This association was noted only among food-insecure participants. This suggests that SNAP may play a different role in individuals who remain food insecure even while receiving SNAP, likely a more vulnerable population than those who are food secure after receiving SNAP. Many individuals who receive SNAP remain food insecure (54 % in 2013, according to the US Department of Agriculture)<sup>(1)</sup>.

The present study is the first one we are aware of to look at the role of food assistance receipt in glucose control among elderly and non-elderly adults. Literature examining food assistance receipt and health has focused on obesity and child health outcomes<sup>(43,44)</sup>. SNAP recipients can spend their benefits on most food items (excluding hot food and alcohol). Research evaluating what SNAP recipients purchase *v.* eligible non-participating counterparts (none of which has focused on individuals with diabetes) has yielded mixed results, from worse diet quality to no difference<sup>(21,22,45)</sup>. In the only study we are aware of focusing on food assistance receipt and diabetes, Nicholas examined glucose control in older Americans and found no difference between the risk of poorly controlled glucose in those receiving Food Stamps and likely eligible non-recipients<sup>(46)</sup>. One explanation for our contrasting finding is our inclusion of

**Table 4** Interaction between food insecurity and SNAP receipt on the risk of poor glucose control among low-income patients with diabetes from an urban US medical centre, June–December 2013

Comparison	OR for poor glucose control	95 % CI	<i>P</i> value
Food insecure and no SNAP receipt <i>v.</i> food secure and no SNAP receipt	5.94	1.98, 17.84	0.001
Food secure and SNAP receipt <i>v.</i> food secure and no SNAP receipt	1.23	0.61, 2.51	0.56
Food insecure and SNAP receipt <i>v.</i> food insecure and no SNAP receipt	0.35	0.13, 0.91	0.03
Food insecure and SNAP receipt <i>v.</i> food secure and SNAP receipt	1.68	0.87, 3.25	0.12

SNAP, Supplemental Nutrition Assistance Program.

The interaction term food insecurity  $\times$  SNAP receipt in our multivariable logistic model had OR = 0.28 (95 % CI 0.09, 0.89);  $P = 0.03$ .

Results are adjusted for age, income, employment status, use of insulin, BMI and other coping strategies.

**Table 5** Stratified analysis: coping strategies and poor glucose control by food security status among low-income patients with diabetes from an urban US medical centre, June–December 2013

Characteristic	Adjusted OR among food insecure*	<i>P</i> value	Adjusted OR among food secure*	<i>P</i> value
Cost-related medication non-adherence	0.76	0.52	0.52	0.08
Foregone medical care in last year	0.48	0.10	2.0	0.20
Used emergency food programmes	0.69	0.36	0.78	0.51
Received SNAP	0.27	0.02	1.26	0.56
Diet				
Variance-adjusted mean daily cup equivalents of fruits and vegetables†	0.38	0.03	0.57	0.07
Variance-adjusted mean daily teaspoons of added sugar	0.83	0.67	1.66	0.16
Mean score on Food Management Practices Scale	0.97	0.48	0.96	0.25

SNAP, Supplemental Nutrition Assistance Program.

\*Odds ratio for poor glucose control from stratified, adjusted logistic regression analysis, including the following covariates: age, income, employment status, use of insulin and BMI.

†Excluding beans and fried potatoes.

non-elderly adults, who face different financial pressures from the elderly. Our study suggests that further research should explore how low-income patients with diabetes who are receiving SNAP differ from those who are not receiving SNAP.

The coping strategies we examined did not account for the difference in glycaemic control observed between food-secure and food-insecure patients (i.e. there are remaining differences even taking into account coping strategies). One interpretation of these findings is that these coping strategies may be adaptive, rather than harmful – that these strategies lead to better control for patients. Second, other unmeasured factors may account for the relationship between food insecurity and glucose control, such as emotional distress related to food insecurity, which in a previous study partially mediated the relationship between food insecurity and control<sup>(9)</sup>. Third, our cross-sectional study does not let us examine the longitudinal effects of the measured coping strategies. For example, a patient classified in our study as food secure and skipping medications because of cost may have been food insecure before deciding to use medication funds for food. If as a consequence of skipping medications their glucose control became worse, it would appear as though this coping mechanism was detrimental to those who were food secure not insecure.

While we found that greater intake of fruits and vegetables was associated with better glucose control, we found no difference between the food-secure and food-insecure groups. One interpretation of this finding is that fruit and vegetable intake was similar between the two groups and other unmeasured factors associated with food insecurity account for differences in glucose control. It is also possible that we did not detect differences due to our measurement tool, a screener limited to assessment of fruit, vegetable and added sugar intake, rather than a measure of full dietary intake. Screeners are subject to systematic error. They can be used to compare different populations as to higher or lower intake, or to examine associations between intake of certain dietary components and other variables, but they cannot be reliably used to characterize individual intake<sup>(47)</sup>. Our findings contrast with a recent study of Puerto Rican adults with diabetes in Boston that found lower overall diet quality and lower fruit and vegetable intake among food-insecure participants<sup>(48)</sup>. Our different findings may relate to differences in dietary patterns between the study populations or to differences in measurement. Further research can more fully characterize the diets of low-income patients with diabetes in additional populations and further examine the relative roles of dietary patterns and other factors important to glucose control.

The present study has several additional limitations. Given our cross-sectional study design, we cannot draw conclusions regarding longitudinal phenomena, including the causal role of food insecurity in glucose control.

We examined patients attending a single medical centre in one US city and thus our findings may not be nationally generalizable. In addition, the patients in our study had all recently received medical care and thus may represent a population with relatively greater access to medical care. These patients may experience different social contexts and utilize different coping strategies from those without regular access, also limiting the generalizability of our findings. Another limitation of the study is our response rate (34 % of the sample, 51 % of those contacted). Those who refused to participate and those with whom we were unable to make contact may differ from those who completed the survey. This also limits our generalizability, as the sample represents a limited subset of all low-income patients with diabetes who are seen in this health system.

Our findings have several important implications. Multiple studies have shown a relationship between food insecurity and glucose control; physicians and health systems should consider screening and addressing food insecurity in patients with diabetes. The coping strategies we studied, apart from SNAP receipt, did not act as modifiers of the relationship between food insecurity and diabetes control, and further research is warranted to elucidate what factors do mediate the relationship, such as diabetes distress or self-efficacy, other known predictors of diabetes control which may be particularly salient in low-income populations<sup>(49–51)</sup>. The association of SNAP receipt with lower risk of poor glucose control in food-insecure individuals suggests that programmes aimed at ameliorating food insecurity are avenues for further research and intervention. Understanding how food assistance affects diabetes self-care, including food purchasing and diet, could help elucidate this relationship. Physicians and health systems can pilot programmes to increase food assistance coverage among patients with diabetes. Finally, our findings indicate that recent cuts to SNAP benefits<sup>(52,53)</sup> may have unintended consequences, such as worse chronic disease control among low-income patients with diabetes.

## Acknowledgements

*Acknowledgements:* The authors would like to acknowledge Esther Kim and Elizabeth Wall for their important work in data collection for this study. They thank Steven Honeywell and Marie Synnstedt for their work developing and executing electronic tools for patient selection. Also, they thank Steve Schachterle for additional statistical consultation. *Financial support:* V.L.M. was supported with funding from the Division of General Internal Medicine Matt Slap Award from the Perelman School of Medicine at the University of Pennsylvania and the National Institutes of Health Institutional Training Grant 5-T32-HP-100296-20-00. She is currently supported by the Empire Clinical Research Investigator Program (Principal

Investigator Dr Carol Horowitz). The funders had no role in the design, analysis, or writing of this article. *Conflict of interest:* None. *Authorship:* V.L.M conceived of and designed the study, researched data and wrote the manuscript. K.M. researched data and reviewed/edited the manuscript. H.S. contributed to study design and reviewed/edited the manuscript. N.M. contributed to researching data and reviewed/edited the manuscript. J.A.L. contributed to study design, researching data and reviewed/edited the manuscript. *Ethics of human subject participation:* The study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the University of Pennsylvania Institutional Review Board. Verbal consent was obtained from all participants and formally recorded.

### Supplementary material

To view supplementary material for this article, please visit <http://dx.doi.org/10.1017/S1368980015002323>

### References

- Coleman-Jensen A, Gregory C & Singh A (2014) Household Food Security in the United States in 2013. [http://www.ers.usda.gov/media/1565410/err173\\_summary.pdf](http://www.ers.usda.gov/media/1565410/err173_summary.pdf) (accessed May 2015).
- Life Sciences Research Office, Federation of American Societies for Experimental Biology (1990) Core indicators of nutritional state for difficult-to-sample populations. *J Nutr* **120**, Suppl. 11, 1559–1600.
- Lee JS & Frongillo EA Jr (2001) Nutritional and health consequences are associated with food insecurity among US elderly persons. *J Nutr* **131**, 1503–1509.
- Seligman HK, Laraia BA & Kushel MB (2010) Food insecurity is associated with chronic disease among low-income NHANES participants. *J Nutr* **140**, 304–310.
- Haering SA & Syed SB (2009) *Community Food Security in the United States: A Survey of the Relevant Scientific Literature*. Baltimore, MD: Johns Hopkins Bloomberg School of Public Health, Center for a Livable Future.
- Gucciardi E, Vogt JA, DeMelo M *et al.* (2009) Exploration of the relationship between household food insecurity and diabetes in Canada. *Diabetes Care* **32**, 2218–2224.
- Seligman HK, Bindman AB, Vittinghoff E *et al.* (2007) Food insecurity is associated with diabetes mellitus: results from the National Health Examination and Nutrition Examination Survey (NHANES) 1999–2002. *J Gen Intern Med* **22**, 1018–1023.
- Seligman HK, Davis TC, Schillinger D *et al.* (2010) Food insecurity is associated with hypoglycemia and poor diabetes self-management in a low-income sample with diabetes. *J Health Care Poor Underserved* **21**, 1227–1233.
- Seligman HK, Jacobs EA, Lopez A *et al.* (2012) Food insecurity and glycemic control among low-income patients with type 2 diabetes. *Diabetes Care* **35**, 233–238.
- Marjerrison S, Cummings EA, Glanville NT *et al.* (2011) Prevalence and associations of food insecurity in children with diabetes mellitus. *J Pediatr* **158**, 607–611.
- Berkowitz SA, Meigs JB, DeWalt D *et al.* (2014) Material need insecurities, control of diabetes mellitus, and use of health care resources: results of the measuring economic insecurity in diabetes study. *JAMA Intern Med* **175**, 257–265.
- World Health Organization, Department of Emergency and Humanitarian Action (1999) Emergency Health Training Programme for Africa. 1. Overview. 1.9. Coping Mechanisms. <http://apps.who.int/disasters/repo/5517.pdf> (accessed October 2014).
- Drewnowski A & Specter SE (2004) Poverty and obesity: the role of energy density and energy costs. *Am J Clin Nutr* **79**, 6–16.
- Drewnowski A & Darmon N (2005) Food choices and diet costs: an economic analysis. *J Nutr* **135**, 900–904.
- Kendall A, Olson CM & Frongillo EA Jr (1996) Relationship of hunger and food insecurity to food availability and consumption. *J Am Diet Assoc* **96**, 1019–1024.
- Rehm CD, Monsivais P & Drewnowski A (2011) The quality and monetary value of diets consumed by adults in the United States. *Am J Clin Nutr* **94**, 1333–1339.
- Tarasuk V, McIntyre L & Li J (2007) Low-income women's dietary intakes are sensitive to the depletion of household resources in one month. *J Nutr* **137**, 1980–1987.
- Dinour LM, Bergen D & Yeh MC (2007) The food insecurity–obesity paradox: a review of the literature and the role food stamps may play. *J Am Diet Assoc* **107**, 1952–1961.
- Larson NI & Story MT (2011) Food insecurity and weight status among US children and families: a review of the literature. *Am J Prev Med* **40**, 166–173.
- Franklin B, Jones A, Love D *et al.* (2012) Exploring mediators of food insecurity and obesity: a review of recent literature. *J Community Health* **37**, 253–264.
- Leung CW, Ding EL, Catalano PJ *et al.* (2012) Dietary intake and dietary quality of low-income adults in the Supplemental Nutrition Assistance Program. *Am J Clin Nutr* **96**, 977–988.
- Nguyen BT, Shuval K, Njike VY *et al.* (2014) The Supplemental Nutrition Assistance Program and dietary quality among US adults: findings from a nationally representative survey. *Mayo Clin Proc* **89**, 1211–1219.
- Billimek J & Sorkin DH (2012) Food insecurity, processes of care, and self-reported medication underuse in patients with type 2 diabetes: results from the California Health Interview Survey. *Health Serv Res* **47**, 2159–2168.
- Philadelphia Research Initiative (2013) Philadelphia 2012: The State of the City. [http://www.pewtrusts.org/our\\_work\\_report\\_detail\\_wide.aspx?id=85899461859](http://www.pewtrusts.org/our_work_report_detail_wide.aspx?id=85899461859) (accessed September 2013).
- Inzucchi SE, Bergenstal RM, Buse JB *et al.* (2012) Management of hyperglycemia in type 2 diabetes: a patient-centered approach: position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* **35**, 1364–1379.
- American Diabetes Association (2014) Standards of medical care in diabetes – 2014. *Diabetes Care* **37**, Suppl. 1, S14–S80.
- Bickel G, Nord M, Price C *et al.* (2000) Measuring Food Security in the United States. Guide to Measuring Household Food Security. Revised 2000. <http://www.fns.usda.gov/fsec/files/fsguide.pdf> (accessed October 2012).
- Pierre-Jacques M, Safran DG, Zhang F *et al.* (2008) Reliability of new measures of cost-related medication nonadherence. *Med Care* **46**, 444–448.
- Ngo-Metzger Q, Sorkin DH, Billimek J *et al.* (2012) The effects of financial pressures on adherence and glucose control among racial/ethnically diverse patients with diabetes. *J Gen Intern Med* **27**, 432–437.
- Centers for Disease Control and Prevention (2010) QuickStats: Delayed or forgone medical care because of cost concerns among adults aged 18–64 years, by disability and health insurance coverage status – National Health Interview Survey, United States, 2009. *MMWR Morb Mortal Wkly Rep* **59**, issue 44, 1456.



31. US Census Bureau (2011) December 2011: Food Security Supplement. <http://www.census.gov/cps/methodology/techdocs.html> (accessed September 2012).
32. Sargeant LA, Khaw KT, Bingham S *et al.* (2001) Fruit and vegetable intake and population glycosylated haemoglobin levels: the EPIC-Norfolk Study. *Eur J Clin Nutr* **55**, 342–348.
33. Evert AB, Boucher JL, Cypress M *et al.* (2014) Nutrition therapy recommendations for the management of adults with diabetes. *Diabetes Care* **37**, Suppl. 1, S120–S143.
34. Johnson RK, Appel LJ, Brands M *et al.* (2009) Dietary sugars intake and cardiovascular health a scientific statement from the American Heart Association. *Circulation* **120**, 1011–1020.
35. Aggarwal A, Monsivais P & Drewnowski A (2012) Nutrient intakes linked to better health outcomes are associated with higher diet costs in the US. *PLoS One* **7**, e37533.
36. National Cancer Institute (2012) The Dietary Screener in the 2009 California Health Interview Survey. <http://appliedresearch.cancer.gov/surveys/chis/dietscreener/2009/> (accessed September 2012).
37. Kempson KM, Keenan DP, Sadani PS *et al.* (2002) Food management practices used by people with limited resources to maintain food sufficiency as reported by nutrition educators. *J Am Diet Assoc* **102**, 1795–1799.
38. Hoisington A, Shultz JA & Butkus S (2002) Coping strategies and nutrition education needs among food pantry users. *J Nutr Educ Behav* **34**, 326–333.
39. Kempson K, Keenan DP, Sadani PS *et al.* (2003) Maintaining food sufficiency: coping strategies identified by limited-resource individuals versus nutrition educators. *J Nutr Educ Behav* **35**, 179–188.
40. Wicks R, Trevena LJ & Quine S (2006) Experiences of food insecurity among urban soup kitchen consumers: insights for improving nutrition and well-being. *J Am Diet Assoc* **106**, 921–924.
41. Smith C & Richards R (2008) Dietary intake, overweight status, and perceptions of food insecurity among homeless Minnesotan youth. *Am J Hum Biol* **20**, 550–563.
42. Baron RM & Kenny DA (1986) The moderator–mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* **51**, 1173–1182.
43. Currie J (2003) US food and nutrition programs. In *Means-Tested Transfer Programs in the United States*, pp. 199–289 [R Moffitt, editor]. Chicago, IL: University of Chicago Press.
44. Ver Ploeg M & Ralston K (2008) Food Stamps and Obesity: What Do We Know? <http://www.ers.usda.gov/publications/eib-economic-information-bulletin/eib34.aspx> (accessed September 2012).
45. Todd JE & Ploeg MV (2014) Caloric beverage intake among adult supplemental nutrition assistance program participants. *Am J Public Health* **104**, E80–E85.
46. Nicholas LH (2011) Can Food Stamps help to reduce Medicare spending on diabetes? *Econ Hum Biol* **9**, 1–13.
47. National Cancer Institute (n.d.) Dietary Assessment Primer: Screeners. <http://dietaassessmentprimer.cancer.gov/profiles/screeners/index.html> (accessed December 2014).
48. Berkowitz SA, Gao X & Tucker KL (2014) Food-insecure dietary patterns are associated with poor longitudinal glycemic control in diabetes: results from the Boston Puerto Rican Health study. *Diabetes Care* **37**, 2587–2592.
49. Aikens JE (2012) Prospective associations between emotional distress and poor outcomes in type 2 diabetes. *Diabetes Care* **35**, 2472–2478.
50. Walker RJ, Gebregziabher M, Martin-Harris B *et al.* (2015) Quantifying direct effects of social determinants of health on glycemic control in adults with type 2 diabetes. *Diabetes Technol Ther* **17**, 80–87.
51. Gao J, Wang J, Zheng P *et al.* (2013) Effects of self-care, self-efficacy, social support on glycemic control in adults with type 2 diabetes. *BMC Fam Pract* **14**, 66.
52. Rampel C (2013) As cuts to Food Stamps take effect, more trims to benefits are expected. *The New York Times*, 31 October; Sect. Politics, p. A20. [https://urldefense.proofpoint.com/v2/url?u=http-3A\\_\\_www.nytimes.com\\_2013\\_11\\_01\\_us\\_as-2Dcuts-2Dto-2Dfood-2Dstamps-2Dtake-2Deffect-2Dmore-2Dtrims-2Dto-2Dbenefits-2Dare-2Dexpected.html-3F-5Fr-3D0&d=AwIF-g&c=4R1YgkJNMyVWjMjneTwN5tJRn8m8VqT SNCjYlg1wNX4&r=ZlJG1qEgJ19wXUK0734EnnjEUeJmDassmLGqKnp5khc&m=mxI0h1YqfwgTdcC3fSK5rJEWmswXVSVlWt7e1HRw\\_Us&s=z-bCl64SeEOR8lOZQVnh\\_ecyzPQXoeb\\_Qgk9EbTRoIU&e=](https://urldefense.proofpoint.com/v2/url?u=http-3A__www.nytimes.com_2013_11_01_us_as-2Dcuts-2Dto-2Dfood-2Dstamps-2Dtake-2Deffect-2Dmore-2Dtrims-2Dto-2Dbenefits-2Dare-2Dexpected.html-3F-5Fr-3D0&d=AwIF-g&c=4R1YgkJNMyVWjMjneTwN5tJRn8m8VqT SNCjYlg1wNX4&r=ZlJG1qEgJ19wXUK0734EnnjEUeJmDassmLGqKnp5khc&m=mxI0h1YqfwgTdcC3fSK5rJEWmswXVSVlWt7e1HRw_Us&s=z-bCl64SeEOR8lOZQVnh_ecyzPQXoeb_Qgk9EbTRoIU&e=) (accessed February 2014).
53. Shear MD (2014) In signing Farm Bill, Obama extols rural growth. *The New York Times*, 7 February; Sect. Politics, p. A12. [https://urldefense.proofpoint.com/v2/url?u=http-3A\\_\\_www.nytimes.com\\_2014\\_02\\_08\\_us\\_politics\\_farm-2Dbill.html&d=AwIF-g&c=4R1YgkJNMyVWjMjneTwN5tJRn8m8VqT SNCjYlg1wNX4&r=ZlJG1qEgJ19wXUK0734EnnjEUeJmDassmLGqKnp5khc&m=mxI0h1YqfwgTdcC3fSK5rJEWmswXVSVlWt7e1HRw\\_Us&s=7Fd-WW0wvNzY-WPiYsqNcY\\_xZTIJbXfcMlTaDT8UX7Q&e=](https://urldefense.proofpoint.com/v2/url?u=http-3A__www.nytimes.com_2014_02_08_us_politics_farm-2Dbill.html&d=AwIF-g&c=4R1YgkJNMyVWjMjneTwN5tJRn8m8VqT SNCjYlg1wNX4&r=ZlJG1qEgJ19wXUK0734EnnjEUeJmDassmLGqKnp5khc&m=mxI0h1YqfwgTdcC3fSK5rJEWmswXVSVlWt7e1HRw_Us&s=7Fd-WW0wvNzY-WPiYsqNcY_xZTIJbXfcMlTaDT8UX7Q&e=) (accessed February 2014).