

STUDY OF ANTARCTIC ICE SHEET MOTION BY MEANS OF DOPPLER EFFECT LASER EQUIPMENT

By V. V. BOGORODSKIY, I. M. BELOUSOVA and I. P. IVANOV

(Arkticheskiy i Antarkticheskiy Nauchno-Issledovatel'skiy Institut, Fontanka 34, Leningrad D104, U.S.S.R.)

ABSTRACT. Doppler-effect laser equipment was successfully employed in Antarctica in 1969–70 to measure the rate of ice-sheet motion. The method is based on measuring differential beat frequency generated by the interaction of the direct pulse and its echo. A He–Ne laser with the wavelength of 6328 Å was used in the 15th Soviet Antarctic Expedition. The paper gives detailed results of the measurements along with a description of the equipment used and of the observational procedures. A new modified version of the equipment was made ready for 1971–72 Antarctic season, and was used in the 18th Soviet Antarctic Expedition in the vicinity of Mirny station. New results are also discussed. Analysis of all available data makes it possible to believe that the movement of the ice sheet is irregular with occasional stops. The progressive motion of the ice sheet is accompanied by strain oscillations.

ELECTROMAGNETIC CHARACTERISTICS OF SEA ICE IN THE METRE AND DECIMETRE RANGES, AND ITS RADIO-ECHO SOUNDING

By V. V. BOGORODSKIY and V. P. TRIPOLNIKOV

(Arkticheskiy i Antarkticheskiy Nauchno-Issledovatel'skiy Institut, Fontanka 34, Leningrad D104, U.S.S.R.)

ABSTRACT. The paper gives measurements of the velocity of electromagnetic wave propagation in sea ice, the tangent loss, and the specific absorption obtained by means of coaxial-line experiments, assuming sea ice to be the dielectric. Measurements have been made at frequencies of 30, 60, 100, 200 and 400 MHz by pulses of 0.05 and 0.5 ms and video pulses of 0.02 ms at a repetition frequency of 10 kHz, with temperatures from -40°C to -10°C and salinities 17, 8.6, 4.4 and 0.56‰. Data obtained on the values of the tangent loss are in a good agreement with those obtained by Addison. The velocity has been found to be constant (170 m/ms) for radio and video pulses in the ice at any salinity value and temperatures below -22°C . At higher temperatures the velocity decreases being more pronounced with the salinity increase. Data on electrical characteristics of sea ice averaged throughout the ice thickness are presented. These measurements have been made by using long lines, frozen into the ice. Experimental radio-echo soundings of sea ice carried out at the drifting stations SP-19 and SP-21 in 1971–72 and 1972–73 by means of pulsed radar systems in the range 200–100 MHz are also discussed.