

Dynamic behaviour of supraglacial lakes on cold polar glaciers: Canada Glacier, McMurdo Dry Valleys, Antarctica

The cold polar glaciers of the McMurdo Dry Valleys (MCM DV), Antarctica, are characterized by low accumulation ($10 \text{ cm w.e. a}^{-1}$), high sublimation ($7\text{--}9 \text{ cm w.e. a}^{-1}$) and low melt rates ($1\text{--}3 \text{ cm w.e. a}^{-1}$) (Fountain and others, 1998). Meltwater production and flow is limited to the near surface of these cold polar glaciers, where the average ice temperature at 15 m depth is $\sim -17^\circ\text{C}$. Meltwater flow, as on other glacier surfaces, is confined to discrete hydrological basins, which consist of series of ice-topped pool and riffle sequences of varying dimensions (Fountain and others, 2004). The larger ice-topped pools, 10–50 m in diameter, are called ‘cryolakes’ (Tranter and others, 2010) and are the focus for supraglacial meltwater flow and sediment transport on Canada Glacier. They function similarly to hydrologically connected cryoconite holes (Fountain and others, 2004). We show below that water flow through this system is highly episodic during the melt season.

The supraglacial drainage system of Canada Glacier, one of the most extensively studied glaciers in the MCM DV, is ice-topped for much of the year, so that flow largely occurs below a veneer of up to 0.5 m of surface ice. The drainage system consists of a series of interconnected channels, riffles and cryoconite holes ($<1 \text{ m}$ diameter), the latter connected to the pool and riffle system by a series of linear passageways

($<10 \text{ cm}$ wide), probably the traces of former crevasses. About half the cryoconite holes are hydrologically connected, while the other half remain isolated (Fountain and others, 2008; Tranter and others, 2010). The drainage system is frozen solid throughout winter and reactivates by melting in November. Increasing solar radiation in spring melts the cryolakes internally due to a solid-state heating effect arising from the relatively low albedo of the lake ice (compared to the glacier ice) and the subsurface sediment concentrated in the lake bottom (Brandt and Warren, 1993; Podgorny and Grenfell, 1996; Fountain and others, 2008). Similar processes occur in the frozen channel beds connecting the cryolakes. The surface ice over the channels, in particular, may rot and melt out completely when air temperatures are $>0^\circ\text{C}$ for a few days. Melt enlargement of the channels also occurs under these same conditions. Drainage system closure due to refreezing usually begins in late January.

We had assumed that the cryolakes steadily melted out as the ablation season progressed (Fountain and others, 2004). Occasional observations of empty cryolake basins have been recorded, but it was presumed that the lake drainage was gradual (Johnston and others, 2005; Tranter and others, 2005). Observations in the 2008/09 ablation season show that, by contrast, episodic filling and drainage of these lakes is both possible and probably frequent. Figure 1a shows a typical cryolake on Canada Glacier on 7 December 2008. Figure 1b shows the cryolake 3 days later, after the lake rapidly flooded and the water level rose by $\sim 2 \text{ m}$ in $<24 \text{ hours}$. At no point during this period did the air temperature reach 0°C (Fig. 2). The mean air temperature prior to the flood (1–6 December) was -5.7°C , with a maximum of -1.2°C . During the flood (8–10 December), the mean air temperature was -2.9°C and the maximum was -0.5°C . We infer that the increase in total diurnal solar radiation during three consecutive cloud-free days (7–9 December) was sufficient to increase production of subsurface meltwater. We further surmise that a number of upstream pools also filled during this period. Their subsequent and progressively rapid drainage, due to channel opening or downcutting, led to the rapid flooding of the cryolake under observation. Meltwater drainage from the monitored cryolake was restricted because the outflow channel was still largely frozen. We observed that smaller cryolakes higher up the same drainage basin did indeed show evidence of having filled and drained back to their former water levels during the same period. This evidence included perched ice ceilings above the cryolakes with clear void space beneath and distinctive downcut and flow-enlarged outflow channels.

We can quantify the relative fluxes of water into and out of the monitored cryolake before, during and after the flood as follows. The estimated volume of the lake increased by $\sim 300\%$ during the flood, from ~ 180 to $\sim 630 \text{ m}^3$. The mean discharge during the peak summer ablation season for a number of different supraglacial streams on Canada Glacier was 2 L s^{-1} (personal communication from M. Hoffman, 2008). By contrast, the maximum discharge flowing into a similar nearby cryolake during the flood, recorded using salt tracing (Moore, 2004), was nearly a factor of 40 higher at 75 L s^{-1} . The monitored cryolake would flood in $\sim 2.3 \text{ hours}$ if this maximum discharge were maintained and outflow were negligible. The flooded lake would drain in 88 hours if the outlet stream flowed at the usual mean rate and no other water inputs occurred. In reality, the cryolake only drained




Fig. 1. A typical cryolake on the surface of Canada Glacier, Taylor Valley (77.61924° N , $162.96803^\circ \text{ E}$), (a) before (7 December) and (b) after (10 December) the flood event in 2008. Note the person on the lake ice in (a) for scale; the ice lid rose by 2 m in $<24 \text{ hours}$.




Fig. 2. Temperature and incoming solar radiation at Canada Glacier throughout the monitoring period, measured by a Campbell 107 probe and a Licor L1200X silicon pyranometer respectively. The red lines indicate the three cloud-free days which triggered the flood, and the associated increase in air temperature. The air temperature prior to the flood rose by 7°C, but did not reach 0°C.

by $\sim 260\text{ m}^3$ after 5 days, as inflows declined and outflows increased due to flow melting and enlargement of the outflow channel.

The episodic filling and emptying of cryolakes has implications for models of streamflow which are based on meteorological variables and assume a proportional response in meltwater discharge (Doran and others, 2008). Summer temperatures can be predicted with reasonable confidence using the dry adiabatic lapse rate and distance from the coast (Doran and others, 2002; Ebnét and others, 2005), and can be used to predict glacier meltwater delivery to proglacial stream, soil and lake ecosystems via simple relationships between air temperature and melt generation (Jaros, 2002; Ebnét and others, 2005). However, the temporary storage and release of meltwater during transit through the supraglacial drainage system adds a layer of complexity to this simple association. Storage of water in pools and cryolakes delays the delivery of meltwater, and hence the release can occur as a series of discrete large-volume pulses or floods to downstream ecosystems.

The role of cryolakes in the mosaic of MCMDV ecosystems in terms of water and nutrient dynamics is also likely to be more stochastic than was previously thought. The interaction of liquid water, sediment and biological material in each basin, coupled with water from the cryoconite hole network, supplies nutrient to glacially fed meltwater streams and into the ice-covered lakes (Foreman and others, 2004). Small-scale flood events, such as that described here, are more likely to send pulses of nutrient through the glacier, stream and lake ecosystems. A number of authors have proposed that the Dry Valley ecosystem operates under a 'pulse-reserve' regime (Moorhead, 2007; Wall, 2007), in which the ecosystem responds to a 'pulse' disturbance, such as an increase in the availability of liquid water, or an influx of labile nutrient from an adjacent part of the ecosystem, generating an accumulation of 'biological

currency' (e.g. biota or biomass). The results of this pulse persist as a 'reserve' during normal conditions. The observed behaviour of the cryolake fits into this model well, by periodically accentuating the magnitude of the water and nutrient delivered to downstream biological communities. The longer residence time of water in the cryolakes potentially maximizes the production of organic carbon and nutrient (Bagshaw and others, 2007) before the water is released. The eventual delivery of the 'pulse' to downstream biological communities may trigger a rapid response in productivity, such as that recorded in the streams (McKnight and others, 2007) and lakes (Foreman and others, 2004).

ACKNOWLEDGEMENTS

This work was supported by US National Science Foundation grant ANT-0423595 and UK Engineering and Physical Sciences Research Council (EPSRC) grant EP/D057620/1. Fieldwork was conducted with the MCMDV Long Term Ecological Research site team whose support is gratefully acknowledged, with logistics provided by Raytheon Polar Services and PHI Helicopters.

*Bristol Glaciology Centre,
School of Geographical Sciences,
University of Bristol,
University Road,
Bristol BS8 1SS, UK
E-mail: liz.bagshaw@bristol.ac.uk*

E.A. BAGSHAW
M. TRANTER
J.L. WADHAM

*Departments of Geography and Geology,
Portland State University,
PO Box 751, Portland,
Oregon 97207-0751, USA*

A.G. FOUNTAIN
H. BASAGIC

6 May 2010

REFERENCES

- Bagshaw, E.A., M. Tranter, A.G. Fountain, K.A. Welch, H. Basagic and W.B. Lyons. 2007. Biogeochemical evolution of cryoconite holes on Canada Glacier, Taylor Valley, Antarctica. *J. Geophys. Res.*, **112**(G4), G04S35. (10.1029/2007JG000442.)
- Brandt, R.E. and S.G. Warren. 1993. Solar-heating rates and temperature profiles in Antarctic snow and ice. *J. Glaciol.*, **39**(131), 99–110.
- Doran, P.T. and 6 others. 2002. Valley floor climate observations from the McMurdo dry valleys, Antarctica, 1986–2000. *J. Geophys. Res.*, **107**(D24), 4772. (10.1029/2001JD002045.)
- Doran, P.T. and 6 others. 2008. Hydrologic response to extreme warm and cold summers in the McMurdo Dry Valleys, East Antarctica. *Antarct. Sci.*, **20**(5), 499–509.
- Ebnet, A.F., A.G. Fountain, T.H. Nylén, D.M. McKnight and C.L. Jaros. 2005. A temperature-index model of stream flow at below-freezing temperatures in Taylor Valley, Antarctica. *Ann. Glaciol.*, **40**, 76–82.
- Foreman, C.M., C.F. Wolf and J.C. Prisco. 2004. Impact of episodic warming events on the physical, chemical and biological relationships of lakes in the McMurdo Dry Valleys, Antarctica. *Aquat. Chem.*, **10**(3), 239–268.
- Fountain, A.G., G.L. Dana, K.J. Lewis, B.H. Vaughn and D.M. McKnight. 1998. Glaciers of the McMurdo Dry Valleys, southern Victoria Land, Antarctica. In Prisco, J.C., ed. *Ecosystem dynamics in a polar desert: the McMurdo Dry Valleys, Antarctica*. Washington, DC, American Geophysical Union, 65–75. (Antarctic Research Series 72.)
- Fountain, A.G., M. Tranter, T.H. Nylén, K.J. Lewis and D.R. Mueller. 2004. Evolution of cryoconite holes and their contribution to meltwater runoff from glaciers in the McMurdo Dry Valleys, Antarctica. *J. Glaciol.*, **50**(168), 35–45.
- Fountain, A.G., T.H. Nylén, M. Tranter and E. Bagshaw. 2008. Temporal variations in physical and chemical features of cryoconite holes on Canada Glacier, McMurdo Dry Valleys, Antarctica. *J. Geophys. Res.*, **113**(G1), G01S92. (10.1029/2007JG000430.)
- Jaros, C. 2002. Climatic controls on interannual variation in streamflow in Fryxell Basin, Taylor Valley. (MSc thesis, University of Colorado.)
- Johnston, R.R., A.G. Fountain and T.H. Nylén. 2005. The origin of channels on lower Taylor Glacier, McMurdo Dry Valleys, Antarctica, and their implication for water runoff. *Ann. Glaciol.*, **40**, 1–7.
- McKnight, D.M. and 7 others. 2007. Reactivation of a cryptobiotic stream ecosystem in the McMurdo Dry Valleys, Antarctica: a long-term geomorphological experiment. *Geomorphology*, **89**(1–2), 186–204.
- Moore, D. 2004. Introduction to salt dilution gauging for stream-flow measurement, Part I. *Streamline, Watershed Manage. Bull.*, **7**(4), 20–23.
- Moorhead, D.L. 2007. Mesoscale dynamics of ephemeral wetlands in the Antarctic Dry Valleys: implications to production and distribution of organic matter. *Ecosystems*, **10**(1), 87–95.
- Podgorny, I.A. and T.C. Grenfell. 1996. Absorption of solar energy in a cryoconite hole. *Geophys. Res. Lett.*, **23**(18), 2465–2468.
- Tranter, M., A.G. Fountain, W.B. Lyons, T.H. Nylén and K.A. Welch. 2005. The chemical composition of runoff from Canada Glacier, Antarctica: implications for glacier hydrology during a cool summer. *Ann. Glaciol.*, **40**, 15–19.
- Tranter, M., E. Bagshaw, A.G. Fountain and C. Foreman. 2010. The biogeochemistry and hydrology of McMurdo Dry valley glaciers: is there life on Martian ice now? In Doran, P.T., W.B. Lyons and D.M. McKnight, eds. *Life in Antarctic deserts and other cold dry environments: astrobiological analogues*. Cambridge, etc., Cambridge University Press.
- Wall, D.H. 2007. Global change tipping points: above- and below-ground biotic interactions in a low diversity ecosystem. *Philos. Trans. R. Soc., Ser. B*, **362**(1488), 2291–2306.