

# ON CERTAIN DUAL INTEGRAL EQUATIONS

by E. T. COPSON

(Received 20 May, 1960)

1. In his book on Fourier Integrals, Titchmarsh [1] gave the solution of the dual integral equations

$$\int_0^\infty \xi^{2\alpha} \psi(\xi) J_\nu(\xi\rho) d\xi = f(\rho) \quad (0 < \rho < 1), \tag{1}$$

$$\int_0^\infty \psi(\xi) J_\nu(\xi) d\xi = 0 \quad (\rho > 1), \tag{2}$$

for the case  $\alpha > 0$ , by some difficult analysis involving the theory of Mellin transforms. Sneddon [2] has recently shown that, in the cases  $\nu = 0$ ,  $\alpha = \pm \frac{1}{2}$ , the problem can be reduced to an Abel integral equation by making the substitution

$$\psi(\xi) = \xi \int_0^1 \phi(t) \cos(\xi t) dt \quad (\alpha = -\frac{1}{2})$$

or

$$\psi(\xi) = \int_0^1 \chi(t) \sin(\xi t) dt, \quad \chi(0) = 0 \quad (\alpha = \frac{1}{2}).$$

It is the purpose of this note to show that the general case can be dealt with just as simply by putting

$$\psi(\xi) = \xi^{1-\alpha} \int_0^1 \phi(t) J_{\nu+\alpha}(\xi t) dt. \tag{3}$$

The analysis is formal: no attempt is made to supply details of rigour.

2. We need the following two lemmas.

LEMMA A. If  $\lambda > \mu > -1$ ,

$$\int_0^\infty J_\lambda(at) J_\mu(bt) t^{1+\mu-\lambda} dt = \begin{cases} 0 & (0 < a < b), \\ \frac{b^\mu(a^2 - b^2)^{\lambda-\mu-1}}{2^{\lambda-\mu-1} a^\lambda \Gamma(\lambda-\mu)} & (0 < b < a). \end{cases}$$

This result is well known [3].

LEMMA B. Let  $f(x), f'(x)$  be continuous in  $0 \leq x \leq a$ . Let  $0 < \kappa < 1$ . Then the solution of

$$\int_0^x g(t) (x^2 - t^2)^{-\kappa} dt = f(x) \quad (0 < x < a)$$

is

$$g(x) = \frac{2 \sin \pi \kappa}{\pi} \frac{d}{dx} \int_0^x f(t) (x-t)^{\kappa-1} dt.$$

This is a simple transformation of the solution of Abel's integral equation [4].

3. In order to determine the function  $\phi(t)$  in equation (3), we have to consider separately the cases when  $\alpha$  is positive and when  $\alpha$  is negative. The case when  $\alpha$  is zero obviously does not arise, since then equations (1) and (2) reduce to a single integral equation in which the right-hand side is zero when  $\rho > 1$ .

If  $\alpha > 0$ , inversion of the order of integration gives

$$\int_0^\infty \psi(\xi) J_\nu(\xi\rho) d\xi = \int_0^1 \phi(t) \int_0^\infty \xi^{1-\alpha} J_{\nu+\alpha}(\xi t) J_\nu(\xi\rho) d\xi dt = 0,$$

when  $\rho > 1$ , by Lemma A, provided that  $\nu > -1$ . Thus (2) is satisfied.

To deal with equation (1), we modify (3) by integration by parts. For we have

$$\begin{aligned} \psi(\xi) &= \xi^{1-\alpha} \int_0^1 \phi(t) J_{\nu+\alpha}(\xi t) dt \\ &= -\xi^{-\alpha} \int_0^1 \phi(t) t^{\nu+\alpha-1} \frac{d}{dt} \{t^{1-\nu-\alpha} J_{\nu+\alpha-1}(\xi t)\} dt \\ &= -\xi^{-\alpha} \left[ \phi(t) J_{\nu+\alpha-1}(\xi t) \right]_0^1 + \xi^{-\alpha} \int_0^1 \Phi(t) J_{\nu+\alpha-1}(\xi t) dt, \end{aligned}$$

where

$$\Phi(t) = t^{1-\nu-\alpha} \frac{d}{dt} \{ \phi(t) t^{\nu+\alpha-1} \}.$$

Hence if

$$\lim_{t \rightarrow +0} t^{\nu+\alpha-1} \phi(t) = 0, \tag{5}$$

we obtain

$$\psi(\xi) = \xi^{-\alpha} \int_0^1 \Phi(t) J_{\nu+\alpha-1}(\xi t) dt - \xi^{-\alpha} \phi(1) J_{\nu+\alpha-1}(\xi).$$

Substitution in (1) then gives

$$f(\rho) = \int_0^1 \Phi(t) \int_0^\infty \xi^\alpha J_\nu(\xi\rho) J_{\nu+\alpha-1}(\xi t) d\xi dt - \phi(1) \int_0^\infty \xi^\alpha J_\nu(\xi\rho) J_{\nu+\alpha-1}(t) dt,$$

where  $0 < \rho < 1$ . By Lemma A, this reduces to

$$f(\rho) = \frac{2^\alpha}{\Gamma(1-\alpha)} \rho^{-\nu} \int_0^\rho t^{\nu+\alpha-1} \Phi(t) (\rho^2 - t^2)^{-\alpha} dt,$$

provided that  $\nu+1 > \nu+\alpha > 0$ , i.e. provided that  $\nu > -\alpha$ ,  $0 < \alpha < 1$ . But this is an integral equation of the type given in Lemma B; its solution is

$$\Phi(\rho) \rho^{\nu+\alpha-1} = \frac{2^{1-\alpha}}{\Gamma(\alpha)} \frac{d}{d\rho} \int_0^\rho t^{\nu+1} f(t) (\rho^2 - t^2)^{\alpha-1} dt,$$

it being assumed that  $t^\nu f(t)$  and its first derivative are continuous in  $0 \leq t \leq 1$ . If we substitute for  $\Phi(\rho)$  from (4), integrate and use condition (5), we find that

$$\rho^{\nu+\alpha-1} \phi(\rho) = \frac{2^{1-\alpha}}{\Gamma(\alpha)} \int_0^\rho t^{\nu+1} f(t) (\rho^2 - t^2)^{\alpha-1} dt, \tag{6}$$

a result which may also be written as

$$\rho^{\nu+\alpha}\phi(\rho) = \frac{2^{-\alpha}}{\Gamma(1+\alpha)} \frac{d}{d\rho} \int_0^\rho t^{\nu+1} f(t) (\rho^2 - t^2)^\alpha dt. \tag{7}$$

The conditions on the parameters are  $0 < \alpha < 1, \nu > -\alpha$ .

When  $\alpha = -\beta$ , where  $\beta > 0$ , we start by integrating the integral in (3) by parts. This gives

$$\begin{aligned} \psi(\xi) &= \xi^{1+\beta} \int_0^1 \phi(t) J_{\nu-\beta}(\xi t) dt \\ &= \xi^\beta \int_0^1 \phi(t) t^{\beta-\nu-1} \frac{d}{dt} \{t^{\nu-\beta+1} J_{\nu-\beta+1}(\xi t)\} dt \\ &= \xi^\beta \left[ \phi(t) J_{\nu-\beta+1}(\xi t) \right]_0^1 - \xi^\beta \int_0^1 \Psi(t) J_{\nu-\beta+1}(\xi t) dt, \end{aligned}$$

where

$$\Psi(t) = t^{\nu-\beta+1} \frac{d}{dt} \{ \phi(t) t^{\beta-\nu-1} \}.$$

Hence

$$\psi(\xi) = \phi(1) \xi^\beta J_{\nu-\beta+1}(\xi) - \xi^\beta \int_0^1 \Psi(t) J_{\nu-\beta+1}(\xi t) dt$$

provided that

$$\lim_{t \rightarrow +0} t^{\nu-\beta+1} \phi(t) = 0.$$

It then follows from Lemma A that condition (2) is satisfied when  $0 < \beta < 1, \nu > -1$ .

If we now substitute from (3) in (1) and invert the order of integration, we obtain

$$\begin{aligned} f(\rho) &= \int_0^1 \phi(t) \int_0^\infty \xi^{1-\beta} J_\nu(\xi \rho) J_{\nu-\beta}(\xi t) dt d\xi \\ &= \frac{2^{1-\beta}}{\Gamma(\beta)} \rho^{-\nu} \int_0^\rho t^{\nu-\beta} \phi(t) (\rho^2 - t^2)^{\beta-1} dt, \end{aligned}$$

by Lemma A. Using Lemma B, we get

$$\rho^{\nu-\beta}\phi(\rho) = \frac{2^\beta}{\Gamma(1-\beta)} \frac{d}{d\rho} \int_0^\rho t^{\nu+1} f(t) (\rho^2 - t^2)^{-\beta} dt, \tag{8}$$

which is, in fact, equation (7) with  $\alpha$  replaced by  $-\beta$ : but the conditions now are  $-1 < \alpha < 0, \nu > -1$ . The limiting condition on  $\phi$  is evidently satisfied.

4. Having obtained formulae for  $\phi(\rho)$ , we can deduce the desired solution of (1) and (2). The simplest form of the solution is given by using (7), viz.

$$\psi(\xi) = \frac{(2\xi)^{1-\alpha}}{2\Gamma(1+\alpha)} \int_0^1 \rho^{-\nu-\alpha} J_{\nu+\alpha}(\xi \rho) \frac{d}{d\rho} \int_0^\rho t^{\nu+1} f(t) (\rho^2 - t^2)^\alpha dt d\rho, \tag{9}$$

valid when  $0 < \alpha < 1, \nu > -\alpha$  or when  $-1 < \alpha < 0, \nu > -1$ . From this, the results given by Sneddon for  $\nu = 0, \alpha = \pm \frac{1}{2}$  readily follow.

To get the solution valid for  $0 < \alpha < 1$ ,  $\nu > -\alpha$ , in the form given by Titchmarsh, we use (6), which gives

$$\psi(\xi) = \frac{(2\xi)^{1-\alpha}}{\Gamma(\alpha)} \int_0^1 J_{\nu+\alpha}(\xi\rho) \rho^{1-\nu-\alpha} \int_0^\rho t^{\nu+1} f(t) (\rho^2 - t^2)^{\alpha-1} dt d\rho. \quad (10)$$

Lastly, if in addition  $\nu + 2\alpha + 2 > 0$ , we may integrate (9) by parts to get

$$\psi(\xi) = \frac{(2\xi)^{1-\alpha}}{2\Gamma(1+\alpha)} \left\{ J_{\nu+\alpha}(\xi) \int_0^1 t^{\nu+1} f(t) (1-t^2)^\alpha dt + \xi \int_0^1 \rho^{-\nu-\alpha} J_{\nu+\alpha+1}(\xi\rho) \int_0^\rho t^{\nu+1} f(t) (\rho^2 - t^2)^\alpha dt d\rho \right\}. \quad (11)$$

#### REFERENCES

1. E. C. Titchmarsh, *Introduction to the theory of Fourier integrals* (Clarendon Press, Oxford, 1937), pp. 334–339.
2. I. N. Sneddon, The elementary solution of dual integral equations, *Proc. Glasgow Math. Assoc.*, **4** (1960), 108–110.
3. G. N. Watson, *A treatise on the theory of Bessel functions* (University Press, Cambridge, 1944), p. 401, equations (1) and (3).
4. E. T. Whittaker and G. N. Watson, *A course of modern analysis* (University Press, Cambridge, 1920), p. 229.

ST. SALVATOR'S COLLEGE  
UNIVERSITY OF ST. ANDREWS