


RESEARCH PAPER

Ground beetles (Coleoptera: Carabidae) of Akimiski Island, Nunavut, Canada

Kaitlyn J. Fleming¹ , James A. Schaefer² , Kenneth F. Abraham³ , Rodney W. Brook³ ,
Glen S. Brown³ , and David V. Beresford^{1,2} 

¹Trent School of the Environment, Trent University, 1600 W Bank Drive, Peterborough, Ontario, K9L 0G2, Canada,

²Department of Biology, Trent University, 1600 W Bank Drive, Peterborough, Ontario, K9L 0G2, Canada, and ³Wildlife Research and Monitoring Section, Ministry of Natural Resources, 2140 East Bank Drive, Peterborough, Ontario, K9L 0G2, Canada

Corresponding author: Kaitlyn J. Fleming; Email: kaitlynfleming@trentu.ca

(Received 22 May 2024; accepted 12 July 2024)

Abstract

In many regions of Canada, knowledge of the distribution of insect species is far from complete. This knowledge gap, known as the Wallacean Shortfall, is often manifest by species records separated by large, often remote areas with no records. Paradoxically, these difficult-to-access areas offer the best opportunity to study unaltered native community assemblages. Such gaps in knowledge are exemplified by ground beetles, a well-known group, yet with record gaps in many unstudied areas of Canada, including Akimiski Island, Nunavut. This postglacial rebound island, located in James Bay, has no permanently occupied human dwellings and almost no human-altered habitat. Using a combination of pitfall-malaise traps, pitfall traps, and hand captures during 2008–2014, we collected 1368 ground beetles (Coleoptera: Carabidae) as part of a larger biodiversity survey. We identified 31 species, 29 of which were first territorial records for Nunavut. Our results almost double the number of Carabidae known from Nunavut and extend the known range of eight other species. Seventeen of the species that we caught cannot fly, evidence for colonists arriving on Akimiski on floating debris. Our study fills substantial range gaps and serves as baseline information to detect future change.

Introduction

Habitat loss represents the principal driver of the decline of biodiversity in Canada (Venter *et al.* 2006) and elsewhere (Maxwell *et al.* 2016), but assessing its effects depends on good baseline information. Knowledge from relatively unaltered habitats is key; it provides insight into biogeographic processes and enables us to quantify the effects of habitat change. Yet, our understanding of species distributions in areas with unaltered habitats remains inadequate for many groups (Whittaker *et al.* 2005; Lomolino *et al.* 2017) and is often short-term (Mihoub *et al.* 2017). This Wallacean shortfall (the paucity of knowledge of species distributions) is particularly acute for insects, especially in geographically isolated areas (Lomolino *et al.* 2017; Vergara-Asenjo *et al.* 2023).

The situation is exemplified by ground beetles (Coleoptera: Carabidae, Latrielle), one of the largest families of insects, with more than 40 000 species worldwide (Lövei and Sunderland 1996) and more than 2000 in North America (Bell 1990). Despite being well known and extensively studied across Canada (*e.g.*, Lindroth 1969; Bousquet 2010, 2012; Bousquet *et al.* 2013;

Subject editor: Delano Lewis

© The Author(s), 2024. Published by Cambridge University Press on behalf of Entomological Society of Canada. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Ernst and Buddle 2015; and many others), there is almost no knowledge of carabids in many remote and difficult-to-access regions, even though these regions offer the best opportunities to study relatively unaltered native community assemblages.

One such remote area is Akimiski Island in the Qikiqtaaluk Region of Nunavut, Canada. The island, located in James Bay, the southernmost part of the Arctic Ocean, is a postglacial rebound island between 3500 and 4000 years old (Martini 1981), with no permanently occupied human dwellings, only seasonal residents, and almost no human-altered habitat. Akimiski Island has a history of biological research. The island's birds (Nguyen *et al.* 2003; Pollock *et al.* 2012; Richards and Gaston 2018; Brook *et al.* 2019, 2021; Gan *et al.* 2019), mammals (Kolenosky and Prevett 1983; Peacock *et al.* 2010; Obbard and Middel 2012), and plants (Blaney and Kotanen 2000; O *et al.* 2005) have been well documented. Although some insect species have been inventoried (*e.g.*, Dytiscidae; DeGasparro *et al.* 2018), little is known about most of the island's ground beetles.

In this paper, we report the results of a seven-year survey of ground beetles (excluding subfamily Elaphrinae; Fleming and Beresford 2019) found on Akimiski Island. Our results almost double the number of Carabidae known from Nunavut, extend the known range of many species, fill many record gaps between western and eastern Canada, and provide an essentially complete account of the relative abundance of the ground beetle community on Akimiski. Our work also highlights the importance of multiyear studies to provide complete inventories, a common challenge in biodiversity studies (Moreno and Halffter 2000).

Materials and methods

Study area

We conducted our surveys on Akimiski Island, Nunavut, from 2008 to 2014. Akimiski Island has an area of about 3000 km², with habitat characteristics of both tundra and boreal forest (Martini 1981; Blaney and Kotanen 2000). The island is 13.7 km east of Ontario's nearest coastline, with three islands in the strait between Akimiski and Ontario. Akimiski Island emerged from the ocean 3500–4000 years ago, after the retreat of the glaciers (Martini 1981). With no permanent inhabitants, Akimiski Island is part of the Omushkego Cree's territory (Tsuji *et al.* 2020) and is a breeding site for waterfowl, shorebirds, and passerines (Brook *et al.* 2021). Human activity is largely restricted to wildlife harvesting and scientific research, and there is almost no human-altered habitat.

A low-lying island, Akimiski Island has tidal mudflats, marshes, and gravel beach ridges along the coast and fens and bogs in the interior. The most common tree is black spruce, *Picea mariana* (Miller) (Pinaceae). The coastal marshes are dominated by two grasses, *Puccinellia phryhanodes* (Trinius) Scribner and Merrill, and *Festuca rubra* Linnaeus (Poaceae) (Blaney and Kotanen 2000; Kotanen and Abraham 2013). The coastal sand and gravel ridges contain gooseberry, *Ribes oxycanthoides* Linnaeus (Grossulariaceae), and juniper, *Juniperus* sp. Linnaeus (Cupressaceae) shrubs, with willow, *Salix* spp. Linnaeus (Salicaceae), occupying the lower areas (Martini and Glooschenko 1984). The nearby coastal area of Ontario, directly west of Akimiski Island, is essentially the same habitat extending for about 40 km inland, which is characteristic of the Hudson Bay Lowlands (Blaney and Kotanen 2000; Crins *et al.* 2009).

Fieldwork for the present study was conducted at two sample sites on the island's north coast (Fig. 1). The first site was located on the coast at an Ontario Ministry of Natural Resources research station (53° 06' 18" N, 80° 57' 25" W); the second site was located in the island's interior, on a dry ridge of gravel and shallow soil 2 km southwest of the research station (53° 06' 00" N, 80° 58' 00" W). The interior site was a small area (~ 30 m²) of open canopy and young poplar trees that was surrounded by wooded bog and fen.




Figure 1. Inset map of Akimiski Island, Nunavut, Canada. The circle represents two sampling sites, 2008–2014. The eastern two-thirds of the island is the Akimiski Island Bird Sanctuary, the border of which is denoted by the meridian at approximately 80° W.

Collecting methods

Specimens were collected by pitfall traps, modified pitfall-malaise traps, and individual hand capture from mid-July to the first of August each year except 2008, when sampling was conducted from mid-June to the end of August. During 2008 and 2009, we used 10 modified pitfall-malaise traps (International Polar Year, or IPY, traps; McKinnon *et al.* 2008), which acted as both intercept traps for flying insects and pitfall traps for crawling insects (Fig. 2). The pitfall part of the trap was constructed using plastic trays (38 cm long \times 7 cm wide and \times 5 cm high) set with the top edge level with the ground. The trap was surmounted by a collecting bottle to capture flying insects. Specimens that flew into the screen would be directed either into the collecting bottle at the top or fall into the tray at the bottom (Gan *et al.* 2009; Bolduc *et al.* 2013). The trays and collecting bottles were partially filled with soapy water to drown trapped insects. We deployed IPY traps 20 m apart along two transects parallel to the coast. One transect comprised five traps in supratidal habitat; the second transect comprised five traps in the intertidal habitat.

In 2010–2013, we used pitfall traps constructed from 500-mL cups (12 cm high \times 9.2 cm diameter) set in the ground with the top of the cup level with the soil to allow crawling beetles to fall in. The traps were partially filled with about 200 mL of nontoxic propylene glycol (nontoxic RV antifreeze) and did not contain any attractant or bait. Our sampling varied by year: four traps in 2010 (all at the research station), eight traps in 2011 (six at the research station and two at the interior site), 12 traps in 2012 (five at the research station and seven at the interior site), and seven traps in 2013 (all at the interior site). In 2014, sampling was done by hand capture only.

Finally, in all years, we collected beetles by hand *ad hoc* when specimens were spotted.



Figure 2. Photograph of International Polar Year (IPY) trap set on the coast of Akimiski Island, Nunavut, Canada. The pitfall portion of this trap is set below the interception screen at ground level. The white cone acts as a rain shield to prevent the pitfall trough from filling with rainwater and as a funnel to direct climbing insects up into the collection head at the top. Photo by Lisa Pollock.

Specimen preparation and identification

We preserved collected specimens in vials containing 80% ethanol. For identification, specimens were pinned, labelled, and identified using dichotomous keys (Lindroth 1969; Bousquet 2010). Our identifications were confirmed by experts Dr. David Maddison (Oregon State Arthropod Collection, Oregon State University, Corvallis, Oregon, United States of America) and Dr. James Liebherr (Cornell University Insect Collection, Cornell University, Ithaca, New York, United States of America). Pinned specimens are housed in the entomology reference collection at Trent University (Peterborough, Ontario, Canada), except for reference specimens of *Bembidion*, which are deposited in the Oregon State Arthropod Collection, and of *Agonum*, which are deposited in the Cornell University Insect Collection.

Analysis. We determined existing known ranges for each species using records from publications and databases (Lindroth 1969; Bousquet 2010, 2012; Bousquet *et al.* 2013; Ernst and Buddle 2015; Canadian National Collection of Insects, Arachnids and Nematodes (CNC) 2022; Fleming *et al.* 2022; Global Biodiversity Information Facility 2024). We categorised any species with records from northeastern Ontario or northern Québec as “expected.” Species with records from further afield – that is, from western or southern Ontario or Québec – were categorised as new range records.

Given the importance of flight in carabid dispersal (Den Boer 1970; As 1984; Venn 2016), we calculated the proportion of flight-capable *versus* flightless species in our collection. Both morphologies are well represented by species of Carabidae (Lindroth 1969; Bousquet 2010).

Diversity. We estimated the number of species of ground beetles using Chao-1 and iChao-1 numbers (Gotelli and Colwell 2010). Chao-1 estimates the expected number of species from sampled abundances using the proportion of species represented by only one or two specimens (singletons and doubletons) in the total catch (Gotelli and Colwell 2010); iChao1 is similar but also includes species with three or four specimens (Chiu *et al.* 2014). We also assessed diversity using Hill's numbers (Jost 2006). Hill's numbers are interpreted as the number of species (H_0), the number of abundant species (H_1), and the number of very abundant species (H_2 ; Ludwig and Reynolds 1988) – that is, H_0 is species richness, and H_1 and H_2 are algebraically related to Shannon's and Simpson's indices (Jost 2006). Finally, we used the rarefaction method to assess how many specimens needed to be caught to adequately inventory the Carabidae community. Rarefaction returns the mean expected number of species that would have been captured at smaller sample sizes (Krebs 1989). These analyses were performed using PAST 4.0 (Hammer *et al.* 2001).

Results

We collected 1368 specimens, which were identified to 31 species from 15 genera. Nineteen species (853 specimens) have Nearctic distributions, and 12 species (515 specimens) have Holarctic distributions. Of the 31 species caught, 29 are first records for the territory of Nunavut (Table 1). Fourteen of the species are able to fly, whereas 17 species do not fly (Lindroth 1969; Bousquet 2010; Table 1).

Of the 31 species ($H_0 = 31$), seven were determined to be abundant ($H_1 = 7.1$) on Akimiski Island and four were determined to be very abundant ($H_2 = 4.1$). These four were *Bembidion bimaculatum* (Kirby) (571 specimens; 41.7% of the total catch), *Carabus maeander* Fischer von Waldheim (316 specimens; 23.1%), *Amara torrida* (Panzer) (132 specimens; 9.6%), and *Pterostichus punctatissimus* (Randall) (59 specimens; 4.3%). Two of the four, *C. maeander* and *A. torrida*, have Holarctic distributions (Table 1). Four species were represented by a single specimen each, and seven species were represented by two specimens each (Table 1). The Chao-1 and iChao-1 estimates of richness were 31.8 and 32.1, respectively, implying that our sampling likely captured the full complement of species in the study area.

Based on the rarefaction analysis, each year of sampling produced fewer species than expected from the number of captured specimens (Fig. 3). This slow accumulation of species continued for the first three years of data collection (Fig. 3).

Discussion

The arrival of species to islands is a long-standing topic in biogeography (Lomolino *et al.* 2017). As a postglacial rebound island, Akimiski Island is a *tabula rasa*, or blank slate, for new colonists (Coope *et al.* 1986; Buckland and Dugmore 1991). Interestingly, our inventory contained a near-equal mix of flight-capable and flightless species (Table 1). Although the 14 flight-capable species could have arrived by air, the 17 flightless species almost certainly were transported by rafting on floating ice, vegetation, or freshwater slicks (Coope *et al.* 1986). Rafting may have carried the flight-capable species, too (As 1984). The distance from the mainland to Akimiski Island, although not insurmountable, poses a considerable barrier to insect species dispersal, emphasising the importance of both aerial and rafting mechanisms in colonisation (Heatwole and Levins 1972; Kotze *et al.* 2000).

Carabid beetles, in general, exhibit a wide range of flight capabilities, which are highly variable between and even within species (Den Boer 1970; Thiele 1977; Desender 2000). Although most carabids can fly short distances well, the prevailing winds in the region could assist weak fliers travelling to Akimiski Island (As 1984). Additionally, the Akimiski Strait islands would likely act

Table 1. Annual and total numbers of specimens of each species of Carabidae captured on Akimiski Island, Nunavut, Canada, in 2008–2014. Flight capability is from Bousquet (2010), except where indicated *

Species	Realm	Flight-capable	Year							Total
			2008	2009	2010	2011	2012	2013	2014	
<i>Agonum affine</i>	Nearctic	yes	5	10						15
<i>Agonum gratiosum</i>	Holarctic	yes		10						10
<i>Agonum superioris</i>	Nearctic	no		2						2
<i>Amara lacustris</i> †	Nearctic	yes				2				2
<i>Amara latior</i>	Nearctic	yes				2				2
<i>Amara lindrothi</i>	Nearctic	no				6				6
<i>Amara quenseli</i>	Holarctic	yes			1	1	1	2		5
<i>Amara sinuosa</i>	Nearctic	no			1	1				2
<i>Amara torrida</i>	Holarctic	no	9	8	13	89	13			132
<i>Bembidion bimaculatum</i>	Nearctic	yes*			144	363	64			571
<i>Bembidion graphicum</i>	Nearctic	yes*	5	1	9	4	2			21
<i>Bembidion morulum</i>	Nearctic	yes*	1	5						6
<i>Calathus ingratus</i>	Nearctic	no		3	10	15	2			30
<i>Carabus maeander</i>	Holarctic	no	94	220		2				316
<i>Carabus taedatus agassii</i>	Nearctic	no	1	1	10	13	5		1	31
<i>Chlaenius alternatus</i>	Holarctic	no		2						2
<i>Cymindis cribricollis</i>	Nearctic	no			3		2	1		6
<i>Loricera pilicornis pilicornis</i>	Holarctic	yes		4						4
<i>Miscodera arctica</i>	Holarctic	yes		1	1	5	1	1		9
<i>Notiophilus aquaticus</i>	Holarctic	no			1					1
<i>Notiophilus semistriatus</i>	Nearctic	no				1				1
<i>Patrobus foveocollis</i>	Holarctic	yes	1	4						5
<i>Patrobus lecontei</i>	Nearctic	yes				2				2
<i>Patrobus stygicus</i>	Nearctic	no	20	20						40
<i>Pelophila rudis</i>	Nearctic	yes		2						2
<i>Platynus mannerheimii</i>	Holarctic	no	3	14						17
<i>Pterostichus adstrictus</i>	Holarctic	no		4	1	8				13
<i>Pterostichus brevicornis</i> †	Holarctic	no	1							1
<i>Pterostichus patruelis</i>	Nearctic	yes	19	35						54
<i>Pterostichus punctatissimus</i>	Nearctic	no	10	48		1				59
<i>Sericoda obsoleta</i>	Nearctic	yes		1						1
Number of species			12	20	11	16	8	3	1	31
Number of species with singletons and doubletons			4	7	5	8	5	3	1	11
Number of new species			12	9	5	5	0	0	0	
Total catch			169	395	194	515	90	4	1	1368

*Based on Lindroth (1969); either not listed (*B. bimaculatum*, *B. graphicum*) or listed as flightless (*B. morulum*) in Bousquet (2010)

† Species previously recorded from Nunavut




Figure 3. Rarefaction analysis of the mean number of Carabidae species expected based on the number of collected specimens, Akimiski Island, Nunavut, Canada, 2008–2014. The dashed lines are the upper and lower 95% confidence limits. White symbols represent individual years; black symbols represent cumulative years. The values for 2008–2012 and 2008–2014 are coincident.

as stepping stones, facilitating colonisation by providing rest points for migrating beetles (MacArthur and Wilson 2001). The establishment of founding populations on Akimiski Island further underscores the importance of these varied dispersal strategies. Overall, the mix of flight-capable and flightless species highlights the complex dynamics of species dispersal to this isolated island (Venn 2016).

Dispersal to Akimiski by flight

Flight facilitates carabid dispersal (Den Boer 1970, 1977). In our study, eight species found on Akimiski Island are undocumented in watersheds in the eastern half of northern Ontario that empty into James Bay: *Amara lacustris* LeConte, *Amara lindrothi* Hieke, *Bembidion bimaclatum*, *Bembidion graphicum* Casey, *Bembidion morulum* LeConte, *Patrobis lecontei* Chaudoir, *Pelophila rudis* (LeConte), and *Sericoda obsoleta* (Say). Only one of these species is flightless, *A. lindrothi*, which was only recently described and is consequently underrepresented in published records. The other seven species are known to fly. The lack of records of these seven flight-capable species from the northeastern Ontario watersheds is consistent with these species not depending on flotsam as a transport mechanism (Buckland 1988; Fleming *et al.* 2021).

Dispersal to Akimiski by rafting

The direction of flow of water in James and Hudson bays is affected by tides, water currents, and wind direction and could be important to the arrival of colonists to the island. The current in southern Hudson Bay flows eastwards along the north shore of Ontario, with some current entering the east side of James Bay (Hachey 1935). In James Bay, water flows south along the western side of the bay past Akimiski Island towards the southern coast of the bay (Martini 1981; Stewart and Lockhart 2005) and then follows the coastline, flowing east and then north along the eastern side (Stewart and Lockhart 2005). The surface water generally flows almost due east (St-Laurent *et al.* 2011); however, wind changes can reverse the direction of surface water movements (Prinsenber 1978).

The closest distance between Akimiski Island and the mainland is approximately 14 km. Akimiski is situated east of Akimiski Strait, opposite the mouth of the Attawapiskat River (Déry *et al.* 2011), south of the outflows from the Ekwan River, and north of the outflow from the Albany and Moose River systems (Déry *et al.* 2011). Debris floating downstream from the Ekwan and Attawapiskat rivers (among others) into James Bay could arrive on the shore of Akimiski Island. Due to the northward current on the east side of James Bay, it is less likely that debris from Québec

rivers would end up on Akimiski Island. However, debris in the eastern or southern parts of James Bay could also be carried by surface water to Akimiski Island (Barber 1972).

New records and abundance

Our study substantially augments the list of carabid beetles in Nunavut. Indeed, Akimiski Island is at the southernmost reaches of this jurisdiction, and species richness is generally higher at lower latitudes (Lomolino *et al.* 2017). Many of these new territorial records are artefacts of political boundaries – a reminder that jurisdiction boundaries, although important to listing and conservation responsibility, are not synonymous with geographical or ecological boundaries.

At present, 41 species of Carabidae (excluding Cicindelidae; Duran and Gough 2020) are recognised from Nunavut: 34 are listed in Bousquet *et al.* (2013) and seven Elaphrinae Nunavut records appear in Fleming and Beresford (2019). The addition of our 29 territorial records for Nunavut increases this total to 70 Carabidae species. All are native species. Moreover, our inventory of the Carabidae on Akimiski Island reveals both similarities and differences with the carabid communities found on the adjacent mainland. For example, many more species are known in Ontario (517), Québec (474), and the Northwest Territories (212; Table 2). Nunavut and Yukon are the only regions in Canada with no known adventive species (Bousquet *et al.* 2013). This pattern is consistent with the global tendency for polar regions to have lower proportions of alien species than elsewhere (Alsos *et al.* 2015).

Two of the 31 species caught in the present study, *A. lacustris* LeConte, 1855 and *Pterostichus brevicornis* (Kirby), had been previously recorded in mainland Nunavut. *Amara lacustris* is transcontinental, but its distribution is predominantly in northern North America (Lindroth 1969). It is flight-capable and is not known from Québec or Ontario. The two specimens in our collection represent an eastward range extension of more than 1000 km. *Pterostichus brevicornis* is flightless, with a Holarctic distribution across northern Eurasia, Alaska, and northern Canada, including Ontario (Fleming *et al.* 2022) and Québec (Bousquet 2012).

We anticipated that northern species documented elsewhere might be part of our collection. Nineteen of the 29 species (comprising 710 specimens) with new records in this study are known from similar latitudes and habitats in the adjacent provinces of Ontario and Québec (Lindroth 1969; Ernst and Buddle 2015; Fleming *et al.* 2022). The species documented at similar latitudes are *Agonum affine* Kirby, *Agonum gratiosum* (Mannerheim), *Agonum superioris* Lindroth, *Amara latior* (Kirby), *Amara torrida*, *Calathus ingratus* Dejean, *Carabus maeander*, *Chlaenius alternatus* Horn, *Cymindis cribricollis* Dejean, *Loricera pilicornis pilicornis* (Fabricius), *Miscodera arctica* (Paykull), *Notiophilus aquaticus* (Linnaeus), *Notiophilus semistriatus* Say, *Patrobus foveocollis* (Eschscholtz), *Platynus mannerheimii* (Dejean), *Pterostichus adstrictus* Eschscholtz, *Pterostichus brevicornis*, *Pterostichus patruelis* (Dejean), and *Pterostichus punctatissimus*. All 19 species were expected from existing records, and our results, of which 18 are first records for Nunavut (*P. brevicornis* having been previously reported in that territory), do not fill substantial range gaps for these species or extend the ecological range of these species. All 19 are described as widespread in northeastern North America by Bousquet (2010) and as circumpolar, transcontinental, or transamerican by Lindroth (1969; Lindroth (1969) reports *P. mannerheimii* as *Agonum mannerheimi* Dejean, 1828). Of these, 13 species are flightless (Table 1). This life-history feature, plus the species' presence in northeastern Ontario or northern Québec, is consistent with the hypothesis of these species being transported on debris to Akimiski Island.

Three other species have southern Ontario and Québec records: *Amara quenseli* (Schönherr), *Amara sinuosa* (Casey), and *Patrobus stygicus* Chaudoir. These three species are also widespread in northeastern North America (Bousquet 2010), and although our records fill range gaps between southern Ontario and Québec, we also expected to find these species on Akimiski Island because of their widespread distributions.

Table 2. Number of Carabidae species in Nunavut and adjacent jurisdictions. Cicindelidae, although listed in Bousquet (2010) and Bousquet *et al.* (2013) as subfamily Cicindelinae, is now considered as a distinct family (Duran and Gough 2020)

Jurisdiction	Number of species						
	Total, including Cicindelidae ¹	Cicindelidae ²	Total, less Cicindelidae	Additional species ³	Total species	Adventive ⁴	Native
Ontario	532	16	516	1	517	21	496
Québec	479	15	464		464	32	432
NWT	218*	6	212		212	1	211
Nunavut	34	0	34	7	41	0	41

¹Bousquet *et al.* (2013, Table 1).

²Bousquet (2010, inferred from pp. 43 and 35).

³Fleming and Beresford (2019).

⁴Bousquet (2012, Table 5). * 218 is from Bousquet (2012); note: Bousquet *et al.* (2013) reported this as 217.

Surprisingly, the list of Carabidae on Akimiksi Island that we anticipated based on nearby occurrences (above) does not include three of our seven most abundant species: *Bembidion bimaculatum* (571; 42% of the specimens captured), *Patrobus stygicus* (40; 2.9%), and *Carabus taedatus agassii* LeConte (31; 2.3%). We expected that more abundant species would have nearby records and that any range extensions would be from those species that were rare in our collections – a reasonable expectation, but wrong in this case. These species' abundance on Akimiksi Island may represent a case of ecological release (Lomolino *et al.* 2017) – that is, the filling of vacant ecological space where insular species are free of competition from mainland species. We discuss these and other individual species below.

Species accounts

The remaining eight species caught during the present study are either new range records or gap infills. All are Nearctic species, and only two are flightless: *A. lindrothi* and *C. taedatus agassii*.

Amara lindrothi. *Amara lindrothi* was first described in 1990 (Bousquet 2012) and is not commonly found; we collected only six individuals. Records are scarce and widespread; most are from the southwestern United States of America. In Canada, the species has been found in the Yukon, northern Alberta, northern Manitoba, and Labrador; it is not known from either Ontario or Québec. In the United States of America, it has been found in Colorado, New Mexico, Nevada, Utah, and Wyoming (Bousquet 2012; Bousquet *et al.* 2013; Global Biodiversity Information Facility 2024). The habitat preferences of *A. lindrothi* are not known other than that the beetle occurs in the Arctic bioclimatic zone (Bousquet 2010). The few records may, in part, be attributed to having been described only 34 years ago. Our record marks the species' presence in the vast distributional gap between Manitoba and Labrador.

Bembidion bimaculatum. We caught 571 *B. bimaculatum*, our most abundant species. It is a predominantly western species (Lindroth 1969), and Bousquet (2012) describes the northern part of its range as “north-central Ontario (CNC 2022) to the Arctic Plains of Alaska.” Current distribution records are scarce east of Manitoba. Ontario records are from Geraldton, about 100 km north of Lake Superior (CNC 2022), as well as Moosonee and the Ottawa area (Global Biodiversity Information Facility 2024). There are also records from New Hampshire (Global Biodiversity Information Facility 2024) and Greenland (Lindroth 1969). We could not find any habitat data for this species; our records add to knowledge of the eastern distribution of this

species, and the abundance of our collections suggests the species may prefer gravel ridge or coastal habitats.

Bembidion graphicum. *Bembidion graphicum* was common in our study: 21 specimens were collected. It is a western North American species (Bousquet 2012). In 2010, two specimens were collected on Manitoulin Island, Ontario, the easternmost records for this species (Paiero 2017). Our collection of 21 specimens of *B. graphicum* from Akimiski Island is 800 km north of Manitoulin Island and is a large range extension in the eastern part of the range. *Bembidion graphicum* prefers habitats along the margins of pools and lakes and saline environments (Lindroth 1969), habitats common on Akimiski Island.

Bembidion morulum. *Bembidion morulum* has been collected in Churchill, Manitoba, and in northern Ontario near the Manitoba border near Fort Severn (CNC 2022). Records exist from Newfoundland (Bousquet 2010, 2012). Records are scarce (only 27 sites are listed in the Global Biodiversity Information Facility 2024 database, and three sites are listed in CNC 2022). With so few records, the present study provides important additional records for this species and partially fills a record gap from Fort Severn to Newfoundland, a stretch of some 2100 km.

Carabus taedatus agassii. *Carabus taedatus agassii* is known from Newfoundland in the east to the Yukon Territory in the northern part of North America (Bousquet 2010, 2012), with intermediate records from Québec (Bousquet 2010). Lindroth (1969) described the range as remarkably disjunct. Ontario records exist for about 100 km north of Lake Superior (Global Biodiversity Information Facility 2024) and at three sites near the Hudson Bay coast (Fleming *et al.* 2022; Global Biodiversity Information Facility 2024). The record from the present study fills a gap between the Ontario and Québec reports.

Patrobis lecontei. *Patrobis lecontei* has an extensive west-to-east distribution across much of southern Canada, with a record gap in Ontario (Bousquet 2012; Global Biodiversity Information Facility 2024). The species is found along standing water among sedges but never in sphagnum moss (Lindroth 1969). Our two specimens are at the northern edge of its known range, 620 km from the closest eastern record in northern Québec, and fill a large record gap between Manitoba and Québec.

Pelophila rudis. The range of *P. rudis* was described by Lindroth (1969) as “a rare and local species, restricted to Canada.” It is known from Newfoundland in the east, with no records until Fort Severn and Cape Henrietta Maria on the Hudson Bay coast in Ontario (Lindroth 1969), and records westward in Manitoba. Our record of this uncommon species helps fill the vast known range gap, placing *P. rudis* east of Cape Henrietta Maria.

Sericoda obsoleta. This species is found across Canada, with records from Newfoundland to the Yukon Territory. Ontario records extend from Toronto in the southern part of the province to Moose Factory in the north (Lindroth 1969). It is known from the southern half of Québec. In eastern North America, our record extends the known range northwards from Moose Factory by about 200 km.

Conclusions and significance

As the rarefaction analysis reveals, four years of sampling were required to capture all 31 species (Table 1), and during the first three years, the aggregate tally of species fell somewhat short of expectations based on the tally of specimens (Fig. 3). Notably, rarefaction is based on resampling being random. Even though our traps were placed in similar areas each year, the years differed in phenology, a major determinant of insect activity. Our results underscore the potential importance of seasonal timing in surveys of insects, and of ground beetles in particular (Niemelä *et al.* 1989; Adlam *et al.* 2017). For example, our mid-summer sampling captured only one *Notiophilus aquaticus*, whereas, in Norway, most have been caught in late August (Erikstad *et al.* 1989).

Transitional areas, such as Akimiski Island, are often the first to show evidence of altered species composition in response to habitat changes (Payette *et al.* 2004). Although all the species in our collection from Akimiski Island were native to eastern Canada, many nonnative ground beetle species are found in the neighbouring regions of Ontario (Fleming *et al.* 2022) and Québec (Bousquet 2012). The absence of nonnative ground beetles highlights the island's importance as a reference point for understanding ecological change. Akimiski Island therefore serves as an important point for comparisons both in space and time – that is, as a contrast with regions harbouring introduced species and as a baseline for shifts in the community, for example, should introduced species disperse to Nunavut and to Akimiski Island.

All ground beetles on Akimiski Island arrived in the last 3500–4000 years, after the island rebounded from the ocean following the retreat of the glaciers (Martini and Glooschenko 1984). A period of a few millennia is not long in ecological time. Ground beetle species in northern temperate regions are still recovering from the last glacial period (Baselga *et al.* 2012; Fleming *et al.* 2021), and there is no reason to assume that ground beetles have completed their dispersal to Akimiski Island. For example, in Greenland, only 17% of the Carabidae species that are able to survive in that climate and habitat are present (Coope *et al.* 1986). Given the relative newness of Akimiski Island, novel species, including nonnative forms, likely will continue to arrive. The present study's results provide a basis for identifying those arrivals, contingent on continued monitoring and long-term investment. In this regard, we believe that the present study may help remedy the truncated baselines that often obscure ecological change (Mihoub *et al.* 2017).

Acknowledgements. We thank the many personnel of the Ontario Ministry of Natural Resources' Far North and waterfowl programmes for their support. Funding was provided by the Ontario Ministry of Natural Resources through its Far North Branch and Wildlife Research and Development Section. We also thank the many dedicated researchers, technicians, volunteers, and pilots who helped collect the IPY and pitfall specimens and whose indefatigable cheerfulness in often harsh conditions made the time at the research station so enjoyable and productive. A special thanks to Lisa Pollock for the photograph of the IPY trap. We thank Dr. David Maddison, Director of the Oregon State Arthropod Collection (Oregon State University, Corvallis, Oregon, United States of America), for confirming the *Bembidion* identifications and Dr. James Liebherr, Professor Emeritus and Emeritus Curator of the Cornell University Insect Collection (Ithaca, New York, United States of America), for confirming the *Agonum* identifications. Finally, we thank the anonymous reviewers of an earlier draft for their careful attention and suggestions.

Competing interests. The authors declare that they have no competing interests.

References

- Adlam, C., Despland, E., and Beaulieu, F. 2017. Spatial and temporal patterns of ground beetle (Coleoptera: Carabidae) activity across habitats at the Mont St. Hilaire Biosphere Reserve, Quebec. *The Journal of the Entomological Society of Ontario*, **148**: 23–38.

- Alsos, I.G., Ware, C., and Elven, R. 2015. Past Arctic aliens have passed away, current ones may stay. *Biological Invasions*, **17**: 3113–3123. <https://doi.org/10.1007/s10530-015-0937-9>.
- As, S. 1984. To fly or not to fly? Colonization of Baltic islands by winged and wingless carabid beetles. *Journal of Biogeography*, **11**: 413–426.
- Barber, F.G. 1972. On the Oceanography of James Bay. Manuscript Report Series 24. Department of the Environment, Marine Sciences Branch, Ottawa, Ontario, Canada. 95 pp.
- Baselga, A., Lobo, J.M., Svenning, J.C., Aragón, P., and Araújo, M.B. 2012. Dispersal ability modulates the strength of the latitudinal richness gradient in European beetles. *Global Ecology and Biogeography*, **21**: 1106–1113. <https://doi.org/10.1111/j.1466-8238.2011.00753.x>.
- Bell, R.T. 1990. Insecta: Coleoptera Carabidae (adults and larvae). *In Soil Biology Guide*. Edited by D.L. Dindal. Wiley, Toronto, Ontario, Canada. Pp. 1053–1092.
- Blaney, C.S. and Kotanen, P.M. 2000. The vascular flora of Akimiski Island, Nunavut Territory, Canada. *The Canadian Field-Naturalist*, **115**: 88–98.
- Bolduc, E., Casajus, N., Legagneux, P., McKinnon, L., Gilchrist, H.G., Leung, M., *et al.* 2013. Terrestrial arthropod abundance and phenology in the Canadian Arctic: modelling resource availability for Arctic-nesting insectivorous birds. *The Canadian Entomologist*, **145**: 155–170. <https://doi.org/10.4039/tce.2013.4>.
- Bousquet, Y. 2010. Illustrated Identification Guide to Adults and Larvae of Northeastern North American Ground Beetles (Coleoptera: Carabidae). PenSoft Publishers, Sofia, Bulgaria.
- Bousquet, Y. 2012. Catalogue of Geadephaga (Coleoptera, Adephaga) of America, north of Mexico. *ZooKeys*, **245**: 1–1722. <https://doi.org/10.3897/zookeys.245.3416>.
- Bousquet, Y., Bouchard, P., Davies, A.E., and Sikes, D. 2013. Checklist of beetles (Coleoptera) of Canada and Alaska. Second edition. *ZooKeys*, **360**: 1–44. <https://doi.org/10.3897/zookeys.360.4742>.
- Brook, R.W., Leafloor, J.O., Abraham, K.F., Ankney, C.D., and Patton, K.A. 2019. Canada goose gosling mortality during pre fledging and early migration on Akimiski Island, Nunavut. *Journal of Fish and Wildlife Management*, **10**: 314–322.
- Brook, R.W., Pollock, L., Abraham, K.F., and Brown, G. 2021. Bird trends from long-term observation data at sites in the Hudson Bay Lowlands. *Avian Conservation and Ecology*, **16**: 10.
- Buckland, P. 1988. North Atlantic faunal connections introduction or endemics? *Entomologica Scandinavica Supplement*, **32**: 7–29.
- Buckland, P. and Dugmore, A. 1991. ‘If this is a refugium, why are my feet so bloody cold?’ The origins of the Icelandic biota in the light of recent research. *In Environmental Change in Iceland: Past and Present*. Edited by J.K. Maizels and C. Caseldine. Kluwer Academic Publishers, Dordrecht, The Netherlands. Pp. 107–125.
- Canadian National Collection of Insect, Arachnids and Nematodes (CNC). 2022. CNC Collection Database [online]. Available from <https://www.cnc.agr.gc.ca/taxonomy/TaxonMain.php> [accessed 15 December 2022].
- Chiu, C.H., Wang, Y.T., Walther, B.A., and Chao, A. 2014. An improved nonparametric lower bound of species richness *via* a modified Good–Turing frequency formula. *Biometrics*, **70**: 671–682.
- Coope, G.R., Moore, P.D., and Gibbs, A. 1986. The invasion and colonization of the North Atlantic islands: a palaeoecological solution to a biogeographic problem. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, **314**: 619–635.
- Crins, W.J., Gray, P.A., Uhlig, P.C., and Wester, M.C. 2009. The Ecosystems of Ontario. Part I: Ecozones and Ecoregions. SIB TER IMA TR-01. Inventory, Monitoring and Assessment, Ontario Ministry of Natural Resources, Peterborough, Ontario, Canada. 71 pp.
- DeGasparro, S.L, Brown, G.S., Alarie, Y., and Beresford, D.V. 2018. Predaceous diving beetle records from Akimiski Island, Nunavut (Coleoptera: Dytiscidae). *The Coleopterists Bulletin*, **72**: 866–869. <https://doi.org/10.1649/0010-065X-72.4.866>.

- Den Boer, P.J. 1970. On the significance of dispersal power for populations of carabid beetles (Coleoptera, Carabidae). *Oecologia*, **4**: 1–28.
- Den Boer, P.J. 1977. Dispersal power and survival: carabids in a cultivated countryside. Communication of the Biological Station Wijster, no 169. H. Veen Man and Zonen B.V., Wageningen, The Netherlands.
- Déry, S.J., Mlynowski, T.J., Hernández-Henríquez, M.A., and Straneo, F. 2011. Interannual variability and interdecadal trends in Hudson Bay streamflow. *Journal of Marine Systems*, **88**: 341–351.
- Desender, K. 2000. Flight muscle development and dispersal in the life cycle of carabid beetles: patterns and processes. *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique*, **70**: 13–31.
- Duran, D.P. and Gough, H.M. 2020. Validation of tiger beetles as a distinct family (Coleoptera: Cicindelidae), review and reclassification of tribal relationships. *Systematic Entomology*, **45**: 723–729.
- Erikstad, K.E., Byrkjedal, I., and Kålås, J.A. 1989. Resource partitioning among seven carabid species on Hardangervidda, southern Norway. *In Annales Zoologici Fennici*. Finnish Zoological Publishing Board, Finnish Academy of Sciences, Societas Scientiarum Fennica, Societas pro Fauna et Flora Fennica and Societas Biologica Fennica Vanamo, Helsinki, Finland. Pp. 113–120.
- Ernst, C.M. and Buddle, C.M. 2015. Drivers and patterns of ground-dwelling beetle biodiversity across northern Canada. *PLOS One*, **10**: e0122163. <https://doi.org/10.1371/journal.pone.0122163>.
- Fleming, K.J. and Beresford, D.V. 2019. Range updates for eight species of the Elaphrinae subfamily (Coleoptera: Carabidae) in Ontario's far north and Akimiski Island, Nunavut. *The Coleopterists Bulletin*, **73**: 433–439.
- Fleming, K.J., Schaefer, J., Abraham, K., Smith, A., and Beresford, D.V. 2021. Evidence for passive dispersal of ground beetles (Coleoptera: Carabidae) in the Nearctic boreal forest. *Ecoscience*, **28**: 93–105. <https://doi.org/10.1080/11956860.2021.1872265>.
- Fleming, K.J., Schaefer, J.A., and Beresford, D.V. 2022. New records and range extensions of Carabidae of Ontario's boreal forest. *The Canadian Entomologist*, **154**: 1–25. <https://doi.org/10.4039/tce.2022.33>.
- Gan, S.K., Abraham, K.F., Brook, R.W., and Murray, D.L. 2019. The influence of habitat selection on Canada Goose, *Branta canadensis*, nest success on Akimiski Island, Nunavut, Canada. *Wildfowl*, **69**: 118–133.
- Gan, S., Jumeau, Z., Beresford, D.V., and Abraham, K.F. 2009. The Terrestrial and Low-flying Arthropods of Akimiski Island, Nunavut: A Bioinventory. Ontario Ministry of Natural Resources, Peterborough, Canada.
- Global Biodiversity Information Facility. 2024. GBIF data portal [online]. Available from <https://www.gbif.org> [accessed May 10 2024].
- Gotelli, N.J. and Colwell, R.K. 2010. Estimating species richness. *In Biological Diversity: Frontiers in Measurement and Assessment*. Edited by A.E. Magurran and B.J. McGill. Oxford University Press, New York, New York, United States of America. Pp. 39–54.
- Hachey, H.B. 1935. The circulation of Hudson Bay water as indicated by drift bottles. *Science*, **82**: 275–276.
- Hammer, Ø., Harper, D.A.T., and Ryan, P.D. 2001. PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica*, **4**: 1–9.
- Heatwole H. and Levins, R. 1972. Biogeography of the Puerto Rican bank: flotsam transport of terrestrial animals. *Ecology*, **53**: 112–117. <https://doi.org/10.2307/1935715>.
- Jost, L. 2006. Entropy and diversity. *Oikos*, **113**: 363–375.
- Kolenosky, G.B. and Prevett, J.P. 1983. Productivity and maternity denning of polar bears in Ontario. *Bears: Their Biology and Management*, **5**: 238–245.
- Kotanen, P.M. and Abraham, K.F. 2013. Decadal changes in vegetation of a subarctic salt marsh used by lesser snow and Canada geese. *Plant Ecology*, **214**: 409–422.

- Kotze, D.J., Niemelä, J., and Nieminen, M. 2000. Colonization success of carabid beetles on Baltic islands. *Journal of Biogeography*, **27**: 807–819.
- Krebs, C.J. 1989. *Ecological Methodology*. Harper & Row, New York, New York, United States of America.
- Lindroth, C.H. 1969. The ground-beetles (Carabidae, excl. Cicindelinae) of Canada and Alaska. Parts 1–6. *Opuscula Entomologica Supplementum*, **20**, **24**, **29**, **33**, **34**, **35**. Berlingska Boktryckeriet, Sweden.
- Lomolino, M., Riddle, B., and Whittaker, R.J. 2017. *Biogeography*. Sinauer, Sunderland, Massachusetts, United States of America.
- Lövei, G.L. and Sunderland, K.D. 1996. Ecology and behavior of ground beetles (Coleoptera: Carabidae). *Annual Review of Entomology*, **41**: 231–236.
- Ludwig, J.A. and Reynolds, J.F. 1988. *Statistical Ecology*. John Wiley and Sons, New York, New York, United States of America.
- MacArthur, R.H. and Wilson, E.O. 2001. *The Theory of Island Biogeography*. Princeton University Press, Princeton, New Jersey, United States of America.
- Martini, I.P. 1981. Morphology and sediments of the emergent Ontario coast of James Bay, Canada. *Geografiska Annaler: Series A, Physical Geography*, **63**: 81–94.
- Martini, I.P. and Glooschenko, W.A. 1984. Emergent coasts of Akimiski Island, James Bay, Northwestern Territories, Canada: geology, geomorphology, and vegetation. *Sedimentary Geology*, **37**: 229–250.
- Maxwell, S.L., Fuller, R.A., Brooks, T.M., and Watson, J.E.M. 2016. Biodiversity: the ravages of guns, nets and bulldozers. *Nature*, **536**: 143–145. <https://doi.org/10.1038/536143a>.
- McKinnon, L., Bety, J., Smith, P., Morrison, G., and Bolduc, E. 2008. Arthropod monitoring: seasonal variation in the abundance of surface active, sub-surface and low-flying arthropods. ArcticWOLVES monitoring protocols. Version 1 [online]. Available from https://science.cen.ulaval.ca/arcticwolves/document/arthropod_monitoring.pdf [accessed 3 July 2024].
- Mihoub, J.B., Henle, K., Titeux, N., Brotons, L., Brummitt, N.A., and Schmeller, D.S. 2017. Setting temporal baselines for biodiversity: the limits of available monitoring data for capturing the full impact of anthropogenic pressures. *Scientific Reports*, **7**: 41591. <https://doi.org/10.1038/srep41591>.
- Moreno, C.E. and Halffter, G. 2000. Assessing the completeness of bat biodiversity inventories using species accumulation curves. *Journal of Applied Ecology*, **37**: 149–158.
- Nguyen, L.P., Nol, E., and Abraham, K.F. 2003. Nest success and habitat selection of the semipalmated plover on Akimiski Island, Nunavut. *The Wilson Bulletin*, **115**: 285–291.
- Niemelä, J., Haila, Y., Halme, E., Pajunen, T., and Punttila, P. 1989. The annual activity cycle of carabid beetles in the southern Finnish taiga. *In* *Annales Zoologici Fennici*. Finnish Zoological Publishing Board, Finnish Academy of Sciences, Societas Scientiarum Fennica, Societas pro Fauna et Flora Fennica and Societas Biologica Fennica Vanamo, Helsinki, Finland. Pp. 35–41.
- O, P.C., Kotanen, P.M., and Abraham, K.F. 2005. Survival and growth of the forage grass *Festuca rubra* in naturally and artificially revegetated sites in a subarctic coastal marsh. *Ecoscience*, **12**: 279–285.
- Obbard, M.E. and Middel, K.R. 2012. Bounding the southern Hudson Bay polar bear subpopulation. *Ursus*, **23**: 134–144.
- Paiero, S. 2017. University of Guelph Insect Collection (DEBU). Version 2.2. Occurrence dataset [online]. University of Guelph, Guelph, Ontario, Canada. Available from <https://doi.org/10.5886/4pp2vt5a> [via GBIF.org; accessed 17 July 2023].
- Payette, S., Delwaide, A., Caccianiga, M., and Beauchemin, M. 2004. Accelerated thawing of subarctic peatland permafrost over the last 50 years. *Geophysical Research Letters*, **31**: L18208.

- Peacock, E., Derocher, A.E., Lunn, N.J., and Obbard, M.E. 2010. Polar bear ecology and management in Hudson Bay in the face of climate change. *In A Little Less Arctic: Top Predators in the World's Largest Northern Inland Sea, Hudson Bay. Edited by S.H. Ferguson, L.L. Loseto, and M.L. Mallory.* Springer, Dordrecht, The Netherlands. Pp. 93–116.
- Pollock, L.A., Abraham, K.F., and Nol, E. 2012. Migrant shorebird use of Akimiski Island, Nunavut: a sub-Arctic staging site. *Polar Biology*, **35**: 1691–1701.
- Prinsenberg, S.J. 1978. Analytical Study of the Circulation of James Bay. Manuscript Report Series Number 9. Fisheries and Environment Canada, Ocean and Aquatic Sciences Central Region, Canada Centre for Inland Waters, Burlington, Ontario, Canada.
- Richards, J.M. and Gaston, A.J. (editors). 2018. *Birds of Nunavut.* University of British Columbia Press, Vancouver, British Columbia, Canada.
- Stewart, D.B. and Lockhart, W.L. 2005. An Overview of the Hudson Bay Marine Ecosystem [online]. Canadian Technical Report of Fisheries and Aquatic Sciences 2586. Fisheries and Oceans Canada, Ottawa, Ontario, Canada. 487 pp. Available from <http://www.dfo-mpo.gc.ca/Library/314704.htm> [accessed 15 May 2024].
- St-Laurent, P., Straneo, F., Dumais, J.F., and Barber, D.G. 2011. What is the fate of the river waters of Hudson Bay? *Journal of Marine Systems*, **88**: 352–361.
- Thiele, H.U. 1977. *Carabid Beetles in Their Environments.* Springer, New York, New York, United States of America.
- Tsuji, L.J., General, Z., Tsuji, S.R., Powell, E., Latychev, K., Clark, J., and Mitrovica, J.X. 2020. Akimiski Island, Nunavut, Canada. *Arctic*, **73**: 421–432.
- Venn, S. 2016. To fly or not to fly: factors influencing the flight capacity of carabid beetles (Coleoptera: Carabidae). *European Journal of Entomology*, **113**: 587–600, <https://doi.org/10.14411/eje.2016.079>.
- Venter, O., Brodeur, N.N., Nemiroff, L., Belland, B., Dolinsek, I.J., and Grant, J.W.A. 2006. Threats to endangered species in Canada. *BioScience*, **56**: 903–910.
- Vergara-Asenjo, G., Alfaro, F.M., and Pizarro-Araya, J. 2023. Linnean and Wallacean shortfalls in the knowledge of arthropod species in Chile: challenges and implications for regional conservation. *Biological Conservation*, **281**: 110027.
- Whittaker, R.J., Araújo, M.B., Jepson, P., Ladle, R.J., Watson, J.E., and Willis, K.J. 2005. Conservation biogeography: assessment and prospect. *Diversity and Distributions*, **11**: 3–23.

Cite this article: Fleming, K.J., Schaefer, J.A., Abraham, K.F., Brook, R.W., Brown, G.S., and Beresford, D.V. 2024. Ground beetles (Coleoptera: Carabidae) of Akimiski Island, Nunavut, Canada. *The Canadian Entomologist*. <https://doi.org/10.4039/tce.2024.26>.