

COMMENTARY

Older people's mental health is more complex than urban-rural differences

Commentary on “Decomposition and Comparative Analysis of Differences in Depressive Symptoms between Urban and Rural Older Adults: Evidence from a National Survey” by Yuan *et al.*

Katarzyna M. Lion¹  and Jenny Murfield² 

¹Menzies Health Institute Queensland, Griffith University, Brisbane, QLD, Australia

²Food & Mood Centre, The Institute for Mental and Physical Health and Clinical Translation (IMPACT), School of Medicine, Deakin University, Geelong, VIC, Australia

Email: k.lion@griffith.edu.au

With the increasing interest in health equity, as highlighted by special issue of *International Psychogeriatrics*, there is a particular need to delineate that not all health differences can be framed as health disparities alone, i.e. differences in health status between groups, such as younger and older people. Health disparities are closely linked to differences in people's economic, social, or environmental disadvantages, which then adversely affect their health outcomes and care, such as being part of a discriminated group (i.e. racial/ethnic group) or due to geographic location (Braveman, 2014). These health inequalities have many faces, exist within and across countries, and require coordinated international and national multi-level changes in policies, government programs, and systems to be reduced (WHO, 2015).

The inequalities between urban and remote/rural areas within the health and social systems are well-known (WHO, 2010). This disparity is mostly caused by the limited availability of health care workers who prefer living and working in urban areas (76% of physicians and 62% of nursing workforce), despite half of the world's population living in remote and rural areas (WHO, 2010). This discrepancy is commonly observed in most countries, including the USA, Australia, and France; however, it has the biggest impact in low-income countries, such as Bangladesh, the Democratic Republic of the Congo and Mali (WHO, 2010). One of the ways to reduce these inequalities, as proposed by the WHO (2010), is by encouraging health professionals to work in more remote/rural areas.

Unfortunately, the general inequality within services across urban-rural areas mostly affects the most vulnerable groups who usually require higher levels of healthcare access due to their increased risk of diseases, such as older people. There is a relatively well-explored rural-urban gap in physical health among older adults. Conversely, research investigating the rural-urban gap in cognitive functioning is scant (Glauber, 2022), with what is available suggesting increased risk of developing dementia or mild cognitive impairment when living in rural regions (Weden *et al.*, 2018).

Yet, there is not much known about how populations of older people are affected by those differences in access to care and if there are other modifiable factors, outside of changes in health systems and providing more services and resources for remote and rural populations, which can play a role in reducing those differences and supporting better healthcare outcomes.

The paper by Yuan and colleagues (2023) in this *International Psychogeriatrics* issue sheds important new light on these factors. The authors investigated factors influencing urban-rural discrepancies in depressive symptoms among older adults in China and measured their contribution as part of the 2018 Chinese Longitudinal Health Longevity Survey. In their paper, the authors present data of $n = 11,245$ older adults from rural ($n = 7,907$) and urban ($n = 3,338$) areas of China (covering 23 provinces). They reported higher levels of depressive symptoms among those living in rural areas (12.41% vs. 10.13%). This finding adds to the research body reporting increased depressive symptoms across

rural areas in older people (i.e. Wang *et al.*, 2021), but sits in contrast to other studies that either show the opposite (i.e. Ziarko *et al.*, 2015) or find no evidence of a rural-urban difference (i.e. Sun *et al.*, 2022).

A novel aspect of the study by Yuan and colleagues (2023) is that they conducted Fairlie decomposition analysis to explore factors that might be influencing the difference in depressive symptoms between the urban and rural older adults in their cohort. Their analyses showed that the difference in depressive symptoms between Chinese people living in urban and rural areas was related to a series of modifiable factors, explaining nearly three-quarters (73.96%) of the observed difference. These modifiable factors include annual income (31.51%), education level (28.05%), sleep time (−25.67%), self-reported health status (24.18%), dysfunction in instrumental activities of daily living (20.73%), exercise (17.72%), living status (−8.31%), dysfunction in daily living activities (−3.29%), and social activity (2.44%). The other 26.04% of the difference in depressive symptoms was related to urban/remote and unobserved factors. The authors also calculated protective and risk factors for developing depressive symptoms and these differed for rural and urban older adults in their study. For example, both higher annual income and exercising were specific protective factors in urban areas, whereas one of the non-modifiable risk factors associated with higher depressive symptoms was 80 years or older (Yuan *et al.*, 2023).

Other research confirms that these modifiable factors, such as coping strategies, lifestyle, and environmental changes, might support better well-being of people living in rural areas (Ziarko *et al.*, 2015). Additionally, Cosco and colleagues (2018) associate adult socioeconomic advantage (i.e. social support and leisure time physical activity) with greater resilience (understood as higher well-being despite declines in physical capabilities). Hopper and colleagues' (2023) large cross-sectional study from Canada ($n = 30,097$ aged between 45 and 85 years old) adds to the understanding of associations between socioeconomic position and mental health resilience (defined as “*the discrepancy between one's reported current mental health and one's predicted mental health based on their physical performance*”). Their results suggest that these links are stronger for people declaring low and middle values of household incomes than those with high values. However, financial status can be mediated by individual resources such as social networks (accounted for 16% of the association) and physical activity (accounted for 6%) (Hopper *et al.*, 2023).

This suggests that recommendations to reduce the inequalities between urban and non-urban areas

should extend to providing more support and services than purely increasing the ratio of healthcare workers. This could start with increasing services providing psychoeducation about lifestyle, including exercise and sleep, as well as general access to education. Similar conclusions are drawn by a study investigating the differences in white matter hyperintensities in socioeconomically deprived groups, suggesting that they are associated with health-related risk factors and should be targeted by prevention programs (Rodríguez *et al.*, 2023; $n = 1,185$).

Therefore, there are limits to addressing the health rural-urban gap by simply increasing the number of healthcare workforce in more remote areas. Rather, what is needed is for more interventions that address lifestyle factors, like facilitating more social activities and psychoeducation about the role of sleep and exercise. Governments and policymakers should also continue to implement long-term plans to reduce the inequalities in education and income levels, which should target the younger generations to reduce health inequalities while they age.

There are also some limitations to Yuan and colleagues' (2023) study. First, the sample size of people living in the rural areas was much greater than those from urban areas ($n = 7,907$ vs. $n = 3,338$). Second, the authors focused on depressive symptoms measured by the Center for Epidemiological Studies Depression Scale. These questions focus on depressive symptoms, loneliness, hopefulness, and sleep, with responses ranging from “0: all of the time” to “3: rarely or never” and a total score between 0 and 30. Despite its well-known high psychometric properties, the cutoff point of 10 does not indicate the clinical diagnosis of depression and does not consider how long the depressive symptoms last. In the future, it would be beneficial to conduct similar studies using more objective measures including a clinical diagnosis of depression. Third, the study was limited in that it focused on a specific set of modifiable factors and did not explore the contribution of other known, influential lifestyle factors, such as diet (Wu *et al.*, 2021).

In conclusion, Yuan and colleagues' (2023) article provides an important insight into the potentially modifiable risk factors for depression, which can help address inequalities in care among older people living in rural-urban areas. Factors like education, better sleep, and exercise can be relatively easily modified and could potentially reduce the levels of depressive symptoms among older people in rural/remote areas. Although other factors like income or living status could be more challenging to address in the short term, they still could be included in longer-term strategies reducing the disparities across different areas of the same

country. This study presents a great opportunity for further research exploring factors influencing other mental health symptoms (i.e. anxiety, cognitive problems) as they often coexist with depressive symptoms. Future work should pilot and evaluate the efficacy of some community-based interventions to reduce the differences and therefore decrease the risk of depressive symptoms in remote/rural populations.

Conflict of interest

None.

References

- Braveman, P.** (2014). What are health disparities and health equity? We need to be clear. *Public Health Reports*, 129, 5–8. <https://doi.org/10.1177/00333549141291S203>
- Cosco, T., Cooper, R., Kuh, D. and Stafford, M.** (2018). Socioeconomic inequalities in resilience and vulnerability among older adults: a population-based birth cohort analysis. *International Psychogeriatrics*, 30, 695–703. <https://doi.org/10.1017/S1041610217002198>
- Glauber, R.** (2022). Rural depopulation and the rural-urban gap in cognitive functioning among older adults. *The Journal of Rural Health*, 38, 696–704. <https://doi.org/10.1111/jrh.12650>
- Hopper, S., Best, J., Wister, A. and Cosco, T.** (2023). Contributors to mental health resilience in middle-aged and older adults: an analysis of the Canadian Longitudinal Study on Aging. *International Psychogeriatrics*, 1–10. <https://doi.org/10.1017/S1041610223000224>
- Rodriguez, F. et al.** (2023). Differences in white matter hyperintensities in socioeconomically deprived groups: results of the population-based LIFE Adult Study. *International Psychogeriatrics*, 1–14. <https://doi.org/10.1017/S104161022300025X>
- Sun, N., Hua, C. L., Qiu, X. and Brown, J. S.** (2022). Urban and rural differences in trajectories of 418 depressive symptoms in later life in the United States. *Journal of Applied Gerontology*, 41, 148–157. <https://doi.org/10.1177/0733464820972527>
- Wang, Y., Li, Z. and Fu, C.** (2021). Urban-Rural differences in the association between social activities and depressive symptoms among older adults in China: a cross-sectional study. *BMC Geriatrics*, 21, 569. <https://doi.org/10.1186/s12877-021-02541-y>
- Weden, M. M., Shih, R. A., Kabeto, M. U. and Langa, K. M.** (2018). Secular trends in dementia and cognitive impairment of US rural and urban older adults. *American Journal of Preventive Medicine*, 54, 164–172.
- World Health Organisation** (2010). Increasing access to health workers in remote and rural areas through improved retention. Global policy recommendations. Available at: <https://www.who.int/publications/i/item/increasing-access-to-health-workers-in-remote-and-rural-areas-through-improved-retention>; last accessed 05 July 2023.
- World Health Organization** (2015). Monitoring health inequality: an essential step for achieving health equity. World Health Organization. Available at: <http://apps.who.int/iris/handle/10665/164530>; last accessed 04 July 2023.
- Wu, P. Y., Chen, K. M. and Belcastro, F.** (2021). Dietary patterns and depression risk in older adults: systematic review and meta-analysis. *Nutrition Reviews*, 79, 976–987. <https://doi.org/10.1093/nutrit/nuaa118>
- Yuan, L. et al.** (2023). Decomposition and comparative analysis of differences in depressive symptoms between urban and rural older adults: evidence from a national survey. *International Psychogeriatrics*, 1–12. <https://doi.org/10.1017/S1041610223000078>
- Ziarko, M. et al.** (2015). Do urban and rural residents living in Poland differ in their ways of coping with chronic diseases? *European Review for Medical and Pharmacological Sciences*, 19, 4227–4234.