Index of Subjects

Age—sex distribution of various diseases with particular reference to toxoplasmic lymphadenopathy, 215

Agglutinin production in mice in the determination of the definitive serotype of Bordetella pertussis, 257

Antibacterial activity of chloroxylenol in combination with EDTA, 11

Antibiotics and the Aberdeen typhoid outbreak in 1964, 191

Antiseptics used for hand disinfection in wards, 75

Bacillus cereus food-poisoning in Finland associated with boiled rice, 319

Bacterial dispersion in relation to operating room clothing, 367

Bacteriological evaluation of a mobile laminar cross-flow unit for surgery, under laboratory circumstances, 1

Bacteriological quality control, preparation of simulated water samples for, 49

Bacteriological survey of washrooms and toilets, 183

Bedpans, the cleaning and disinfection of by heat, 341

Bioassay results, point and interval estimation in the combination of, 147

Bordetella pertussis, the importance of agglutinin production in mice in the determination of the definitive serotype, 257

Bordetella pertussis, serotypes of, isolated in Great Britain between 1941 and 1968 and a comparison with the serotypes observed in other countries during this period, 265

Bordetella pertussis, the stability of the serotypes of, with particular reference to serotype 1,2,3,4,277

Bovine brucellosis, serological tests for: sensitivity of to different antibody classes, 287

Brucella abortus agglutinins in whey, the effect of chloroform on, 425

Brucella abortus, the immunogenic activity of ribosomal fractions derived from, 65

Characteristics of a virus causing pox disease in sheep and goats in Kenya, with observations on the epidemiology and control, 163

Chlamydia A (TRIC agents), the sensitivity of replicating McCoy cells in the isolation and growth of, 441

Chloroform, the effect of on Brucella abortus agglutinins in whey, 425

Chloroxylenol, antibacterial activity in combination with EDTA, 11

Chocolate, survival of Salmonella in, 41

Cleaning and disinfection by heat of bedpans in automatic and semi-automatic machines, 341

Clothing in the operating room, its relation to bacterial dispersion, 367

Congenital malformations in India, role of rubella in, 33

Cow-pats, the serotypes of $Escherichia\ coli$ from, compared with serotypes from human sources, 403

Cross-infection in isolation wards of different design, 23

Cytomegalovirus, porcine, in commercial pig herds, 125

Dairy creamery effluent sludge, isolation of mycobacteria from, 407

Disinfection of hands in wards, evaluation of antiseptics used for, 75

Echovirus 19, epidemic of, in the north-east of England, 307

Enteric fever in Scotland, 1967-1974, 83

Environmental contamination in a microbiological laboratory, 91

Epidemic of echovirus 19 in the north-east of England, 307

Epidemiology and control of a virus causing pox-disease in sheep and goats in Kenya, 163 Escherichia coli serotypes from cow-pats, compared with those from human sources, 403

Ethylenediaminetetra-acetic acid, anti-bacterial activity of chloroxylenol in combination with, 11

Food-poisoning in Finland by *Bacillus cereus*, associated with boiled rice, 319 Foot-and-mouth disease virus, growth of in respiratory tract of cattle, 467

Foot-and-mouth disease in Kenya, the role of sheep and goats in the epizootiology of, 395 Frozen broiler chickens, estimation of the hygienic condition of, 57

Goats and sheep, the role of in the epizootiology of foot-and-mouth disease in Kenya, 395

Hamster as a secondary reservoir host of lymphocytic choriomeningitis virus, 299 Hand disinfection in wards, evaluation of antiseptics used for, 75

HBV infections in Athens, Greece, sero-epidemiological study of, 229

Infected patients, a source isolator for, 355

Influenza vaccine, WRL 105 strain live attenuated, used for comparing four methods of intranasal vaccination, 459

Iodophor disinfectants, the laboratory evaluation of with yeast suspensions, 349 Isolation wards of different design, cross-infection in, 23

Laboratory bench, isolation of salmonellas and Shigella sonnei from, 337

Laminar cross-flow unit for surgery, bacteriological evaluation of, 1

Low temperature denaturation injury in T4Bo phage, prevention of by low concentrations of traditional cryoprotective additives, 453

Lymphocytic choriomeningitis virus, the hamster as a secondary reservoir host of, 299

Major gene controlling warfarin-resistance in the house mouse, 173

Marmosets and tamarins in the laboratory, an outbreak of shigellosis in, 415

McCoy cells, factors affecting the sensitivity of in the isolation and growth of chlamydia (TRIC agents), 441

Meat products, methods for detection of vegetable proteins in, 329

Mouse or man? Which are pertussis vaccines to protect? 249

Mycobacteria, the effect of contact with on the pattern of sensitivity to a range of new Tuberculins amongst Ugandan adults, 205

Mycobacteria, isolation of from dairy creamery effluent sludge, 407

Oral poliovaccine, response to in persons aged 16-18 years, 235

Pertussis vaccines, to protect mouse or man? 249

Phage T4Bo, prevention of low temperature denaturation injury in by low concentrations of traditional cryoprotective additives, 453

Point and interval estimation in the combination of bioassay results, 147

Polar isolation, studies of experimental rhinovirus type 2 infections in, compared with in England, 379

Porcine cytomegalovirus in commercial pig herds, 125

Pseudomonas, a new vaccine against: preliminary trial on human volunteers, 429

Response to oral poliovaccine in persons aged 16-18 years, 235

Rhinoviruses in Britain 1963-1973, 137

Rhinovirus type 2 infections studied experimentally in polar isolation and in England, 379

Ribosomal fractions derived from Brucella abortus, the immunogenic activity of, 65

Rice, boiled, associated with an outbreak of Bacillus cereus food-poisoning in Finland, 319

Rubella, role of in congenital malformations in India, 33

Rubella syndrome (congenital), specific immunoglobins in infants with, 109

 $Salmonella\ east bourne\ {\it and}\ S.\ typhimurium\ {\it in\ chocolate},\ 41$

Salmonella, the virulence of trimethoprim-resistant thymine-requiring strains of, 97

Salmonellas and Shigella sonnei isolated from a laboratory bench, 337

Seroepidemiological study of HBV infections in Athens, Greece, 229

Serological tests for bovine brucellosis, a quantitative comparison of the sensitivity of to different antibody classes, 287

Serotypes of Bordetella pertussis isolated in Great Britain between 1941 and 1968 and a comparison with serotypes observed in other countries during the period, 265
Sheep and goats, the role of in the epizootiology of foot-and-mouth disease in Kenya, 395
Shigella sonnei and salmonellas isolated from a laboratory bench, 337
Shigellosis in laboratory marmosets and tamarins, 415

Stability of the serotypes of *Bordetella pertussis* with particular reference to serotype 1,2,3,4, 277

Toxoplasmic lymphadenopathy, age–sex distribution of various diseases with particular reference to, 215

Tuberculins, new, sensitivity of Uganda adults, to 205

Typhoid outbreak in Aberdeen, 1964, the use of antibiotics in, 191

Vaccination, intranasal, a comparison of four methods in a clinical trial of WRL 105 strain live attenuated influenza vaccine, 459

Vaccine against Pseudomonas. Preliminary trial on human volunteers, 429

Vegetable proteins in meat products, extraction methods and test techniques for their detection, 329

Virulence of trimethoprim-resistant thymine-requiring strains of Salmonella, 97

Viruses: echo, 307; foot-and-mouth, 395, 467; hepatitis B, 229; lymphocytic choriomeningitis, 299; porcine cytomegalo, 125; pox, 163; rhino, 137, 379; rubella, 33, 109.

Warfarin-resistance in the house mouse, a major gene controlling, 173 Washrooms and toilets, a bacteriological survey of, 183 Water samples, simulated, for bacteriological quality control, 49

Yeast suspensions used in the laboratory evaluation of iodophor disinfectants, 349