

ON MAHLER'S CLASSIFICATION OF p -ADIC NUMBERS

YANN BUGEAUD[✉] and GÜLCAN KEKEÇ

(Received 21 April 2018; accepted 14 May 2018; first published online 30 July 2018)

Abstract

We give transcendence measures for p -adic numbers ξ , having good rational (respectively, integer) approximations, that force them to be either p -adic S -numbers or p -adic T -numbers.

2010 *Mathematics subject classification*: primary 11J61; secondary 11J82.

Keywords and phrases: Mahler's classification of p -adic numbers, p -adic S -number, p -adic T -number, transcendence measure.

1. Introduction

In 1955, Roth [12] proved that irrational real algebraic numbers cannot be approximable by rational numbers at an order greater than 2.

THEOREM 1.1 (Roth [12], 1955). *Let ξ be a real number and ε be a positive real number. Suppose that there exists a sequence $(p_n/q_n)_{n=1}^{\infty}$ of rational numbers such that $2 \leq q_1 < q_2 < \dots$ and*

$$0 < \left| \xi - \frac{p_n}{q_n} \right| < q_n^{-2-\varepsilon} \quad (n = 1, 2, \dots).$$

Then ξ is transcendental.

In 1964, under an additional assumption on the growth of the sequence $(q_n)_{n \geq 1}$ in Theorem 1.1, Baker [2] obtained a more precise conclusion than the simple transcendence of ξ . Before stating his result, we shall recall the classifications of transcendental real numbers defined by Mahler [9] in 1932 and by Koksma [7] in 1939. Let d be a positive rational integer and ξ a real number. Then $w_d(\xi)$ is defined as the supremum of the real numbers w_d for which there exist infinitely many polynomials $P(X)$ with rational integral coefficients and of degree at most d satisfying the inequalities

$$0 < |P(\xi)| \leq H(P)^{-w_d},$$

This work is supported by the Scientific Research Projects Coordination Unit of Istanbul University, project number IRP-52249.

© 2018 Australian Mathematical Publishing Association Inc.

where $H(P)$ denotes the height of the polynomial $P(X)$, that is, $H(P)$ is the maximum of the absolute values of the coefficients of $P(X)$. On the other hand, $w_d^*(\xi)$ is defined as the supremum of the real numbers w_d^* for which there exist infinitely many real algebraic numbers α of degree at most d satisfying the inequalities

$$0 < |\xi - \alpha| \leq H(\alpha)^{-w_d^*-1},$$

where $H(\alpha)$ denotes the height of α , that is, $H(\alpha)$ is the height of the minimal polynomial of α over \mathbb{Z} . Setting $w(\xi) = \limsup_{d \rightarrow \infty} (w_d(\xi)/d)$, Mahler [9] called ξ :

- an S -number if $0 < w(\xi) < \infty$;
- a T -number if $w(\xi) = \infty$ and $w_d(\xi) < \infty$ for all positive rational integers d ;
- a U -number if $w(\xi) = \infty$ and $w_d(\xi) = \infty$ from some d onward.

Exactly in the same manner, setting $w^*(\xi) = \limsup_{d \rightarrow \infty} (w_d^*(\xi)/d)$, and using $w^*(\xi)$ and $w_d^*(\xi)$ instead of $w(\xi)$ and $w_d(\xi)$, Koksma [7] defined the classes of S^* -, T^* - and U^* -numbers and proved that they coincide with those of S -, T - and U -numbers, respectively. Thus, the real transcendental numbers are divided into three disjoint classes. (See Bugeaud [3] for details of the classifications of Mahler and Koksma.) Now we can state the result of Baker.

THEOREM 1.2 (Baker [2], 1964). *Let ξ be a real number and ε be a positive real number. Suppose that there exists a sequence $(p_n/q_n)_{n=1}^\infty$ of rational numbers with $\gcd(p_n, q_n) = 1$ ($n = 1, 2, \dots$) such that $2 \leq q_1 < q_2 < \dots$ and*

$$0 < \left| \xi - \frac{p_n}{q_n} \right| < q_n^{-2-\varepsilon} \quad (n = 1, 2, \dots).$$

If

$$\limsup_{n \rightarrow \infty} \frac{\log q_{n+1}}{\log q_n} < \infty,$$

then there exists a real number c , depending only on ξ and ε , such that

$$w_d^*(\xi) \leq \exp \exp \{cd^2\} \quad (d = 1, 2, \dots).$$

In particular, ξ is either an S -number or a T -number.

Recently, in 2010, Adamczewski and Bugeaud [1, Théorème 3.1] gave a new proof of Theorem 1.2, based on a new application (Théorème EL in [1]) of a quantitative version of Theorem 1.1 obtained by Evertse [6] and Locher [8]. Moreover, the result of Adamczewski and Bugeaud [1, Théorème 3.1] improves the transcendence measure given by Theorem 1.2. In Theorem 1.4, we give a p -adic analogue of Theorem 1.2 by following the method of proof of Théorème 3.1 in [1]. Before stating our new result, we recall the classifications of p -adic numbers in analogy with the classifications of Mahler and Koksma of real numbers.

Throughout the present paper, p denotes a fixed prime number and $|\cdot|_p$ denotes the p -adic absolute value on the field \mathbb{Q} of rational numbers, normalised such that $|p|_p = p^{-1}$. We also denote the unique extension of $|\cdot|_p$ to the field \mathbb{Q}_p of p -adic numbers, the completion of \mathbb{Q} with respect to $|\cdot|_p$, by the same notation $|\cdot|_p$.

In 1958, Ridout [11] proved the p -adic analogue of Theorem 1.1. For coprime nonzero integers x, y , write $|x, y|$ for the maximum of $|x|$ and $|y|$, that is, for the height of the rational number x/y .

THEOREM 1.3 (Ridout [11], 1958). *Let ξ be a p -adic number and ε be a positive real number. Suppose that there exists a sequence $(x_n/y_n)_{n=1}^\infty$ of rational numbers with $\gcd(x_n, y_n) = 1$ ($n = 1, 2, \dots$) such that $2 \leq |x_1, y_1| < |x_2, y_2| < \dots$ and*

$$0 < \left| \xi - \frac{x_n}{y_n} \right|_p < |x_n, y_n|^{-2-\varepsilon} \quad (n = 1, 2, \dots).$$

Then ξ is transcendental.

Unlike the real case, we cannot replace $|x_n, y_n|$ by $|y_n|$ in the statement of Theorem 1.3 since, for any irrational p -adic number ξ and any positive real number ε , there exists an integer x such that $|\xi - x/y|_p$ is less than ε .

Mahler [10], in 1935, proposed a classification of transcendental p -adic numbers in analogy with his classification of transcendental real numbers. Let d be a positive rational integer and ξ be a transcendental p -adic number. Then $w_d(\xi)$ is defined as the upper limit of the real numbers w_d for which there exist infinitely many polynomials $P(X)$ with rational integral coefficients and of degree at most d satisfying the inequalities

$$0 < |P(\xi)|_p \leq H(P)^{-w_d-1}.$$

Setting $w(\xi) = \limsup_{d \rightarrow \infty} (w_d(\xi)/d)$, we then call ξ :

- a p -adic S -number if $0 < w(\xi) < \infty$;
- a p -adic T -number if $w(\xi) = \infty$ and $w_d(\xi) < \infty$ for all positive rational integers d ;
- a p -adic U -number if $w(\xi) = \infty$ and $w_d(\xi) = \infty$ from some d onward.

The p -adic transcendental numbers are divided into the three disjoint classes S, T and U . (See Bugeaud [3] for more information about Mahler's classification in \mathbb{Q}_p .) On the other hand, in analogy with Koksma's classification of real numbers, we define $w_d^*(\xi)$ as the upper limit of the real numbers w_d^* for which there exist infinitely many p -adic algebraic numbers α of degree at most d satisfying the inequalities

$$0 < |\xi - \alpha|_p \leq H(\alpha)^{-w_d^*-1}.$$

Setting $w^*(\xi) = \limsup_{d \rightarrow \infty} (w_d^*(\xi)/d)$, we define the p -adic S^* -numbers, T^* -numbers and U^* -numbers, respectively, exactly as in the real case. Again, the classes S, T and U are the same as the classes S^*, T^* and U^* , respectively. (See Bugeaud [3] and Schlickewei [13].)

A first goal of the present paper is to establish the following p -adic analogue of Baker's Theorem 1.2.

THEOREM 1.4. *Let ξ be a p -adic number and ε be a positive real number. Suppose that there exists a sequence $(x_n/y_n)_{n=1}^\infty$ of rational numbers with $\gcd(x_n, y_n) = 1$ (for $n = 1, 2, \dots$) such that $2 \leq |x_1, y_1| < |x_2, y_2| < \dots$ and*

$$0 < \left| \xi - \frac{x_n}{y_n} \right|_p < |x_n, y_n|^{-2-\varepsilon} \quad (n = 1, 2, \dots). \tag{1.1}$$

If

$$\limsup_{n \rightarrow \infty} \frac{\log |x_{n+1}, y_{n+1}|}{\log |x_n, y_n|} < \infty, \tag{1.2}$$

then ξ is transcendental and there exists a real number c , depending only on ξ and ε , such that

$$w_d^*(\xi) \leq (2d)^{c \log \log 3d} \quad (d = 1, 2, \dots).$$

In particular, ξ is either a p -adic S -number or a p -adic T -number.

We point out that the bound on $w_d^*(\xi)$ in Theorem 1.4 does not depend on p . It has the same shape as the bound obtained in the real case in [1, Theorem 3.1]. The same remarks hold for our next result.

THEOREM 1.5. *Let ξ be a p -adic number and let $\xi = \sum_{k \geq -k_0} a_k p^k = \sum_{j \geq 1} a_{n_j} p^{n_j}$ denote its Hensel expansion, where $k_0 \geq 0$, $a_{-k_0} \neq 0$ if $k_0 > 0$, $(n_j)_{j \geq 1}$ is strictly increasing and a_{n_j} is in $\{1, \dots, p - 1\}$ for $j \geq 1$. Let ε be a positive real number. Suppose that there exists an increasing sequence $\mathcal{J} = (j_k)_{k \geq 1}$ of positive integers such that $n_{j_{k+1}} \geq (1 + \varepsilon)n_j$ for j in \mathcal{J} and*

$$\limsup_{k \rightarrow \infty} \frac{j_{k+1}}{j_k} < \infty.$$

Then ξ is transcendental and there exists a real number c , depending only on ξ and ε , such that

$$w_d^*(\xi) \leq (2d)^{c \log \log 3d} \quad (d = 1, 2, \dots).$$

In particular, ξ is either a p -adic S -number or a p -adic T -number.

We display a straightforward application of Theorem 1.5.

COROLLARY 1.6. *For any real number $c > 1$, the p -adic number $\sum_{i=0}^{+\infty} p^{\lfloor c^i \rfloor}$ is either an S -number or a T -number.*

The proofs of Theorems 1.4 and 1.5 follow a method introduced in [1] and depend on a quantitative version of Ridout’s theorem given in Theorem 2.1 in the next section.

We take this opportunity to give, in addition, an application of Theorem 2.1 to the number of digit changes in the Hensel expansion of irrational algebraic p -adic numbers, thereby improving [4, Theorem 2].

Let ξ be a p -adic number and denote by

$$\xi = \sum_{k=-k_0}^{+\infty} a_k p^k \quad (a_k \in \{0, 1, \dots, p - 1\}, k_0 \geq 0, a_{-k_0} \neq 0 \text{ if } k_0 > 0)$$

its Hensel expansion. For a positive integer n , set

$$\text{nbdc}(n, \xi, p) = \text{Card}\{1 \leq k \leq n : a_k \neq a_{k+1}\}$$

and

$$\mathcal{S}(n, \xi, p) = \sum_{k=1}^n a_k.$$

THEOREM 1.7. *Let p be a prime number. Let ξ be an algebraic irrational number in \mathbb{Q}_p . For any positive real number δ with $\delta < 1/2$ and any sufficiently large integer n ,*

$$\text{nbd}c(n, \xi, p) \geq (\log n)^{1+\delta},$$

there are at least $(\log n)^{1+\delta}$ nonzero digits among the first n digits of the Hensel expansion of ξ and, moreover,

$$(\log n)^{1+\delta} \leq \mathcal{S}(n, \xi, p) \leq n(p - 1) - (\log n)^{1+\delta}.$$

The proof of Theorem 1.7 follows the same lines as that of [5, Theorem 3.1]. We omit the details. The good approximations to ξ are obtained by truncating its Hensel expansion and repeating the last digit. They are rational numbers, whose denominators divide $p - 1$, and we apply Theorem 2.1 to $(p - 1)\xi$.

COROLLARY 1.8. *For all real numbers $c > 1$ and $\eta > 2/3$, the p -adic number*

$$\sum_{j \geq 1} p^{\lfloor c^j \rfloor}$$

is transcendental.

2. Auxiliary result

The following theorem is a consequence of [5, Proposition A.1] and is the key point in the proof of Theorems 1.4 and 1.5. It can be regarded as a p -adic analogue of Théorème EL in [1].

THEOREM 2.1 (Bugeaud and Evertse [5], 2008). *Let α be a p -adic algebraic number of degree d and height H , and let ε be a positive real number. Then the inequality*

$$\left| \alpha - \frac{x}{y} \right|_p \leq |x, y|^{-2-\varepsilon}$$

has at most

$$2^{26}(1 + 2/\varepsilon)^3 \log(2d + 4) \log((1 + 2/\varepsilon) \log(2d + 4))$$

solutions $(x, y) \in \mathbb{Z}^2$ with $\text{gcd}(x, y) = 1$ and

$$|x, y| \geq \max((2\sqrt{d+1}H)^{1/((d+1)(d+2))}, 4^{2/\varepsilon}).$$

Likewise, the inequality

$$|\alpha - x|_p \leq |x|^{-1-\varepsilon}$$

has at most

$$2^{26}(1 + 1/\varepsilon)^3 \log(2d + 4) \log((1 + 1/\varepsilon) \log(2d + 4))$$

solutions $x \in \mathbb{Z}$ with

$$|x| \geq \max((2\sqrt{d+1}H)^{1/((d+1)(d+2))}, 4^{1/\varepsilon}).$$

Theorem 2.1 follows from [5, Proposition A.1] in the same way as [5, Corollary 5.2] eventually follows from [5, Proposition A.1]. We omit the details.

3. Proof of Theorem 1.4

Let all the hypotheses of Theorem 1.4 be satisfied. It follows from Theorem 1.3 that ξ is a p -adic transcendental number. By (1.2), there exists a real number $c > 1$ such that

$$|x_n, y_n| < |x_{n+1}, y_{n+1}| \leq |x_n, y_n|^c \quad (n = 1, 2, \dots). \tag{3.1}$$

Let d be a positive rational integer and α be a p -adic algebraic number of degree d . We choose α with sufficiently large height $H(\alpha)$ so that

$$|x_1, y_1| \leq (2\sqrt{d+1}H(\alpha))^{1/((d+1)(d+2))}$$

is satisfied. Let $j \geq 2$ be the unique integer satisfying

$$|x_{j-1}, y_{j-1}| \leq (2\sqrt{d+1}H(\alpha))^{1/((d+1)(d+2))} < |x_j, y_j|. \tag{3.2}$$

We suppose that we choose α with sufficiently large height $H(\alpha)$ so that the inequality

$$|x_j, y_j| > \max(4^{2/\varepsilon}, 4^c(d+1)^c) \tag{3.3}$$

is satisfied. Let χ be the real number defined by

$$|\xi - \alpha|_p = H(\alpha)^{-\chi}.$$

We suppose that $\chi > 1$ and we will bound χ from above.

Let K be the largest of the positive integers h satisfying $|x_{j+h}, y_{j+h}|^{2+\varepsilon} < H(\alpha)^\chi$. Then

$$|\xi - \alpha|_p = H(\alpha)^{-\chi} < |x_{j+h}, y_{j+h}|^{-2-\varepsilon} \quad (h = 1, 2, \dots, K). \tag{3.4}$$

By (1.1) and (3.4),

$$\left| \alpha - \frac{x_{j+h}}{y_{j+h}} \right|_p \leq \max \left(\left| \xi - \frac{x_{j+h}}{y_{j+h}} \right|_p, |\xi - \alpha|_p \right) < |x_{j+h}, y_{j+h}|^{-2-\varepsilon}$$

for $h = 1, 2, \dots, K$. As a result, the inequality

$$\left| \alpha - \frac{x}{y} \right|_p < |x, y|^{-2-\varepsilon}$$

has at least K rational solutions x/y with $\gcd(x, y) = 1$ and $|x, y| > |x_j, y_j|$. Hence, by (3.2), (3.3) and Theorem 2.1,

$$K \leq 2^{26}(1 + 2/\varepsilon)^3 \log(2d + 4) \log((1 + 2/\varepsilon) \log(2d + 4)). \tag{3.5}$$

On the other hand, by $\chi > 1$ and the choice of K , the inequalities (3.1), (3.2) and (3.3) imply that

$$|x_j, y_j|^{(2+\varepsilon)c^{K+1}} \geq |x_{j+K+1}, y_{j+K+1}|^{2+\varepsilon} \geq H(\alpha)^\chi \geq 2^{-\chi}(d+1)^{-\chi/2} |x_j, y_j|^{\chi/c} \geq |x_j, y_j|^{\chi/(2c)}$$

and so

$$\chi \leq 2(2 + \varepsilon)c^{K+2}. \tag{3.6}$$

It follows from (3.5) and (3.6) that there exists a real number c' , depending only on ξ and ε , such that

$$w_d^*(\xi) \leq (2d)^{c' \log \log 3d} \quad (d = 1, 2, \dots).$$

Then ξ is either a p -adic S -number or a p -adic T -number. This completes the proof of Theorem 1.4.

4. Proof of Theorem 1.5

Let all the hypotheses of Theorem 1.5 be satisfied. It follows from Theorem 2.1 that ξ is a p -adic transcendental number. For $J \geq 1$, set $\xi_J := \sum_{j=1}^J a_n_j p^{n_j}$. By assumption, there exists a real number $c > 1$ such that

$$\xi_j < \xi_{j+1} \leq \xi_j^c \quad (j = 1, 2, \dots). \tag{4.1}$$

Without loss of generality, we assume that $k_0 = 0$. Consequently,

$$1 \leq \xi_j < p^{1+n_j} \quad (j = 1, 2, \dots)$$

and there exists j_0 such that

$$|\xi - \xi_j|_p = p^{-n_{j+1}} < \xi_j^{-n_{j+1}/(1+n_j)} < \xi_j^{-1-\varepsilon/2} \quad (j \geq j_0). \tag{4.2}$$

Let d be a positive rational integer and α be a p -adic algebraic number of degree d . We choose α with sufficiently large height $H(\alpha)$ so that

$$\xi_1 \leq (2\sqrt{d+1}H(\alpha))^{1/((d+1)(d+2))}$$

is satisfied. Let $j \geq 2$ be the unique integer satisfying

$$\xi_{j-1} \leq (2\sqrt{d+1}H(\alpha))^{1/((d+1)(d+2))} < \xi_j. \tag{4.3}$$

We suppose that we choose α with sufficiently large height $H(\alpha)$ so that the inequality

$$\xi_j > \max(4^{2/\varepsilon}, 4^c(d+1)^c) \tag{4.4}$$

is satisfied. Let χ be the real number defined by

$$|\xi - \alpha|_p = H(\alpha)^{-\chi}.$$

We suppose that $\chi > 1$ and we will bound χ from above.

Let K be the largest of the positive integers h satisfying $\xi_{j+h}^{1+\varepsilon/2} < H(\alpha)^\chi$. Then

$$|\xi - \alpha|_p = H(\alpha)^{-\chi} < \xi_{j+h}^{-1-\varepsilon/2} \quad (h = 1, 2, \dots, K). \tag{4.5}$$

By (4.2) and (4.5),

$$|\alpha - \xi_{j+h}|_p \leq \max(|\xi - \xi_{j+h}|_p, |\xi - \alpha|_p) < \xi_{j+h}^{-1-\varepsilon/2}$$

for $h = 1, 2, \dots, K$. As a result, the inequality

$$|\alpha - x|_p < x^{-1-\varepsilon/2}$$

has at least K solutions in positive integers x with $x > \xi_j$. Hence, by (4.3), (4.4) and Theorem 2.1,

$$K \leq 2^{26}(1 + 2/\varepsilon)^3 \log(2d + 4) \log((1 + 2/\varepsilon) \log(2d + 4)). \tag{4.6}$$

On the other hand, by $\chi > 1$ and the choice of K , the inequalities (4.1), (4.3) and (4.4) imply that

$$\xi_j^{(1+\varepsilon/2)c^{K+1}} \geq \xi_{j+K+1}^{1+\varepsilon/2} \geq H(\alpha)^\chi \geq 2^{-\chi}(d+1)^{-\chi/2} \xi_j^{\chi/c} \geq \xi_j^{\chi/(2c)}$$

and so

$$\chi \leq 2(1 + \varepsilon/2)c^{K+2}. \quad (4.7)$$

It follows from (4.6) and (4.7) that there exists a real number c' , depending only on ξ and ε , such that

$$w_d^*(\xi) \leq (2d)^{c' \log \log 3d} \quad (d = 1, 2, \dots).$$

Then ξ is either a p -adic S -number or a p -adic T -number. This completes the proof of Theorem 1.5.

Acknowledgements

We would like to thank Jan-Hendrik Evertse for his valuable advice and the referee for a careful reading.

References

- [1] B. Adamczewski and Y. Bugeaud, 'Mesures de transcendance et aspects quantitatifs de la méthode de Thue–Siegel–Roth–Schmidt', *Proc. Lond. Math. Soc.* **101** (2010), 1–26.
- [2] A. Baker, 'On Mahler's classification of transcendental numbers', *Acta Math.* **111** (1964), 97–120.
- [3] Y. Bugeaud, *Approximation by Algebraic Numbers*, Cambridge Tracts in Mathematics, 160 (Cambridge University Press, Cambridge, 2004).
- [4] Y. Bugeaud, 'On the b -ary expansion of an algebraic number', *Rend. Semin. Mat. Univ. Padova* **118** (2007), 217–233.
- [5] Y. Bugeaud and J.-H. Evertse, 'On two notions of complexity of algebraic numbers', *Acta Arith.* **133** (2008), 221–250.
- [6] J.-H. Evertse, 'The number of algebraic numbers of given degree approximating a given algebraic number', in: *Analytic Number Theory (Kyoto, 1996)*, London Mathematical Society Lecture Note Series, 247 (Cambridge University Press, Cambridge, 1997), 53–83.
- [7] J. F. Koksma, 'Über die Mahlersche Klasseneinteilung der transzendenten Zahlen und die Approximation komplexer Zahlen durch algebraische Zahlen', *Monatsh. Math. Phys.* **48** (1939), 176–189.
- [8] H. Locher, 'On the number of good approximations of algebraic numbers by algebraic numbers of bounded degree', *Acta Arith.* **89** (1999), 97–122.
- [9] K. Mahler, 'Zur Approximation der Exponentialfunktionen und des Logarithmus. I, II', *J. reine angew. Math.* **166** (1932), 118–150.
- [10] K. Mahler, 'Über eine Klasseneinteilung der p -adischen Zahlen', *Mathematica (Leiden)* **3** (1935), 177–185.
- [11] D. Ridout, 'The p -adic generalization of the Thue–Siegel–Roth theorem', *Mathematika* **5** (1958), 40–48.
- [12] K. F. Roth, 'Rational approximations to algebraic numbers', *Mathematika* **2** (1955), 1–20; corrigendum, 168.
- [13] H. P. Schlickewei, ' p -adic T -numbers do exist', *Acta Arith.* **39** (1981), 181–191.

YANN BUGEAUD, Université de Strasbourg, CNRS, IRMA UMR 7501,
7, rue René Descartes, 67000 Strasbourg, France
e-mail: bugeaud@math.unistra.fr

GÜLCAN KEKEÇ, Department of Mathematics, Faculty of Science,
Istanbul University, 34134, Vezneciler-Istanbul, Turkey
e-mail: gulkecec@istanbul.edu.tr