

when they come in contact with warm wash solution, the processor must allow time for the eggs to warm before processing. The result would be that eggs, like every other raw food of animal origin, are held under strict time-temperature control to ensure a safe, wholesome product reaches the consumer.

In summary, *S enteritidis* in poultry does not appear to be a problem that is a function of high production, controlled environment husbandry methods, but is an evolving problem throughout the industry, complicated by transovarian transmission.

Accordingly, institutional food service operations, such as hospitals, should recognize the special susceptibility to their patients and adhere to published guidelines of "Safe Handling of Eggs in Quantity." Four key guidelines from this document are:

- Avoid serving raw eggs and foods containing raw eggs; institutionally prepared Caesar salad and Hollandaise sauce, for example. Products such as homemade ice cream, homemade eggnog and homemade mayonnaise also should be avoided, but commercial forms of these products are safe to serve since they are made with pasteurized eggs.

- Cook eggs thoroughly until both the yolk and white are firm, not runny, in order to kill any bacteria that may be present. There may be some risk in serving eggs lightly cooked; soft-cooked, soft-scrambled or sunny-side-up, for example.

- Realize that serving lightly cooked foods containing eggs, such as soft custards, meringues and french toast, may be particularly risky for people with weakened immune systems and other high-risk groups.

- Review recipes and food-handling practices to consider using pasteurized egg products instead of shell eggs whenever possible.

Russell W. Currier, DVM  
Susan C. Trock, DVM

#### REFERENCES

1. Hargrett-Bean NJ, Pavia AT, Tauxe RV: *Salmonella* isolates from humans in the United States. *MMWR* 1988; SS-2:25-31.
2. St. Louis ME, Morse DL, Potter ME, et al: The emergence of grade A eggs as a major source of *Salmonella enteritidis* infections. *JAMA* 1988; 259:2103-2107.
3. British ad warns against using uncooked eggs. *Food Chemical News* 1988; 43:47-48.
4. Incidence of *Salmonella*-contaminated eggs reported low. *Food Chemical News* 1988; 12:6-7.
5. Conner MW, Crouch EA, Green LC, et al: Assessment of safety of grade A shell eggs. In a risk assessment sponsored by the Egg Nutrition Center, Washington, DC, 1988.
6. AMS to announce model program for egg testing. *Food Chemical News* 1988; 26:36.
7. Baker RC, Hogartv S, Poon W, et al: Survival of *Salmonella typhimurium* and *Staphylococcus aureus* in eggs cooked by different methods. *Poultry Science* 1983; 62:1211-1216.
8. Code of Federal Regulations; Agriculture. Federal Register 1987; 56(Jan 1):133-134.
9. Safe handling of eggs in quantity. *Food Service Institutions Bulletin*, Food and Drug Administration; September 1988: 1.

## Diamidine Use in Treatment of *Pneumocystis carinii*

#### To the Editor:

The reintroduction of pentamidine into therapy for pneumocystosis was preceded by an unusual chapter in pharmacologic investigation. The basic studies formed a romantic search for a compound that would influence the metabolism of glucose, an essential metabolite of trypanosomes. In 1935, an investigation of hypoglycemic-producing guanidine derivatives was initiated. Synthalin (decamethylene diguanidine hydrochloride) was determined to have trypanocidal activity, an action later found not due to hypoglycemia. Subsequently, an exploration of compounds in which aromatic carrier chains were substituted for the alkyl chains and a shift to terminal amidine groups was made. The fundamental chemotherapeutic constituent was determined to be the aromatic diamidine group. Oxygen, nitrogen and sulfur linkages were also introduced into the alkane chain. Four promising compounds were isolated (stilbamidine,

pentamidine, propamidine and phenamidine). The last three compounds contained an oxygen link in the alkane chain.

These aromatic diamidines have therapeutic activity in human trypanosomiasis, leishmaniasis and gram-positive bacterial infections. Although stilbamidine seemingly was the most effective drug, its neurotoxicity made pentamidine the agent of choice for such therapy. At a later time, stilbamidine was demonstrated to be satisfactory treatment for blastomycosis.<sup>4</sup> Advantage was taken of its neurotoxicity in the treatment of tic douloureux.<sup>5</sup> The use of the diamidines for such purposes is a far cry from the agent sought as a hypoglycemic drug for trypanosomiasis, and the story is a tribute to the many investigators who by perseverance and perhaps, serendipity, brought these compounds to modern medicine for the treatment of *Pneumocystis carinii*.

Joseph M. Miller, MD  
Baltimore, Maryland

#### REFERENCES

1. von Jancso II, von Jancso H: Chemotherapeutische Wirkung und Kohlehydratstoffwechsel. Die Heilwirkung von Guaninderivaten auf die Trypanosomeninfektion. *Zschr f. Immunitätsforsch. u exper Therap* 1935; 86: 1-30.
2. Lourie EM, Yorke W: Studies in chemotherapy. XVI. The trypanocidal action of synthalin. *Ann Trop Med Parasitol* 1937; 31:435-445.
3. Ashley JN, Barber IJ, Ewins AJ, et al: A chemotherapeutic comparison of the trypanocidal action of some aromatic diamidines. *J Chem Soc Pat-t I*, 1942; 20:103-116.
4. Schoenbach EB, Miller JM, Ginsberg AJ, et al: Systemic blastomycosis treated with stilbamidine (preliminary report). *JAMA* 1951; 146:1317-1318.
5. Smith GW, Miller JM: The relief of tic douloureux with stilbamidine. *Ann Intern Med* 1953; 38:335-338.

Letters to the Editor should be addressed to INFECTION CONTROL AND HOSPITAL EPIDEMIOLOGY Editorial Offices, CAI General Hospital, University of Iowa Hospitals and Clinics, Iowa City, IA 52242. All letters must be typed, double spaced, and may not exceed four pages nor include more than one figure or table. The editors reserve the right to edit for purposes of clarity or brevity.