

is a consultation service within the Division of Medicine and CL Psychiatry. Staffed by Medical Registrars and Mental Health Nurses, the collaboration provides a unique healthcare response to acute general wards. The BoC RRT has been implemented to address the rising number of incidences whereby staff and patient safety are compromised. Using evidence-based skills the team aimed to: respond to episodes of clinical agitation that require an internal security response, assist ward referrals by exploring biopsychosocial contributors to behaviour, develop individual patient support plans and review and reduce restrictive intervention practices.

Objectives: To determine if the rapid response model has influenced:

- The impact on staff/patient safety
- Frequency of emergency responses for aggression
- Frequency of restrictive intervention use

Methods: This project was approved as a quality assurance project (QA2022018). The patients within scope of the BoC RRT include inpatients in medical and surgical wards. It excludes patients in Emergency Departments, mental health units, outpatient clinics, and visitors. The evaluation of the pilot has used a PDSA (Plan, Do, Study, Act) cycle when implementing new improvements. A mixed methods approach explored the impact of the BoC RRT. Staff consultation will identify challenges in responding to scenarios whereby there is risk of harm to staff and patients. Staff feedback and the emergency response data was monitored.

Results: In 2021, there was approx. 720 code greys per month, requiring a security response. Since the implementation of BoC RRT, these numbers have reduced to 527. Reviewing restrictive intervention practices has identified areas for policy review and need for education. Staff consultation found that nurses were confident caring for those patients exhibiting clinical agitation associated with delirium and dementia. However, caring for people with mental health or substance use disorders were more challenging.

Conclusions: These interim results indicate that BoC RRT has been generally well received by clinical staff. The decline in code grey responses indicates that it is likely having a positive impact in early identification and management of clinical agitation for hospital inpatients. There is support for this response model to continue beyond the pilot phase and further area for research.

Disclosure of Interest: None Declared

EPP0561

Descriptive analysis of unfavorable mental health opinions of candidates for bariatric surgery

R. C. Pereira^{1*}, A. Delgado² and F. M. Ferreira²

¹Hospital Magalhães Lemos and ²Unidade Local de Saúde de Matosinhos, Porto, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.862

Introduction: In recent years, there has been an increase in the number of candidates for bariatric surgery and a decrease in psychiatric contraindications.

Objectives: We aim to make a descriptive evaluation of unfavorable feedback concerning mental health of the candidates for bariatric surgery of the Local Health Unit of Matosinhos (Porto, Portugal).

Methods: Descriptive analysis of unfavorable feedback of mental health of candidates for surgical treatment of obesity.

Results: From March 2017 to August 2022, the Mental Health Service of the Local Health Unit of Matosinhos issued 347 pre-

surgical feedback. In 63 cases the initial opinion issued was unfavorable: 11 cases due to a psychiatric contraindication (not meeting conditions for intervention) and 52 cases had a conditional opinion (requiring pre-surgical interventions in order to become eligible for the intervention). Regarding contraindications, these were due to alcohol use disorder (n=3), binge eating (n=3), intellectual development disorder (n=2), purgative behavior (n=1), psychotic disorder (n=1) and mood disturbance (n=1). In terms of conditional opinions, the issues mentioned were lack of motivation for surgery (n=22), psychopathology (n=20), doubts about informed consent (n=8) and need for multidisciplinary discussion/coordination (n= 7).

Conclusions: There was an increase in eligibility of candidates for surgical treatment as most of the initial unfavorable opinions were conditional. This could be explained by the decline of complications associated with bariatric surgery, but also because psychiatric disorders are now being viewed as treatable. Notably people with eating disorders are now fit for surgery after a medical or psychotherapeutic intervention.

Disclosure of Interest: None Declared

EPP0562

Delirium, Antipsychotics and Death in the time of COVID-19

R. Softic* and N. Becarevic

of psychiatry, University Clinical Center Tuzla, Tuzla, Bosnia and Herzegovina

*Corresponding author.

doi: 10.1192/j.eurpsy.2023.863

Introduction: Delirium is an acute, transient, global organic disorder of CNS functioning resulting in impaired consciousness, attention, and other cognitive functions. Causes of delirium are multifactorial and can be unrecognized in 2/3 of cases. It is recommended to use as few psychotropic medications as possible because many of them can worsen delirium. Antipsychotics are not recommended as a drug of first choice.

Objectives: To present rate of delirium regarding to treatment and outcome.

Methods: A retrospective observational study was conducted in the department of consultative psychiatry of the University Clinical Center Tuzla during the one-year period of the COVID-19 pandemic.

Results: 761 calls from different clinics of the University Clinical Center Tuzla were received in one year period. Delirium was diagnosed in 213 patients (28%). The total number of deaths was 147 (19.3%), the number of deaths in patients with delirium was 88 (41.3%). Antipsychotics were used in 137 (64%) patients with delirium. Death as an outcome was more common in patients treated with antipsychotics (64%) $p < 0.05$. The most used antipsychotic was Promazine 94 (44.1%). Number of deaths in patients with delirium treated with Promazine was 42 (44.7%) $p < 0.05$.

Conclusions: In patients with delirium mortality is significantly higher in those treated with antipsychotics, especially when treated with Promazine. The choice of antipsychotic medications should be made according to pharmacological properties and the clinical context.

Disclosure of Interest: None Declared