

Temporal patterns of loneliness and their associations with mental health outcomes:

Observations from a longitudinal study

Short title: Temporal patterns of loneliness and mental health

Błażej Misiak^{1*}, Paweł Liśkiewicz², Jerzy Samochowiec²

¹Department of Psychiatry, Wrocław Medical University, Wrocław, Poland

²Department of Psychiatry, Pomeranian Medical University, Szczecin, Poland

*Corresponding author:

Błażej Misiak, MD, PhD, Professor of Psychiatry, e-mail: blazej.misiak@umw.edu.pl

Department of Psychiatry, Wrocław Medical University, Pasteura 10 Street, 50-367 Wrocław,
Poland

This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI.

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

21

22

23

24

25 **Abstract**

26 **Background:** Loneliness is a global public health concern. Investigating loneliness in the
27 general population offers a greater generalizability over various levels of health-related
28 impairments, identification of at-risk individuals, detection of various loneliness severity
29 levels, and broader insights into social determinants. Previous studies have shown that
30 loneliness might be a transient or chronic experience depending on how consistently it is
31 reported across at least two timepoints. This study aimed to assess differential associations of
32 chronic and transient with various domains of psychopathology.

33 **Methods:** Participants were enrolled from the general population and assessed at two
34 timepoints spanning 6-7 months. Depressive symptoms, generalized anxiety, social anxiety,
35 and paranoid thoughts were measured using self-reports. The data were analysed using binary
36 logistic regressions.

37 **Results:** Altogether, 3,275 completed both assessments with the retention rate of 64.2%.
38 Chronic loneliness was associated with higher baseline and follow-up scores across all
39 symptom domains. The strongest association was observed for social anxiety. Transient
40 loneliness was not robustly associated with symptom scores. It was not significantly associated
41 with depressive symptoms (at neither of timepoints) and paranoid ideation (at baseline). The
42 strongest association was observed for generalized anxiety. Chronic loneliness, compared to
43 transient loneliness, was associated with significantly higher odds of social anxiety, depressive
44 symptoms, and paranoid ideation, but not generalized anxiety.

45 **Conclusions:** Both transient and chronic loneliness are associated with mental health
46 outcomes with the latter one showing generally stronger associations. Risk stratification and
47 early intervention among individuals experiencing loneliness might be needed to prevent the
48 development of more severe psychopathology.

49 **Keywords:** perceived social isolation; social disconnection; depression; anxiety disorder;
50 paranoia

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73 Introduction

74 The perception of social bonds has evolved significantly over time and was shaped by
75 cultural, social, and historical contexts [1]. Pre-modern perspectives posited social isolation
76 (currently conceptualized as an objective measure denoting that individual social bonds are
77 lower than average) as a choiceful or even esteemed experience associated with spiritual
78 reflection, creativity, or even philosophical inquiry [2, 3]. The term “loneliness”, understood
79 today as a painful self-perception of social disconnection or the discrepancy between actual
80 and desired social [4], was not commonly used or conceptualized by ancient and medieval
81 societies [3]. The concept of loneliness, as a distressing experience, emerged more
82 prominently in the 20th century and appeared to be influenced by societal shifts related to
83 industrialization, urbanization, and the rise of individualism [5]. According to the Bauman’s
84 “liquid modernity” concept, these processes are largely unpredictable and coincide with the
85 “virtualization of reality” manifesting in an ongoing digitalization of daily life and social
86 interactions [6, 7]. The 20th century brought a growing academic interest in the psychological
87 aspects of loneliness with early studies that began to link loneliness with depression and
88 anxiety [8, 9].

89 Currently, loneliness is perceived as a global public health concern due to high
90 prevalence rates and a negative impact on all aspects of health, wellbeing, and development
91 [10, 11]. Experiencing loneliness has been associated with 26% increased risk of mortality over
92 time [12]. Taking into account the mental health perspective, it is needed to note bidirectional

associations of loneliness with symptoms across a variety of mental health outcomes, including mood and anxiety disorders [13-15], psychosis [16, 17], substance use disorders [18, 19], and problematic internet use [20]. It is also important to note that contextual factors underlying the emergence of loneliness might show some interindividual variability [21]. For instance, factors related to social identity formation, social rejection, life transition, and digital overconnectivity might be of greater importance in younger populations [22-24]. In turn, contexts related to the loss of significant social connections, retirement, institutional care, physical illness and related disability might play important roles in older adults [25-27].

Studying loneliness in the general population offers several advantages over focusing on clinical samples. Although clinical samples provide valuable insights into loneliness among people with poor health status, population-based studies allow for a more comprehensive understanding of loneliness as a public health concern. Studies based on the general population include a wide range of participants providing the opportunity to apply the findings to the whole society. Given that these studies offer to study loneliness alongside a variety of demographic, social, behavioural, and environmental factors, potential risk and protective factors might be identified. Another benefit is related to the whole spectrum of loneliness severity and thus the opportunity for risk stratification. At this point, it is needed to note that population-based studies hold the potential to detect at-risk individuals who show a lower severity of loneliness. Therefore, it might be foreseen that these studies hold the potential to contribute to the development of early intervention strategies applicable to primary care. Finally, existing evidence indicates that the effects of psychotherapeutic interventions on loneliness might be limited as reported effect size estimates are mostly small-to-medium [28, 29]. These observations might indicate the necessity to adopt more comprehensive

approaches that also cover top-down strategies at the level of public health interventions. However, their development might require broader insights into population-wide processes.

Following the considerations about the spectrum of loneliness severity that might be covered by population-based studies, it is needed to note that previous studies have found loneliness to be either a transient or chronic experience [30-36]. Although the exact threshold duration of loneliness defining its chronic temporal pattern has not been established so far, previous studies have conceptualized chronic loneliness as its occurrence across at least two timepoints spanning between 1 and 6 years [30-36]. Investigating differences between chronic and transient loneliness might be of importance to understand interindividual variability of underlying mechanisms and outcomes. According to the cognitive and evolutionary model [4, 37, 38], the experience of loneliness might be adaptive as it triggers cognitions and behaviours that aim to restore social connections. However, some individuals, especially those remaining lonely over time, focus on threats related to social interactions and thus remain socially disconnected [39]. In agreement with this model, it has been observed that transient loneliness is associated with a smaller interpersonal distance while chronic loneliness is related to a greater interpersonal distance [40]. A recent population-based study further demonstrated that chronic and transient loneliness show important differences in terms of underlying risk factors [41]. In this study, female gender, not being married, poor educational attainment, poor mental and physical health, being limited in activities, poor social network, and living in a culturally individualistic country are risk factors for chronic loneliness. For transient loneliness, the effects of some risk factors including gender, physical health, the level of education, and social network size were either not significant or not robust. It has also been suggested that transient loneliness tends to occur after stressful events (e.g., retirement and

loss of close social interactions), while chronic loneliness is more closely related to poor social cognition, low social support, and a lack of intimate relationships [42].

Little is known about the differential associations of chronic and transient loneliness with mental health outcomes. To date, temporal patterns of loneliness have not been tested with respect to mental health outcomes beyond those related to depressive symptoms. It has recently been observed that both chronic and transient loneliness, assessed over the course of one year, are significantly associated with a higher likelihood of reporting a history of depression diagnosis and psychiatric distress [30]. However, stronger associations were found for chronic loneliness. Similar observations have been obtained in older populations [33, 36, 43, 44] and college students [45]. To bridge existing research gaps, the present study aimed to assess differential associations of chronic and transient loneliness with a variety of mental health outcomes represented by depressive symptoms, generalized anxiety, social anxiety, and paranoid thoughts in a large, general population sample. The hypothesis behind the present study was that both temporal patterns are associated with poor mental health outcomes; however, these associations might be stronger for chronic loneliness.

Methods

Recruitment procedures

The cohort reported in this study was developed using the quota sampling method to provide sample representativeness with respect to age, gender, education, employment status, and place of residence. All assessments were carried out using self-reports implemented in an internet-based survey. Baseline data were collected between July and August, 2024. The follow-up assessment took place in February, 2025. At both timepoints, after

the first invitation, potential participants received up to two reminders. Participants were enrolled by a research company using its own online access panel of registered and verified participants. The panel includes over 70,000 participants residing all administrative regions of Poland. It is continuously being developed by means of regular campaigns. Individuals with underrepresented backgrounds (e.g., ethnic and social minorities) are continuously invited to register by additional campaigns initiated through trusted channels (e.g., ethnic media, cultural festivals, and local events). These campaigns are implemented by the staff trained on cultural sensitivity that approaches community leaders and organizations, and disseminates culturally appropriate contents (in terms of language, imagery, and messaging) with transparent information about research activities. In this study, the panel members aged at least 18 years were eligible for participation. For completing both surveys, the participants received incentives equivalent to 10 EUR. To optimize the reliability of responses, the accuracy checks were used. Specifically, respondents were excluded while violating any of the following accuracy checks: (1) short survey completion time (i.e., below 30% of the median completion time); (2) failure to pass attention checks (i.e., participants were asked to respond to items requesting them to select a specific answer); (3) inconsistent responses to repeated items, and (4) responses with random strings of characters. The study received approval from the Bioethics Committee at Wroclaw Medical University, Wroclaw, Poland (approval number: 553/2024) and all participants provided the online version of informed consent.

Measures

Depressive symptoms: The Patient Health Questionnaire-9 (PHQ-9) was administered to measure depressive symptoms [46]. It is a 9-item questionnaire measuring the frequency of various depressive symptoms over preceding two weeks with a 4-point scale (0 - "not at all"

to 3 - "nearly every day"). The total PHQ-9 score are between 0 and 27 (higher scores reflect a greater level of depressive symptoms). The PHQ-9 total score of ≥ 10 has optimal sensitivity and specificity in detecting depression and was used in this study [47]. The Cronbach's alpha of the PHQ-9 was 0.897 in this study.

Generalized anxiety: Anxiety symptoms were recorded using the Generalized Anxiety Disorder-7 (GAD-7) [48]. It is a 7-item questionnaire (rated on 4-point scale: 0 - "not at all", 3 - "nearly every day") that refers to generalized anxiety symptoms in the preceding 2 weeks (a 4-point scale; 0 - "not at all", 3 - "nearly every day"). The total score is 0 - 21, where higher scores indicate a greater level of anxiety symptoms. The optimal cut-off score of GAD-7 to detect generalized anxiety disorder has been estimated at ≥ 10 and thus it was used in our study [48]. The Cronbach's alpha of GAD-7 was 0.944 in this study.

Social anxiety: The Social Interaction Anxiety Scale (SIAS) was used [49]. It is a 20-item questionnaire that covers various of social anxiety. Respondents are instructed to rate the level each item is typical for them on a 5-point scale (0 – "not at all", 4 – "extremely"). The cut-off score of ≥ 36 was used [50]. The Cronbach's alpha was 0.947 in this study.

Paranoid ideation: Participants were asked to fill in the Revised Green et al. Paranoid Thoughts Scale (R-GPTS) [51]. It is based on two subscales measuring ideas of reference (part A, 8 items) and ideas of persecution (part B, 10 items) over the period of preceding month. Items are rated on a 5-point scale (0 – "not at all", 4 – "totally"). Respondents are asked to assess experiences that did not appear as the consequence of substance use. For the persecution scale, the recommended cut-off to detect clinical levels of persecutory ideation was estimated at ≥ 11 [51]. This cut-off corresponds with moderately severe levels of ideas of reference (part A, ≥ 16 points). Therefore, in our study, the participants showing threshold scores reaching at least one of these cut-offs were classified as those showing paranoid

ideation. In this study, the Cronbach's alphas for parts A and B were 0.930 and 0.964, respectively.

Social network size: To measure the social network size, the 6-item version of Lubben Social Network Scale (LSNS-6) was used [52]. The LSNS-6 items record the number of family members and friends who are seen or heard at least once a month, with whom the respondent can talk about private matters, and who can be called on for help (a 6-point scale). The total LSNS-6 score is 0 - 20. Higher scores correspond with a greater social network size. The Cronbach's alpha of the LSNS-6 was 0.876 in this study.

Loneliness: To record the level of loneliness, the 11-item version of the De Jong Gierveld Loneliness Scale (DJGLS) was used [53, 54]. Each item is rated using a 5-point scale (potential responses are as follows: "yes!", "yes", "more or less", "no", and "no!"). There are two DJGLS subscales: the first one measures emotional loneliness (6 items) and the second one refers to social loneliness (5 items). The total score is estimated by counting positive and neutral responses ("yes!", "yes", and "more or less") to items for emotional loneliness as well as negative and neutral responses ("no!", "no", and "more or less") to items developed for social loneliness. The total DJGLS score ranges between 0 and 11 (higher scores reflect a greater level of loneliness). We used the cut-off proposed by the authors of DJGLS of > 2 to classify participants as those experiencing loneliness [53]. In the present study, the Cronbach's alphas for emotional loneliness subscale and social loneliness subscale were 0.874 and 0.810, respectively.

Data analysis

In the first step, descriptive characteristics of participants who completed assessments at both timepoints and follow-up non-completers were compared using the chi-square test

(categorical variables) and t-tests (continuous variables). Next, participants who completed assessments at both timepoints were divided into three groups according to threshold scores of the DJGLS [53] as proposed by previous studies [30, 32-36, 43, 55]. These groups were as follows: (1) individuals with the DJGLS score of < 3 at both time points (i.e., participants without loneliness at neither of timepoints); (2) individuals with the DJGLS score of ≥ 3 at one timepoint (i.e., participants with transient loneliness), and (3) individuals with the DJGLS score of ≥ 3 at both timepoints (i.e., participants with chronic loneliness). Finally, binary logistic regression models were analysed to assess differential associations of loneliness temporal patterns with symptom scores. Separate models were analysed for baseline and follow-up symptoms taking into consideration three specific comparisons (i.e., chronic loneliness vs. no loneliness, transient loneliness vs. no loneliness, and chronic loneliness vs. transient loneliness). Covariates included age, gender, the level of education, place of residence, employment status, mean monthly income, social network size, substance use in the preceding month (except for alcohol and nicotine), and psychiatric treatment history (in the preceding month). There were no missing data among individuals who completed assessments at both timepoints. Results were interpreted as significant while obtaining the p-value lower than 0.05. All analyses were carried out in the SPSS software, version 28.

Results

Descriptive characteristics of the sample

Altogether, 10,985 individuals were approached for participation (Supplementary Figure 1). Baseline assessment was completed by 5,099 individuals (aged 44.9 ± 15.4 years, 47.7% men, 46.4% of individuals invited to participate) while the follow-up assessment

included 3,275 individuals resulting the retention rate of 64.2%. Individuals who completed both assessments did not differ significantly from individuals who did not participate in the follow-up assessment with respect to sociodemographic and clinical characteristics (Table 1). Chronic loneliness was observed in 58.1% participants while transient loneliness was found in 16.2% individuals.

Baseline symptoms and temporal patterns of loneliness

Transient loneliness was associated with significantly higher odds of baseline generalized anxiety, social anxiety, and paranoid ideation, but not depressive symptoms. This observation remained significant after adjustment for covariates (Table 2, Figure 1A). The highest effect size estimate was found for generalized anxiety (unadjusted analysis: OR = 1.94, 95%CI: 1.16 – 3.27, $p = 0.012$; adjusted analysis: OR = 1.92, 95%CI: 1.11 – 3.33, $p = 0.020$). Among all tested covariates, only a lower social network size at baseline was significantly associated with transient loneliness.

Chronic loneliness was associated with significantly higher baseline levels of all symptom scores before and after adjustment for covariates (Table 2, Figure 1B). The highest effect size estimate was observed for social anxiety (unadjusted analysis: OR = 4.29, 95%CI: 3.35 – 5.50, $p < 0.001$; adjusted analysis: OR = 3.18, 95%CI: 2.38 – 4.24, $p < 0.001$). Chronic loneliness was also significantly associated with female gender, higher levels of education, higher odds of substance use, and a lower social network size.

Follow-up symptoms and temporal patterns of loneliness

Transient loneliness was associated with significantly higher odds of follow-up generalized anxiety and social anxiety. This observation was significant before and after

adjustment for covariates (Table 3, Figure 1C). The highest effect size estimate was found for generalized anxiety (unadjusted analysis: OR = 2.57, 95%CI: 1.46 – 4.53, $p = 0.001$; adjusted analysis: OR = 2.51, 95%CI: 1.39 – 4.53, $p = 0.002$). Among all tested covariates, only a lower social network size at the follow-up was significantly associated with transient loneliness.

Chronic loneliness was associated with significantly higher odds of all symptoms before and after adjustment for covariates (Table 3, Figure 1D). The highest effect size estimate was found for social anxiety (unadjusted analysis: OR = 4.29, 95%CI: 3.35 – 5.50, $p < 0.001$; adjusted analysis: OR = 3.18, 95%CI: 2.38 – 4.24, $p < 0.001$). Chronic loneliness was also significantly associated with female gender, higher levels of education, higher odds of substance use, and a lower social network size.

Differences between temporal patterns of loneliness in their associations with mental health outcomes

Chronic loneliness, compared to transient loneliness, showed stronger associations with baseline and follow-up symptoms of depression, social anxiety, and paranoid thoughts, but not generalized anxiety (Table 4, Figure 2). These differences remained significant after adjustment for covariates. With respect to baseline symptoms, the highest effect size estimate was found for depressive symptoms (unadjusted analysis: OR = 2.46, 95%CI: 1.74 – 3.49, $p < 0.001$; adjusted analysis: OR = 2.20, 95%CI: 1.53 – 3.16, $p < 0.001$). In turn, for follow-up symptoms, the highest effect size estimate was observed for social anxiety (unadjusted analysis: OR = 2.68, 95%CI: 2.10 – 3.44, $p < 0.001$; adjusted analysis: OR: 2.30, 95%CI: 1.77 – 2.99, $p < 0.001$).

Discussion

307

308 *Main findings*

309 Findings from the present study indicate that chronic loneliness, compared to its
310 transient temporal pattern, shows stronger and robust associations with the symptoms of
311 social anxiety, generalized anxiety, depression, and paranoia that are likely to be observed in
312 a short-term perspective. In case of chronic loneliness, these associations are likely to be
313 bidirectional with the largest effect size estimates observed for social anxiety. In case of
314 transient loneliness, the associations with psychopathological symptoms might be more
315 complex. First, it is important to note that our study did not demonstrate any significant
316 associations of transient loneliness with depressive symptoms. Second, transient loneliness
317 was significantly associated with higher baseline, but not follow-up odds of paranoid ideation.
318 Third, the strongest effect driven by transient loneliness was found for generalized anxiety and
319 this effect was comparable to the one found for chronic loneliness.

320 In general, the findings about quantitative differences in the strength of associations
321 with symptom domains between chronic and transient loneliness are in agreement with our
322 expectations and observations from previous studies measuring the level of depressive
323 symptoms [30, 33, 36, 43-45]. Various processes might explain observed associations between
324 loneliness and mental health outcomes. There is evidence that individuals experiencing
325 loneliness tend to show a number of social information processing biases that might increase
326 the risk of developing mental disorders. These include attention for threat bias, negative and
327 hostile intent attributions, increased rejection sensitivity, negative evaluation of self and
328 others, endorsement of less promotion- and more prevention-oriented goals, and low
329 perceived self-efficacy [39]. These cognitive biases might be bidirectionally associated with

depression [56], generalized anxiety disorder [57], social anxiety disorder [58], and psychosis [59].

It is also important to discuss some discrepancies between our findings and previous observations. Indeed, our findings, to some extent, contradict the evolutionary model of loneliness positing its transient temporal pattern as an adaptive experience that signals the need to reconnect with others and indicating a lack of unfavourable associations with mental health. However, observations from previous studies also do not support these considerations. For instance, a longitudinal study of adolescents with the observation period of 12 years identified three trajectories of loneliness, i.e., “stable low”, “high decreasing”, and “low increasing” [60]. Both “high decreasing” and “low increasing” temporal patterns were associated with elevated risk of depression. In addition, the “low increasing” pattern was associated with a higher risk of anxiety. Another study, based on a 14-year observation period of older adults revealed two frequency patterns of loneliness (i.e., moderate and high frequency) and both of them were associated with increased risk of depression [61]. Also, the study which spanned one year and included two waves of data demonstrated that both chronic and transient loneliness are related to a higher risk of depression [30]. Notably, the present study did not demonstrate any significant associations of transient loneliness with depressive symptoms while observing significant associations with social and generalized anxiety as well as follow-up paranoid ideation. This difference might originate from the fact that our study, as opposed to previous studies, analysed the associations with depressive symptoms while accounting for co-occurring psychopathology.

Methodological considerations

To our knowledge, this study has various strengths related to broad insights into symptom domains associated with loneliness, a relatively large sample size, and a longitudinal design. However, the findings also need to be interpreted in light of some limitations.

Recruitment procedures were implemented among the online panel users. While this approach provided representativeness of the sample in terms of sociodemographic characteristics, the risk of selection bias should be considered. Indeed, the study was performed among internet users and initiated during the summer period when many people spend holidays. Moreover, we did not use any measures of problematic internet use that has been associated with loneliness [20]. Following these considerations, it cannot be excluded that individuals with problematic internet use and those with higher levels of loneliness were overrepresented in the present cohort. Another limitation is that we did not record the information about the social or ethnic minority status. Also, the response and retention rates (46.4% and 64.2%, respectively) were relatively low. However, a recent meta-analysis estimated the mean response rate across online surveys at 44.1% [62]. Moreover, response rates are not always directly related to the validity of findings [63]. For instance, it has been found that studies with low response rate, even those with the response rate of 20%, are able to provide more accurate results than those with the response rate of 60 – 70% [64].

Another limitation is related to a lack of the general consensus on how loneliness should be operationalized as a categorical construct. Therefore, prevalence rates of loneliness show relatively high variability across previous studies that is further influenced by the use of specific questionnaires. In general, studies using single-item measures tend to report lower prevalence rates and some authors argue that simple measures may not thoroughly capture the construct of loneliness [65, 66]. A recent meta-analysis estimated the prevalence of loneliness at 55.4% in community-dwelling older adults while pooling the studies using DJGLS

[65]. For studies based on single-item questions, the prevalence rate was found to be 21.2%. Single-item measures often use direct indicators of loneliness (e.g., “I feel lonely” and “how often do you feel lonely?”). Indirect indicators, represented by the DJGLS that was used in the present study, are not based on the reference to loneliness. Therefore, they do not evoke negative stereotypes and limit socially desirable responses [66, 67]. Indeed, individuals experiencing loneliness are likely to hold self-stigmatizing perceptions that might be defined as “the shame for being lonely and inclination to conceal loneliness” [68]. In this regard, the use of single-item and direct indicators might lead to underreporting of loneliness. Findings from our study may also provide limited insights into the temporal patterns of loneliness due to a low number of assessments over time and a relatively short observation period. However, we believe that approaching shorter observation periods may also be important as it informs about early consequences of loneliness and processes that directly precede its occurrence. Considering our approach to data analysis, it is needed to note that while we controlled for the effects of various sociodemographic characteristics, we did not assess as to whether these variables moderate the associations of loneliness with mental health outcomes.

It should also be noted that while the general population provides opportunities for studying loneliness as a public health concern taking into consideration the heterogeneity of social determinants, this approach limits the potential to translate findings over individuals with mental disorders. At this point, it is also important to highlight that the measures of mental health outcomes were based on screening questionnaires and a comprehensive clinical validation was not performed. Finally, it should be noted that the study does enable to draw conclusions about causality due to observational design.

Directions for future studies

The most important direction for future studies is related to the need of a consensus-driven operationalization of loneliness temporal patterns. In this regard, two research directions need to be discussed. First, it should be considered to develop questionnaires recording duration of loneliness. Second, it might be needed to perform intensive longitudinal studies that record the experience of loneliness and mental health characteristics over multiple timepoints in order to find the threshold duration defining temporal patterns of loneliness. Another important direction is related to the need of better understanding of factors moderating the associations of loneliness with mental health in order to define populations with the highest vulnerability. For instance, it has consistently been shown that the prevalence of loneliness is the highest in young adulthood (up to 30 years of age), decreases in middle adulthood and early old age, and then increases during the oldest old age [69-71]. To date, it has been suggested that contextual factors underlying loneliness in various age groups might differ [22-27]. However, as shown by a recent meta-analysis, interventions targeting loneliness in various age groups might show comparable efficacy [29].

Finally, with an ongoing digitalization and frequent involvement in remote social interactions, it might be needed to better understand on how various problematic online behaviours are interrelated with loneliness. A recent meta-analysis demonstrated that problematic internet use is bidirectionally associated with loneliness [20]. This association was found to be stronger in samples from Eastern countries, with more men and young adults, and those studied over recent years. On one side, problematic internet use might be perceived as the strategy to cope with social disconnection (either loneliness or social isolation) and boredom [72-74]. On the other side, problematic internet use may increase the level of loneliness through the mechanisms related to social displacement, a lack of adequate sensory cues and bodily feedback [75, 76]. However, it is needed to point out that problematic internet

use is now believed to serve as the spectrum of various problematic online behaviours that show differential associations with mental health outcomes [77]. Addressing their differential associations with loneliness is warranted. Also, the need to explore on how digital advancements might be reframed to tackle loneliness should be highlighted. For instance, there is some evidence that enhancing video communication between care home residents and family members might be feasible in supporting person-centred care, social interactions, and wellbeing [78].

Clinical implications

Although the present study was carried out in the general population, some implications for public health and clinical practice might be formulated. The findings imply the necessity to consider early intervention and risk stratification taking into consideration the temporal pattern of loneliness. It should also be considered which interventions need to be prioritized among individuals showing specific temporal patterns of loneliness. According to the cognitive-evolutionary model, transient loneliness might motivate individuals to seek social interactions [4, 37, 38]. A recent study revealed that while transient loneliness is associated with a smaller preferred interpersonal distance, chronic loneliness was found to be linked with a greater preferred interpersonal distance [40]. In this regard, it might be concluded that specific therapeutic interventions targeting transient loneliness, especially those related to motivating social involvement, may not be necessary. In this group, awareness of transition to chronicity of loneliness, improvement of social skills maintaining close relationships, community engagement, and clinical assessment should be considered [79]. In turn, the initiation of therapeutic interventions might be the priority for individuals with chronic loneliness. To date, a variety of approaches targeting chronic loneliness have been

developed. A recent meta-analysis revealed that the priority should be given to the reminiscence intervention, interventions promoting social identity formation, and cognitive-behavioural therapy [29]. Another meta-analysis revealed the greatest effect size for cognitive interventions and a lack of benefits related to the interventions enhancing social skills [80].

Conclusions

Taken together, the findings indicate that chronic and transient loneliness show qualitative and quantitative differences in their associations with psychopathological symptoms that might already be observed in a short-term perspective. Chronic loneliness shows robust and bidirectional associations across various domains of psychopathology covering depressive symptoms, social anxiety, generalized anxiety, and paranoid ideation. In turn, transient loneliness shows bidirectional associations with generalized and social anxiety. These observations hold some implications referring to the public health and clinical perspectives by indicating the rationale to detect loneliness in its early development. However, additional studies approaching longer observation periods and a thorough clinical assessment are also needed to provide broader insights into the mechanisms, consequences, and temporal patterns of loneliness.

Figure legend

Figure 1. Temporal patterns of loneliness and their associations with mental health outcomes. The associations with baseline symptoms are shown in Figure 1A (transient loneliness vs. no loneliness) and Figure 1B (chronic loneliness vs. no loneliness), while the associations with follow-up symptoms are visualized in Figure 1C (transient loneliness vs. no loneliness) and

Figure 1D (chronic loneliness vs. no loneliness). Results are adjusted for age, gender, education, place of residence, employment status, monthly income, social network size, substance use (preceding month), and psychiatric treatment history (preceding month).

Figure 2. Differences between temporal patterns of loneliness in their associations with baseline (A) and follow-up (B) symptoms. Chronic loneliness is a reference category. Results are adjusted for age, gender, education, place of residence, employment status, monthly income, social network size, substance use (preceding month), and psychiatric treatment history (preceding month).

Financial support. The study was funded from “the Initiative of Excellence – Research University” provided by Wroclaw Medical University, Wroclaw, Poland (task number: IDUB.C230.24.001).

Acknowledgments. None to report.

Conflicts of interest. There is no conflict of interest to report regarding the present article.

Data availability statement. Due to sensitivity reasons, the data supporting the present study are available upon reasonable request sent to the corresponding author.

References

- 495 [1] Casabianca E, Kovacic M. Historical roots of loneliness and its impact on second-
 496 generation immigrants' health. *J Econ Behav Organ.* 2024;224:407-37.
 497 <https://doi.org/10.1016/j.jebo.2024.06.011>.
- 498 [2] Ahmed MZ. Introductory Chapter: Unraveling Loneliness – Historical Perspectives and
 499 Modern Realities. In: Ahmed MZ, editor *Determinants of Loneliness*, Rijeka:
 500 IntechOpen; 2024. <https://doi.org/10.5772/intechopen.1005761>.
- 501 [3] Sebo E. Exiles: Medieval Experiences of Isolation. *Neophilologus.* 2023;107(1):83-4.
 502 <https://doi.org/10.1007/s11061-022-09741-w>.
- 503 [4] Cacioppo JT, Hawkley LC, Bernston GG. The anatomy of loneliness. *Curr Dir Psychol Sci.*
 504 2003;12(3):71-4. <https://doi.org/10.1111/1467-8721.01232>.
- 505 [5] Luhmann M, Buecker S, Rusberg M. Loneliness across time and space. *Nat Rev Psychol.*
 506 2023;2(1):9-23. <https://doi.org/10.1038/s44159-022-00124-1>.
- 507 [6] Franklin A. A lonely society? Loneliness and liquid modernity in Australia. *Aust J Soc*
 508 *Issues.* 2012;47(1):11-28. <https://doi.org/10.1002/j.1839-4655.2012.tb00232.x>.
- 509 [7] Bauman Z. *Liquid Modernity*. Oxford, England: Polity Press; 2000.
- 510 [8] Ortega MJ. Depression, loneliness, and unhappiness. *Int Psychiatry Clin.* 1969;6(2):143-
 511 53.
- 512 [9] Mark FR. America's major diseases: loneliness, anxiety, boredom. *Rehabil Rec.*
 513 1972;13(3):5-8.
- 514 [10] Ng IKS. The 'loneliness' epidemic: a new social determinant of health? *Intern Med J.*
 515 2024;54(3):365-7. <https://doi.org/10.1111/imj.16356>.
- 516 [11] Park C, Majeed A, Gill H, Tamura J, Ho RC, Mansur RB, et al. The Effect of Loneliness on
 517 Distinct Health Outcomes: A Comprehensive Review and Meta-Analysis. *Psychiatry Res.*
 518 2020;294:113514. <https://doi.org/10.1016/j.psychres.2020.113514>.

- 519 [12] Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D. Loneliness and social
520 isolation as risk factors for mortality: a meta-analytic review. *Perspect Psychol Sci.*
521 2015;10(2):227-37. <https://doi.org/10.1177/1745691614568352>.
- 522 [13] Chen Z, Song X, Lee TMC, Zhang R. The robust reciprocal relationship between
523 loneliness and depressive symptoms among the general population: Evidence from a
524 quantitative analysis of 37 studies. *J Affect Disord.* 2023;343:119-28.
525 <https://doi.org/10.1016/j.jad.2023.09.035>.
- 526 [14] Maes M, Nelemans SA, Danneel S, Fernandez-Castilla B, Van den Noortgate W,
527 Goossens L, et al. Loneliness and social anxiety across childhood and adolescence:
528 Multilevel meta-analyses of cross-sectional and longitudinal associations. *Dev Psychol.*
529 2019;55(7):1548-65. <https://doi.org/10.1037/dev0000719>.
- 530 [15] Domenech-Abella J, Mundo J, Haro JM, Rubio-Valera M. Anxiety, depression, loneliness
531 and social network in the elderly: Longitudinal associations from The Irish Longitudinal
532 Study on Ageing (TILDA). *J Affect Disord.* 2019;246:82-8.
533 <https://doi.org/10.1016/j.jad.2018.12.043>.
- 534 [16] Michalska da Rocha B, Rhodes S, Vasilopoulou E, Hutton P. Loneliness in Psychosis: A
535 Meta-analytical Review. *Schizophr Bull.* 2018;44(1):114-25.
536 <https://doi.org/10.1093/schbul/sbx036>.
- 537 [17] Chau AKC, Zhu C, So SH. Loneliness and the psychosis continuum: a meta-analysis on
538 positive psychotic experiences and a meta-analysis on negative psychotic experiences.
539 *Int Rev Psychiatry.* 2019;31(5-6):471-90.
540 <https://doi.org/10.1080/09540261.2019.1636005>.
- 541 [18] Bonar EE, Walton MA, Carter PM, Lin LA, Coughlin LN, Goldstick JE. Longitudinal within-
542 and between-person associations of substance use, social influences, and loneliness

- 543 among adolescents and emerging adults who use drugs. *Addict Res Theory*.
 544 2022;30(4):262-7. <https://doi.org/10.1080/16066359.2021.2009466>.
- 545 [19] Gutkind S, Gorfinkel LR, Hasin DS. Prospective effects of loneliness on frequency of
 546 alcohol and marijuana use. *Addict Behav*. 2022;124:107115.
 547 <https://doi.org/10.1016/j.addbeh.2021.107115>.
- 548 [20] Zhang Y, Li J, Zhang M, Ai B, Jia F. Bidirectional associations between loneliness and
 549 problematic internet use: A meta-analytic review of longitudinal studies. *Addict Behav*.
 550 2024;150:107916. <https://doi.org/10.1016/j.addbeh.2023.107916>.
- 551 [21] Barjakova M, Garnero A, d'Hombres B. Risk factors for loneliness: A literature review.
 552 *Soc Sci Med*. 2023;334:116163. <https://doi.org/10.1016/j.socscimed.2023.116163>.
- 553 [22] Matthews T, Arseneault L, Bryan BT, Fisher HL, Gray R, Henchy J, et al. Social media use,
 554 online experiences, and loneliness among young adults: A cohort study. *Ann N Y Acad*
 555 *Sci*. 2025. <https://doi.org/10.1111/nyas.15370>.
- 556 [23] Haslam SA, Haslam C, Cruwys T, Jetten J, Bentley SV, Fong P, et al. Social identity makes
 557 group-based social connection possible: Implications for loneliness and mental health.
 558 *Curr Opin Psychol*. 2022;43:161-5. <https://doi.org/10.1016/j.copsyc.2021.07.013>.
- 559 [24] Watson J, Nesdale D. Rejection sensitivity, social withdrawal, and loneliness in young
 560 adults. *J Appl Soc Psychol*. 2012;42:1984-2005. <https://doi.org/10.1111/j.1559-1816.2012.00927.x>.
- 561
 562 [25] Carrasco PM, Crespo DP, Garcia AIR, Ibanez ML, Rubio BM, Montenegro-Pena M.
 563 Predictive factors and risk and protection groups for loneliness in older adults: a
 564 population-based study. *BMC Psychol*. 2024;12(1):238.
 565 <https://doi.org/10.1186/s40359-024-01708-7>.

- 566 [26] Hutten E, Jongen EMM, Hajema KJ, Ruiter RAC, Hamers F, Bos AER. Risk factors of
567 loneliness across the life span. *Soc Pers Relat.* 2021;39(5):1482-507.
- 568 [27] Sheftel MG, Margolis R, Verdery AM. Life Events and Loneliness Transitions Among
569 Middle-Aged and Older Adults Around the World. *J Gerontol B Psychol Sci Soc Sci.*
570 2024;79(1). <https://doi.org/10.1093/geronb/gbad149>.
- 571 [28] Duffner LA, Janssen N, Deckers K, Schroyen S, de Vugt ME, Kohler S, et al. Facing the
572 Next "Geriatric Giant"-A Systematic Literature Review and Meta-Analysis of
573 Interventions Tackling Loneliness and Social Isolation Among Older Adults. *J Am Med*
574 *Dir Assoc.* 2024;25(9):105110. <https://doi.org/10.1016/j.jamda.2024.105110>.
- 575 [29] Hickin N, Kall A, Shafran R, Sutcliffe S, Manzotti G, Langan D. The effectiveness of
576 psychological interventions for loneliness: A systematic review and meta-analysis. *Clin*
577 *Psychol Rev.* 2021;88:102066. <https://doi.org/10.1016/j.cpr.2021.102066>.
- 578 [30] Wolska K, Creaven AM. Associations between transient and chronic loneliness, and
579 depression, in the understanding society study. *Br J Clin Psychol.* 2023;62(1):112-28.
580 <https://doi.org/10.1111/bjc.12397>.
- 581 [31] Cutrona CE. Transition to college: Loneliness and the process of social adjustment. In:
582 Peplau LA, Perlman D, editors. *Loneliness: A sourcebook of current theory, research*
583 *and therapy*, New York: John Wiley; 1982, p. 291 - 309.
- 584 [32] Ojagbemi A, Bello T, Gureje O. The roles of depression and social relationships in the
585 onset and course of loneliness amongst Nigerian elders. *Int J Geriatr Psychiatry.*
586 2021;36(4):547-57. <https://doi.org/10.1002/gps.5451>.
- 587 [33] Theeke LA. Sociodemographic and health-related risks for loneliness and outcome
588 differences by loneliness status in a sample of U.S. older adults. *Res Gerontol Nurs.*
589 2010;3(2):113-25. <https://doi.org/10.3928/19404921-20091103-99>.

- 590 [34] Martin-Maria N, Caballero FF, Miret M, Tyrovolas S, Haro JM, Ayuso-Mateos JL, et al.
 591 Differential impact of transient and chronic loneliness on health status. A longitudinal
 592 study. Psychol Health. 2020;35(2):177-95.
 593 <https://doi.org/10.1080/08870446.2019.1632312>.
- 594 [35] Gerst-Emerson K, Jayawardhana J. Loneliness as a public health issue: the impact of
 595 loneliness on health care utilization among older adults. Am J Public Health.
 596 2015;105(5):1013-9. <https://doi.org/10.2105/AJPH.2014.302427>.
- 597 [36] Martin-Maria N, Caballero FF, Lara E, Domenech-Abella J, Haro JM, Olaya B, et al. Effects
 598 of transient and chronic loneliness on major depression in older adults: A longitudinal
 599 study. Int J Geriatr Psychiatry. 2021;36(1):76-85. <https://doi.org/10.1002/gps.5397>.
- 600 [37] Cacioppo JT, Hawkley LC, Ernst M, Burleson M, Bernston GG, Nouriani B, et al.
 601 Loneliness within a nomological net: An evolutionary perspective. J Res Personality.
 602 2006;40:1054-85. <https://doi.org/10.1016/j.jrp.2005.11.007>.
- 603 [38] Cacioppo JT, Hawkley LC. Perceived social isolation and cognition. Trends Cogn Sci.
 604 2009;13(10):447-54. <https://doi.org/10.1016/j.tics.2009.06.005>.
- 605 [39] Spithoven AWM, Bijttebier P, Goossens L. It is all in their mind: A review on information
 606 processing bias in lonely individuals. Clin Psychol Rev. 2017;58:97-114.
 607 <https://doi.org/10.1016/j.cpr.2017.10.003>.
- 608 [40] Saporta N, Scheele D, Lieberz J, Stuhr-Wulff F, Hurlemann R, Shamay-Tsoory SG.
 609 Opposing Association of Situational and Chronic Loneliness with Interpersonal
 610 Distance. Brain Sci. 2021;11(9). <https://doi.org/10.3390/brainsci11091135>.
- 611 [41] Domenech-Abella J, Gabarrell-Pascuet A, Mundo J, Haro JM, Varga TV. Chronic and
 612 Transient Loneliness in Western Countries: Risk Factors and Association With

- 613 Depression. A 2-Year Follow-Up Study. *Am J Geriatr Psychiatry*. 2024;32(4):412-23.
 614 <https://doi.org/10.1016/j.jagp.2023.11.001>.
- 615 [42] Heinrich LM, Gullone E. The clinical significance of loneliness: a literature review. *Clin*
 616 *Psychol Rev*. 2006;26(6):695-718. <https://doi.org/10.1016/j.cpr.2006.04.002>.
- 617 [43] Shiovitz-Ezra S, Ayalon L. Situational versus chronic loneliness as risk factors for all-
 618 cause mortality. *Int Psychogeriatr*. 2010;22(3):455-62.
 619 <https://doi.org/10.1017/S1041610209991426>.
- 620 [44] Zhang Y, Kuang J, Xin Z, Fang J, Song R, Yang Y, et al. Loneliness, social isolation,
 621 depression and anxiety among the elderly in Shanghai: Findings from a longitudinal
 622 study. *Arch Gerontol Geriatr*. 2023;110:104980.
 623 <https://doi.org/10.1016/j.archger.2023.104980>.
- 624 [45] Hojat M. Comparison of transitory and chronic loners on selected personality variables.
 625 *Br J Psychol*. 1983;74(Pt 2):199-202. [https://doi.org/10.1111/j.2044-](https://doi.org/10.1111/j.2044-8295.1983.tb01855.x)
 626 [8295.1983.tb01855.x](https://doi.org/10.1111/j.2044-8295.1983.tb01855.x).
- 627 [46] Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity
 628 measure. *J Gen Intern Med*. 2001;16(9):606-13. [https://doi.org/10.1046/j.1525-](https://doi.org/10.1046/j.1525-1497.2001.016009606.x)
 629 [1497.2001.016009606.x](https://doi.org/10.1046/j.1525-1497.2001.016009606.x).
- 630 [47] Negeri ZF, Levis B, Sun Y, He C, Krishnan A, Wu Y, et al. Accuracy of the Patient Health
 631 Questionnaire-9 for screening to detect major depression: updated systematic review
 632 and individual participant data meta-analysis. *BMJ*. 2021;375:n2183.
 633 <https://doi.org/10.1136/bmj.n2183>.
- 634 [48] Spitzer RL, Kroenke K, Williams JB, Lowe B. A brief measure for assessing generalized
 635 anxiety disorder: the GAD-7. *Arch Intern Med*. 2006;166(10):1092-7.
 636 <https://doi.org/10.1001/archinte.166.10.1092>.

- 637 [49] Mattick RP, Clarke JC. Social Interaction Anxiety Scale (SIAS). In: APA PsycTests; 1998.
- 638 [50] Peters L. Discriminant validity of the Social Phobia and Anxiety Inventory (SPAI), the
639 Social Phobia Scale (SPS) and the Social Interaction Anxiety Scale (SIAS). Behav Res Ther.
640 2000;38(9):943-50. [https://doi.org/10.1016/s0005-7967\(99\)00131-x](https://doi.org/10.1016/s0005-7967(99)00131-x).
- 641 [51] Freeman D, Loe BS, Kingdon D, Startup H, Molodynski A, Rosebrock L, et al. The revised
642 Green et al., Paranoid Thoughts Scale (R-GPTS): psychometric properties, severity
643 ranges, and clinical cut-offs. Psychol Med. 2021;51(2):244-53.
644 <https://doi.org/10.1017/S0033291719003155>.
- 645 [52] Lubben J, Blozik E, Gillmann G, Iliffe S, von Renteln Kruse W, Beck JC, et al. Performance
646 of an abbreviated version of the Lubben Social Network Scale among three European
647 community-dwelling older adult populations. Gerontologist. 2006;46(4):503-13.
648 <https://doi.org/10.1093/geront/46.4.503>.
- 649 [53] de Jong-Gierveld J, Kamphuls F. The Development of a Rasch-Type Loneliness Scale.
650 Appl Psychol Meas. 1985;9(3):289-99. <https://doi.org/10.1177/014662168500900307>.
- 651 [54] Grygiel P, Humenny G, Rebisz S, Świtaj P, Sikorska J. Validating the Polish Adaptation of
652 the 11-Item De Jong Gierveld Loneliness Scale. Eur J Psychol Assess. 2013;29(2):129-
653 39. <https://doi.org/10.1027/1015-5759/a000130>.
- 654 [55] Vanhalst J, Luyckx K, Van Petegem S, Soenens B. The Detrimental Effects of Adolescents'
655 Chronic Loneliness on Motivation and Emotion Regulation in Social Situations. J Youth
656 Adolesc. 2018;47(1):162-76. <https://doi.org/10.1007/s10964-017-0686-4>.
- 657 [56] Vos LMW, Nieto I, Amanvermez Y, Smeets T, Everaert J. Do cognitive biases
658 prospectively predict anxiety and depression? A multi-level meta-analysis of
659 longitudinal studies. Clin Psychol Rev. 2025;116:102552.
660 <https://doi.org/10.1016/j.cpr.2025.102552>.

- 661 [57] Krahe C, Whyte J, Bridge L, Loizou S, Hirsch CR. Are different forms of repetitive
 662 negative thinking associated with interpretation bias in generalized anxiety disorder
 663 and depression? Clin Psychol Sci.7(5):969-81.
 664 <https://doi.org/10.1177/2167702619851808>.
- 665 [58] Liu H, Li X, Han B, Liu X. Effects of cognitive bias modification on social anxiety: A meta-
 666 analysis. PLoS One. 2017;12(4):e0175107.
 667 <https://doi.org/10.1371/journal.pone.0175107>.
- 668 [59] Gaweda L, Kowalski J, Aleksandrowicz A, Bagrowska P, Dabkowska M, Pionke-Ubych R.
 669 A systematic review of performance-based assessment studies on cognitive biases in
 670 schizophrenia spectrum psychoses and clinical high-risk states: A summary of 40 years
 671 of research. Clin Psychol Rev. 2024;108:102391.
 672 <https://doi.org/10.1016/j.cpr.2024.102391>.
- 673 [60] Hutten E, Jongen EMM, Verboon P, Bos AER, Smeekens S, Cillessen AHN. Trajectories
 674 of Loneliness and Psychosocial Functioning. Front Psychol. 2021;12:689913.
 675 <https://doi.org/10.3389/fpsyg.2021.689913>.
- 676 [61] Liu I, Huang YJ, Wang LK, Tsai YH, Hsu SL, Chang CJ, et al. Dual trajectories of loneliness
 677 and depression and their baseline correlates over a 14-year follow-up period in older
 678 adults: Results from a nationally representative sample in Taiwan. Int J Older People
 679 Nurs. 2021;16(6):e12410. <https://doi.org/10.1111/opn.12410>.
- 680 [62] Wu MJ, Zhao K, Fils-Aime F. Response rates of online surveys in published research: A
 681 meta-analysis. Computers in Human Behavior Reports. 2022;7:100206.
 682 <https://doi.org/https://doi.org/10.1016/j.chbr.2022.100206>
 683 .

- 684 [63] Morton SM, Bandara DK, Robinson EM, Carr PE. In the 21st Century, what is an
 685 acceptable response rate? Aust N Z J Public Health. 2012;36(2):106-8.
 686 <https://doi.org/10.1111/j.1753-6405.2012.00854.x>.
- 687 [64] Visser PS, Krosnick JA, Marquette J, Curtin M. Publ Opin Quart. 1996;60(2):181-227.
 688 <https://doi.org/10.1086/297748>.
- 689 [65] Stegen H, Duppen D, Savieri P, Stas L, Pan H, Aartsen M, et al. Loneliness prevalence of
 690 community-dwelling older adults and the impact of the mode of measurement, data
 691 collection, and country: A systematic review and meta-analysis. Int Psychogeriatr.
 692 2024;36(9):747-61. <https://doi.org/10.1017/S1041610224000425>.
- 693 [66] Mund M, Maes M, Drewke PM, Gutzeit A, Jaki I, Qualter P. Would the Real Loneliness
 694 Please Stand Up? The Validity of Loneliness Scores and the Reliability of Single-Item
 695 Scores. Assessment. 2023;30(4):1226-48.
 696 <https://doi.org/10.1177/10731911221077227>.
- 697 [67] Russell D. The measurement of loneliness. In: Pervin LA, Perlman D, editors. Loneliness:
 698 A sourcebook of current theory, research and therapy, John Wiley; 1982, p. 81 - 104.
- 699 [68] Barreto M, van Breen J, Victor C, Hammond C, Eccles A, Richins MT, et al. Exploring the
 700 nature and variation of the stigma associated with loneliness. J Soc Pers Relat.
 701 2022;39(9):2658-79. <https://doi.org/10.1177/02654075221087190>.
- 702 [69] Hawkley LC. Loneliness and health. Nat Rev Dis Primers. 2022;8(1):22.
 703 <https://doi.org/10.1038/s41572-022-00355-9>.
- 704 [70] Hawkley LC, Buecker S, Kaiser T, Luhmann M. Loneliness from Young Adulthood to Old
 705 Age: Explaining Age Differences in Loneliness. Int J Behav Dev. 2022;46(1):39-49.
 706 <https://doi.org/10.1177/0165025420971048>.

- 707 [71] Graham EK, Beck ED, Jackson K, Yoneda T, McGhee C, Pieramici L, et al. Do We Become
708 More Lonely With Age? A Coordinated Data Analysis of Nine Longitudinal Studies.
709 Psychol Sci. 2024;35(6):579-96. <https://doi.org/10.1177/09567976241242037>.
- 710 [72] Orsolini L, Longo G, Volpe U. The Mediatory Role of the Boredom and Loneliness
711 Dimensions in the Development of Problematic Internet Use. Int J Environ Res Public
712 Health. 2023;20(5). <https://doi.org/10.3390/ijerph20054446>.
- 713 [73] Orsolini L, Volpe U, Albert U, Carmassi C, Carra G, Cirulli F, et al. Use of social network
714 as a coping strategy for depression among young people during the COVID-19
715 lockdown: findings from the COMET collaborative study. Ann Gen Psychiatry.
716 2022;21(1):44. <https://doi.org/10.1186/s12991-022-00419-w>.
- 717 [74] Volpe U, Orsolini L, Salvi V, Albert U, Carmassi C, Carra G, et al. COVID-19-Related Social
718 Isolation Predispose to Problematic Internet and Online Video Gaming Use in Italy. Int
719 J Environ Res Public Health. 2022;19(3). <https://doi.org/10.3390/ijerph19031539>.
- 720 [75] Costa RM, Patrao I, Machado M. Problematic internet use and feelings of loneliness.
721 Int J Psychiatry Clin Pract. 2019;23(2):160-2.
722 <https://doi.org/10.1080/13651501.2018.1539180>.
- 723 [76] Kim J, LaRose R, Peng W. Loneliness as the cause and the effect of problematic Internet
724 use: the relationship between Internet use and psychological well-being. Cyberpsychol
725 Behav. 2009;12(4):451-5. <https://doi.org/10.1089/cpb.2008.0327>.
- 726 [77] Bloch M, Misiak B. Problematic online behaviors and their early patterns of co-
727 occurrence in young adults: insights from directed and undirected networks. Front
728 Psychiatry. 2025;16:1446338. <https://doi.org/10.3389/fpsyt.2025.1446338>.

- 729 [78] Austbo LBH, Testad I, Gjesten MT. Using a Robot to Address the Well-Being, Social
730 Isolation, and Loneliness of Care Home Residents via Video Calls: Qualitative Feasibility
731 Study. JMIR Form Res. 2025;9:e59764. <https://doi.org/10.2196/59764>.
- 732 [79] Meehan DE, Clare P, Grunseit A, Merom D. Community-level determinants of loneliness
733 and social isolation: a population-based cohort study across younger and older adults.
734 Front Public Health. 2025;13:1526166. <https://doi.org/10.3389/fpubh.2025.1526166>.
- 735 [80] Masi CM, Chen HY, Hawkley LC, Cacioppo JT. A meta-analysis of interventions to reduce
736 loneliness. Pers Soc Psychol Rev. 2011;15(3):219-66.
737 <https://doi.org/10.1177/1088868310377394>.
- 738
- 739

740 **Table 1.** The general characteristics of the cohort.

	Total sample (<i>n</i> = 5,099)	Completers (<i>n</i> = 3,275)	Non- completers (<i>n</i> = 1,824)	<i>p</i>
Age, years	44.9 ± 15.4	45.2 ± 15.7	44.8 ± 15.2	0.189
Gender, men	2431 (47.7)	1568 (47.9)	863 (47.3)	0.718
Education				
Primary	94 (1.8)	53 (1.6)	41 (2.2)	0.858
Vocational	411 (8.1)	273 (8.3)	138 (7.6)	
Secondary	2098 (41.1)	1345 (41.1)	753 (41.3)	
Higher	2496 (49.0)	1604 (49.0)	892 (48.9)	
Employment				
Unemployed	435 (8.5)	265 (8.1)	170 (9.3)	0.406
Retired/disability pension	1002 (19.6)	632 (19.3)	370 (20.3)	
Student	234 (4.6)	156 (4.8)	78 (4.3)	
Part-time work	482 (9.5)	317 (9.7)	165 (9.0)	
Full-time work	2946 (57.8)	1905 (58.1)	1041 (57.1)	
Place of residence				
Rural	1650 (32.4)	1065 (32.5)	585 (32.1)	0.991
Urban, < 50,000 inhabitants	1166 (22.9)	750 (22.9)	416 (22.8)	
Urban, 50,000 – 150,000 inhabitants	816 (16.0)	525 (16.0)	291 (16.0)	
Urban, 150,000 – 500,000 inhabitants	714 (14.0)	457 (14.0)	257 (14.0)	
Urban, > 500,000 inhabitants	753 (14.7)	478 (14.6)	275 (15.1)	
Monthly income				
< 750 USD	1082 (21.1)	695 (21.2)	387 (21.2)	0.170
750 – 1500 USD	2485 (48.7)	1621 (49.6)	864 (47.4)	
1,500 – 2,500 USD	722 (14.2)	468 (14.3)	254 (13.9)	
2,500 – 3,750 USD	133 (2.6)	80 (2.4)	53 (2.9)	
> 3,750 USD	44 (0.9)	31 (0.9)	13 (0.7)	
Refused to answer	633 (12.4)	380 (11.6)	253 (13.9)	
Social network size, baseline	15.9 ± 6.0	16.0 ± 6.0	15.8 ± 5.8	0.222
Social network size, follow-up	–	15.7 ± 5.9	–	–
Psychiatric treatment				
Lifetime	1108 (21.7)	705 (21.5)	403 (22.1)	0.638
Preceding month	539 (10.6)	327 (10.0)	512 (28.1)	0.068
Substance use, preceding month*	353 (6.9)	213 (6.5)	140 (7.7)	0.353
Depression, baseline	1362 (26.7)	860 (26.3)	502 (27.5)	0.329
Depression, follow-up	–	823 (25.1)	–	–
Generalized anxiety, baseline	1084 (21.3)	683 (20.9)	401 (22.0)	0.345
Generalized anxiety, follow-up	–	664 (20.3)	–	–
Social anxiety, baseline	1890 (37.1)	1206 (36.8)	684 (37.5)	0.632
Social anxiety, follow-up	–	1219 (37.2)	–	–
Paranoid ideation, baseline	1618 (31.7)	1017 (31.1)	601 (32.9)	0.163
Paranoid ideation, follow-up	–	1018 (31.1)	–	–
Loneliness				
Chronic	–	1904 (58.1)	–	–
Transient	–	532 (16.2)	–	
None	–	839 (25.6)	–	

741 Data are reported as mean ± SD or *n* (%)

742 *Except for nicotine and alcohol

743

744 **Table 2.** The associations of baseline mental health measures with temporal patterns of loneliness.

Model	Independent variable	Analysis					
		Unadjusted			Adjusted		
		OR	95%CI	p	OR	95%CI	p
Transient loneliness vs. no loneliness	Depressive symptoms	1.20	0.74 – 1.94	0.455	1.16	0.69 – 1.92	0.578
	Generalized anxiety	1.94	1.16 – 3.27	0.012	1.92	1.11 – 3.33	0.020
	Social anxiety	1.87	1.36 – 2.56	< 0.001	1.65	1.18 – 2.31	0.004
	Paranoid ideation	1.62	1.16 – 2.24	0.004	1.64	1.16 – 2.32	0.005
	Age	–	–	–	0.99	0.98 – 1.00	0.134
	Gender (women)	–	–	–	1.26	0.97 – 1.62	0.084
	Education	–	–	–	1.18	0.98 – 1.43	0.081
	Urbanicity	–	–	–	0.96	0.87 – 1.04	0.311
	Unemployment ^a	–	–	–	1.01	0.76 – 1.34	0.967
	Income	–	–	–	1.04	0.88 – 1.23	0.626
	Psychiatric treatment ^b	–	–	–	1.54	0.82 – 2.90	0.176
	Substance use ^c	–	–	–	1.87	0.90 – 3.89	0.094
	Social network size	–	–	–	0.89	0.87 – 0.92	< 0.001
Chronic loneliness vs. no loneliness	Depressive symptoms	3.39	2.35 – 4.90	< 0.001	2.96	1.94 – 4.50	< 0.001
	Generalized anxiety	1.85	1.20 – 2.86	0.006	1.86	1.14 – 3.03	0.013
	Social anxiety	4.29	3.35 – 5.50	< 0.001	3.18	2.38 – 4.24	< 0.001
	Paranoid ideation	2.17	1.67 – 2.82	< 0.001	2.14	1.59 – 2.89	< 0.001
	Age	–	–	–	0.99	0.98 – 1.05	0.326
	Gender (women)	–	–	–	1.39	1.10 – 1.76	0.005
	Education	–	–	–	1.26	1.07 – 1.49	0.007
	Urbanicity	–	–	–	0.95	0.88 – 1.03	0.252
	Unemployment ^a	–	–	–	0.93	0.72 – 1.20	0.569
	Income	–	–	–	0.97	0.83 – 1.13	0.661
	Psychiatric treatment ^b	–	–	–	1.02	0.62 – 1.66	0.950
	Substance use ^c	–	–	–	2.27	1.22 – 4.22	0.009
	Social network size	–	–	–	0.80	0.78 – 0.82	< 0.001

745 Significant associations ($p < 0.05$) are bolded

746 ^aRefers to the lack of any employment and student status

747 ^bRefers to the prior-month history

748 ^cRefers to the prior-month history; nicotine and alcohol use were not assessed

749

750

Table 3. The associations of follow-up mental health measures with temporal patterns of loneliness.

Model	Independent variable	Analysis					
		Unadjusted			Adjusted		
		OR	95%CI	p	OR	95%CI	p
Transient loneliness vs. no loneliness	Depressive symptoms	1.12	0.69 – 1.82	0.656	1.11	0.67 – 1.85	0.690
	Generalized anxiety	2.57	1.46 – 4.53	0.001	2.51	1.39 – 4.53	0.002
	Social anxiety	1.77	1.28 – 2.45	< 0.001	1.52	1.08 – 2.15	0.017
	Paranoid ideation	1.30	0.93 – 1.81	0.126	1.37	0.96 – 1.94	0.081
	Age	–	–	–	0.99	0.98 – 1.00	0.147
	Gender (women)	–	–	–	1.26	0.97 – 1.63	0.080
	Education	–	–	–	1.21	1.00 – 1.46	0.050
	Urbanicity	–	–	–	0.97	0.89 – 1.06	0.485
	Unemployment ^a	–	–	–	0.99	0.74 – 1.31	0.929
	Income	–	–	–	1.06	0.90 – 1.24	0.517
	Psychiatric treatment ^b	–	–	–	1.39	0.74 – 2.60	0.303
	Substance use ^c	–	–	–	2.03	0.98 – 4.23	0.058
	Social network size	–	–	–	0.89	0.86 – 0.91	< 0.001
Chronic loneliness vs. no loneliness	Depressive symptoms	2.26	1.57 – 3.24	< 0.001	2.11	1.39 – 3.21	< 0.001
	Generalized anxiety	2.92	1.84 – 4.62	< 0.001	2.38	1.43 – 3.95	< 0.001
	Social anxiety	5.00	3.89 – 6.41	< 0.001	3.39	2.56 – 4.50	< 0.001
	Paranoid ideation	2.34	1.81 – 3.03	< 0.001	2.58	1.92 – 3.47	< 0.001
	Age	–	–	–	0.99	0.98 – 1.01	0.639
	Gender (women)	–	–	–	1.36	1.08 – 1.72	0.009
	Education	–	–	–	1.25	1.05 – 1.48	0.010
	Urbanicity	–	–	–	0.97	0.90 – 1.05	0.475
	Unemployment ^a	–	–	–	0.90	0.70 – 1.16	0.422
	Income	–	–	–	0.97	0.83 – 1.13	0.690
	Psychiatric treatment ^b	–	–	–	0.97	0.60 – 1.58	0.915
	Substance use ^c	–	–	–	2.42	1.31 – 4.47	0.005
	Social network size	–	–	–	0.79	0.77 – 0.81	< 0.001

Significant associations ($p < 0.05$) are bolded^aRefers to the lack of any employment and student status^bRefers to the prior-month history^cRefers to the prior-month history; nicotine and alcohol use were not assessed


Table 4. Differences between chronic and transient loneliness with respect to mental health measures.

Model	Independent variable	Analysis					
		Unadjusted			Adjusted		
		OR	95%CI	p	OR	95%CI	p
Baseline measures of mental health	Depressive symptoms	2.46	1.74 – 3.49	< 0.001	2.20	1.53 – 3.16	< 0.001
	Generalized anxiety	1.05	0.72 – 1.52	0.808	1.07	0.73 – 1.57	0.731
	Social anxiety	2.29	1.80 – 2.91	< 0.001	1.95	1.51 – 2.53	< 0.001
	Paranoid ideation	1.33	1.03 – 1.72	0.030	1.36	1.03 – 1.78	0.028
	Age	–	–	–	1.00	0.99 – 1.01	0.718
	Gender (women)	–	–	–	1.21	0.96 – 1.53	0.111
	Education	–	–	–	1.06	0.89 – 1.25	0.520
	Urbanicity	–	–	–	0.98	0.90 – 1.06	0.566
	Unemployment ^a	–	–	–	0.95	0.73 – 1.22	0.675
	Income	–	–	–	0.94	0.82 – 1.09	0.440
	Psychiatric treatment ^b	–	–	–	0.74	0.46 – 1.19	0.215
	Substance use ^c	–	–	–	0.85	0.51 – 1.42	0.536
	Social network size	–	–	–	0.89	0.87 – 0.91	< 0.001
Follow-up measures of mental health	Depressive symptoms	1.82	1.28 – 2.59	< 0.001	1.78	1.22 – 2.58	0.003
	Generalized anxiety	1.16	0.79 – 1.70	0.444	1.10	0.73 – 1.63	0.657
	Social anxiety	2.68	2.10 – 3.44	< 0.001	2.30	1.77 – 2.99	< 0.001
	Paranoid ideation	1.78	1.37 – 2.32	< 0.001	1.98	1.50 – 2.61	< 0.001
	Age	–	–	–	1.00	0.99 – 1.01	0.727
	Gender (women)	–	–	–	1.18	0.93 – 1.50	0.166
	Education	–	–	–	1.06	0.89 – 1.26	0.495
	Urbanicity	–	–	–	0.98	0.91 – 1.07	0.668
	Unemployment ^a	–	–	–	0.95	0.74 – 1.23	0.695
	Income	–	–	–	0.95	0.82 – 1.10	0.481
	Psychiatric treatment ^b	–	–	–	0.73	0.46 – 1.17	0.195
	Substance use ^c	–	–	–	0.83	0.49 – 1.39	0.474
	Social network size	–	–	–	0.88	0.86 – 0.90	< 0.001

Chronic loneliness was included as reference


Significant associations ($p < 0.05$) are bolded^aRefers to the lack of any employment and student status^bRefers to the prior-month history^cRefers to the prior-month history; nicotine and alcohol use were not assessed

766 Figure 1
767



768
769

770 Figure 2



771