

## NOTES FOR AUTHORS

Papers to be considered for publication should be sent to the Editorial Secretary, The Royal Society of Edinburgh, 22 George Street, Edinburgh EH2 2PQ, Scotland.

A paper by more than one author must be submitted with a statement, signed by each author, to the effect that the paper in its entirety is approved by the joint authors and naming the author who will be responsible for correspondence with the Society.

Authors will receive fifty (50) offprints free of charge, this number to be shared between joint authors. Additional offprints may be obtained, in units of fifty, at a fixed scale of prices given on a form which will be attached to the proof.

In view of the high cost of publication, authors must prepare their papers as concisely as possible. Manuscripts should be submitted in triplicate and preferably should be typewritten on one side of A4 paper, double spaced with adequate margins. Authors are advised to retain a copy of their papers as the Society cannot accept responsibility for any loss.

Every paper must be accompanied by a Synopsis, in general not exceeding two hundred words, which will be printed in small type at the beginning of the paper.

References within the text should be indicated by bold numbers in square brackets, e.g. [2] or [3, p. 167]. For style of references at end of text, see recent issues of *Proceedings A*.

Authors should ensure that punctuation carries through the mathematics in the proper manner. The use of hyphens should be consistent. In the text avoid such abbreviations as: iff, w.r.t., a.e.,  $\forall$ ,  $\exists$ , and thm.

Footnotes should be avoided. Headings should not be underlined. Every effort should be made to avoid complicated subscripts, superscripts, ranges of summation and integration. Horizontal fraction signs should normally be avoided: use either solidus signs / or negative exponents. Replace  $e^{(\dots)}$  by  $\exp[. . .]$  if the expression in parenthesis is complicated. Simple formulae should *not* be displayed unless they require a formula number. Use the prime ' or  $d/dx$ , but preferably not a dot, to denote ordinary differentiation. If possible use subscripts to denote partial differentiation of  $\partial/\partial x$  etc. Bars reaching over several letters should be avoided: use  $\sqrt{()}$  or the exponent  $1/2$  for the square root. Sub-subscripts and super-superscripts should be avoided if possible: bars and other devices over indices cannot be supplied.

Note that confusion very often arises between 1 (one) and  $l$  (ell): 0 (zero) and O (Capital oh):  $\circ$  (composition) and  $o$  (lower case oh):  $x$  and  $\times$ :  $U$  and  $\cup$ :  $c$  and  $\subset$ :  $\in$  (belongs to) and  $\epsilon$  (epsilon):  $\emptyset$  (empty set) and  $\phi$  (phi):  $;$  and comma  $,$ : prime ' and  $^1$ :  $K$  and  $\kappa$ :  $p$  and  $\rho$ :  $w$  and  $\omega$ :  $\sum$  (summation) and  $\Sigma$  (capital sigma):  $\prod$  (product) and  $\Pi$  (capital pi):  $v$  (lower case vee) and  $\nu$  (Greek nu):  $a$  (lower case a) and  $\alpha$  (Greek alpha):  $y$  (lower case y) and  $\gamma$  (Greek gamma). Please provide pencilled indicators in the margin where necessary. Where capitals and lower case of the same shape have to be printed, please indicate accordingly. Show italics by single underlining (except in the formulae which are set up normally in italics), bold face/Clarendon by wavy underlining and Greek by red underlining.

The statement of theorems, lemmas, et cetera, will be printed in italics and should be underlined. In definitions key words only should be in italics.

Equations should be indicated by numbers in parentheses in the right-hand margin.

Proofs of papers will be sent to the author. The cost of *authors' corrections in excess of five per cent* of the printers' charge for the setting of a particular paper will be charged to the author.

### Copyright

© 1981 The Royal Society of Edinburgh and the authors of individual papers.

It is the policy of the Royal Society of Edinburgh not to charge any royalty for the production of a single copy of any one article made for private study or research. Requests for the copying or reprinting of any article for any other purpose should be sent to the Royal Society of Edinburgh, 22/24 George Street, Edinburgh EH2 2PQ

CONTENTS

W. D. EVANS and B. J. HARRIS	
Bounds for the point spectra of separated Dirac operators	1
HEINRICH VOSS	
Positive solutions of superlinear boundary value problems	17
R. W. DICKEY	
Stability theory for quasi-linear wave equations with linear damping	25
RANDY TULER	
Subgroups of $SL_2\mathbb{Z}$ generated by elementary matrices	43
ROBERT M. KAUFFMAN	
A differentiability property and its application to the spectral theory of differential operators	49
CATHERINE BUNDLE	
<i>A priori</i> bounds for a class of nonlinear elliptic equations and applications to physical problems	75
HELMUT HOFER	
Existence and multiplicity result for a class of second order elliptic equations	83
C. M. ELLIOTT and V. JANOVSKÝ	
A variational inequality approach to Hele-Shaw flow with a moving boundary	93
PATRIZIA DONATO, LUCIA MIGLIACCIO and ROSANNA SCHIANCHI	
Semilinear elliptic equations in unbounded domains of $R^n$	109
MAN KAM KWONG and A. ZETTL	
Weighted norm inequalities of sum form involving derivatives	121
C. C. TRAVIS and NORIO YOSHIDA	
Oscillation criteria for third order hyperbolic characteristic initial value problems	135
COLIN MACLACHLAN	
Groups of units of zero ternary quadratic forms	141
J. M. HOWIE	
Some subsemigroups of infinite full transformation semigroups	159
J. M. HOWIE	
A class of bisimple, idempotent-generated congruence-free semigroups	169
B. D. SLEEMAN	
Boundary value problems for the vibrating string equation	185