

New insights on the conservation status of the Endangered coastal endemic plant *Astragalus berytheus* (Fabaceae) in Lebanon

FARAH ABDEL SAMAD, FREDERIC MEDAIL, ALEX BAUMEL
SONJA SILJAK YAKOVLEV, ANTHONY ROUKOZ, GEORGES TOHME
HENRIETTE TOHME and MAGDA BOU DAGHER KHARRAT

Abstract Coastal vegetation includes unique species with adaptations to the specific conditions of this environment, and in the Mediterranean region urban development of coastal areas has severely affected coastal ecosystems. *Astragalus berytheus* Boiss. & Blanche is a narrow endemic plant of the eastern Mediterranean coast, occurring in Lebanon and Palestine/Israel; its Red List status has not been previously assessed. As a result of the effects of urban sprawl in coastal areas *A. berytheus* is one of the most threatened plant taxa in Lebanon. We assess the conservation status of this taxon and propose conservation measures. Only one population of *A. berytheus* is extant in Lebanon, in the Tyre Coast Nature Reserve. We recommend that *A. berytheus* should be categorized as Endangered globally and as Critically Endangered at the national level in Lebanon. Unless adequate conservation measures are implemented *A. berytheus* is likely to go extinct in Lebanon in the near future.

Keywords *Astragalus berytheus*, coastal ecosystem, conservation, Critically Endangered, Eastern Mediterranean, endemic, extinction, habitat degradation

Supplementary material for this article is available at doi.org/10.1017/S0030605320000149

MAGDA BOU DAGHER KHARRAT (Corresponding author, orcid.org/0000-0001-7969-1673), FARAH ABDEL SAMAD*† and ANTHONY ROUKOZ Laboratoire Biodiversité et Génomique Fonctionnelle, Faculté des Sciences, Campus Sciences et Technologies, Université Saint-Joseph, Beirut, Lebanon
E-mail magda.boudagher@usj.edu.lb

FREDERIC MEDAIL† (orcid.org/0000-0002-1429-6661) and ALEX BAUMEL (orcid.org/0000-0003-4245-197X) Institut Méditerranéen de Biodiversité et d'Écologie Marine et Continentale, Aix Marseille Université, CNRS, IRD, Faculté des Sciences et Techniques St-Jérôme, Marseille, France

SONJA SILJAK YAKOVLEV (orcid.org/0000-0002-6247-3351) Université Paris-Sud, Écologie, Systématique, Évolution, CNRS-PSU-AgroParisTech, Orsay Cedex, France

GEORGES TOHME and HENRIETTE TOHME Conseil National de Recherches Scientifiques, CNRS, Beirut, Lebanon

*Also at: Aix Marseille Université, CNRS, IRD, Institut Méditerranéen de Biodiversité et d'Écologie Marine et Continentale

†Also at: Technopôle de l'Environnement Arbois-Méditerranée, Aix-en-Provence, France

Received 3 September 2019. Revision requested 13 November 2019.

Accepted 25 February 2020. First published online 24 November 2020.

Coastal vegetation includes unique species with adaptations to the specific conditions of this environment (Vorontkova et al., 2008). In the Mediterranean region urban development of coastal areas has severely affected coastal ecosystems and led to loss of biodiversity (Doxa et al., 2017). In Lebanon, one of the 10 regional biodiversity hotspots in this region (Médail & Myers, 2004), the increase in the human population and intensive urbanization has destroyed one-third of the 220 km coastline (MOE, 2004).

Astragalus berytheus Boiss. & Blanche (Plate 1) is a coastal species endemic to Lebanon and Palestine/Israel (Tohmé & Tohmé, 2014), and one of the most threatened taxa in Lebanon (MOE, 2004). It occurs in prime tourist areas, such as sandy beaches, which are targeted for the establishment of resorts. These habitats are categorized as highly threatened because of extensive tourism throughout the Mediterranean (IUCN, 1994). Here we assess the conservation status of *A. berytheus*, using the IUCN Red List criteria (IUCN, 2012), and make appropriate recommendations for its conservation.

Information on the historical distribution of *A. berytheus* was gathered from regional floras (Mouterde, 1984; Tohmé & Tohmé, 2014) and herbaria (Royal Botanic Garden Edinburgh Herbarium, Herbarium WU at University of Vienna, and MNHN Paris). During 1994–2018, all sandy shores along the coast of Lebanon were visited by GT and HT to determine the species' occurrence. Historically the species was known from five localities in Lebanon (Fig. 1, Supplementary Table 1).

Three of the five Lebanese populations (Beirut, Ouzai and Khaldeh) were already extinct when we began this study in 2009. The coastal strip was severely damaged by the construction of the highway connecting Beirut to southern Lebanon and the construction of Beirut airport in 1997. The last population in the Beirut area (Bir Hassan) became extinct following construction of a residential building in 2015. This population was in an area of 1,000 m² and was characterized by annual fluctuation in the number of individuals: eight in 2010, 14 in 2013 and five in 2014.

The only extant population of *A. berytheus* in Lebanon is in an area of c. 2,000 m² in Tyre Coast Nature Reserve. In 2013 there were > 20 individuals, bearing healthy fruits and seeds. These were, however, affected by the invasion of camphor-weed *Heterotheca subaxillaris* (Lam.) Britton & Rusby

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

Oryx, 2021, 55(4), 519–521 © The Author(s), 2020. Published by Cambridge University Press on behalf of Fauna & Flora International doi:10.1017/S0030605320000149




PLATE 1 *Astragalus berytheus* in Tyre Nature Reserve: (a) plant photographed in May, (b) flowers and leaves photographed in March, and (c) immature seed pods photographed in April.



FIG. 1 Present and historical distribution of *Astragalus berytheus* on the eastern Mediterranean coast (see Supplementary Table 1 for additional details). The only extant population in Lebanon is in Tyre Coast Nature Reserve (see text for details).

(Family Asteraceae), which originated from the USA; it is an invader of coastal sand dunes (Sternberg, 2016) and was introduced into the Middle East in 1975 (di Castri et al., 1990). It was first observed in the Reserve in 1999 (Tohmé & Tohmé, 2009), and in 2015 it invaded *A. berytheus* habitat. Eradication has, however, allowed *A. berytheus* to recover

(Bou Dagher Kharrat et al., 2016), with 32 individuals recorded in 2016 and 76 individuals bearing fruits in 2019.

We calculated the AOO (area of occupancy) and the EOO (extent of occurrence) of *A. berytheus* using the *Geospatial Conservation Assessment Tool* (Bachman et al., 2011), considering the Lebanese population and the populations in Palestine/Israel (Danin & Fragman-Sapir, 2016; Supplementary Table 1). With an EOO of c. 945 km² and an AOO of c. 36 km², *A. berytheus* should be categorized as Endangered based on criteria B2ab(ii); i.e. with AOO < 500 km² (B2), severely fragmented (a) and projected decline of quality of habitat b(ii).

Nationally in Lebanon, *A. berytheus* should be categorized as Critically Endangered based on criteria A2ac; B1B2ab(i, ii, iii, iv); C2b; i.e. with a population reduction observed of > 80% and where the causes of reduction may not be reversible (A2) and direct observation (a) of decline in AOO and EOO (c), with an EOO < 100 km² (B1) and an AOO < 10 km² (B2) showing only one location (a) and a continuing decline of EOO (i), AOO (ii), quality of habitat (iii), and number of subpopulations (iv), and with < 250 mature individuals in the wild (C2) and extreme fluctuations in the number of mature individuals (b).

Passive habitat protection is not a guarantee of long-term plant conservation (Martinell et al., 2011), and the last *A. berytheus* population in Lebanon would probably have been extirpated if encroachment of the invasive *H. subaxillarisi* had not been halted. Plant species known from only one or two populations, growing in areas with some protection, are known to have gone extinct (Bañares et al., 2003). Appropriate planning of local management actions, with recovery plans for individual species (Fenu et al., 2011; Ballesteros et al., 2013; Cambrollé et al., 2018), are required for successful plant conservation. Environmental monitoring, population viability analysis and population genetics and studies of reproductive biology are required to support the conservation of *A. berytheus*. The micro-reserve concept (Laguna et al., 2001), for the monitoring and in situ conservation of threatened plants in small protected areas, could be appropriate for *A. berytheus* and for other narrow endemic species of Lebanon, such as *Matthiola crassifolia* Boiss. & Gaill. (Bou Dagher Kharrat et al., 2018).

In 2013, 28 pods of *A. berytheus* were collected from the Bir Hassan population (before it vanished), containing a total of 112 seeds, but only nine seeds escaped infestation by a chalcid wasp *Bruchophagus* sp. (Hymenoptera: Eurytomidae) often associated with the seeds of legumes (Burks, 1957). These seeds, together with some seeds collected from the Tyre Coast Nature Reserve in 2011 were stored in the Laboratory of Seed Germination and Conservation–Jouzour Loubnan at Saint-Joseph University, Beirut, Lebanon.

In 2020, 50 plantlets grown from ex situ germination of these seeds will be introduced to the Tyre archaeological

site, 3 km from Tyre Nature Reserve, with similar ecological conditions. As the archaeological site is a UNESCO world heritage site this will facilitate protection and monitoring. The presence of endemic plants on archaeological sites is an opportunity to raise the conservation awareness of visitors (Siljak-Yakovlev et al., 2011).

A national legal framework is required to halt the loss of biodiversity in Lebanon, especially of narrow endemics plant species (Médail & Baumel, 2018), working from the principle that the narrower the range of a taxon the greater the local responsibility (Gauthier et al., 2010). An additional imperative to conserve *A. berytheus* is that it bears the name of Beirut, the nation's capital. If the results of the pilot introduction of the species to the Tyre archaeological site are favourable, future introductions could be considered in other suitable coastal locations.

Acknowledgements This research was supported by a PhD grant from the National Council of Scientific Research of Lebanon and from Critical Ecosystem Partnership Fund grant #108497.

Author contributions Fieldwork: FAS, GT, HT; writing: FAS, FM, MBDK, AB, SSY.

Conflicts of interest None.

Ethical standards This research complied with the *Oryx* guidelines on ethical standards.

References

- BACHMAN, S., MOAT, J., HILL, A., DE LA TORRE, J. & SCOTT, B. (2011) Supporting Red List threat assessments with *GeoCAT*: geospatial conservation assessment tool. In *e-Infrastructures for Data Publishing in Biodiversity Science* (eds V. Smith & L. Penev), pp. 117–126. ZooKeys, Pensoft Publishers, Sofia, Bulgaria.
- BALLESTEROS, M., FORONDA, A., CAÑADAS, E.M., PEÑAS, J. & LORITE, J. (2013) Conservation status of the narrow endemic gypsophile *Ononis tridentata* subsp. *crassifolia* in southern Spain: effects of habitat disturbance. *Oryx*, 47, 199–202.
- BAÑARES, Á., MARRERO, M., CARQUÉ, E. & FERNÁNDEZ, Á. (2003) Plan de recuperación de la flora amenazada del Parque Nacional de Garajonay. La Gomera (Islas Canarias). Germinación y restituciones de *Pericallis hansenii*, *Gonospermum gomerae* e *Ilex Perado* ssp. *Lopezilloi*. *Botanica Macaronésica*, 24, 3–16.
- BOU DAGHER KHARRAT, M., EL ZEIN, H. & ROUHAN, G. (2018) Setting conservation priorities for Lebanese flora—Identification of important plant areas. *Journal for Nature Conservation*, 43, 85–94.
- BOU DAGHER KHARRAT, M., FARHAT, P., DAKIK, N., KAHALE, R., CHAHINE, T. & POTASH, L. (2016) Invasion and management of *Heterotheca subaxillaris* in the Coast Nature Reserve of Tyre, Lebanon. *Conference Poster: 1st Mediterranean Plant Conservation Week “Building a Regional Network to Conserve Plants and Cultural Diversity”* Ulcinj–Montenegro. [dx.doi.org/10.13140/RG.2.2.21409.40808](https://doi.org/10.13140/RG.2.2.21409.40808) [accessed 15 July 2020].
- BURKS, B.D. (1957) A new Bruchophagus from a liliaceous plant with a host plant list for the genus (Hymenoptera: Eurytomidae). *Proceedings of the Entomological Society of Washington*, 59, 273–277.
- CAMBROLLÉ, J., MANCILLA-LEYTÓN, J.M., MUÑOZ-VALLÉS, S., MORENZA-BARRERA, B. & FIGUEROA, M.E. (2018) Exploring the conservation status of an Endangered coastal plant over the last nine years: a case study in Andalusia, Spain. *Journal of Coastal Conservation* 22, 1201–1207.
- DANIN, A. & FRAGMAN-SAPIR, O. (2016) *Flora of Israel Online*. flora.org.il/en/plants [accessed 6 March 2020].
- DI CASTRI, F., HANSEN, A.J. & DEBUSSCHE, M.E. (1990) *Biological Invasions in Europe and the Mediterranean Basin*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- DOXA, A., ALBERT, C.H., LERICHE, A. & SAATKAMP, A. (2017) Prioritizing conservation areas for coastal plant diversity under increasing urbanization. *Journal of Environmental Management*, 201, 425–434.
- FENU, G., MATTANA, E. & BACCHETTA, G. (2011) Conservation of endemic insular plants: the genus *Ribes* L. (Grossulariaceae) in Sardinia. *Oryx*, 46, 219–222.
- GAUTHIER, P., DEBUSSCHE, M. & THOMPSON, J. (2010) Regional priority setting for rare species based on a method combining three criteria. *Biological Conservation*, 143, 1501–1509.
- IUCN (1994) *Guidelines for Protected Area Management Categories*. IUCN, Gland, Switzerland and Cambridge, UK.
- IUCN (2012) *IUCN Red List Categories and Criteria: Version 3.1*. 2nd edition. Gland, Switzerland and Cambridge, UK.
- LAGUNA, E., BALLESTER, G., FABREGAT, C., OLIVARES, A., SERRA, L., DELTORO, V. et al. (2001) Plant micro-reserves: a new model of micro protected areas, Spain. *Re-Introduction News*, 20, 19–21.
- MARTINELL, M.C., LOPEZ-PUJOL, J., BLANCHÉ, C., MOLERO, J. & SÁEZ, L. (2011) Conservation assessment of *Aquilegia paui* (Ranunculaceae): a case study of an extremely narrow endemic. *Oryx*, 45, 187–190.
- MÉDAIL, F. & BAUMEL, A. (2018) Using phylogeography to define conservation priorities: the case of narrow endemic plants in the Mediterranean Basin hotspot. *Biological Conservation*, 224, 258–266.
- MÉDAIL, F. & MYERS, N. (2004) Mediterranean basin. In *Hotspots Revisited: Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions, CEMEX (Monterrey)* (eds R.A. Mittermeier, P. Robles Gil, M. Hoffmann, J. Pilgrim, T. Brooks, C.G. Mittermeier, J. Lamoreaux & G.A.B. da Fonseca), pp. 144–147. Conservation International, Washington, DC, USA and Agrupación Sierra Madre, Mexico.
- MOE (2004) *Final Report Biodiversity Assessment and Monitoring*. The Protected Areas/Lebanon LEB/95/G31, Tyre Beach Nature Reserve, Tyre, Lebanon.
- MOUTERDE, P. (1984) *Nouvelle Flore du Liban et de la Syrie*. Editions de l'Imprimerie Catholique, Beirut, Lebanon.
- SILJAK-YAKOVLEV, S., TAN, K., TSOUNIS, G. & TSOUNIS, L. (2011) Genome size and chromosome number of *Micromeria acropolitana* (Lamiaceae), a steno-endemic from Greece. *Phytologia Balcanica*, 17, 201–204.
- STERNBERG, M. (2016) From America to the Holy Land: disentangling plant traits of the invasive *Heterotheca subaxillaris* (Lam.) Britton & Rusby. *Plant Ecology*, 217, 1307–1314.
- TOHMÉ, G. & TOHMÉ, H. (2009) Espèces nouvelles du Liban et redescription d'espèces endémiques. *Lebanese Science Journal*, 10, 125–130.
- TOHMÉ, G. & TOHMÉ, H. (2014) *Illustrated Flora of Lebanon, National Council for Scientific Research*. National Council for Scientific Research, Beirut, Lebanon.
- VORONKOVA, N.M., BURKOVSKAYA, E.V., BEZDELEVA, T.A. & BURUNDUKOVA, O.L. (2008) Morphological and biological features of plants related to their adaptation to coastal habitats. *Russian Journal of Ecology*, 39, 1–7.