



Original Article

Cite this article: Redfern JM, Hawkes S, Bryan A, Cullington D, and Ashrafi R (2024) The oxygen uptake efficiency slope in adults with CHD: group validity. *Cardiology in the Young* **34**: 1285–1294. doi: [10.1017/S1047951123004365](https://doi.org/10.1017/S1047951123004365)

Received: 26 February 2023

Revised: 2 October 2023

Accepted: 2 December 2023

First published online: 15 January 2024

Keywords:

Oxygen uptake efficiency slope; CHD; adults; exercise testing

Corresponding author:

J. M. Redfern; Email: james.redfern4@nhs.net

¹Countess of Chester Hospital NHS Foundation Trust, Chester, UK; ²Liverpool Heart and Chest Hospital NHS Foundation Trust, Liverpool, LP, UK and ³Manchester University NHS Foundation Trust, Manchester, GM, UK

Abstract

The maximal oxygen uptake ($V_{O2\max}$) is a well-validated measure of cardiorespiratory function that is calculated during a maximal cardiopulmonary exercise test. $V_{O2\max}$ enables physicians to objectively assess cardiopulmonary function to aid in decision-making for patients with CHD. A significant proportion of these patients however are unable to achieve a maximal exercise test, and as such, there is a need for reliable submaximal predictors of cardiorespiratory reserve.

The oxygen uptake efficiency slope represents a measure of how effectively oxygen is extracted from the lungs and taken into the body and can be calculated from a submaximal exercise test. Its reliability as a predictor of cardiorespiratory reserve has been validated in various patient populations, but there is limited evidence for its validity in adult patients with CHD.

Retrospective analysis of cardiopulmonary exercise test data in 238 consecutive patients with CHD who completed a maximal cardiopulmonary exercise test at our tertiary cardiology centre demonstrated a strong correlation between peak V_{O2} and the oxygen uptake efficiency slope (0.936). A strong correlation with peak V_{O2} was also demonstrated when oxygen uptake efficiency slope was calculated at ventilatory anaerobic threshold ($OUES_{VAT}$), 75% ($OUES_{75}$), and 90% ($OUES_{90}$) of the test (0.833, 0.905, 0.927 respectively).

In adult patients with CHD who are unable to complete a maximal cardiopulmonary exercise test, the oxygen uptake efficiency slope is a reliable indicator of cardiopulmonary fitness which correlates strongly with peak V_{O2} at or beyond the ventilatory anaerobic threshold. Further research is required to validate the findings in patients with less common anatomies and to assess the relationship between the oxygen uptake efficiency slope and mortality.

CHD represents around 0.8% of all live births in developed countries.¹ As a result of improving surgical techniques, post-operative care, and medical therapy, over 80% now survive to adulthood.² Unfortunately, these patients still may have a greater degree of morbidity and exercise limitation than their healthy counterparts.³ They may also have a distorted perception of their functional limitation and can struggle to recognise further deterioration as a result of long-term adaptation to reduced physical activity.^{3–5}

Physicians caring for patients with CHD are often faced with the dilemma of trying to predict untoward events in patients who remain clinically well despite deteriorating imaging results, or conversely, trying to explain worsening symptoms despite stable investigation results. Cardiopulmonary exercise testing enables physicians to objectively assess and monitor cardiopulmonary function and has been used for risk stratification to aid decision-making in a non-invasive, reproducible manner since the 1930s.⁶ It has also been demonstrated that in patients with CHD, lower exercise capacity is associated with a worse prognosis.^{7,8} Maximal oxygen uptake, defined as the point at which oxygen uptake plateaus despite increasing workload is a well-validated measure of cardiorespiratory function,⁹ and in various patient groups predicts prognosis as an aid to decision-making.¹⁰ A true plateau however is difficult to achieve, is effort dependant, and may be influenced by patient motivation and by the team performing the test.¹¹ There is variation in achievement by lesion as evidenced by the study from Buys et al where an RER of > 1.10 (indicative of a maximal test)¹² was achieved in 78.7% of patients with coarctation, and around 60% in those with tetralogy of Fallot, transposition of the great arteries, and univentricular heart.¹³ The inability to achieve a maximal exercise test is a significant issue in patients with CHD, and as such, there is a need for a reliable predictor of cardiorespiratory reserve that can be achieved using a submaximal test.

Several submaximal indices to predict maximal oxygen uptake have been investigated but are associated with various limitations. Ventilatory anaerobic threshold is not always identifiable and is subject to interobserver and intraobserver variability.^{14,15} The regression line showing the relationship between minute ventilation and carbon dioxide production has been demonstrated to be inversely correlated with the maximal oxygen uptake; however, the correlation is weak ($r = -0.56$).¹⁶ Buller et al proposed an additional method to predict cardiorespiratory functional

reserve independent of exercise duration using a quadratic function of the carbon dioxide production and oxygen uptake termed the extrapolated maximal oxygen consumption;¹⁷ however, this has not been validated by other investigators and so remains of limited clinical validity.

In 1996, Baba et al described the oxygen uptake efficiency slope as a novel method of estimating cardiorespiratory functional reserve from a submaximal exercise test and demonstrated a strong positive correlation between the oxygen uptake efficiency slope and the maximal oxygen uptake in a group of young patients.¹¹

The oxygen uptake efficiency slope is the slope of the regression line of the oxygen uptake on a logarithmically adjusted minute ventilation as measured during incremental exercise.¹⁸ It represents a measure of how effectively oxygen is extracted from the lungs and taken into the body. Oxygen uptake efficiency slope is represented as a slope whereby a steeper gradient represents a greater uptake in oxygen for any given increase in ventilation which is indicative of a more efficient cardiorespiratory system (Fig. 1). Put simply, oxygen uptake efficiency slope is the absolute increase in oxygen uptake associated with a 10-fold increase in ventilation. The benefit of the oxygen uptake efficiency slope over other submaximal indices may be that the slope incorporates cardiovascular, musculoskeletal, and respiratory function into a single measure as physiologically it is based on both the development of metabolic acidosis and lung perfusion.^{11,19} The oxygen uptake efficiency slope is best described by a single exponential function:

$$\dot{V}O_2 = a \log_{10}(\dot{V}_E / \dot{V}_{E_{rest}}) + \dot{V}O_{2_{rest}}$$

The reliability of the oxygen uptake efficiency slope as a predictor of cardiorespiratory reserve in submaximal exercise tests has been validated in healthy adults,^{20,21} children with CHD,^{11,22} and a variety of adult cardiac patients including those with heart failure,^{23,24} coronary artery disease,²⁵ and pulmonary hypertension.^{26–28} To date, there is limited and equivocal evidence for the role of the oxygen uptake efficiency slope as a submaximal predictor of exercise capacity in adult patients with CHD.^{13,29} Giardini et al in 2008 demonstrated good correlation of the with the last 50% of the entire exercise duration (oxygen uptake efficiency slope 50–100) and maximal oxygen uptake in a group of Fontan ($n = 23$) and Mustard/Senning patients ($n = 30$). In the Fontan group, the oxygen uptake efficiency slope 50 (first 50% of the test) differed significantly from the oxygen uptake efficiency slope 50–100, with the difference appearing most marked in the cyanotic Fontan group.²⁹ Buys et al published a larger study including patients with tetralogy of Fallot, aortic coarctation, univentricular heart and transposition of the great arteries and reported good correlation between oxygen uptake efficiency slope and oxygen uptake efficiency slope 75 (where data points from only the first 75% of the exercise test were analysed) in the aortic coarctation group, but poor correlation in all other patient groups.¹³ Previously, there has been discrepancy about the decimalisation of the y -intercept and the unit of oxygen uptake efficiency slope. We suggest calculating oxygen uptake efficiency slope using oxygen uptake expressed in litres/minute and that the oxygen uptake efficiency slope itself is unitless.¹⁸

This study seeks to assess the validity of the oxygen uptake efficiency slope as a submaximal predictor of maximal oxygen uptake in patients with all the major adult CHD diagnosis groups and establish a reproducible method by which it can be calculated and thus be used as the standard industry-wide method.

Additionally, we will evaluate the correlation between oxygen uptake efficiency slope at ventilatory anaerobic threshold and maximal oxygen uptake to add more direct clinical application of the oxygen uptake efficiency slope in this group of patients.

Methods

Participants

As part of a service development project to establish a departmental standard, we retrospectively analysed the exercise date of all patients with CHD referred for routine cardiopulmonary exercise testing in the Department of Pulmonary Function, Liverpool Heart, and Chest Hospital between 2019–2020. CHD diagnoses were split into 11 subgroups as defined by Kempny et al, 2012.³⁰ Ethics approval for this project as a service development project was based on retrospective analysis of data acquired in routine care.

Anthropometric

Anthropometric measurements were taken in all participants including body mass (kg) and height (m) using electronic scales and a stadiometer (Seca GmbH & CO, Germany). Body mass index and body surface area were calculated.

Spirometry

All patients performed resting spirometry in accordance with American Thoracic Society guidelines for acceptability and repeatability.³¹ Measurements of forced vital capacity, forced expiratory volumes in 1 s, and the forced expiratory volumes in 1 s / forced vital capacity ratio were calculated. Predicted values for spirometry used were derived from the Global Lung Index 2012.³²

Cardiopulmonary exercise testing procedure

Patients were familiarised with equipment and protocol. Prior to each test, equipment was calibrated as per manufacturer's guidelines using 3L reference syringe for volume calibration and a two-point calibration of gases. All patients performed cardiopulmonary exercise testing using an electronically braked cycle ergometer (Ergoselect 100, ergoline GmbH, Germany). Cardiopulmonary exercise test followed standardised protocol¹² consisting of 3 minutes of rest, 3 minutes unloaded cycling at 60 rpm, followed by incremental uniformed increase in load 5–30 W/min, load calculated using Wasserman formula³³ with the aim of eliciting maximum performance in 8–12 minutes of exercise. During cardiopulmonary exercise testing, patients were required to maintain revolutions of 60–80 r.p.m. and were given verbal encouragement throughout. The test ceased when patients achieved volitional exhaustion and were either unable to maintain the required cadence or all the identifiers of a maximal exercise test were present (respiratory exchange ratio > 1.10, heart rate > 90% age-predicted maximum and a sustained plateau of oxygen uptake was achieved). As a maximal test was required to develop a correlation, tests whereby the respiratory exchange ratio did not meet the threshold for maximal exercise were excluded from this study (respiratory exchange ratio < 1.10), $n = 7$.

Standard 12 lead electrocardiograms (AMEDTEC Medizintechnik GmbH, Germany) and pulse oximetry were continuously monitored throughout cardiopulmonary exercise testing. Blood pressure (automatic cuff) was recorded regularly throughout exercise, at peak and into recovery for assessment of haemodynamic recovery. Carbon

dioxide (carbon dioxide production ml/min), oxygen uptake (ml/min), minute ventilation (L/min), and respiratory rate were measured continuously via breath-by-breath gas analysis. Patients breathed through a face mask (Hans Rudolph, Kansas City, MO) connected to a metabolic cart system (Geratherm GmbH, Germany). Expired gas was passed through a flowmeter and rapidly responding oxygen and carbon dioxide analysers connected to a computer, and derived variables such as respiratory exchange rate, oxygen uptake/heart rate, and ventilatory equivalents (minute ventilation/carbon dioxide production, minute ventilation/oxygen uptake) were obtained throughout. Maximal oxygen uptake was determined as the highest 30 s average of oxygen uptake. Remaining peak parameters were calculated during this time period. Ventilatory anaerobic threshold was calculated by the V-slope method, and minute ventilation/carbon dioxide production was determined at ventilatory anaerobic threshold.¹² Cardiopulmonary exercise testing resting, predicted and peak data were recorded against known predicted equations³⁴ and reported into electronic patient record (Allscripts, Chicago, IL), whereby additional calculations of breathing reserve (forced expiratory volumes in 1 s x40 – peak minute ventilation), ventilatory anaerobic threshold as a percentage of achieved and predicted workload were calculated.^{31,34}

Oxygen uptake efficiency slope calculation

Oxygen uptake efficiency slope is best expressed by the following exponential function:

$$VO_2 = a \log_{10}(V_E/V_{E_{rest}}) + VO_{2_{rest}}$$

Semi-log transformation of the X-axis demonstrates a linear relationship between oxygen consumption and minute ventilation whereby the steeper slope represents improved oxygen uptake with exercise (Fig. 1).

To evaluate its usefulness in the CHD population, oxygen uptake efficiency slope was calculated using breath-by-breath data (including the last minute of resting data, having allowed for a mask familiarisation period). To further assess its utility as an index derived from submaximal cardiopulmonary exercise testing, the oxygen uptake efficiency slope was also calculated from data collected during the first 75 and 90% of exercise duration and correlated against maximal oxygen uptake (L/min). This was achieved by removing all data points beyond 75 and 90% of the total exercise time, respectively, and recalculating the oxygen uptake efficiency slope based on the remaining data. Ventilatory anaerobic threshold was calculated using the V-slope method and then used to calculate oxygen uptake efficiency slope at ventilatory anaerobic threshold by removing all subsequent data points and recalculating the oxygen uptake efficiency slope.

Statistical analysis

The relationships between oxygen uptake efficiency slope from the first 75% of the test, the oxygen uptake efficiency slope from the first 90% of the test, the oxygen uptake efficiency slope from the entire test, and the oxygen uptake efficiency slope at ventilatory anaerobic threshold with each of maximal oxygen uptake, oxygen uptake at lactate threshold, and ventilatory equivalents for carbon dioxide at lactate threshold and slope were investigated using either Pearson's product-moment correlation coefficient or Spearman's rho correlation coefficient. These tests were performed for the whole patient cohort and the subset patient cohorts of tetralogy of




Figure 1. Relationship between the oxygen uptake and the minute ventilation during a cardiopulmonary exercise test in a healthy 27-year-old woman, and in sex and age-matched patients with transposition of the great arteries and Fontan circulation. The values of the oxygen uptake efficiency slope are 2.49, 1.50, and 0.96, respectively, and the data are presented as linear (top graph) and semi-log plots of the x-axis (bottom graph). TGA: transposition of the great arteries, VE: minute ventilation, VO₂: oxygen uptake.

Fallot, valvular, transposition of the great arteries and Fontan. Preliminary analyses were performed to ensure no violation of the assumptions of linearity, homoscedasticity and normality. Partial correlation coefficients were used to explore the relationships as above while controlling for body surface area. Differences between the groups were investigated using Student's t-test or Mann-Whitney U tests after checking for normality. All analyses were done using IBM SPSS Statistics v23.

Results

Patients

We analysed the results of 238 consecutive patients (age 33.5 ± 11.7 , 44% female) who completed a maximal cardiopulmonary exercise test. The cohort included 10 distinct anatomical groups of patients plus a further mixed group of complex anatomy that included Eisenmenger's, anomalous pulmonary venous drainage, truncus, and atrioventricular septal defects. Anthropometric data are summarised in Table 1.

Table 1. Characteristics of study participants.

	Sex (m/f)	Age (years)	Height (cm)	Weight (kg)	BMI (kg/m ²)	BSA (m ²)
Total n = 238	125 / 113	33.2 (11.4)	169 (10)	75.3 (18.1)	26.2 (5.6)	1.87 (0.25)
TGA Arterial Switch n = 10	9 / 1	25.9 (4.7)	173.8 (10.2)	74.7 (11.7)	24.9 (4.8)	1.89 (0.2)
Valvular n = 55	25 / 30	33.0 (11.5)	169.3 (10.4)	77.41 (17.5)	26.9 (5.5)	1.9 (0.2)
CoA n = 10	9 / 1	34.1 (8.5)	174.8 (6.7)	78.8 (19.1)	25.8 (6.6)	1.94 (0.2)
ASD n = 7	1 / 6	38 (13.3)	164.1 (5.1)	76.1 (18.3)	28.1 (6.1)	1.85 (0.2)
VSD n = 3	2 / 1	44.0 (14.9)	169.67 (12.7)	94.0 (24.9)	32.0 (4.7)	2.09 (0.3)
ToF n = 53	24 / 29	32.55 (12.4)	167.2 (10.8)	74.7 (20.3)	26.5 (5.8)	1.85 (0.3)
TGA Atrial Switch n = 30	23 / 7	37.6 (4.4)	171.62 (6.1)	80.83 (16.0)	27.38 (5.1)	1.96 (0.2)
ccTGA n = 9	7 / 2	43.4 (13.5)	171.1 (10.6)	84.0 (17.1)	28.6 (5.0)	1.9 (0.2)
Ebstein's Anomaly n = 10	5 / 5	40.1 (13.9)	168.70 (7.4)	67.10 (9.2)	23.49 (2.3)	1.77 (0.1)
Fontan n = 31	18 / 13	26.5 (5.6)	167.8 (9.9)	66.6 (14.9)	23.6 (4.7)	1.75 (0.2)
Complex n = 20	10 / 10	31.43 (12.0)	168.21 (9.3)	73.40 (17.1)	25.76 (5.1)	1.84 (0.2)

Exercise test characteristics

All patients achieved a maximal test (respiratory exchange ratio > 1.10 and HR > 90% age-predicted maximum). The results are shown in Table 2. Four minor adverse events were recorded. Three patients experienced vasovagal syncope on completion of the test, and another was admitted overnight for treatment of sustained atrial arrhythmia developed post-test.

Relationship of maximal oxygen uptake with oxygen uptake efficiency slope and minute ventilation/carbon dioxide production

In total, 238 patient tests were included in the full patient cohort and n = 53,55,30,31, respectively, for the tetralogy of Fallot, valvular, transposition of the great arteries and Fontan sub-patient groups. There was a strong positive correlation between maximal oxygen uptake and each of oxygen uptake efficiency slope from the first 75% of the test, the oxygen uptake efficiency slope from the first 90% of the test, the oxygen uptake efficiency slope from the entire test, and the oxygen uptake efficiency slope at ventilatory anaerobic threshold for all patient groups as shown in Table 3 and Figure 2. The only exception was a moderate correlation between maximal oxygen uptake and oxygen uptake efficiency slope at ventilatory anaerobic threshold in the Fontan patient subgroup (+0.678). Higher maximal oxygen uptakes were associated with high levels of oxygen uptake efficiency slope from the first 75% of the test, the oxygen uptake efficiency slope from the first 90% of the test, the oxygen uptake efficiency slope from the entire test, and the oxygen uptake efficiency slope at ventilatory anaerobic threshold for the total patient cohort (Fig. 3). The maximum Pearson correlation was the oxygen uptake efficiency slope from the entire test of + 0.943 for the tetralogy of Fallot cohort and the lowest + 0.678 in the Fontan group, oxygen uptake efficiency slope at ventilatory anaerobic threshold. The partial correlation used to explore the relationship between maximal oxygen uptake and each of oxygen uptake efficiency slope from the first 75% of the test, the oxygen uptake efficiency slope from the first 90% of the test, the oxygen uptake efficiency slope from the entire test, and the oxygen uptake efficiency slope at ventilatory anaerobic threshold

when controlling for body surface area was lower in each case but retained the strong positive correlations for the total cohort and each of the patient subgroups (Fig. 4). This was supported when the analyses were repeated using Bongers' method which divided the variables by body surface area rather than using the partial correlation method to examine the covariate.²²

For comparison with the above, the analyses were repeated to compare the oxygen uptake efficiency slope variables with each of the cardiopulmonary exercise testing variables, ventilatory anaerobic threshold, and minute ventilation/carbon dioxide production (both slope and at anaerobic threshold) in all the patient groups. Similar to the above, there was a strong positive correlation with ventilatory anaerobic threshold and each of oxygen uptake efficiency slope from the first 75% of the test, the oxygen uptake efficiency slope from the first 90% of the test, the oxygen uptake efficiency slope from the entire test, and the oxygen uptake efficiency slope at ventilatory anaerobic threshold in all groups albeit lower coefficients than when comparing with maximal oxygen uptake. Controlling for body surface area had little effect on the strength of the relationship with ventilatory anaerobic threshold in the groups, it caused slightly more of a reduction in the coefficients in the tetralogy of Fallot group than the others. When comparing minute ventilation/carbon dioxide production at anaerobic threshold and the oxygen uptake efficiency slope variables the correlation was a weak, negative association being 0.501,0.503,0.502,0.425, respectively, for oxygen uptake efficiency slope from the first 75% of the test, the oxygen uptake efficiency slope from the first 90% of the test, the oxygen uptake efficiency slope from the entire test, and the oxygen uptake efficiency slope at ventilatory anaerobic threshold in the total patient cohort, which was improved slightly when replaced with minute ventilation/carbon dioxide production slope and with a similar effect of the covariate body surface area (Table 3). The subgroups showed a similar pattern with the latter.

In the total patient cohort, there were no significant differences between the means of any one of the variables, oxygen uptake efficiency slope from the first 75% of the test, the oxygen uptake efficiency slope from the first 90% of the test,

Table 2. Exercise test characteristics.

	Total n = 238	TGA arterial switch n = 10	Valvular n = 55	CoA n = 10	ASD n = 7	VSD n = 3	ToF n = 53	TGA atrial switch n = 30	ccTGA n = 9	Ebstein's anomaly n = 10	Fontan n = 31	Complex n = 20
Peak work (watts)	136.12 (43.7)	181.6 (34.8)	149.51 (49.0)	160.9 (39.8)	105.4 (18.5)	164.67 (48.6)	133.32 (37.9)	132.03 (33.4)	131.89 (44.6)	125.00 (28.6)	117.5 (35.1)	136.71 (40.9)
VO ₂ /WR	9.76 (1.9)	10.49 (1.0)	10.23 (1.6)	10.02 (2.0)	10.57 (1.5)	11.63 (1.6)	10.01 (1.4)	9.51 (2.1)	9.96 (2.4)	8.36 (2.0)	9.17 (2.0)	9.11 (1.8)
RER peak (VCO ₂ /VO ₂)	1.28 (0.1)	1.32 (0.1)	1.29 (0.1)	1.31 (0.1)	1.25 (0.1)	1.25 (0.1)	1.31 (0.1)	1.26 (0.1)	1.34 (0.1)	1.27 (0.1)	1.24 (0.1)	1.28 (0.1)
SPO ₂ Rest (%)	98.48 (3.4)	99.4 (1.3)	99.75 (0.6)	99.8 (0.6)	99.29 (1.4)	99.67 (0.5)	99.58 (0.9)	99.10 (1.2)	99.33 (1.3)	98.40 (3.3)	92.35 (5.3)	98.86 (2.6)
SPO ₂ peak (%)	96.62 (5.6)	98.8 (1.3)	98.7 (1.9)	97.9 (2.7)	98 (1.6)	98.0 (0.8)	98.60 (2.5)	96.30 (4.6)	99.22 (1.6)	94.90 (8.5)	88.45 (7.3)	96.86 (4.7)
VO ₂ peak (ml/min/kg)	22.9 (6.9)	30.51 (6.4)	24.7 (7.4)	26.41 (7.4)	20.43 (7.2)	24.13 (2.0)	23.24 (5.7)	20.55 (5.8)	20.57 (6.0)	21.71 (6.0)	21.52 (6.0)	21.79 (6.1)
VO ₂ Peak % predicted	70.1 (16.1)	75.3 (8.7)	78.6 (15.0)	72.7 (17)	76 (5.7)	93.67 (6.0)	73.02 (13.9)	63.53 (13.9)	68.33 (13.1)	68.20 (19.6)	56.97 (13)	65.36 (12.6)
VO ₂ peak (ml/min)	1696.44 (549.1)	2232.3 (382.2)	1883.29 (607.2)	2051.1 (633.2)	1459.43 (327.8)	2313.67 (758.3)	1690 (456.7)	1629.37 (431.4)	1696.33 (487)	1486.70 (433.1)	1399.03 (373.6)	1559.07 (458.2)
AT (ml/min)	1114.41 (338.4)	1360 (309.5)	1195.49 (404.9)	1292.4 (390.9)	985.14 (202.1)	1499.33 (572.5)	1085.34 (280.7)	1122.33 (252.7)	1077.44 (296.1)	1027.10 (230.6)	970.1 (245.3)	1074.07 (282.7)
VE (L/min)	85.79 (25.8)	112.7 (13.6)	87.78 (29.0)	98 (23.6)	78.43 (19.6)	113.33 (36.0)	83.21 (22.6)	89.27 (24.7)	91 (25.9)	78.20 (15.8)	77.52 (18.0)	84.93 (30.2)
VE/VCO ₂	30.8 (6.3)	27.71 (2.4)	28.00 (3.7)	27.93 (5.4)	31.59 (5.4)	29.03 (4.2)	28.80 (3.3)	33.20 (6.7)	29.43 (4.1)	34.18 (8.3)	34.8 (5.0)	31.52 (4.6)
HR peak (beats/min)	160.63 (27.0)	175.9 (10.5)	166.47 (23.3)	166.6 (10.0)	148.14 (32.8)	170.0 (19.3)	164.09 (23.8)	158.80 (28.9)	144.56 (22.5)	160.70 (28.0)	151.42 (30.6)	163.07 (29.0)
OUES 100 (L/min)	1.65 (0.5)	2.12 (0.5)	1.82 (0.6)	1.96 (0.6)	1.40 (0.3)	2.04 (0.8)	1.63 (0.4)	1.57 (0.4)	1.68 (0.4)	1.43 (0.4)	1.46 (0.4)	1.54 (0.3)
OUES 90 (L/min)	1.69 (0.5)	2.15 (0.4)	1.85 (0.6)	2.01 (0.7)	1.43 (0.3)	2.01 (0.8)	1.65 (0.4)	1.61 (0.4)	1.73 (0.5)	1.48 (0.4)	1.51 (0.4)	1.58 (0.3)
OUES 75 (L/min)	1.65 (0.5)	2.06 (0.3)	1.82 (0.6)	1.97 (0.6)	1.39 (0.3)	1.99 (0.8)	1.60 (0.4)	1.62 (0.4)	1.70 (0.4)	1.45(0.3)	1.49 (0.4)	1.56 (0.4)
OUESVAT (L/min)	1.48 (0.5)	1.83 (0.4)	1.62 (0.6)	1.74 (0.5)	1.26 (0.3)	1.81 (0.8)	1.40 (0.4)	1.43 (0.4)	1.41 (0.3)	1.44 (0.4)	1.34 (0.4)	1.34 (0.3)

Table 3. Relationships between cardiopulmonary exercise testing and oxygen uptake efficiency slope variables for the combined and separate patient cohorts.

Patient group	CPET parameter for correlation	OUES100 (R, p) (Partial correlation control for BSA)	OUES90 (R,p) (partial correlation control for BSA)	OUES75 (R,p) (partial correlation control for BSA)	OUESVAT (R,p) (partial correlation control for BSA)
Combined (n = 238)					
	VO2 peak	+0.937 (+.914) ¹	+0.926 (+ 0.90) ¹	+0.901 (+.865) ¹	+0.830 (+.768) ¹
	VE/VCO2 LT	-0.501 (-.496) ²	-.503 (-.486) ²	-.502 (-.489) ²	-0.425 (-.403) ²
	VAT	+0.892 (+.852) ¹	+.90 (+.864) ¹	+.903 (+.868) ¹	+.900 (+.863) ¹
	VE/VCO2 Slope	-0.559 (-.540) ²	-.571 (-0.5582)	-0.575 (-.564) ²	-0.500 (-.484) ²
TOF (n = 53)					
	VO2 peak	+0.943 (+.904) ¹	+0.941 (+0.901) ¹	+.905 (+.840) ¹	+.824 (+.686) ¹
	VE/VCO2 LT	-0.2254, p = 0.07 (-0.361, p = .009) ¹	-0.282, p = 0.044 (-0.409, p = .003) ¹	-0.276, p = 0.045 (-0.424, p = .002) ¹	-0.216, p = .121 (-0.339, p = .014) ¹
	VAT	+.871 (+.767) ¹	+0.863 (+.750) ¹	+0.852 (+.727) ¹	+0.808 (+.636) ¹
	VE/VCO2 Slope	-.416, p = 0.002 (-0.448, p = .001) ¹	-0.453, p = .001 (-.503) ¹	-0.443, p = .001 (-.508) ¹	-0.377, p = .005 (- 0.413, p = .002) ¹
Valvular (n = 55)					
	VO2 peak	+0.928 (+.899) ¹	+0.916 (+.882) ¹	+0.902 (+.861) ¹	+.840 (+.776) ¹
	VE/VCO2 LT	-0.411, p = 0.002 (-0.405, p = .002) ¹	-0.435, p = .001(-0.439, p = .001) ¹	-0.451, p = .001 (-0.452, p = .001) ¹	-0.450, p = 0.001 (-0.445, p = .001) ¹
	VAT	+0.919 (+.884) ¹	+0.927 (+.897) ¹	+0.932 (+.903) ¹	+0.939 (+.914) ¹
	VE/VCO2 Slope	-0.486 (-0.439, p = .001) ¹	-0.533 (-.501) ¹	-0.550 (-.519) ¹	-.542 (-.505) ¹
TGA (n = 30)					
	VO2 peak	+.894 (+.874) ¹	+0.856 (+0.834) ¹	+.822 (+.799) ¹	+0.730 (.694) ¹
	VE/VCO2 LT	-0.586, p = 0.001, (-0.465, p = .011) ²	-0.60, p = 0.001 (-0.468,p = .01) ²	-0.492, p = 0.006 (-0.479, p = .009) ²	-0.410, p = 0.024 (-0.419, p = 0.023) ²
	VAT	+0.842 (+.826) ¹	+0.871 (+.858) ¹	+0.876 (+.863) ¹	+0.852 (+.837) ¹
	VE/VCO2 Slope	-0.629 (-0.548, p = .002) ²	-0.598 (-0.547, p = .002) ²	-0.526, p = 0.003 (-0.563, p = .001) ²	-0.463, p = 0.010 (-0.517, p = .004) ²
Fontan (n = 31)					
	VO2 peak	+0.938 (+.930) ¹	+0.896 (+.834) ¹	+0.806 (+.782) ¹	+0.678 (+.648) ¹
	VE/VCO2 LT	-0.371, p = .04 (- 0.363, p = .05) ¹	-0.355, p = 0.05 (0.345, n.s) ¹	-0.419, p = 0.019 (-0.414, p = .023) ¹	-0.314, n.s (-0.301, n.s) ¹
	VAT	+0.812 (+.792) ¹	+0.851 (+.836) ¹	+0.866 (+.852) ¹	+0.862 (+.851) ¹
	VE/VCO2 Slope	-0.479, p = 0.006 (-0.471, p = .009) ¹	-0.486, p = 0.006 (-0.476, p = .008) ¹	-0.547, p = 0.001 (-0.542, p = .002) ¹	-0.463, p = 0.009 (-0.452, p = .012) ¹

CPET = cardiopulmonary exercise testing; OUES100 = oxygen uptake efficiency slope from the entire test; OUES90 = oxygen uptake efficiency slope from 90% of the test; OUES75 = oxygen uptake efficiency slope from 75% of the test; OUESVAT = oxygen uptake efficiency slope at ventilatory anaerobic threshold; BSA = body surface area; TOF = tetralogy of Fallot; TGA = transposition of the great arteries; VO2 peak = maximum oxygen uptake; VE = minute ventilation; CO2 = carbon dioxide production; LT = lactate threshold; VAT = ventilatory anaerobic threshold.

The partial correlation control for body surface area is shown in brackets after each association.

The coefficients of correlation are shown for each comparison.

¹Pearson's correlation coefficient or

²Spearman's Rho coefficient, p < 0.0005 unless otherwise stated.




Figure 2. Graphical representation of oxygen uptake efficiency slope calculation from the entire test, from first 90%, from first 75% and at ventilatory anaerobic threshold. VE; minute ventilation, VO₂; oxygen uptake, OUES₁₀₀; oxygen uptake efficiency slope from the entire test, OUES₉₀; oxygen uptake efficiency slope from first 90%, OUES₇₅; oxygen uptake efficiency slope from first 75%, OUES_{VAT}; oxygen uptake efficiency slope at ventilatory anaerobic threshold.




Figure 3. Relationship between peak VO₂ and OUES₁₀₀ (top left); peak VO₂ and OUES₉₀ (top right); peak VO₂ and OUES₇₅ (bottom left); peak VO₂ and OUES_{VAT} (bottom right). Peak VO₂; maximum oxygen uptake, OUES₇₅; oxygen uptake efficiency slope from the first 75% of the test, OUES₉₀; oxygen uptake efficiency slope from the first 90% of the test, OUES₁₀₀; the oxygen uptake efficiency slope from the entire test, OUES_{VAT}; oxygen uptake efficiency slope at ventilatory anaerobic threshold.




Figure 4. Correlation of OUES₁₀₀ with OUES₉₀ and OUES₇₅. OUES₇₅; oxygen uptake efficiency slope from the first 75% of the test, OUES₉₀; oxygen uptake efficiency slope from the first 90% of the test, OUES₁₀₀; oxygen uptake efficiency slope from the entire test.

the oxygen uptake efficiency slope from the entire test, and the oxygen uptake efficiency slope at ventilatory anaerobic threshold and the other two (Student's *t*-test).

Discussion

The aim of our study was to investigate the reliability of the oxygen uptake efficiency slope as a submaximal marker of cardiorespiratory reserve in adult patients with CHD and build upon the existing evidence. In this retrospective assessment of all adult congenital patients who achieved a maximal exercise test at a single tertiary cardiology centre, there was a strong correlation between the maximum oxygen uptake and the oxygen uptake efficiency slope from the entire test (0.936), the oxygen uptake efficiency slope from the first 90% of the test (0.927), and the oxygen uptake efficiency slope from the first 75% of the test (0.905). Importantly, as it can be calculated from a submaximal test, there was also a strong correlation between the maximum oxygen uptake and the oxygen uptake efficiency slope at ventilatory anaerobic threshold (0.833). In our cohort, the mean percentage of oxygen uptake at which anaerobic threshold was reached was 68%. No significant difference was seen in the oxygen uptake efficiency slope at these four different exercise durations. Analysis of the larger subgroups, namely patients with tetralogy of Fallot, transposition of the great arteries, valvular abnormalities, and those with Fontan circulations demonstrated similarly strong correlation of the maximum oxygen uptake with the oxygen uptake efficiency slope from the first 75% of the test, the oxygen uptake efficiency slope from the first 90% of the test, the oxygen uptake efficiency slope from the entire test, and the oxygen uptake efficiency slope at ventilatory anaerobic threshold other than the slightly weaker correlation between the maximum oxygen uptake and the oxygen uptake efficiency slope at ventilatory anaerobic threshold in the Fontan circulation subgroup.

A greater correlation was demonstrated between the maximum oxygen uptake and the oxygen uptake efficiency slope than that of the maximum oxygen uptake with either ventilatory anaerobic threshold (0.875) or minute ventilation/carbon dioxide production slope (0.524). These findings are consistent with those of previous studies. Baba demonstrated the strongest correlation between the maximum oxygen uptake and the oxygen uptake efficiency slope (0.941), with weaker correlations comparable to our own seen with ventilatory anaerobic threshold (0.861) and the minute ventilation/carbon dioxide production slope (-0.452)⁽¹¹⁾. Buys et al also demonstrated a strong correlation between the maximum oxygen uptake and the oxygen uptake efficiency slope (0.853), with weaker correlations of the maximum oxygen uptake with ventilatory anaerobic threshold (0.840) and the minute ventilation/carbon dioxide production slope (-0.421).¹³

In addition to the strong correlation between the oxygen uptake efficiency slope and the maximum oxygen uptake, the oxygen uptake efficiency slope also showed an association with ventilatory anaerobic threshold (0.893) and a moderate association with the minute ventilation/carbon dioxide production slope (0.511) which further validates the results.

The explanation for the superior correlation of the oxygen uptake efficiency slope with the maximum oxygen uptake may be related to the underlying physiology on which it is based. According to the modified alveolar gas equation, the carbon dioxide set-point, metabolic carbon dioxide production, and the ratio of the pulmonary dead space to tidal volume all affect the

relationship between minute ventilation and oxygen uptake.¹¹ The ventilatory anaerobic threshold is dependent on metabolic carbon dioxide production. The minute ventilation/carbon dioxide production slope is predominantly affected by the ratio of the pulmonary dead space to tidal volume. The oxygen uptake efficiency slope by contrast is impacted by both and as such reflects blood distribution to skeletal muscles as well as perfusion to the lungs.

Two previous studies have investigated the correlation between the maximum oxygen uptake and oxygen uptake efficiency slope in the CHD population. Giardini et al in 2008 demonstrated good correlation of the oxygen uptake efficiency slope during the last 50% of exercise duration and oxygen uptake efficiency slope from the entire test with the maximum oxygen uptake in their patients with previous Fontan procedure or atrial switch. They did however report a substantial difference between the oxygen uptake efficiency slope from the first 50% of the test and the maximum oxygen uptake in cyanotic Fontan patients. It was felt that the mechanism for the non-linear relationship between the Log of the minute ventilation and oxygen uptake in this group may be related to right-left shunting and resulting hyperventilation of blood passing through the lungs during exercise.²⁹ It should be noted however that a study by Bongers et al²² identified a linear relationship between the Log of minute ventilation and oxygen uptake throughout the cardiopulmonary exercise test in their group of paediatric cyanotic Fontan patients. Subsequently, in 2011, Buys et al also demonstrated a correlation between the maximum oxygen uptake and the oxygen uptake efficiency slope in adult patients with transposition of the great arteries and univentricular physiology, as well as those with tetralogy of Fallot and coarctation of the aorta. Good correlation of the maximum oxygen uptake with oxygen uptake efficiency slope from the first 75% of the test, the oxygen uptake efficiency slope from the first 90% of the test, and the oxygen uptake efficiency slope from the entire test was seen in all groups except those following atrial switch procedure for transposition of the great arteries.¹³ As above, we did not identify any significant drop off in the strength of the correlation across all groups including that of the Fontan patients and the patients following atrial switch repair.

As has been previously recommended,²² we also feel that it is important to normalise the oxygen uptake efficiency slope for body surface area. Oxygen uptake is traditionally expressed in ml/min/kg; however, the calculation of oxygen uptake efficiency slope does not account for weight, and as such, changes in weight that are often seen with progressive disease may result in erroneous variation in the calculated oxygen uptake efficiency slope. Our analysis did not demonstrate a significant difference when adjusted for body surface area or body mass index; however, we feel that on an individual basis this could account for significant discrepancies.

The linear correlation seen between the logarithm of minute ventilation and oxygen uptake at different exercise intensities is in keeping with data from various patient groups in suggesting that the relationship is effort-independent.^{11,20-28} This is an important finding in the adult CHD population in whom the rate of submaximal testing is high,¹³ and objective assessment of cardiorespiratory function can be key to interventional decisions. Perhaps most importantly, the strong correlation between the maximum oxygen uptake and the oxygen uptake efficiency slope at ventilatory anaerobic threshold will give clinicians some insight into the cardiorespiratory reserve of patients with CHD who are unable to achieve a maximum oxygen uptake.

Strengths and limitations

As all patients who achieved a maximal test were included this is a truly representative sample of the case mix of our tertiary adult CHD service, and although the heterogeneity of the patients makes some comparisons difficult, we feel that the results are generalisable and the numbers are greater than the two previous similar studies. Retrospective data collection comes with inherent limitations; however, this does remove some potential bias and we made every effort to reduce data collection errors. The clinical utility of the oxygen uptake efficiency slope from the first 90% of the test and the oxygen uptake efficiency slope from the first 75% of the test is limited; however, knowing that there is a strong correlation of the maximum oxygen uptake with the oxygen uptake efficiency slope once the patient achieved ventilatory anaerobic threshold allows conclusions to be drawn from the oxygen uptake efficiency slope in future submaximal tests where this threshold is achieved.

Conclusion

In adult patients with CHD who are unable to complete a maximal cardiopulmonary exercise test, the oxygen uptake efficiency slope is a reliable indicator of cardiopulmonary fitness which correlates strongly with the maximum oxygen uptake irrespective of exercise intensity. A strong correlation has been shown at or beyond ventilatory anaerobic threshold. Further research is required to further validate the findings in patients with less common anatomies and to assess the relationship between the oxygen uptake efficiency slope and mortality.

Competing interests. None.

References

- Ferrer-Sargues FJ, Peiro-Molina E, Salvador-Coloma P, et al. Cardiopulmonary rehabilitation improves respiratory muscle function and functional capacity in children with congenital heart disease. A prospective cohort study. *Int J Environ Res Public Health* 2020; 17: 4328.
- Wren C, O'Sullivan JJ. Survival with congenital heart disease and need for follow up in adult life. *Heart* 2001; 85: 438–443.
- Engelfriet P, Boersma E, Oechslin E, et al. The spectrum of adult congenital heart disease in Europe: morbidity and mortality in a 5 year follow-up period. The Euro Heart Survey on adult congenital heart disease. *Eur Heart J* 2005; 26: 2325–2333.
- Buber J, Rhodes J. Exercise physiology and testing in adult patients with congenital heart disease. *Heart Fail Clin* 2014; 10: 23–33.
- Gratz A, Hess J, Hager A. Self-estimated physical functioning poorly predicts actual exercise capacity in adolescents and adults with congenital heart disease. *Eur Heart J* 2009; 30: 497–504.
- Hallock P. Lactic acid production during rest and after exercise in subjects with various types of heart disease with special reference to congenital heart disease. *J Clin Invest* 1939; 18: 385–394.
- Fernandes SM, Alexander ME, Graham DA, et al. Exercise testing identifies patients at increased risk for morbidity and mortality following Fontan surgery. *Congenit Heart Dis* 2011; 6: 294–303.
- Inuzuka R, Diller GP, Borgia F, et al. Comprehensive use of cardiopulmonary exercise testing identifies adults with congenital heart disease at increased mortality risk in the medium term. *Circulation* 2012; 125: 250–259.
- Taylor HL, Buskirk E, Henschel A. Maximal oxygen intake as an objective measure of cardio-respiratory performance. *J Appl Physiol* 1955; 8: 73–80.
- Mancini DM, Eisen H, Kussmaul W, Mull R, Edmunds LH Jr., Wilson JR. Value of peak exercise oxygen consumption for optimal timing of cardiac transplantation in ambulatory patients with heart failure. *Circulation* 1991; 83: 778–786.
- Baba R, Nagashima M, Goto M, et al. Oxygen uptake efficiency slope: a new index of cardiorespiratory functional reserve derived from the relation between oxygen uptake and minute ventilation during incremental exercise. *J Am Coll Cardiol* 1996; 28: 1567–1572.
- American Thoracic S. American college of chest P. ATS/ACCP statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med* 2003; 167: 211–277.
- Buys R, Cornelissen V, Van De Bruaene A, et al. Measures of exercise capacity in adults with congenital heart disease. *Int J Cardiol* 2011; 153: 26–30.
- Yeh MP, Gardner RM, Adams TD, Yanowitz FG, Crapo RO. Anaerobic threshold: problems of determination and validation. *J Appl Physiol Respir Environ Exerc Physiol* 1983; 55: 1178–1186.
- Shimizu M, Myers J, Buchanan N, et al. The ventilatory threshold: method, protocol, and evaluator agreement. *Am Heart J* 1991; 122: 509–516.
- Metra M, Dei Cas L, Panina G, Visioli O. Exercise hyperventilation chronic congestive heart failure, and its relation to functional capacity and hemodynamics. *Am J Cardiol* 1992; 70: 622–628.
- Buller NP, Poole-Wilson PA. Mechanism of the increased ventilatory response to exercise in patients with chronic heart failure. *Br Heart J* 1990; 63: 281–283.
- Uchida K. Unit of oxygen uptake efficiency slope. *J Sports Med Phys Fitness* 2018; 7: 171–175.
- Mezzani A, Agostoni P, Cohen-Solal A, et al. Standards for the use of cardiopulmonary exercise testing for the functional evaluation of cardiac patients: a report from the exercise physiology section of the european association for cardiovascular prevention and rehabilitation. *Eur J Cardiovasc Prev Rehabil* 2009; 16: 249–267.
- Mollard P, Woorons X, Antoine-Jonville S, et al. 'Oxygen uptake efficiency slope' in trained and untrained subjects exposed to hypoxia. *Respir Physiol Neurobiol* 2008; 161: 167–173.
- Hollenberg M, Tager IB. Oxygen uptake efficiency slope: an index of exercise performance and cardiopulmonary reserve requiring only submaximal exercise. *J Am Coll Cardiol* 2000; 36: 194–201.
- Bongers BC, Hulzebos HJ, Blank AC, van Brussel M, Takken T. The oxygen uptake efficiency slope in children with congenital heart disease: construct and group validity. *Eur J Cardiovasc Prev Rehabil* 2011; 18: 384–392.
- Toste A, Soares R, Feliciano J, et al. Prognostic value of a new cardiopulmonary exercise testing parameter in chronic heart failure: oxygen uptake efficiency at peak exercise - comparison with oxygen uptake efficiency slope. *Rev Port Cardiol* 2011; 30: 781–787.
- Woods PR, Bailey KR, Wood CM, Johnson BD. Submaximal exercise gas exchange is an important prognostic tool to predict adverse outcomes in heart failure. *Eur J Heart Fail* 2011; 13: 303–310.
- Coeckelberghs E, Buys R, Goetschalckx K, Cornelissen VA, Vanhees L. Prognostic value of the oxygen uptake efficiency slope and other exercise variables in patients with coronary artery disease. *Eur J Prev Cardiol* 2016; 23: 237–244.
- Woods PR, Frantz RP, Taylor BJ, Olson TP, Johnson BD. The usefulness of submaximal exercise gas exchange to define pulmonary arterial hypertension. *J Heart Lung Transplant* 2011; 30: 1133–1142.
- Tang Y, Luo Q, Liu Z, et al. Oxygen uptake efficiency slope predicts poor outcome in patients with idiopathic pulmonary arterial hypertension. *J Am Heart Assoc* 2017; 6(7): e005037.
- Tan X, Yang W, Guo J, et al. Usefulness of decrease in oxygen uptake efficiency to identify gas exchange abnormality in patients with idiopathic pulmonary arterial hypertension. *PLoS One* 2014; 9: e98889.
- Giardini A, Specchia S, Gargiulo G, Sangiorgi D, Picchio FM. Accuracy of oxygen uptake efficiency slope in adults with congenital heart disease. *Int J Cardiol* 2009; 133: 74–79.
- Kempny A, Dimopoulos K, Uebing A, et al. Reference values for exercise limitations among adults with congenital heart disease. Relation to activities of daily life—single centre experience and review of published data. *Eur Heart J* 2012; 33: 1386–1396.

31. Graham BL, Steenbruggen I, Miller MR, et al. Standardization of spirometry 2019 Update. An official American thoracic society and european respiratory society technical statement. *Am J Respir Crit Care Med* 2019; 200: e70–e88.
32. Quanjer PH, Stanojevic S, Cole TJ, et al. Multi-ethnic reference values for spirometry for the 3-95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012; 40: 1324–1343.
33. Wasserman K. *Principles of Exercise Testing and Interpretation : Including Pathophysiology and Clinical Applications*, vol. xvi, 4th edn. Lippincott Williams & Wilkins, Philadelphia, 2005, 585–p.
34. Hansen JE, Sue DY, Wasserman K. Predicted values for clinical exercise testing. *Am Rev Respir Dis* 1984; 129: S49–55.