



Letter to the Editor

Vegetation and climate changes during the late Pliocene and early Pleistocene in SW Turkey – Response to comments by Elitez et al., Quaternary Research, 84, pp. 448–456



Keywords:

Pollen analysis
Paleovegetation
Paleoclimate
Pliocene
Pleistocene
SW Anatolia

We found the comments by Elitez et al. irrelevant to the scientific study on the vegetation history of the Çameli Basin in SW Anatolia, as presented by Jiménez-Moreno et al. (2015). The comments are far from the scientific debate but instead deal with technical issues raised after looking for the studied sites on Google Earth. Elitez et al. argue about the geographic coordinates of the localities and the thickness of the studied successions, appear to have a stationary view of a fossil site as a single spot on the Earth' surface, and forget that a sedimentary layer containing fossils can outcrop laterally with changes in thickness.

Elitez et al. also have problems with the stratigraphic position and the age given for the Ericcek and Bicakci localities. Jiménez-Moreno et al. (2015) show that they are different in age; small changes in the dates in the literature are normal as science progresses with further scientific work. For example, notice how many times the Geological Time Scale has changed/evolved.

In order to justify their criticism towards our knowledge of the geology of the area, Elitez et al. discuss a landslide issue that they illustrated with a Google Earth image. They also show a calculation on present-day sediment accumulation on an artificially dammed lake and delta plain far out to the south as a proxy for sedimentary rates in the past. However, they do not take into account that climate has changed since then, and that human impact on the environment (e.g., enhanced erosion due to deforestation) is an important factor today when estimating erosion rates and thus sedimentation. Biostratigraphy is therefore fundamental when trying to learn about the age of old sedimentary sequences and cannot be replaced by simple calculations such as this one.

DOIs of original article: <http://dx.doi.org/10.1016/j.yqres.2016.03.004>, <http://dx.doi.org/10.1016/j.yqres.2015.09.005>.

<http://dx.doi.org/10.1016/j.yqres.2016.03.009>

0033-5894/© 2016 Published by Elsevier Inc. on behalf of University of Washington.

They present the geological map as their own work, but it was extracted from the Geological Map Series of the Geological Survey of Turkey (MTA) compiled by Akdeniz (2011). They also did some computer mapping on a Google Earth image without citation. Elitez et al. failed to present any field evidences and instead showed logging and lithological observations on an old photo. They claim their unpublished evidence is stronger than international published data, and to justify their comments they cite personal communications with us that we are not aware of, and they cite non-scientific sources such as Research Gate. Such sites are unreliable personal domains, like Facebook, and the holder can change them arbitrarily.

Comments such as these do not contain any data refuting our arguments on Plio-Quaternary vegetation history of the Çameli Basin in SW Anatolia and do not stimulate discussion within the scientific community. Regardless, we direct them to read carefully pre-existed studies on the Late Miocene-early Pleistocene extensional graben-type basins in SW Anatolia, including the Çameli Basin (Şenel, 1997a,b,c; Alçiçek et al., 2005, 2006, 2012, 2013a,b, 2015; Alçiçek, 2007, 2009, 2010, 2015; Alçiçek and Ten Veen, 2008; Ten Veen et al., 2009; Over et al., 2010, 2013a,b; Akdeniz, 2011; Helvacı et al., 2013; Alçiçek and Jiménez-Moreno, 2013; Alçiçek and Alçiçek, 2014; Jiménez-Moreno et al., 2015, 2016; van den Hoek Ostende et al., 2015a,b; Ozsayın, 2016).

Here we summary the stratigraphy of the Çameli Basin and its sedimentary infill to better inform the reader:

The Çameli Basin, ca. 40 km wide and 60 km long, is delimited by NE-trending basin-bounding normal faults (see Fig. 1 in Jiménez-Moreno et al., 2016). The basin resides on the Lycian nappes and consists of a series of NE-trending interconnected tilt-block compartments resulting from the NW–SE directed regional crustal extension. The basin formation in SW Anatolia was followed by final southeastward emplacement of the Lycian Nappes, which was sealed by Langhian shallow marine reef carbonates, and its nappe front was eventually covered by Serravalian shallow marine clastics to the southeast (Hayward and Robertson, 1982; Hayward, 1984; Collins and Robertson, 1997; Alçiçek and Ten Veen, 2008). Final movement of the nappes was followed by regional uplift, which caused NE–SW trending extension and resulted in graben-type depressions parallel to the nappe front. The age of basin-fill succession has been well determined as late Miocene (biozones MN9–12; Vallesian-Turolian chronozones) to early Pleistocene (biozone MN17; latest Villanyian chronozone) based on terrestrial macro- and micro-mammal associations (Alçiçek, 2001; Saraç, 2003; Alçiçek et al., 2005; Van den Hoek-Ostende et al., 2015a,b).

The basin-fill succession of the Çameli Basin has been identified as Çameli Formation and grouped into three lithostratigraphic

subunits referred to as the Derindere, Kumafşarı and Değne members, consisting of alluvial, fluvial and lacustrine deposits, respectively. In the central part of the basin these members overlie each other in a 500-m-thick sequence, but they are laterally equivalent along the basin margins. The Derindere Member is composed of coarse-grained alluvial deposits and occurs typically in the lower- and also uppermost parts of the basin fill along the basin margins. It is about 60 m thick with dark-red colored matrix-supported conglomerates and mudstones, and passes laterally and vertically into the fluvial deposits. The Kumafşarı Member is widespread in the northern part of the basin and in the middle stratigraphic level of the basin fill, and it consists of up to 146 m of stacked fluvial deposits characterized by a light yellow color. This member passes laterally and vertically into lacustrine deposits. The Değne Member is composed of lacustrine deposits that vary from 75 to 300 m in thickness. The unit is common in southern parts of the basin and mainly constitutes the upper part of the basin succession, passing laterally and vertically into the fluvial deposits.

References

- Alçiçek, M.C., 2001. Sedimentological investigation of the Çameli Basin (Late Miocene-Late Pliocene, Denizli, SW Anatolia). Unpubl. Ph. D. Thesis. Ankara University, p. 101.
- Akdeniz, N., 2011. Geological Maps of Turkey in 1:100,000 Scale: Denizli N22 Sheet. Mineral Research and Exploration Directorate of Turkey, Ankara, 44 pp.
- Alçiçek, M.C., Kazancı, N., Özkul, M., 2005. Multiple rifting pulses and sedimentation pattern in the Çameli Basin, southwestern Anatolia, Turkey. *Sedimentary Geology* 173, 409–431.
- Alçiçek, M.C., Ten Veen, J.H., Özkul, M., 2006. Neotectonic development of the Çameli Basin, southwestern Anatolia, Turkey. In: Robertson, A.H.F., Mountrakis, D. (Eds.), *Tectonic Development of the Eastern Mediterranean Region*. Geological Society of London, Special Publication, vol. 260, pp. 591–611.
- Alçiçek, M.C., 2007. Tectonic development of an orogen-top rift recorded by its terrestrial sedimentation pattern: the Neogene Eşen Basin of southwestern Anatolia, Turkey. *Sedimentary Geology* 200, 117–140.
- Alçiçek, M.C., Ten Veen, J.H., 2008. The late Early Miocene Acipayam piggy-back basin: refining the last stages of Lycian nappe emplacement in SW Turkey. *Sedimentary Geology* 208, 101–113.
- Alçiçek, H., 2009. Late Miocene nonmarine sedimentation and formation of magnetites in the Acıgöl Basin, southwestern Anatolia, Turkey. *Sedimentary Geology* 219, 115–135.
- Alçiçek, H., 2010. Stratigraphic correlation of the Neogene basins in southwestern Anatolia: regional palaeogeographical, palaeoclimatic and tectonic implications. *Palaeogeography, Palaeoclimatology, Palaeoecology* 291, 297–318.
- Alçiçek, M.C., Mayda, S., Alçiçek, H., 2012. Faunal and palaeoenvironmental changes in the Çal Basin, SW Anatolia: implications for regional stratigraphic correlation of late Cenozoic basins. *Comptes Rendus Geoscience* 344, 89–98.
- Alçiçek, H., Jiménez-Moreno, G., 2013. Late Miocene to Pliocene fluvio-lacustrine system in Karacasu Basin (SW Anatolia, Turkey): depositional, palaeogeographic and palaeoclimatic implications. *Sedimentary Geology* 291, 62–83.
- Alçiçek, M.C., Mayda, S., Titov, V.V., 2013a. Lower Pleistocene stratigraphy of the Burdur Basin of SW Anatolia. *Comptes Rendus Palevol* 12, 1–11.
- Alçiçek, M.C., Brogi, A., Capezzuoli, E., Liotta, D., Meccheri, M., 2013b. Superimposed basin formation during Neogene–Quaternary extension in SW-Anatolia (Turkey): insights from the kinematics of the Dinar fault zone. *Tectonophysics* 608, 713–727.
- Alçiçek, H., Alçiçek, M.C., 2014. Palustrine carbonates and pedogenic calcretes in the Çal basin of SW Anatolia: implication on the Plio-Pleistocene regional climatic pattern in the eastern Mediterranean. *Catena* 112, 48–55.
- Alçiçek, H., Wesselingh, F., Alçiçek, M.C., 2015. Palaeoenvironmental evolution of the late Pliocene-early Pleistocene fluvio-deltaic sequence of the Denizli Basin (SW Turkey). *Palaeogeography, Palaeoclimatology, Palaeoecology* 437, 98–116.
- Alçiçek, M.C., 2015. Comment on The Fethiye–Burdur Fault Zone: a component of upper plate extension of the subduction transform edge propagator fault linking Hellenic and Cyprus Arcs, Eastern Mediterranean. *Tectonophysics* 635, 80–99 by J. Hall, A.E. Aksu, İ. Elitez, C. Yaltrak, G. Çiftçi *Tectonophysics* 664, 1–4.
- Collins, A.S., Robertson, A.H.F., 1997. Lycian mélange, southwestern Turkey: an emplaced Late Cretaceous accretionary complex. *Geology* 25, 255–258.
- Hayward, A.B., Robertson, A.H.F., 1982. Direction of ophiolite emplacement inferred from Cretaceous and Tertiary sediments of an adjacent autochthon, the Bey Dağları, SW Turkey. *Geological Society of America Bulletin* 93, 68–75.
- Hayward, A.B., 1984. Sedimentation and basin formation related to ophiolite emplacement, Miocene, SW Turkey. *Sedimentary Geology* 40, 105–129.
- Helvacı, C., Alçiçek, M.C., Gündoğan, İ., Gemici, Ü., 2013. Tectonosedimentary development and palaeoenvironmental changes in the Acıgöl shallow-perennial playa-lake basin, SW Anatolia, Turkey. *Turkish Journal of Earth Sciences* 22, 173–190.
- Jiménez-Moreno, G., Alçiçek, H., Alçiçek, M.C., van den Hoek Ostende, L.W., Wesselingh, F.P., 2015. Vegetation and climatic cycles during the late Pliocene and early Pleistocene in SW Anatolia, Turkey. *Quaternary Research* 84, 448–456.
- Jiménez-Moreno, G., Alçiçek, H., Alçiçek, M.C., van den Hoek Ostende, L.W., Wesselingh, F.P., 2016. Corrigendum to “Vegetation and climatic cycles during the late Pliocene and early Pleistocene in SW Anatolia”. *Quaternary Research* 85, 185–186.
- Over, S., Yılmaz, H., Pınar, A., Özden, S., Ünlügöç, U.C., Kamacı, Z., 2013a. Plio-Quaternary stress state in the Burdur Basin, SW Turkey. *Tectonophysics* 588, 56–68.
- Over, S., Özden, S., Yılmaz, H., Pınar, A., Ünlügöç, U.C., Kamacı, Z., 2013b. Plio-Quaternary stress regime in the Eşen Çay Basin, SW Turkey. In: Robertson, A.H.F., Parlak, O., Unlugöç, U.C. (Eds.), *Geological Development of Anatolia and the Easternmost Mediterranean Region*. Geological Society, London, pp. 547–560. Special Publications 372.
- Over, S., Pınar, A., Özden, S., Yılmaz, H., Ünlügöç, U.C., Kamacı, Z., 2010. Late Cenozoic stress field in the Çameli Basin, SW Turkey. *Tectonophysics* 492, 60–72.
- Ozsayın, E., 2016. Relative tectonic activity assessment of the Çameli Basin, Western Anatolia, using geomorphic indices. *Geodinamica Acta*. <http://dx.doi.org/10.1080/09853111.2015.1128180>.
- Saraç, G., 2003. Türkiye Omurgalı Fossil Yatakları (Vertebrate Fossil Localities of Turkey). Scientific Report No. 10609. General Directorate of the Mineral Research and Exploration of Turkey (MTA), Ankara, p. 208.
- Şenel, M., 1997a. Geological Maps of Turkey in 1:100,000 Scale: Fethiye L8 Sheet. Mineral Research and Exploration Directorate of Turkey (MTA), Ankara, Turkey, 22 pp.
- Şenel, M., 1997b. Geological Maps of Turkey in 1:100,000 Scale: Fethiye M8 Sheet. Mineral Research and Exploration Directorate of Turkey (MTA), Ankara, Turkey, 15 pp.
- Şenel, M., 1997c. Geological Maps of Turkey in 1:100,000 Scale: Denizli K9 Sheet. Mineral Research and Exploration Directorate of Turkey (MTA), Ankara, Turkey, 17 pp.
- Ten Veen, J.H., Boulton, S.J., Alçiçek, M.C., 2009. From palaeotectonics to neotectonics in the Neotethys realm: the importance of kinematic decoupling and inherited structural grain in SW Anatolia (Turkey). *Tectonophysics* 473, 261–281.
- van den Hoek Ostende, L.W., Diepeveen, F., Tesakov, A., Saraç, G., Mayhew, D., Alçiçek, M.C., 2015a. On the brink: micromammals from the latest Villanyian from Bıçakçı (Anatolia). *Geological Journal* 50, 230–245.
- van den Hoek Ostende, L.W., Gardner, J.D., van Bennekom, L., Alçiçek, M.C., Murray, A.M., Wesselingh, F.P., Alçiçek, H., Tesakov, A.S., 2015b. Eriçek, a new Pliocene vertebrate locality from the Çameli Basin (SW Anatolia, Turkey). *Palaeobiodiversity and Palaeoenvironments* 95, 305–320.

Gonzalo Jiménez-Moreno*

Departamento de Estratigrafía y Paleontología, Universidad de Granada, Fuente Nueva s/n, 18002, Granada, Spain

Hülya Alçiçek, Mehmet Cihat Alçiçek
Pamukkale University, Dept. of Geology, 20070, Denizli, Turkey
E-mail addresses: halcicek@pau.edu.tr (H. Alçiçek),
alcicek@pau.edu.tr (M.C. Alçiçek).

Lars van den Hoek Ostende, Frank P. Wesselingh
Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA, Leiden, The Netherlands
E-mail addresses: lars.vandenhoekostende@naturalis.nl (L. van den Hoek Ostende), frank.wesselingh@naturalis.nl (F.P. Wesselingh).

* Corresponding author.

E-mail address: gonzaloj@ugr.es (G. Jiménez-Moreno).

15 March 2016