

108.09 A visual proof that $b^e < e^b$ when $b > e$

In a recent visual proof ([1]), the author provided a visual proof of the inequality $\pi^e < e^\pi$. However, their visual proof can be used to show the more general inequality $b^e < e^b$, where $e < b$.




FIGURE 1

$$\ln b - 1 = \int_e^b \frac{dx}{x} < \frac{1}{e} (b - e) = \frac{b}{e} - 1$$

and so $b^e < e^b$.

Reference

1. Bikash Chakraborty, A visual proof that $\pi^e < e^\pi$, *Mathematical Intelligencer* **41** (2019) p. 60.

10.1017/mag.2024.26 © The Authors, 2024

Published by Cambridge University Press

on behalf of The Mathematical Association

BIKASH CHAKRABORTY

Department of Mathematics,

Ramakrishna Mission

Vivekananda Centenary College,

Rahara, West Bengal 700 118, India

e-mail: *bikashchakraborty.math@yahoo.com,*

bikash@rkmvccrahara.org


108.10 Proof without words: $\tan \frac{\pi}{12} = 2 - \sqrt{3}$, $\tan \frac{5\pi}{12} = 2 + \sqrt{3}$

The standard proof of $\tan \frac{\pi}{12} = 2 - \sqrt{3}$ is to use the less well-known formula

$$\tan \alpha = \frac{-1 + \sqrt{1 + \tan^2 2\alpha}}{\tan 2\alpha}$$

for $\alpha = \frac{\pi}{12}$ and the well-known value $\tan \frac{\pi}{6} = \frac{1}{\sqrt{3}}$. Using only the last fact,





from the diagram the readers can readily see that

$$\tan \frac{5\pi}{12} = 2 + \sqrt{3} \quad \text{and} \quad \tan \frac{\pi}{12} = \frac{1}{2 + \sqrt{3}} = 2 - \sqrt{3}.$$

There is another PWW from Garcia Capitan Francisco Javier [1]. Paul Stephenson [2] and Nick Lord [3] have offered other demonstrations of the identity of $\tan \frac{\pi}{12} = 2 - \sqrt{3}$, for which Nick Lord gives four Proofs without words, with quite different ideas. For many useful principles and comments about Proofs without words, see [4].

References

1. F. J. Garcia Capitan, Proof without Words: tangents 15 and 17 degrees, *Coll. Math. J.* **48:1** (2017) p. 35.
2. P. Stephenson, Feedback: On what makes a good Proof without Words, *Math. Gaz.* **107** (March 2023) p. 165.
3. Nick Lord, Feedback, *Math. Gaz.* **107** (July 2023) p. 356.
4. G. Leversha, What makes a good Proof without Words, *Math. Gaz.* **105** (July 2021) pp. 271-281.

10.1017/mag.2024.27 © The Authors, 2024

Published by Cambridge University Press
on behalf of The Mathematical Association

MARTIN LUKAREVSKI
*Department of Mathematics
and Statistics,*

*University "Goce Delcev" - Stip,
North Macedonia*

e-mail: *martin.lukarevski@ugd.edu.mk*

108.11 Euler's limit—revisited

Let $e_n = (1 + \frac{1}{n})^n$ for $n \in \mathbb{N}$. It is well known that the sequence (e_n) is monotone increasing and bounded, hence it is convergent. The limit of this sequence is the famous Euler number e . Here we establish a generalisation of this limit.

Theorem: Let $\{a_n\}$ and $\{b_n\}$ be two sequences of positive real numbers such that $a_n \rightarrow +\infty$ and b_n satisfies the asymptotic formula $b_n \sim k \cdot a_n$, where $k > 0$. Then

$$\lim_{n \rightarrow \infty} \left(1 + \frac{1}{a_n}\right)^{b_n} = e^k.$$