

Medical News

GINA PUGLIESE, RN, MS, AND MARTIN S. FAVERO, PhD

HIV Transmission by a Human Bite

The transmission of HIV-1 infection by a human bite is assumed to be biologically possible, but has remained unproven. To date, there have been only two case reports in the literature suggesting transmission by this route. Recently, however, researchers from the Department of Infectious Disease, Clinical Center in Ljubljana, Slovenia, reported a documented HIV seroconversion that occurred after a bite.

The case involved a 47-year-old male with AIDS who first presented with esophageal candidiasis in June 1994 and a CD4 lymphocyte count of 40×10^6 L and a Western blot that showed consistently strong reactivity to gp120 and gp160 only. The HIV-1 p24 antigen test was positive, and a high level of HIV-1 RNA was present in his plasma, all suggesting late-stage HIV-1 infection. He was started on zidovudine, and his condition improved.

Eleven months later, the patient had a grand mal seizure at home. His neighbor, who was unaware of his diagnosis, attended to him during his seizure and was bitten while placing his fingers in the patient's mouth trying to prevent airway obstruction. The neighbor reported that the bite resulted in a small crack and a shallow wound on his finger with teeth imprints but without visible bleeding. Right after the incident, the bitten man noticed blood in the biter's saliva.

One hour later, a physician confirmed the neighbor's description of the wound. Serum taken from the bitten man on the day of the injury was negative for HIV-1 and HIV-2 antibodies, p24 antigen, and HIV-1 RNA. Ten hours after the injury, zidovudine prophylaxis was started. Five weeks after the incident, the bitten man reported symptoms consistent with acute primary HIV-1 infection (fatigue, headache, and sore throat). One week after his symptoms developed, serologic tests showed anti-p24 and anti-p55 antibodies only; p24 antigen test was negative. The bitten man seroconverted completely 54 days after the incident. The bitten man was married for 23 years, his wife consistently tested negative for HIV-1 antibodies, and he denied any risk factors for HIV.

The authors comment that blood in the biter's saliva most probably was crucial for HIV transmission in this case. The bite occurred 13 days before the death of the patient, in his terminal stage of HIV infection, so a high

viral load may explain the relatively short time from exposure to seroconversion in the bitten man.

FROM: Vidmar L, Poljak M, Tomazic J, et al. Transmission of HIV-1 by human bite. *Lancet* 1996; 347:1762-1763.

Nationwide Drop in Percutaneous Injuries

Data from 56 hospitals participating in the EPINet data-sharing network have shown a 32% reduction in percutaneous injuries reported in 1995 as compared to 1994 and 1993. The hospitals vary in size, teaching status, and geographic diversity.

In 1993 and 1994, the overall annual injury rate, calculated by dividing the total number of annual injuries by the average daily census, remained constant at 32 injuries per 100 occupied beds. However, in 1995, the injury rate dropped to 22 injuries per 100 occupied beds. The overall decrease in the number of sharp object injuries was consistent among hospitals, regardless of location, teaching status, or size.

Dr. Janine Jagger, who leads the Health Care Worker Safety Center that coordinates the EPINet data-sharing network, noted that the uniformity of this national decline among different types of hospitals in different geographic regions was surprising. In a recent article in *Advances in Exposure Prevention*, Jagger explains that the decline is not due to a decline in the length of hospital stay or reduced patient census, because these factors are controlled for when percutaneous injury rates are reported per 100 occupied beds. The possible reasons for this decline are still being investigated.

FROM: Jagger J, Bentley B. Substantial nationwide drop in percutaneous injury rates detected for 1995. *Advances in Exposure Prevention* May/June 1996.

Additional news items in this issue: Repeat Fit Testing Not Routinely Required, page 635; Community-Acquired and Nosocomial Bacteremia Trends, page 640; Drug-Resistant Acinetobacter baumannii, page 659; OSHA and JCAHO Form Partnership, page 671; Computer Reminders Did Not Improve Inpatient Preventive Care, page 686; Labor Department Launches Worker Safety Program for Nursing Homes, page 693.

EDITED BY MURRAY D. BATT, MD

CHARLES E. EDMISTON, PhD

JOHN HEGGERS, PhD

SHEA News

THE SOCIETY FOR HEALTHCARE EPIDEMIOLOGY OF AMERICA

PRESIDENT	William J. Martone, MD/Bethesda, Maryland	TREASURER	Alan I. Hartstein, MD/Indianapolis, Indiana
PRESIDENT-ELECT	Robert A. Weinstein, MD/Chicago, Illinois	COUNCILOR	John W. Froggatt, III, MD/Asheville, North Carolina
VICE PRESIDENT	Elias Abrutyn, MD/Philadelphia, Pennsylvania	COUNCILOR	Edward S. Wong, MD/Richmond, Virginia
PAST PRESIDENT	Bryan P. Simmons, MD/Memphis, Tennessee	COUNCILOR	Barry M. Farr, MD, MSc/Charlottesville, Virginia
SECRETARY	Lindsay E. Nicolle, MD/Winnipeg, Manitoba	COUNCILOR	Martha Tanner, MD/Idaho Falls, Idaho

SHEA Membership Continues to Grow

As the premier organization of advanced-degree epidemiologists, SHEA continues toward the strategic membership goals that were outlined at the January 1996 strategic planning meeting. SHEA membership exceeded 1,200 for the first time in August, prior to the recruitment opportunity represented by the annual SHEA presence at the ICAAC/IDSA meeting in September 1996. The highest recorded degree was an MD or DO

for more than 1,100 members; a PhD for more than 50; and a DVM for 1. SHEA now has 13 masters-degree members, each nominated by a SHEA member because of their particular epidemiologic accomplishments (this class of membership is limited by the bylaws to 5% of total membership). Many of our members have dual degrees, particularly MD or RN, plus PhD or MPH.

SHEA Position Papers in the Literature and in the Pipeline

Lindsay Nicolle's group's position paper on "Antimicrobial Use in Long-Term-Care Facilities," and Larry Strausbaugh's group's summary of the issues of "Antimicrobial Resistance in Long-Term-Care Facilities," which were both published earlier this year in *Infection Control and Hospital Epidemiology*, are the 13th and 14th SHEA position

papers to be published in our journal since 1988. In the pipeline, under the guidance of the indicated individual, are the following additional topics: "Selection and Interpretation of Molecular Methods for Microbial Typing for Healthcare Epidemiologists," Fred C. Tenover; "Guidelines for the Prevention of Antimicrobial Resistance in Hospitals," David

New Executive Director of SHEA

With a growing organization exceeding 1,200 members, SHEA has determined the need for a new full-time executive director. She is Stephanie Dickenson. Her background is in the association management field with professional societies and trade associations. As part of the Talley Management Group Team, she will provide business leadership to SHEA officers and committees, as well as managing the day-to-day membership, financial, and administrative functions of the Society. Keep tuned. We have solicited an article for the Fall 1996 *SHEA Newsletter* to reflect her views on SHEA's future.

Shlaes; "Revision of Management of Health Care Workers Who are Infected by Bloodborne Pathogens," Michael Tapper; and "Infrastructure and Essential Activities of Hospital Epidemiology and Infection Control Programs," William Scheckler.

Brief items of interest for the SHEA News or Newsletter may be sent to Murray D. Batt, MD, SHEA News Editor, Washoe Medical Center, 77 Pringle Way, Reno NV 89520; fax (702) 328-4797.