



Cumulants of the q -semicircular Law, Tutte Polynomials, and Heaps

Matthieu Josuat-Vergès

Abstract. The q -semicircular distribution is a probability law that interpolates between the Gaussian law and the semicircular law. There is a combinatorial interpretation of its moments in terms of matchings, where q follows the number of crossings, whereas for the free cumulants one has to restrict the enumeration to connected matchings. The purpose of this article is to describe combinatorial properties of the classical cumulants. We show that like the free cumulants, they are obtained by an enumeration of connected matchings, the weight being now an evaluation of the Tutte polynomial of a so-called crossing graph. The case $q = 0$ of these cumulants was studied by Lassalle using symmetric functions and hypergeometric series. We show that the underlying combinatorics is explained through the theory of heaps, which is Viennot’s geometric interpretation of the Cartier–Foata monoid. This method also gives a general formula for the cumulants in terms of free cumulants.

1 Introduction

Let us consider the sequence $\{m_n(q)\}_{n \geq 0}$ defined by the generating function

$$\sum_{n \geq 0} m_n(q)z^n = \frac{1}{1 - \frac{[1]_q z^2}{1 - \frac{[2]_q z^2}{1 - \dots}}},$$

where $[i]_q = \frac{1-q^i}{1-q}$. For example, $m_0(q) = m_2(q) = 1$, $m_4(q) = 2 + q$, and the odd values are 0. The generating function being a Stieltjes continued fraction, $m_n(q)$ is the n -th moment of a symmetric probability measure on \mathbb{R} (at least when $0 \leq q \leq 1$). An explicit formula for the density $w(x)$ such that $m_n(q) = \int x^n w(x) dx$ is given by Szegő [22]:

$$w(x) = \begin{cases} \frac{1}{\pi} \sqrt{1-q} \sin \theta \prod_{n=1}^{\infty} (1-q^n) |1 - q^n e^{2i\theta}|^2 & \text{if } -2 \leq x\sqrt{1-q} \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where $\theta \in [0, \pi]$ is such that $2 \cos \theta = x\sqrt{1-q}$. At $q = 0$, it is the semicircular distribution with density $(2\pi)^{-1} \sqrt{4-x^2}$ supported on $[-2, 2]$, whereas at the

Received by the editors March 16, 2012.

Published electronically November 13, 2012.

This work was supported by the Austrian Research Foundation (FWF) via the grant Y463.

AMS subject classification: 05A18, 05C31, 46L54.

Keywords: moments, cumulants, matchings, Tutte polynomials, heaps.

limit $q \rightarrow 1$ it becomes the Gaussian distribution with density $(2\pi)^{-1/2}e^{-x^2/2}$. This law is therefore known either as the q -Gaussian or the q -semicircular law. It can be conveniently characterized by its orthogonal polynomials, defined by the relation $xH_n(x|q) = H_{n+1}(x|q) + [n]_q H_{n-1}(x|q)$ together with $H_1(x|q) = x$ and $H_0(x|q) = 1$, and called the continuous q -Hermite polynomials (but we do not insist on this point of view, since the notion of cumulant is not particularly relevant for orthogonal polynomials).

The semicircular law is the analogue in free probability of the Gaussian law [15, 19]. More generally, the q -semicircular measure plays an important role in noncommutative probability theories [3, 5, 7, 8, 24, 25]. This was initiated by Bożejko and Speicher [7, 8] who used creation and annihilation operators on a twisted Fock space to build generalized Brownian motions.

The goal of this article is to examine the combinatorial meaning of the classical cumulants $k_n(q)$ of the q -semicircular law (we recall the definition in the next section). The first values lead to the observation that

$$\tilde{k}_{2n}(q) = \frac{k_{2n}(q)}{(q-1)^{n-1}}$$

is a polynomial in q with nonnegative coefficients. For example:

$$\tilde{k}_2(q) = \tilde{k}_4(q) = 1, \quad \tilde{k}_6(q) = q + 5, \quad \tilde{k}_8(q) = q^3 + 7q^2 + 28q + 56.$$

We actually show in Theorem 3.5 that this $\tilde{k}_{2n}(q)$ can be given a meaning as a generating function of connected matchings, *i.e.*, the same objects that give a combinatorial meaning to the free cumulants of the q -semicircular law. However, the weight function that we use here on connected matching is not as simple as in the case of free cumulants, it is given by the value at $(1, q)$ of the Tutte polynomial of a graph attached to each connected matching, called the crossing graph.

There are various points where the evaluation of Tutte polynomials has combinatorial meaning, in particular $(1, 0)$, $(1, 1)$, and $(1, 2)$. In the first and third case ($q = 0$ and $q = 2$), they can be used to give an alternative proof of Theorem 3.5. These will be provided in Sections 5 and 6 respectively. The integers $\tilde{k}_{2n}(0)$ were recently considered by Lassalle [17] who defines them as a sequence simply related with Catalan numbers, and further studied in [2]. Being the (classical) cumulants of the semicircular law, it might seem unnatural to consider this quantity, since this law belongs to the world of free probability, but on the other hand the free cumulants of the Gaussian have numerous properties (see [4]). The interesting feature is that this particular case $q = 0$ can be proved via the theory of heaps [10, 26]. As for the case $q = 2$, even though the q -semicircular is only defined when $|q| < 1$ its moments and cumulants and the link between still exist, because (2.1) can be seen as an identity between formal power series in z . The particular proof for $q = 2$ is an application of the exponential formula.

2 Preliminaries

Let us first make precise some terms used in the introduction. Besides the moments $\{m_n(q)\}_{n \geq 0}$, the q -semicircular law can be characterized by its *cumulants* $\{k_n(q)\}_{n \geq 1}$,

formally defined by

$$(2.1) \quad \sum_{n \geq 1} k_n(q) \frac{z^n}{n!} = \log \left(\sum_{n \geq 0} m_n(q) \frac{z^n}{n!} \right),$$

or by its *free cumulants* $\{c_n(q)\}_{n \geq 1}$ [19], formally defined by $C(zM(z)) = M(z)$, where

$$M(z) = \sum_{n \geq 0} m_n(q) z^n, \quad C(z) = 1 + \sum_{n \geq 1} c_n(q) z^n.$$

These relations can be reformulated using set partitions.

For any finite set V , let $\mathcal{P}(V)$ denote the lattice of set partitions of V , and let $\mathcal{P}(n) = \mathcal{P}(\{1, \dots, n\})$. We will denote by $\hat{1}$ the maximal element and by μ the Möbius function of these lattices, without mentioning V explicitly. Although we will not use it, let us mention that $\mu(\pi, \hat{1}) = (-1)^{\#\pi-1} (\#\pi - 1)!$, where $\#\pi$ is the number of blocks in π . See [21, Chapter 3] for details. When we have some sequence $(u_n)_{n \geq 1}$, for any $\pi \in \mathcal{P}(V)$ we will use the notation $u_\pi = \prod_{b \in \pi} u_{\#b}$. Then the relations between moments and cumulants read

$$(2.2) \quad m_n(q) = \sum_{\pi \in \mathcal{P}(n)} k_\pi(q), \quad k_n(q) = \sum_{\pi \in \mathcal{P}(n)} m_\pi(q) \mu(\pi, \hat{1}).$$

These are equivalent via the Möbius inversion formula, and both can be obtained from (2.1) using Faà di Bruno’s formula. When $V \subset \mathbb{N}$, let $\mathcal{NC}(V) \subset \mathcal{P}(V)$ denote the subset of *noncrossing partitions*, which form a sublattice with Möbius function μ^{NC} . Then we have ([15, 19])

$$(2.3) \quad m_n(q) = \sum_{\pi \in \mathcal{NC}(n)} c_\pi(q), \quad c_n(q) = \sum_{\pi \in \mathcal{NC}(n)} m_\pi(q) \mu^{NC}(\pi, \hat{1}).$$

Equations (2.2) and (2.3) can be used to compute the first non-zero values:

$$\begin{aligned} k_2(q) &= 1, & k_4(q) &= q - 1, & k_6(q) &= q^3 + 3q^2 - 9q + 5, \\ c_2(q) &= 1, & c_4(q) &= q, & c_6(q) &= q^3 + 3q^2. \end{aligned}$$

Let $\mathcal{M}(V) \subset \mathcal{P}(V)$ denote the set of *matchings*, i.e., set partitions into blocks of size 2. As is customary, a block of $\sigma \in \mathcal{M}(V)$ will be called an *arch*. When $V \subset \mathbb{N}$, a *crossing* [16] of $\sigma \in \mathcal{M}(V)$ is a pair of arches $\{i, j\}$ and $\{k, \ell\}$ such that $i < k < j < \ell$. Let $\text{cr}(\sigma)$ denote the number of crossings of $\sigma \in \mathcal{M}(V)$. Let $\mathcal{N}(V) = \mathcal{M}(V) \cap \mathcal{NC}(V)$ denote the set of *noncrossing matchings*, i.e., those such that $\text{cr}(\sigma) = 0$. Let also $\mathcal{M}(2n) = \mathcal{M}(\{1, \dots, 2n\})$ and $\mathcal{N}(2n) = \mathcal{N}(\{1, \dots, 2n\})$. Let $\mathcal{P}^c(n) \subset \mathcal{P}(n)$ denote the set of *connected set partitions*, i.e., π such that no proper interval of $\{1, \dots, n\}$ is a union of blocks of π , and let $\mathcal{M}^c(2n) = \mathcal{M}(2n) \cap \mathcal{P}^c(2n)$ denote the set of *connected matchings*.

It is known [16] that for any $n \geq 0$, the moment $m_{2n}(q)$ counts matchings on $2n$ points according to the number of crossings:

$$(2.4) \quad m_{2n}(q) = \sum_{\sigma \in \mathcal{M}(2n)} q^{\text{cr}(\sigma)}.$$

Lehner [18] showed that (2.3) and (2.4) give a combinatorial meaning for the free cumulants:

$$c_{2n}(q) = \sum_{\sigma \in \mathcal{M}^c(2n)} q^{\text{cr}(\sigma)}.$$

See [4] for various properties of connected matchings in the context of free probability. Let us also mention that both quantities $m_{2n}(q)$ and $c_{2n}(q)$ are considered in an article by Touchard [23].

3 A Combinatorial Formula for $k_n(q)$

We will use the Möbius inversion formula in equation (2.2), but we first need to consider the combinatorial meaning of the products $m_\pi(q)$.

Lemma 3.1 For any $\sigma \in \mathcal{M}(2n)$ and $\pi \in \mathcal{P}(2n)$ such that $\sigma \leq \pi$, let $\text{cr}(\sigma, \pi)$ be the number of crossings $(\{i, j\}, \{k, \ell\})$ of σ such that $\{i, j, k, \ell\} \subset b$ for some $b \in \pi$. Then we have:

$$(3.1) \quad m_\pi(q) = \sum_{\substack{\sigma \in \mathcal{M}(2n) \\ \sigma \leq \pi}} q^{\text{cr}(\sigma, \pi)}.$$

Proof Denoting $\sigma|_b = \{x \in \sigma : x \subset b\}$, the map $\sigma \mapsto (\sigma|_b)_{b \in \pi}$ is a natural bijection between the set $\{\sigma \in \mathcal{M}(2n) : \sigma \leq \pi\}$ and the product $\prod_{b \in \pi} \mathcal{M}(b)$, in such a way that $\text{cr}(\sigma, \pi) = \sum_{b \in \pi} \text{cr}(\sigma|_b)$. This allows us to factor the right-hand side in (3.1) and obtain $m_\pi(q)$. ■

From equation (2.2) and the previous lemma, we have:

$$(3.2) \quad \begin{aligned} k_{2n}(q) &= \sum_{\pi \in \mathcal{P}(2n)} m_\pi(q) \mu(\pi, \hat{1}) = \sum_{\pi \in \mathcal{P}(2n)} \sum_{\substack{\sigma \in \mathcal{M}(2n) \\ \sigma \leq \pi}} q^{\text{cr}(\sigma, \pi)} \mu(\pi, \hat{1}) \\ &= \sum_{\sigma \in \mathcal{M}(2n)} \sum_{\substack{\pi \in \mathcal{P}(2n) \\ \pi \geq \sigma}} q^{\text{cr}(\sigma, \pi)} \mu(\pi, \hat{1}) = \sum_{\sigma \in \mathcal{M}(2n)} W(\sigma), \end{aligned}$$

where for each $\sigma \in \mathcal{M}(2n)$ we have introduced:

$$(3.3) \quad W(\sigma) = \sum_{\substack{\pi \in \mathcal{P}(2n) \\ \pi \geq \sigma}} q^{\text{cr}(\sigma, \pi)} \mu(\pi, \hat{1}).$$

A key point is to note that $W(\sigma)$ only depends on how the arches of σ cross with respect to each other, which can be encoded in a graph. This leads to the following definition.

Definition 3.2 Let $\sigma \in \mathcal{M}(2n)$. The *crossing graph* $G(\sigma) = (V, E)$ is as follows. The vertex set V contains the arches of σ (i.e., $V = \sigma$), and the edge set E contains the crossings of σ (i.e., there is an edge between the vertices $\{i, j\}$ and $\{k, \ell\}$ if and only if $i < k < j < \ell$).

See Figure 1 for an example. Note that the graph $G(\sigma)$ is connected if and only if σ is a connected matching in the sense of the previous section.




Figure 1: A matching σ and its crossing graph $G(\sigma)$.

We also need the following definition for a general graph.

Definition 3.3 Let $G = (V, E)$ be a graph, and let $\pi \in \mathcal{P}(V)$. Then we denote by $i(E, \pi)$ the number of elements in the edge set E such that both endpoints are in the same block of π .

Lemma 3.4 Let $\sigma \in \mathcal{M}(2n)$ and let $G(\sigma) = (V, E)$ be its crossing graph. Then we have

$$(3.4) \quad W(\sigma) = \sum_{\pi \in \mathcal{P}(V)} q^{i(E, \pi)} \mu(\pi, \hat{1}).$$

Proof There is a natural bijection between the interval $[\sigma, \hat{1}]$ in $\mathcal{P}(2n)$ and the set $\mathcal{P}(V)$, in such a way that $\text{cr}(\sigma, \pi) = i(E, \pi)$. Hence equation (3.4) is just a rewriting of (3.3) in terms of the graph $G(\sigma)$. ■

Now we can use Proposition 4.1. It allows to recognize $(q - 1)^{-n+1}W(\sigma)$ as an evaluation of the Tutte polynomial $T_{G(\sigma)}$, except that it is 0 when the graph is not connected.

Gathering equations (3.2), (3.4), and Proposition 4.1, we have proved the following theorem.

Theorem 3.5 For any $n \geq 1$,

$$\tilde{k}_{2n}(q) = \sum_{\sigma \in \mathcal{M}^c(2n)} T_{G(\sigma)}(1, q).$$

In particular, $\tilde{k}_{2n}(q)$ is a polynomial in q with nonnegative coefficients.

4 The Tutte Polynomial of a Connected Graph

For any graph $G = (V, E)$, let $T_G(x, y)$ denote its Tutte polynomial; we give a short definition here and refer the reader to [1, Chapter 9] for details. This graph invariant can be computed recursively via edge deletion and edge contraction. Let $e \in E$, let $G \setminus e = (V, E \setminus e)$ and $G/e = (V/e, E \setminus e)$ where V/e is the quotient set where both endpoints of the edge e are identified. Then the recursion is

$$(4.1) \quad T_G(x, y) = \begin{cases} xT_{G/e}(x, y) & \text{if } e \text{ is a bridge,} \\ yT_{G \setminus e}(x, y) & \text{if } e \text{ is a loop,} \\ T_{G/e}(x, y) + T_{G \setminus e}(x, y) & \text{otherwise.} \end{cases}$$

The initial case is that $T_G(x, y) = 1$ if the graph G has no edge. Here, a *bridge* is an edge e such that $G \setminus e$ has one more connected component than G , and a *loop* is an edge with identical endpoints.

Proposition 4.1 *Let $G = (V, E)$ be a graph (possibly with multiple edges and loops). Let $n = \#V$. We have:*

$$(4.2) \quad \frac{1}{(q-1)^{n-1}} \sum_{\pi \in \mathcal{P}(V)} q^{i(E, \pi)} \mu(\pi, \hat{1}) = \begin{cases} T_G(1, q) & \text{if } G \text{ is connected,} \\ 0 & \text{otherwise.} \end{cases}$$

Proof Denote by U_G the left-hand side in (4.2) and let e be an edge of G . Suppose that $e \in E$ is a loop; it is then clear that $i(E \setminus e, \pi) = i(E, \pi) - 1$, so $U_G = qU_{G \setminus e}$. Then suppose that e is not a loop, and let x and y be its endpoints. We have

$$U_G - U_{G \setminus e} = \frac{1}{(q-1)^{n-1}} \sum_{\pi \in \mathcal{P}(V)} \left(q^{i(E, \pi)} - q^{i(E \setminus e, \pi)} \right) \mu(\pi, \hat{1}).$$

In this sum, all terms where x and y are in different blocks of π vanish. So we can keep only π such that x and y are in the same block, and these can be identified with elements of $\mathcal{P}(V/e)$ and satisfy $i(E \setminus e, \pi) = i(E, \pi) - 1$. We obtain

$$U_G - U_{G \setminus e} = \frac{1}{(q-1)^{n-2}} \sum_{\pi \in \mathcal{P}(V/e)} q^{i(E \setminus e, \pi)} \mu(\pi, \hat{1}) = U_{G/e}.$$

This is a recurrence relation that determines U_G , and it remains to describe the initial case. So, suppose the graph G has n vertices and no edge, i.e., $G = (V, \emptyset)$. We have $i(\emptyset, \pi) = 0$. By the definition of the Möbius function, we have

$$\sum_{\pi \in \mathcal{P}(V)} \mu(\pi, \hat{1}) = \delta_{n1},$$

hence $U_G = \delta_{n1}$ as well in this case.

We have thus a recurrence relation for U_G , and it remains to show that the right-hand side of (4.2) satisfies the same relation. This is true, because when $x = 1$, and when we consider a variant of the Tutte polynomial that is 0 for a non-connected graph, then the first case of (4.1) becomes a particular case of the third case. ■

Remark 4.2 Proposition 4.1 can also be derived from results of Burman and Shapiro [9], at least in the case where G is connected. More precisely, in the light of [9, Theorem 9] we can recognize the sum in the left-hand side of (4.2) as the *external activity polynomial* $C_G(w)$, where all edge variables are specialized to $q-1$. It is known to be related to $T_G(1, q)$; see for example [20, Section 2.5].

5 The Case $q = 0$, Lassalle’s Sequence and Heaps

In the case $q = 0$, the substitution $z \rightarrow iz$ recasts equation (2.1) as

$$(5.1) \quad -\log \left(\sum_{n \geq 0} (-1)^n C_n \frac{z^{2n}}{(2n)!} \right) = \sum_{n \geq 1} \tilde{k}_{2n}(0) \frac{z^{2n}}{(2n)!},$$

where $C_n = \frac{1}{n+1} \binom{2n}{n}$ is the n -th Catalan number, known to be the cardinal of $\mathcal{N}(2n)$; see [21]. The integer sequence $\{\tilde{k}_{2n}(0)\}_{n \geq 1} = (1, 1, 5, 56, \dots)$ was previously defined by Lassalle [17] via an equation equivalent to (5.1), and [17, Theorem 1] states that the integers $\tilde{k}_{2n}(0)$ are positive and increasing (stronger results are also true; see [2, 17]).

The goal of this section is to give a meaning to (5.1) in the context of the theory of heaps [26], [10, Appendix 3]. This will give an alternative proof of Theorem 3.5 for the case $q = 0$, based on a classical result on the evaluation $T_G(1, 0)$ of a Tutte polynomial in terms of some orientations of the graph G .

Definition 5.1 A graph $G = (V, E)$ is *rooted* when it has a distinguished vertex $r \in V$, called the *root*. An orientation of G is *root-connected* if for any vertex $v \in V$ there exists a directed path from the root to v .

Proposition 5.2 (Greene and Zaslavsky [14]) *If G is a rooted and connected graph, $T_G(1, 0)$ is the number of its root-connected acyclic orientations.*

The notion of heap was introduced by Viennot [26] as a geometric interpretation of elements in the Cartier–Foata monoid [10], and has various applications in enumeration. We refer the reader to [10, Appendix 3] for a modern presentation of this subject (including a comprehensive bibliography).

Let M be the monoid built on the generators $(x_{ij})_{1 \leq i < j}$ subject to the relations $x_{ij}x_{kl} = x_{kl}x_{ij}$ if $i < j < k < \ell$ or $i < k < \ell < j$. We call it the Cartier–Foata monoid, but in other contexts it could be called a partially commutative free monoid or a trace monoid as well. Following [26], we call an element of M a *heap*.

Any heap can be represented as a “pile” of segments, as in the left part of Figure 2 (this is remindful of [6]). This pile is described inductively: the generator x_{ij} corresponds to a single segment whose extremities have abscissas i and j , and multiplication $m_1 m_2$ is obtained by placing the pile of segments corresponding to m_2 above the one corresponding to m_1 . In terms of segments, the relation $x_{ij}x_{kl} = x_{kl}x_{ij}$ if $i < j < k < \ell$ has a geometric interpretation: segments are allowed to move vertically as long as they do not intersect (this is the case of x_{34} and x_{67} in Figure 2). Similarly, the other relation $x_{ij}x_{kl} = x_{kl}x_{ij}$ if $i < k < \ell < j$ can be treated by thinking of each segment as the projection of an arch as in the central part of Figure 2. In

this three-dimensional representation, all the commutation relations are translated in terms of arches that are allowed to move along the dotted lines as long as they do not intersect.

A heap can also be represented as a poset. Consider two segments s_1 and s_2 in a pile of segments, then the relation is defined by saying that $s_1 < s_2$ if s_1 is always below s_2 , after any movement of the arches (along the dotted lines and as long as they do not intersect, as above). This way, a heap can be identified with a poset where each element is labeled by a generator of M and two elements whose labels do not commute are comparable. See the right part of Figure 2 for an example and [10, Appendice 3] for details.




Figure 2: The heap $m = x_{46}x_{67}x_{34}x_{16}x_{47}$ as a pile of segments and the Hasse diagram of the associated poset.

Definition 5.3 For any heap $m \in M$, let $|m|$ denote its length as a product of generators. Moreover, $m \in M$ is called a *trivial heap* if it is a product of pairwise commuting generators. Let $M^\circ \subset M$ denote the set of trivial heaps.

Let $\mathbb{Z}[[M]]$ denote the ring of formal power series in M , i.e., all formal sums $\sum_{m \in M} \alpha_m m$ with multiplication induced by the one of M . A fundamental result of Cartier and Foata [10] is the identity in $\mathbb{Z}[[M]]$ as follows:

$$(5.2) \quad \left(\sum_{m \in M^\circ} (-1)^{|m|} m \right)^{-1} = \sum_{m \in M} m.$$

Note that M° contains the neutral element of M , so that the sum in the left-hand side is invertible, being a formal power series with constant term equal to 1.

Definition 5.4 An element $m \in M$ is called a *pyramid* if the associated poset has a unique maximal element. Let $P \subset M$ denote the subset of pyramids.

A fundamental result of the theory of heaps links the generating function of pyramids with the one of all heaps [10,26]. It essentially relies on the exponential formula

for labeled combinatorial objects and reads

$$(5.3) \quad \log \left(\sum_{m \in M} m \right) =_{\text{comm}} \sum_{p \in P} \frac{1}{|p|} p,$$

where the sign $=_{\text{comm}}$ means that the equality holds in any commutative quotient of $\mathbb{Z}[[M]]$. Combining (5.2) and (5.3), we obtain

$$(5.4) \quad -\log \left(\sum_{m \in M^\circ} (-1)^{|m|} m \right) =_{\text{comm}} \sum_{p \in P} \frac{1}{|p|} p.$$

Now, let us examine how to apply this general equality to the present case.

The following lemma is a direct consequence of the definitions and permits us to identify trivial heaps with noncrossing matchings.

Lemma 5.5 *The map*

$$(5.5) \quad \Phi: x_{i_1 j_1} \cdots x_{i_n j_n} \mapsto \{ \{i_1, j_1\}, \dots, \{i_n, j_n\} \}$$

defines a bijection between the set of trivial heaps M° and the disjoint union of $\mathcal{N}(V)$, where V runs through the finite subsets (of even cardinal) of $\mathbb{N}_{>0}$.

For a general heap $m \in M$, we can still define $\Phi(m)$ via (5.5), but it may not be a matching; for example $\Phi(x_{1,2}x_{2,3}) = \{ \{1, 2\}, \{2, 3\} \}$. Let us first consider the case of $m \in M$ such that $\Phi(m)$ is really a matching.

Lemma 5.6 *Let $\sigma \in \mathcal{M}(V)$ for some $V \subset \mathbb{N}_{>0}$. Then the heaps $m \in M$ such that $\Phi(m) = \sigma$ are in bijection with acyclic orientations of $G(\sigma)$. Thus, such a heap $m \in M$ can be identified with a pair (σ, r) , where r is an acyclic orientation of the graph $G(\sigma)$.*

Proof An acyclic orientation r on $G(\sigma)$ defines a partial order on σ by saying that two arches x and y satisfy $x < y$ if there is a directed path from y to x . In this partial order, two crossing arches are always comparable, since they are adjacent in $G(\sigma)$. We recover the description of heaps in terms of posets, as described above, so each pair (σ, r) corresponds to a heap $m \in M$ with $\Phi(m) = \sigma$. ■

To treat the case where $m \in M$ such that $\Phi(m)$ is not a matching, such as $x_{12}x_{23}$, we are led to introduce a set of commuting variables $(a_i)_{i \geq 1}$ such that $a_i^2 = 0$ and consider the specialization $x_{ij} \mapsto a_i a_j$ that defines a morphism of algebras $\omega: \mathbb{Z}[[M]] \rightarrow \mathbb{Z}[[a_1, a_2, \dots]]$. This way, for any $m \in M$ we have either $\omega(m) = 0$ or $\Phi(m) \in \mathcal{M}(V)$ for some $V \subset \mathbb{N}_{>0}$.

Let $m \in M$ such that $\omega(m) \neq 0$. As seen in Lemma 5.6, it can be identified with the pair (σ, r) , where $\sigma = \Phi(m)$, and r is an acyclic orientation of $G(\sigma)$. Then the condition defining pyramids is easily translated in terms of (σ, r) . Indeed we have $m \in P$ if and only if the acyclic orientation r has a unique source (where a *source* is a vertex having no ingoing arrows).

Under the specialization ω , the generating function of trivial heaps is

$$(5.6) \quad \omega \left(\sum_{m \in M^\circ} (-1)^{|m|} m \right) = \sum_{n \geq 0} (-1)^n C_n e_{2n},$$

where e_{2n} is the $2n$ -th elementary symmetric function in the a_i 's. Indeed, let $V \subset \mathbb{N}_{>0}$ with $\#V = 2n$. Then the coefficient of $\prod_{i \in V} a_i$ in the left-hand side of (5.6) is $(-1)^{\#V} \#N(V) = (-1)^n C_n$, as can be seen using Lemma 5.5. In particular, it only depends on n , so that this generating function can be expressed in terms of the e_{2n} . Moreover, since the variables a_i have vanishing squares, their elementary symmetric functions satisfy

$$e_{2n} = \frac{1}{(2n)!} e_1^{2n},$$

so that the right-hand side of (5.6) is actually the exponential generating of the Catalan numbers (evaluated at e_1). It remains to explain the meaning of taking the logarithm of the left-hand side of (5.6) using pyramids and equation (5.4).

Note that the relation $=_{\text{comm}}$ becomes a true equality after the specialization $x_{ij} \mapsto a_i a_j$. So taking the image of (5.4) under ω and using (5.6), this gives

$$-\log \left(\sum_{n \geq 0} (-1)^n C_n e_{2n} \right) = \sum_{p \in P} \frac{1}{|p|} \omega(p).$$

The argument used to obtain (5.6) shows as well that the right-hand side of the previous equation is $\sum \frac{x_n}{n} e_{2n}$, where $x_n = \#\{p \in P : \omega(p) = a_1 \cdots a_{2n}\}$. So we have

$$-\log \left(\sum_{n \geq 0} (-1)^n C_n e_{2n} \right) = \sum_{n \geq 0} \frac{x_n}{n} e_{2n},$$

and comparing this with (5.1), we obtain $\tilde{k}_{2n}(0) = \frac{x_n}{n}$.

Clearly, a graph with an acyclic orientation always has a source, and it has a unique source only when it is root-connected (for an appropriate root, viz. the source). So a pyramid p such that $\omega(p) \neq 0$ can be identified with a pair (σ, r) where r is a root-connected acyclic orientation of $G(\sigma)$. Then using Proposition 5.2, it follows that

$$x_n = n \sum_{\sigma \in \mathcal{M}^c(2n)} T_{G(\sigma)}(1, 0).$$

Here, the factor n in the right-hand side accounts for the n possible choices of the source in each graph $G(\sigma)$. Eventually, we obtain

$$(5.7) \quad \tilde{k}_{2n}(0) = \sum_{\sigma \in \mathcal{M}^c(2n)} T_{G(\sigma)}(1, 0);$$

i.e., we have proved the particular case $q = 0$ of Theorem 3.5.

Let us restate the result in an equivalent form. We can consider that if $\sigma \in \mathcal{M}(2n)$, the graph $G(\sigma)$ has a canonical root which is the arch containing 1. Then equation (5.7) gives a combinatorial model for the integers $\tilde{k}_{2n}(0)$.

Theorem 5.7 *The integer $\tilde{k}_{2n}(0)$ counts pairs (σ, r) , where $\sigma \in \mathcal{M}^c(2n)$, and r is an acyclic orientation of $G(\sigma)$ whose unique source is the arch of σ containing 1.*

From this, it is possible to give a combinatorial proof that the integers $\tilde{k}_{2n}(0)$ are increasing, as suggested by Lassalle [17] who gave an algebraic proof. Indeed, we can check that pairs (σ, r) where $\{1, 3\}$ is an arch of σ are in bijection with the same objects but of size one less, hence $\tilde{k}_{2n}(0) \leq \tilde{k}_{2n+2}(0)$.

Before ending this section, note that the left-hand side of (5.1) is $-\log(\frac{1}{z}J_1(2z))$ where J_1 is the Bessel function of order 1. There are quite a few other cases where the combinatorics of Bessel functions are related to the theory of heaps; see the articles of Fédou [11, 12], Bousquet-Mélou and Viennot [6].

6 The Case $q = 2$, the Exponential Formula

The specialization at $(1, 2)$ of a Tutte polynomial has combinatorial significance in terms of connected spanning subgraphs (see [1, Chapter 9]), so it is natural to consider the case $q = 2$ of Theorem 3.5. This case is special, because the factor $(q - 1)^{n-1}$ disappears, so that $\tilde{k}_{2n}(2) = k_{2n}(2)$. We can then interpret the logarithm in the sense of combinatorial species, by showing that $\tilde{k}_{2n}(2)$ counts some *primitive* objects and $m_{2n}(2)$ counts *assemblies* of those, just like permutations that are formed by assembling cycles (this is the exponential formula for labeled combinatorial objects; see [1, Chapter 3]). What we obtain is another more direct proof of Theorem 3.5, based on an interpretation of $T_G(1, 2)$ as follows.

Proposition 6.1 (Gioan [13]) *If G is a rooted and connected graph, $T_G(1, 2)$ is the number of its root-connected orientations.*

This differs from the more traditional interpretation of $T_G(1, 2)$ in terms of connected spanning subgraphs mentioned above, but it is what naturally appears in this context.

Definition 6.2 Let $\mathcal{M}^+(2n)$ be the set of pairs (σ, r) , where $\sigma \in \mathcal{M}(2n)$ and r is an orientation of the graph $G(\sigma)$. Such a pair is called an *augmented matching* and is depicted with the convention that the arch $\{i, j\}$ lies above the arch $\{k, \ell\}$ if there is an oriented edge $\{i, j\} \rightarrow \{k, \ell\}$, and behind it if there is an oriented edge $\{k, \ell\} \rightarrow \{i, j\}$.

See Figure 3 for example. Clearly, $\#\mathcal{M}^+(2n) = m_{2n}(2)$. Indeed, each graph $G(\sigma) = (V, E)$ has $2^{\#E}$ orientations, and $\#E = \text{cr}(\sigma)$, so this follows from (2.4).

Notice that if there is no directed cycle in the oriented graph $(G(\sigma), r)$, the augmented matching (σ, r) can be identified with a heap $m \in M$ as defined in the previous section. The one in Figure 3 would be $x_{3,5}x_{4,11}x_{10,12}x_{1,6}x_{7,9}x_{2,8}$. Actually, the application of the exponential formula in the present section is quite reminiscent of the link between heaps and pyramids as seen in the previous section.

Definition 6.3 Recall that each graph $G(\sigma)$ is rooted with the convention that the root is the arch containing 1. Let $\mathcal{J}(2n) \subset \mathcal{M}^+(2n)$ be the set of augmented matchings




Figure 3: An augmented matching (σ, r) and the corresponding orientation of $G(\sigma)$.

(σ, r) such that σ is connected and r is a root-connected orientation of $G(\sigma)$. The elements of $\mathcal{J}(2n)$ are called *primitive augmented matchings*. For any $V \subset \mathbb{N}_{>0}$ with $\#V = 2n$, we also define the set $\mathcal{J}(V)$, with the same combinatorial description as $\mathcal{J}(2n)$ except that matchings are based on the set V instead of $\{1, \dots, 2n\}$.

Using Proposition 6.1, we have

$$\#\mathcal{J}(2n) = \sum_{\sigma \in \mathcal{M}^+(2n)} T_{G(\sigma)}(1, 2),$$

so that the particular case $q = 2$ of Theorem 3.5 is the equality $\#\mathcal{J}(2n) = k_{2n}(2)$. To prove this from (2.1) using the exponential formula, we have to see how an augmented matching can be decomposed into an assembly of primitive ones, as stated in Proposition 6.4. This decomposition thus proves the case $q = 2$ of Theorem 3.5. Note also that the bijection given below is equivalent to the first identity in (2.2).

Proposition 6.4 *There is a bijection*

$$\mathcal{M}^+(2n) \longrightarrow \bigsqcup_{\pi \in \mathcal{P}(n)} \prod_{V \in \pi} \mathcal{J}(V).$$

Proof Let $(\sigma, r) \in \mathcal{M}^+(2n)$; the bijection is defined as follows. Consider the vertices of $G(\sigma)$ that are accessible from the root. This set of vertices defines a matching on a subset $V_1 \subset \{1, \dots, 2n\}$. For example, in the case of Figure 3, the root is $\{1, 6\}$ and the only other accessible vertex is $\{2, 8\}$, so $V_1 = \{1, 2, 6, 8\}$. Together with the restriction of the orientation r on this subset of vertices, this defines an augmented matching $(\sigma_1, r_1) \in \mathcal{M}^+(V_1)$ that is primitive by construction. By repeating this operation on the set $\{1, \dots, 2n\} \setminus V_1$, we find $V_2 \subset \{1, \dots, 2n\} \setminus V_1$ and $(\sigma_2, r_2) \in \mathcal{J}(V_2)$, and so on. See Figure 4 for the result, in the case of the augmented matching in Figure 3.

The inverse bijection is easily described. If $(\sigma_i, r_i) \in \mathcal{J}(V_i)$ for any $1 \leq i \leq k$ where $\pi = \{V_1, \dots, V_k\}$, let $\sigma = \sigma_1 \cup \dots \cup \sigma_k$, and the orientation r of $G(\sigma)$ is as follows. Let e be an edge of $G(\sigma)$ and x_1, x_2 be its endpoints, with $x_1 \in \sigma_{j_1}$ and $x_2 \in \sigma_{j_2}$. If $j_1 = j_2$, the edge e is oriented in accordance with the orientation $r_{j_1} = r_{j_2}$. Otherwise, say $j_1 < j_2$, then the edge e is oriented in the direction $x_1 \leftarrow x_2$. ■




Figure 4: Decomposition of an augmented matching into primitive ones.

7 Cumulants in Terms of Free Cumulants

In this section, $\{\mu_n\}_{n \geq 0}$ is any sequence of moments, and $\{k_n\}_{n \geq 1}$ and $\{c_n\}_{n \geq 1}$ are the corresponding cumulants and free cumulants defined by (2.2) and (2.3). It was shown by Lehner [18] that (2.2) and (2.3) imply

$$(7.1) \quad c_n = \sum_{\pi \in \mathcal{P}^c(n)} k_\pi.$$

By a triangularity argument, this can clearly be inverted to have k_n in terms of $c_1, c_2,$ etc. But it cannot be done using a Möbius function as in (2.2) and (2.3), because in this case we have that

$$c_\pi \neq \sum_{\substack{\rho \in \mathcal{P}^c(n) \\ \rho \leq \pi}} k_\rho$$

in general, contrary to

$$m_\pi = \sum_{\substack{\rho \in \mathcal{P}(n) \\ \rho \leq \pi}} k_\rho \text{ if } \pi \in \mathcal{P}(n), \quad m_\pi = \sum_{\substack{\rho \in \mathcal{NC}(n) \\ \rho \leq \pi}} c_\rho \text{ if } \pi \in \mathcal{NC}(n).$$

Still, we obtain an explicit inverse formula for (7.1) in the theorem below.

For any set partition $\pi \in \mathcal{P}(V)$ for some $V \subset \mathbb{N}$, we can define a crossing graph $G(\pi)$, whose vertices are the blocks of π , and there is an edge between $b, c \in \pi$ if $\{b, c\}$ is not a noncrossing partition. Note that π is connected if and only if the graph $G(\pi)$ is connected. The two different proofs for the semicircular cumulants show the following theorem as well.

Theorem 7.1 For any $n \geq 1$, we have

$$k_n = \sum_{\pi \in \mathcal{P}^c(n)} c_\pi (-1)^{1+\#\pi} T_{G(\pi)}(1, 0).$$

Let us sketch the proofs. If $\pi \in \mathcal{P}(n)$, similar to Lemma 3.1, we have

$$m_\pi = \prod_{b \in \pi} \left(\sum_{\rho \in \mathcal{NC}(b)} c_\rho \right) = \sum_{\substack{\rho \in \mathcal{P}(n) \\ \rho \trianglelefteq \pi}} c_\rho,$$

where the relation $\rho \trianglelefteq \pi$ means that $\rho \leq \pi$ and $\rho|_b$ is a noncrossing partition for each $b \in \pi$. Indeed the map $\rho \mapsto (\rho|_b)_{b \in \pi}$ is a bijection between $\{\rho \in \mathcal{P}(n) : \rho \trianglelefteq \pi\}$ and

$\prod_{b \in \pi} \mathcal{N}\mathcal{C}(b)$. The same computation as in (3.2) and (3.3) gives

$$(7.2) \quad k_n = \sum_{\pi \in \mathcal{P}(n)} m_\pi \mu(\pi, \widehat{1}) = \sum_{\substack{\rho, \pi \in \mathcal{P}(n) \\ \rho \trianglelefteq \pi}} c_\rho \mu(\pi, \widehat{1}) = \sum_{\rho \in \mathcal{P}(n)} c_\rho W(\rho),$$

where

$$W(\rho) = \sum_{\substack{\pi \in \mathcal{P}(n) \\ \rho \trianglelefteq \pi}} \mu(\pi, \widehat{1}).$$

Denoting $G(\rho) = (V, E)$ the crossing graph of ρ , the previous equality is rewritten $W(\rho) = \sum \mu(\pi, \widehat{1})$, where the sum is over $\pi \in \mathcal{P}(V)$ such that for any $b \in \pi, e \in E$, the block b does not contain both endpoints of the edge e . Then the case $q = 0$ of Proposition 4.1 shows that

$$W(\rho) = \begin{cases} (-1)^{1+\#\rho} T_{G(\rho)}(1, 0) & \text{if } \rho \in \mathcal{P}^c(n), \\ 0 & \text{otherwise.} \end{cases}$$

Together with (7.2), this completes the first proof of Theorem 7.1. ■

As for the second proof, we follow the outline of Section 5, but with another definition for M, M°, P , and ω . Let M be the monoid with generators (x_V) , where V runs through finite subsets of $\mathbb{N}_{>0}$, and with relations $x_V x_W = x_W x_V$ if $\{V, W\}$ is a non-crossing partition. We also denote $M^\circ \subset M$ the corresponding set of trivial heaps, i.e., products of pairwise commuting generators, and an element $x_{b_1} \cdots x_{b_i} \in M^\circ$ is identified with $\{b_1, \dots, b_i\} \in \mathcal{N}\mathcal{C}(\cup_{j=1}^i b_j)$. The subset $P \subset M$ is characterized by Definition 5.4. Now, we consider the morphism ω defined on $\mathbb{Z}[[M]]$ by

$$\omega(x_V) = -c_{\#V} \prod_{i \in V} a_i.$$

We have

$$\begin{aligned} \omega\left(\sum_{m \in M^\circ} (-1)^{|m|} m\right) &= \sum_V \sum_{\pi \in \mathcal{N}\mathcal{C}(V)} (-1)^{\#\pi} \prod_{b \in \pi} \omega(x_b) \\ &= \sum_V \sum_{\pi \in \mathcal{N}\mathcal{C}(V)} c_\pi \prod_{i \in V} a_i \\ &= \sum_{n \geq 0} m_n e_n = \sum_{n \geq 0} m_n \frac{e_1^n}{n!}. \end{aligned}$$

We still understand that $V \subset \mathbb{N}_{>0}$ is finite, $(a_i)_{i \geq 1}$ are commuting variables with vanishing squares, and e_n is the n -th elementary symmetric function in the a_i 's. Equation (5.4) is still valid as such with the new definition of M° and P , and taking the image by ω gives

$$-\log\left(\sum_{n \geq 0} m_n \frac{e_1^n}{n!}\right) = \sum_{p \in P} \frac{1}{|p|} \omega(p).$$

By extracting the coefficient of $a_1 \cdots a_n$ on both sides, we get

$$(7.3) \quad -k_n a_1 \cdots a_n = \sum \frac{1}{|p|} \omega(p)$$

where the sum is over $p \in P$ that can be written in the form $p = x_{V_1} \cdots x_{V_k}$, where $\pi = \{V_1, \dots, V_k\} \in \mathcal{P}(n)$. We can gather the terms corresponding to a same π . Note that

$$\omega(p) = (-1)^{\#\pi} c_\pi a_1 \cdots a_n.$$

From (7.3), we obtain

$$(7.4) \quad -k_n a_1 \cdots a_n = \sum_{\pi \in \mathcal{P}(n)} \frac{x_\pi}{\#\pi} (-1)^{\#\pi} c_\pi a_1 \cdots a_n,$$

where x_π is the number of $p \in P$ that are a product of x_b , where b runs through the blocks of π . Following the ideas in Section 5, such a p is characterized by π together with an acyclic orientation of the graph $G(\pi)$ having a unique source. So x_π is the number of such orientations, which is equal to $\#\pi \cdot T_{G(\pi)}(1, 0)$ if π is connected and 0 otherwise. Knowing the value of x_π , from equation (7.4), we complete the second proof of Theorem 7.1. ■

8 Final Remarks

It would be interesting to explain why the same combinatorial objects appear for both $c_{2n}(q)$ and $k_{2n}(q)$. This suggests that there exists some quantity that interpolates between the classical and free cumulants of the q -semicircular law; however, building a noncommutative probability theory that encompasses the classical and free ones appears to be elusive (see [25] for a precise statement). It means that building such an interpolation would rely not only on the q -semicircular law and its moments, but on its realization as a noncommutative random variable. This might be feasible using q -Fock spaces [7, 8] but is beyond the scope of this article.

Acknowledgment This work was initiated during the trimester “Bialgebras in Free Probability” at the Erwin Schrödinger Institute in Vienna. In particular I thank Franz Lehner, Michael Anshelevich, and Natasha Blitvić for their conversations.

References

- [1] M. Aigner, *A course in enumeration*. Graduate Texts in Mathematics, 238, Springer, Berlin, 2007.
- [2] T. Amdeberhan, V. H. Moll, and C. Vignat, *A new proof of a conjecture by D. Zeilberger about Catalan numbers*. [arxiv:1202.1203v1](https://arxiv.org/abs/1202.1203v1).
- [3] M. Anshelevich, S. T. Belinschi, M. Bozejko, and F. Lehner, *Free infinite divisibility for q -Gaussians*. *Math. Res. Lett.* **17**(2010), no. 5, 905–916.
- [4] S. Belinschi, M. Bozejko, F. Lehner, and R. Speicher, *The normal distribution is \boxplus -infinitely divisible*. *Adv. Math.* **226**(2011), no. 4, 3677–3698. <http://dx.doi.org/10.1016/j.aim.2010.10.025>
- [5] N. Blitvić, *On the norm of q -circular operators*. [arxiv:1102.0748](https://arxiv.org/abs/1102.0748)
- [6] M. Bousquet-Mélou and X. G. Viennot, *Empilements de segments et q -énumération de polyominos convexes dirigés*. *J. Combin. Theory Ser. A* **60**(1992), no. 2, 196–224. [http://dx.doi.org/10.1016/0097-3165\(92\)90004-E](http://dx.doi.org/10.1016/0097-3165(92)90004-E)

- [7] M. Bozejko, B. Kümmerer, and R. Speicher, *q-Gaussian processes: non-commutative and classical aspects*. *Comm. Math. Phys.* **185**(1997), no. 1, 129–154. <http://dx.doi.org/10.1007/s002200050084>
- [8] M. Bozejko and R. Speicher, *An example of a generalized brownian motion*. *Comm. Math. Phys.* **137**(1991), no. 3, 519–531. <http://dx.doi.org/10.1007/BF02100275>
- [9] Y. Burman and B. Shapiro, *Around matrix-tree theorem*. *Math. Res. Lett.* **13**(2006), no. 5–6, 761–774.
- [10] P. Cartier and D. Foata, *Problèmes combinatoires de commutation et réarrangements*. *Lecture Notes in Mathematics*, 85, Springer-Verlag, Berlin-New York, 1969.
- [11] J.-M. Fédou, *Sur les fonctions de Bessel*. *Discrete Math.* **139**(1995), no. 1–3, 473–480. [http://dx.doi.org/10.1016/0012-365X\(94\)00150-H](http://dx.doi.org/10.1016/0012-365X(94)00150-H)
- [12] ———, *Combinatorial objects enumerated by q-Bessel functions*. *Rep. Math. Phys.* **34**(1994), no. 1, 57–70. [http://dx.doi.org/10.1016/0034-4877\(94\)90017-5](http://dx.doi.org/10.1016/0034-4877(94)90017-5)
- [13] E. Gioan, *Enumerating degree sequences in digraphs and a cycle-cocycle reversing system*. *European J. Combin.* **28**(2007), no. 4, 1351–1366. <http://dx.doi.org/10.1016/j.ejc.2005.11.006>
- [14] C. Greene and T. Zaslavsky, *On the interpretation of Whitney numbers through arrangements of hyperplanes, zonotopes, non-Radon partitions, and orientations of graphs*. *Trans. Amer. Math. Soc.* **280**(1983), no. 1, 97–126. <http://dx.doi.org/10.1090/S0002-9947-1983-0712251-1>
- [15] F. Hiai and D. Petz, *The semicircle law, free random variables and entropy*. *Mathematical Surveys and Monographs*, 77, American Mathematical Society, Providence, RI, 2000.
- [16] M. E. H. Ismail, D. Stanton, and G. Viennot, *The combinatorics of q-Hermite polynomials and the Askey-Wilson integral*. *European J. Combin.* **8**(1987), no. 4, 379–392.
- [17] M. Lassalle, *Two integer sequences related to Catalan numbers*. *J. Combin. Theory Ser. A* **119**(2012), no. 4, 923–935. <http://dx.doi.org/10.1016/j.jcta.2012.01.002>
- [18] F. Lehner, *Free cumulants and enumeration of connected partitions*. *European J. Combin.* **23**(2002), no. 8, 1025–1031. <http://dx.doi.org/10.1006/eujc.2002.0619>
- [19] A. Nica and R. Speicher, *Lectures on the combinatorics of free probability*. *London Mathematical Society Lecture Note Series*, 335, Cambridge University Press, Cambridge, 2006.
- [20] A. D. Sokal, *The multivariate Tutte polynomial (alias Potts model) for graphs and matroids*. In: *Surveys in combinatorics 2005*, *London Math. Soc. Lecture Note Ser.*, 327, Cambridge University Press, Cambridge, 2005, pp. 173–226.
- [21] R. P. Stanley, *Enumerative combinatorics*. Vol. 2, *Cambridge Studies in Advanced Mathematics*, 62, Cambridge University Press, Cambridge, 1999.
- [22] G. Szegő, *Ein Beitrag zur Theorie der Thetafunktionen*. *Sitz. Preuss. Akad. Wiss. Phys. Math. Kl.* **19**(1926), 242–252.
- [23] J. Touchard, *Sur un problème de configurations et sur les fractions continues*. *Canad. J. Math.* **4**(1952), 2–25. <http://dx.doi.org/10.4153/CJM-1952-001-8>
- [24] H. van Leeuwen and H. Maassen, *A q-deformation of the Gauss distribution*. *J. Math. Phys.* **36**(1995), no. 9, 4743–4756. <http://dx.doi.org/10.1063/1.530917>
- [25] ———, *An obstruction for q-deformation of the convolution product*. *J. Phys. A* **29**(1996), no. 15, 4741–4748. <http://dx.doi.org/10.1088/0305-4470/29/15/036>
- [26] G. X. Viennot, *Heaps of pieces. I. Basic definitions and combinatorial lemmas*. In: *Combinatoire énumérative* (Montreal, Que., 1985) *Lecture Notes in Math.*, 1234, Springer, Berlin, 1986, pp. 321–350.

CNRS and Institut Gaspard Monge, Université Paris-Est Marne-la-Vallée, 5 Boulevard Descartes, Champs-sur-Marne, 77454 Marne-la-Vallée cedex 2, France
 e-mail: matthieu.josuat-verges@univ-mlv.fr