

Constraints to cosmological parameters through cluster evolution

A. Del Popolo and N. Ercan

Boğaziçi University, Physics Department, 80815 Bebek, Istanbul, Turkey

Abstract. In this paper, we revisit the constraints obtained by several authors (Reichart et al. 1999; Eke et al. 1998; Henry 2000) on the estimated values of Ω_m , n and σ_8 in the light of recent theoretical developments: 1) new theoretical mass functions (Sheth & Tormen 1999, Sheth, Mo & Tormen 2001, Del Popolo 2002b); 2) a more accurate mass-temperature relation, also determined for arbitrary Ω_m and Ω_Λ (Del Popolo 2002a).

Firstly, using the quoted improvements, we re-derive an expression for the X-ray Luminosity Function (XLF), similarly to Reichart et al. (1999), and then we get some constraints to Ω_m and n , by using the *ROSAT* BCS and EMSS samples and maximum-likelihood analysis. Then we re-derive the X-ray Temperature Function (XTF), similarly to Henry (2000), re-obtaining the constraints on Ω_m , n , σ_8 . In the case of both the XLF and the XTF, the changes in the mass function and M-T relation produces an increase in Ω_m of $\simeq 20\%$ and similar results in σ_8 and n .

1. Introduction

It is well known that clusters are strong X-ray emitters whose study can put constraints on fundamental cosmological parameters. There are different methods to trace the evolution of the cluster number density: a) The X-ray temperature function (XTF) has been presented for local (e.g. Henry & Arnaud 1991) and distant clusters (Eke et al. 1998; Henry 2000). b) The evolution of the X-ray luminosity function (XLF).

The results obtained for Ω_m and other cosmological parameters are in many cases discrepant the one with the other. Several studies in literature, show that the parameters values span the entire range of acceptable solutions: $0.2 \leq \Omega_m \leq 1$ (see Reichart et al. 1999). The reasons leading to the quoted discrepancies has been studied in several papers (Eke et al. 1998; Borgani et al. 2001): 1) The inadequate approximation given by the PS (e.g., Bryan & Norman 1997). 2) Inadequacy in the structure formation as described by the spherical model leading to changes in the threshold parameter δ_c (e.g., Governato et al. 1998). 3) Inadequacy in the M-T relation obtained from the virial theorem (see Del Popolo 2002a). 4) Effects of cooling flows. 5) Determination of the X-ray cluster catalog's selection function. 6) Evolution of the L-T relation. 7) Optimization methods used in the analysis. These reasons lead us to re-calculate the constraints on Ω_m , n and σ_8 , using the XLF and XTF.

2. Constraints to Ω_m and n from the XLF

Similarly to Reichart et al. (1999), we re-derived an expression for the XLF, using an improved version of the mass function and $M - T$ relation, obtained in Del Popolo (2000a,b), taking account of the effects of asphericity and tidal interaction with neighbors. Then we got some constraints to Ω_m and n , by using the *ROSAT* BCS and EMSS samples.




Figure 1. (a) The 68% confidence contours for the parameters n , k_0 and Ω_m for the open model. The dashed lines are lines of constant σ_8 . (b) Δ (likelihood) for the parameter Ω_m . The solid line is obtained from the model of this paper while the dotted line is that calculated by Henry (2000). The dashed lines represent various confidence levels (65%, 90%, 95%, 99%). (c) The 68% confidence contours for the parameters σ_8 , and Ω_m for the open model (see also Henry 2000, Fig. 9).

As described in Del Popolo (2000b), the mass function can be approximated by:

$$n(m, z) \simeq 1.21 \frac{\bar{p}}{m^2} \frac{d \log(\nu)}{d \log m} \left(1 + \frac{0.06}{(a\nu)^{0.585}} \right) \sqrt{\frac{a\nu}{2\pi}} \exp \left\{ -a\nu \left[1 + \frac{0.57}{(a\nu)^{0.585}} \right]^2 / 2 \right\} \quad (2.1)$$

where $a = 0.707$. Eq. (2.1) can be converted from a mass function to a luminosity function using an $L - T$ relation (we use that of Mathiesen & Evrard 1998), and a $T - M$ relation. This last is the one obtained in Del Popolo (2000a), and is based on the merging-halo formalism of Lacey & Cole (1993), accounting for the fact that massive clusters accrete matter quasi-continuously, and again take account of angular momentum acquisition by protostructures:

$$kT \simeq 8\text{keV} \left(\frac{M^{\frac{2}{3}}}{10^{15} h^{-1} M_{\odot}} \right) \frac{\left[\frac{1}{m_1} + \left(\frac{t_{\Omega}}{t} \right)^{\frac{2}{3}} + \frac{K(m_1, x)}{M^{8/3}} \right]}{\left[\frac{1}{m_1} + \left(\frac{t_{\Omega}}{t_0} \right)^{\frac{2}{3}} + \frac{K_0(m_1, x)}{M_0^{8/3}} \right]} \quad (2.2)$$

(see Del Popolo 2000a for a derivation of the previous equation and the definition of the terms in it). The luminosity function is obtained as in Reichart et al. (1999), using the mass function and the $M - T$ relation previously introduced (see Del Popolo 2003 for a detailed analysis). A Bayesian inference analysis used to constrain the model parameters

shows that: $\Omega_m = 1.15_{-0.33}^{+0.40}$ and $n = -1.55_{-0.41}^{+0.42}$. The previous result shows that the change in the mass function and M-T relation gives rise to an increase of Ω_m and n of $\simeq 20\%$ with respect to Reichart's results. The lesson from the previous calculation is that taking account of non-sphericity in collapse and the fact that massive clusters accrete matter quasi-continuously gives rise to a noteworthy change in the prediction of cosmological parameters, as Ω_m . In order to check the previous trend, we have also estimated the value of Ω_m following Borgani et al. (2001). Analyzing the ROSAT Deep Cluster Survey (RDCS) and using the XLF to obtain constraints on cosmological parameters, Borgani et al. (2001) found that $\Omega_m = 0.35_{-0.10}^{+0.13}$. Using their method and data, but our mass function and M-T relation, one obtains larger values of Ω_m ($\Omega_m \simeq 0.4 \pm 0.1$) that, differently from the previous analysis (Reichart et al. 1999), exclude an Einstein-de Sitter model.


3. Constraints to Ω_m , n , and σ_8 from the XTF

As previously reported, the mass function (MF) is a critical ingredient in putting strong constraints on cosmological parameters (e.g. Ω_m). In the following, we will recalculate the constraints obtained by Henry (2000), by using the mass function and the M-T relation modified as described in the previous section eq. (2.1) and eq. (2.2) (see Del Popolo 2003). we use a maximum likelihood fit to the unbinned data in order to determine various model parameters. The method is described in Marshall et al (1983). The likelihood function is given by their eq. (2), adapted to our present situation. At this point, we can fit the data described in Section. 2 of Henry (2000) to the theory previously described using the quoted maximum likelihood method. The most general description of the results requires the three parameters of the fit (Ω_m , σ_8 and n). These values shows that the correction introduced by the new form of the mass function and M-T relation gives rise to higher values of Ω_m ($\Omega_m = 0.6 \pm 0.13$, while it is $\Omega_m = 0.49 \pm 0.12$ for Henry 2000) and $n = -1.5 \pm 0.32$ ($n = -1.72 \pm 0.34$ in Henry 2000). Constraints are relatively tight when considering this single parameter. We find that $\Omega_m = 0.6_{-0.11}^{+0.12}$ at the 68% confidence level and $\Omega_m = 0.6_{-0.2}^{+0.23}$ at the 95% confidence level for the open model.

Concluding, our analysis shows that improvements in the mass function and M-T relation increases the value of Ω_m and that even small correction in the physics of the collapse can induce noteworthy effects on the constraints obtained.

References

- Borgani S., et al. 2001, ApJ 561, 13
 Bryan G.L., Norman M.L., 1997, ApJ 495, 80
 Del Popolo A., Gambera M., 1998, A&A 337, 96
 Del Popolo A., 2002a, MNRAS 336, 81
 Del Popolo A., 2002b, MNRAS 337, 529
 Del Popolo A., 2003, ApJ 599, 723
 Eke V. R., Cole S, Frenk C. S., Henry J.P., 1998, MNRAS 298, 1145
 Governato F., et al., 1999, MNRAS 307, 949
 Henry J.P., 2000, ApJ 534, 565
 Lacey C., Cole S., 1993, MNRAS 262, 627
 Marshall H. L., et al., 1983, ApJ 269, 35
 Reichart D. E., et al., 1999, ApJ 518, 521
 Sheth R. K., Tormen G., 1999, MNRAS 308, 119
 Sheth R. K., Mo H. J., Tormen G., 2001, MNRAS 323, 1 (SMT)



Roman Juskiewicz reminds when the low-density universe was born (*by J.M. Colberg*).



Brent Tully, Volker Müller and Gustavo Yepes at Palazzo Barolo (*by J.M. Colberg*).



Giuseppe Tormen and Tommaso Treu at the conference dinner (*by J.M. Colberg*).



Rien van de Weygaert toasts to the success of both the Colloquium and the Barolo wine (*by J.M. Colberg*).