

months as a senior house officer and the experiences described mirrored mine exactly.

The most enjoyable aspects were being able to work without the interruption of a bleep, enabling one to concentrate wholeheartedly, and becoming part of a new team whose approach was so different from any in a hospital setting. What struck me was that although each person worked autonomously he or she also contributed significantly to the whole team, either in the weekly referral meeting or during daily work.

My experiences have made me more aware of patients' family and social circumstances which can easily be forgotten in a busy acute unit and I would strongly recommend that all trainees spend at least six months working in a child guidance clinic.

HELEN TUCKER, *Torbay Psychiatric Rotation Scheme, Edith Morgan Unit, Torbay Hospital, Torquay TQ2 7AA*

Sir: I was interested to read Jan Hermsen's description of work in a child guidance clinic. (*Psychiatric Bulletin*, 1993, 17, 626). I am currently working as a registrar in a child and adolescent unit and would agree that an experience in this area is essential for all trainees. The shift to a totally different patient population with their own peculiar set of disorders requires a major rethink of approach, method and technique which can be revealing and, at times, painful.

In this area of psychiatry it is acknowledged that the patient comes as part of a package, i.e. the family. You cannot treat the one without the other. In the clinic in which I work family therapy is a major part of the treatment offered. It has opened my eyes to the importance of an approach neglected or ignored in other areas of psychiatry.

Child and adolescent psychiatry involves the treatment of largely unwilling and involuntary patients who are frequently unable to articulate their problems. The predominantly psychotherapeutic basis of this treatment can be unappealing or alarming to the practitioner. However, trainees can gain vital skills which may not be possible in other specialities.

S. C. CARVILL, *14 Woodloes Road, Shirley, Solihull, West Midlands B90 2RP*

Psychiatric emergencies

Sir: I read with interest Kohen's paper on psychiatric emergencies in people with a mental handicap (*Psychiatric Bulletin*, 1993, 17, 587-589). We have just completed a survey of all psychiatric emergencies in the Borders region over one year (June 1992 to May 1993). In contrast to Kohen's

survey of emergencies seen by a consultant during the daytime in an urban area, most of our cases were seen out of hours by junior doctors and came from a dispersed rural population of 104,000.

During the year there were 13 emergencies with a diagnosis of learning difficulty. This was 2.7% of all psychiatric emergencies. The mean age was 36 with a range of 20-69 years. Eleven were male. Only one of the 13 was seen by a consultant, the others being seen by junior doctors, usually with a co-therapist. Eleven were seen out of hours and all were seen within four hours of referral. The commonest source of referral was from the GP (six cases), and the commonest site of assessment was at home (six cases). Referral was precipitated by self-harm in only one case. Most of the patients had never been married (ten cases), but only one still lived with his parents. The others lived alone (six cases) or with other people (six cases). Just over half were employed full or part-time (seven cases).

Nine patients had another diagnosis apart from learning difficulties. There were five cases of neurotic and stress-related disorders, three of affective psychosis and one of personality disorder. Unlike Kohen's study, none were referred for epilepsy. All the referrals had a history of previous admissions and ten were active cases. The consultant was contacted in only two cases, both times to discuss the decision to admit. Six patients were admitted, the reasons given including risk to self, risk to others, social factors, for assessment and for treatment. Eleven cases were thought to be appropriate emergency referrals.

Only two cases were referred on Saturday and there was no evidence of a 'Saturday afternoon syndrome' as described by Spenser (*Psychiatric Bulletin*, 1993, 17, 565). This may reflect better local support from the social work department or other differences in local practice. We would not dispute her conclusion that it is important to have provision of emergency short-term beds for assessment and treatment of people with learning difficulties. We had only 13 in-patient beds at the time of this study and would also support Kohen's conclusion that anxiety about emergencies should not deter the implementation of community care. Most of the referrals were appropriate and the actual demand was small and, to a large extent, met by existing out-of-hours arrangements.

JOHN TAYLOR, *Dingleton Hospital, Melrose, Roxburghshire TD6 9HN*

Defeat Depression Campaign

Sir: The thesis of the Defeat Depression Campaign is that patients are reluctant to take